

Calculating the spawn index for Pacific Herring (*Clupea pallasii*) in British Columbia, Canada

Matthew H. Grinnell, Jake F. Schweigert, Matt Thompson, Sarah Hawkshaw, and Jaclyn S. Cleary

Fisheries and Oceans Canada
Science Branch, Pacific Region
Pacific Biological Station
3190 Hammond Bay Road
Nanaimo, BC V9T 6N7

2023

**Canadian Technical Report of
Fisheries and Aquatic Sciences 3489**



Fisheries and Oceans
Canada

Pêches et Océans
Canada

Canada

Canadian Technical Report of Fisheries and Aquatic Sciences

Technical reports contain scientific and technical information that contributes to existing knowledge but which is not normally appropriate for primary literature. Technical reports are directed primarily toward a worldwide audience and have an international distribution. No restriction is placed on subject matter and the series reflects the broad interests and policies of Fisheries and Oceans Canada, namely, fisheries and aquatic sciences.

Technical reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in the data base *Aquatic Sciences and Fisheries Abstracts*.

Technical reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page.

Numbers 1--456 in this series were issued as Technical Reports of the Fisheries Research Board of Canada. Numbers 457--714 were issued as Department of the Environment, Fisheries and Marine Service, Research and Development Directorate Technical Reports. Numbers 715--924 were issued as Department of Fisheries and Environment, Fisheries and Marine Service Technical Reports. The current series name was changed with report number 925.

Rapport technique canadien des sciences halieutiques et aquatiques

Les rapports techniques contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui ne sont pas normalement appropriés pour la publication dans un journal scientifique. Les rapports techniques sont destinés essentiellement à un public international et ils sont distribués à cet échelon. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques de Pêches et Océans Canada, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports techniques peuvent être cités comme des publications à part entière. Le titre exact figure au-dessus du résumé de chaque rapport. Les rapports techniques sont résumés dans la base de données *Résumés des sciences aquatiques et halieutiques*.

Les rapports techniques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre.

Les numéros 1 à 456 de cette série ont été publiés à titre de Rapports techniques de l'Office des recherches sur les pêcheries du Canada. Les numéros 457 à 714 sont parus à titre de Rapports techniques de la Direction générale de la recherche et du développement, Service des pêches et de la mer, ministère de l'Environnement. Les numéros 715 à 924 ont été publiés à titre de Rapports techniques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 925.

Canadian Technical Report of
Fisheries and Aquatic Sciences 3489

2023

CALCULATING THE SPAWN INDEX FOR PACIFIC HERRING (*CLUPEA PALLASII*) IN BRITISH COLUMBIA, CANADA

by

Matthew H. Grinnell¹, Jake F. Schweigert², Matt Thompson¹, Sarah Hawkshaw³,
and Jaclyn S. Cleary¹

¹ Pacific Biological Station, Fisheries and Oceans Canada,
3190 Hammond Bay Road, Nanaimo, British Columbia V9T 6N7

² Emeritus, Pacific Biological Station, Fisheries and Oceans Canada,
3190 Hammond Bay Road, Nanaimo, British Columbia V9T 6N7

³ Institute of Ocean Sciences, Fisheries and Oceans Canada,
9860 W. Saanich Road, Sidney, British Columbia V8L 5T5

© His Majesty the King in Right of Canada, as represented by the
Minister of the Department of Fisheries and Oceans, 2023
Cat. No. Fs97-6/3489E-PDF ISBN 978-0-660-44033-0 ISSN 1488-5379

Correct citation for this publication:

Grinnell, M.H., Schweigert, J.F., Thompson, M., Hawkshaw, S., and Cleary, J.S.
2023. Calculating the spawn index for Pacific Herring (*Clupea pallasii*) in British Columbia, Canada. Can. Tech. Rep. Fish. Aquat. Sci. 3489: viii + 45 p.

CONTENTS

LIST OF FIGURES	v
LIST OF TABLES	vi
ABSTRACT	vii
RÉSUMÉ	viii
1 INTRODUCTION	1
2 STATISTICAL FRAMEWORK	6
2.1 SURFACE SPAWN FRAMEWORK	6
2.2 DIVE SPAWN FRAMEWORK	6
2.2.1 UNDERSTORY SPAWN FRAMEWORK	6
2.2.2 MACROCYSTIS SPAWN FRAMEWORK	9
3 SAMPLING PROTOCOL	10
3.1 SURFACE SPAWN PROTOCOL	12
3.1.1 SPAWN INTENSITY CATEGORIES	13
3.2 MACROCYSTIS SPAWN PROTOCOL	14
3.3 UNDERSTORY SPAWN PROTOCOL	14
4 CONVERT EGGS TO BIOMASS	14
5 SURFACE SPAWN CALCULATIONS	15
5.1 SAMPLE OBSERVATIONS AND CALCULATIONS	16
5.2 SPAWN OBSERVATIONS AND CALCULATIONS	17
6 MACROCYSTIS SPAWN CALCULATIONS	18
6.1 PLANT OBSERVATIONS	18
6.2 TRANSECT OBSERVATIONS AND CALCULATIONS	18
6.3 SPAWN OBSERVATIONS AND CALCULATIONS	19
7 UNDERSTORY SPAWN CALCULATIONS	21
7.1 QUADRAT OBSERVATIONS AND CALCULATIONS	21
7.2 TRANSECT OBSERVATIONS AND CALCULATIONS	26
7.3 SPAWN OBSERVATIONS AND CALCULATIONS	26

8	TOTAL SPAWN CALCULATIONS	27
8.1	SCALED BIOMASS	28
9	SPAWN-ON-KELP CALCULATIONS	28
10	SOURCES OF UNCERTAINTY	30
11	FUTURE RESEARCH	32
12	CAVEATS	35
13	R PACKAGE	35
14	ACKNOWLEDGEMENTS	36
15	REFERENCES	36
A	SPAWN WIDTH ADJUSTMENTS	40
A.1	SURFACE SPAWN WIDTH	40
A.2	UNDERSTORY SPAWN WIDTH	41
B	SURFACE SPAWN UPDATES	42
C	R PACKAGE CROSSWALK	44

LIST OF FIGURES

1	Pacific Herring spawn index by Location, as well as Sections and Statistical Areas in the Strait of Georgia stock assessment region	3
2	Spatial boundaries for Pacific Herring stock assessment regions in British Columbia	4
3	Sequence of steps for Pacific Herring spawn index calculations	5
4	Aerial view of Pacific Herring spawn	11
5	Number of Pacific Herring egg layers	12
6	Egg density for Pacific Herring surface spawn surveys	17
7	Number of eggs per plant for Pacific Herring <i>Macrocystis</i> spawn surveys	22
8	Substrate egg density for Pacific Herring underwater spawn surveys .	24
9	Algae egg density for Pacific Herring underwater spawn surveys . . .	25

LIST OF TABLES

1	Index notation for Pacific Herring spawn index calculations	7
2	Notation for Pacific Herring spawn survey statistical framework	9
3	Spawn intensity categories and number of egg layers for Pacific Herring surface spawn surveys	13
4	Notation for converting Pacific Herring eggs to spawning biomass	15
5	Notation for Pacific Herring surface spawn index calculations	16
6	Notation for Pacific Herring Macrocystis spawn index calculations	19
7	Notation for Pacific Herring understory spawn index calculations	23
8	Algae types and coefficients for Pacific Herring understory spawn surveys	24
9	Notation for Pacific Herring total spawn index calculations	28
10	Notation for Pacific Herring spawn-on-kelp calculations	30
11	Details for parameters and prediction models used in Pacific Herring spawn index calculations	33
A.1	Spawn width correction factors for Pacific Herring understory spawn surveys	43
A.2	Notation for Pacific Herring understory spawn width adjustments	43
C.1	Crosswalk table for equations and functions in the <code>SpawnIndex R</code> package	45

ABSTRACT

Grinnell, M.H., Schweigert, J.F., Thompson, M., Hawkshaw, S., and Cleary, J.S. 2023. Calculating the spawn index for Pacific Herring (*Clupea pallasii*) in British Columbia, Canada. Can. Tech. Rep. Fish. Aquat. Sci. 3489: viii + 45 p.

The spawn index is one component of Pacific Herring (*Clupea pallasii*) stock assessments in British Columbia, Canada. This document describes how we calculate the spawn index from spawn survey observations (e.g., spatial extent, number of egg layers, substrate type). There are three types of spawn survey observations: (1) observations of spawn taken from the surface usually at low tide; (2) underwater observations of spawn on giant kelp, *Macrocystis* (*Macrocystis pyrifera*); and (3) underwater observations of spawn on other types of algae and the substrate, which we refer to as ‘understory.’ We calculate the spawn index in four steps. First, we develop a statistical framework and sampling protocol to estimate the number of eggs in a given area. Second, we develop a conversion factor to convert Pacific Herring eggs to biomass. Third, we calculate the spawn index for each spawn survey observation type: surface, *Macrocystis*, and understory. Finally, we add the spawn indices from the three types of survey observations, and aggregate by stock assessment region and year to produce a relative index of combined sex spawning biomass. We also provide an approach to calculate biomass for spawn-on-kelp fisheries. In addition, we identify uncertainties in calculations, suggest future research, and list caveats to using the spawn index. Last, we describe how users can install the **SpawnIndex R** package to calculate the spawn index. Although we transform the spawn survey data from egg density to biomass in tonnes, the resulting data is a relative index of spawning biomass.

RÉSUMÉ

Grinnell, M.H., Schweigert, J.F., Thompson, M., Hawkshaw, S., and Cleary, J.S. 2023. Calculating the spawn index for Pacific Herring (*Clupea pallasii*) in British Columbia, Canada. Can. Tech. Rep. Fish. Aquat. Sci. 3489: viii + 45 p.

L'indice de frai est une composante des évaluations des stocks de hareng du Pacifique (*Clupea pallasii*) en Colombie-Britannique, au Canada. Ce document décrit comment nous calculons l'indice de frai à partir des observations du relevé du frai (par ex., l'étendue spatiale, le nombre de couches d'oeufs, le type de substrat). Il existe trois types d'observations du relevé des frayères: (1) observations des frayères effectuées à la surface, généralement à marée basse; (2) observations sous-marines des frayères sur le varech géant, *Macrocystis* (*Macrocystis pyrifera*); et (3) observations sous-marines des frayères sur les autres algues et le substrat, que nous appelons "sous-étage." Nous calculons l'indice de frai en quatre étapes. Premièrement, nous développons un cadre statistique et un protocole d'échantillonnage pour estimer le nombre d'oeufs dans une zone donnée. Deuxièmement, nous développons un facteur de conversion pour convertir les oeufs en biomasse. Troisièmement, nous calculons l'indice de frai pour chaque types d'observations du relevé des frayères: surface, *Macrocystis* et sous-étage. Enfin, nous additionnons les indices de frai provenant des trois types d'observations de relevés et nous les agrégeons par région d'évaluation des stocks et par année pour produire un indice relatif de la biomasse de reproducteurs de sexe combiné. Nous fournissons également une approche pour calculer la biomasse pour les pêcheries de frai sur varech. En outre, nous identifions les incertitudes dans les calculs, suggérons des recherches futures, et énumérons les mises en garde relatives à l'utilisation de l'indice de frai. Enfin, nous décrivons comment les utilisateurs peuvent installer le paquet **R** **SpawnIndex** pour calculer l'indice de frai. Bien que nous transformions les données de l'enquête sur le frai de la densité des oeufs en biomasse en tonnes, les données résultantes sont un indice relatif de la biomasse de frai.

1 INTRODUCTION

Statistical age-structured stock assessment models rely on an indicator of relative population abundance or biomass to reconstruct a time series of estimated abundance (or biomass). For Pacific Herring (*Clupea pallasii*) in British Columbia (BC), Canada, an index of relative biomass is provided by monitoring the extent and intensity of spawn (i.e., egg) deposition throughout coastal BC (DFO 2022). This document describes our calculations to convert spawn survey observations (e.g., spatial extent, number of egg layers, substrate type) to the Pacific Herring spawn index. These calculations have been described elsewhere, in either published or informal, internal documents. The objective of this document is to collate the calculations in their order of application. Spawn index calculations have been updated over the years as more data and analyses justified improvements; we restrict this document to describing the current method.

Hart and Tester (1934) first demonstrated that an estimate of Pacific Herring biomass could be determined by counting egg deposition in a small set of sampling quadrats. Based on their work, annual spawn surveys collect data used to calculate the spawn index. There are three types of spawn survey observations:

1. Observations of spawn taken from the surface usually at low tide,
2. Underwater observations of spawn on giant kelp, *Macrocystis* (*Macrocystis pyrifera*), and
3. Underwater observations of spawn on other types of algae and the substrate, which we refer to as ‘understory’.


Surface spawn surveys are believed to be the least accurate of the three survey types due to challenges in estimating spawn width (Hay and Kronlund 1987), but they have the greatest temporal and spatial extent (Schweigert 1993). For example, surface spawn surveys were the only survey type prior to 1988, and they are still used extensively for minor spawns, remote spawns (e.g., outside stock assessment region boundaries; see below), unusually early or late spawns, and during exceptional circumstances such as the COVID-19 pandemic. *Macrocystis* and understory spawn surveys are conducted under water using SCUBA gear and have been used for all major spawns since 1988. Thus, we describe the spawn index as having two survey periods based on the predominant spawn survey observation type: the surface survey period from 1951 to 1987 and the dive survey period from 1988 to present.

The intent of Pacific Herring spawn surveys is to monitor spawn throughout the BC coast. Pacific Herring spawn surveys began in 1928, but are considered incomplete for indexing purposes prior to 1937 because many potential areas were not surveyed (Hay and Kronlund 1987). In addition, survey records prior to 1951 are

unreliable for indexing purposes and are excluded from stock assessments. The introduction of dive surveys in 1988 (Schweigert and Haegele 2021) makes it challenging to compare the spawn index between these two periods. For example, surface surveys are less accurate than dive surveys (section A.1) and spawn surveyors used subjective intensity categories instead of direct egg layer estimates until 1978 (section 3.1.1). In addition, Pacific Herring spawn survey effort has been inconsistent over time due to variation in available resources and departmental priorities. For example, prior to 1988, surveyors often dedicated several months each year to spawn surveys: they used small vessels to search for spawn and the surface survey method to estimate egg deposition. Since 1988 surveyors primarily use aircraft to identify spawning activity and underwater SCUBA surveys to estimate egg deposition. Starting in 2018 surveyors have used satellite imagery to identify spawns in remote areas. It typically takes less time to search for spawn using aircraft and more effort to measure egg deposition using SCUBA surveys. Thus, widespread effort (both spatially and temporally) in the past has been replaced with intense effort in the present.

Pacific Herring spawn survey observations have a nested hierarchical structure: samples, *Macrocystis* plants, and quadrats are nested within transects, transects are nested within spawns, and spawns are nested within Locations. To develop spawn indices, Locations are nested within Sections (Figure 1a), Sections are nested within Statistical Areas (Figure 1b), and Statistical Areas are nested within stock assessment regions (SARs; Figure 1c). There are seven SARs in BC, which we categorize as either ‘major’ or ‘minor’ (Figure 2; Haist and Rosenfeld 1988). The terms ‘major’ and ‘minor’ describe relative differences in geographic and biomass scales. The major SARs are Haida Gwaii (formerly known as Queen Charlotte Islands), Prince Rupert District (formerly known as North Coast), Central Coast, Strait of Georgia, and West Coast of Vancouver Island. The minor SARs are Area 27 (A27) and Area 2 West (A2W).

We calculate the spawn index in four main steps. First, we develop a statistical framework (section 2) and sampling protocol (section 3) to estimate the number of eggs in each spawn. Second, we develop a conversion factor to convert Pacific Herring eggs to spawning biomass (section 4), which is critical to spawn index calculations. Third, we calculate the spawn index for each of the three aforementioned spawn survey types: surface (section 5), *Macrocystis* (section 6), and understory (section 7). Note that in this report, we use subsections within sections to separate levels of spatial aggregation (e.g., calculations at the quadrat, or transect level; Figure 3). Finally, we add the spawn indices from the three types of survey observations, and aggregate by SAR and year to produce a relative index of combined sex spawning biomass (section 8).



(a) Spawn index in tonnes (t) in 2021 by Location in Section 142, Statistical Area 14, SoG SAR.
 (b) Sections (Sec) in Statistical Area 14, SoG SAR. Section 142 is emphasized with a thick boundary line.
 (c) Statistical Areas (SA) in the SoG SAR. Statistical Area 14 is emphasized with a thick boundary line.

Figure 1. Pacific Herring spawn index in 2021 by Location (a), as well as Sections (b) and Statistical Areas (c) in the Strait of Georgia (SoG) stock assessment region (SAR; Figure 2).

This document accompanies our **SpawnIndex R** package (RCT 2021) which has functions to calculate the spawn index (section 13). Previously, spawn index calculations were implemented in a **Microsoft Access** database which presented some challenges:

1. The **Access** database has been used for various purposes over the last two decades and has incidental calculations that make it overly complex,
2. The **Access** database is difficult to troubleshoot because it is hard to differentiate between input (i.e., data) and derived values, and
3. It is difficult to determine the equations and parameter values in the **Access** database.

Because of these challenges, we updated the calculations to an **R** package which has several benefits:

1. The **R** package is open and transparent; researchers can view the script and




Figure 2. Spatial boundaries for Pacific Herring stock assessment regions (SARs) in British Columbia. There are five major SARs: Haida Gwaii (HG), Prince Rupert District (PRD), Central Coast (CC), Strait of Georgia (SoG), and West Coast of Vancouver Island (WCVI). There are two minor SARs: Area 27 (A27) and Area 2 West (A2W).

download the package with an example spawn survey database to run functions which implement calculations described in this document,

2. The **R** package clearly distinguishes data from analyses,
3. The **R** package will facilitate future research (e.g., quantify spawn index uncertainty), and
4. The **R** package will allow reproducible research using **knitr** (Xie 2015; Marwick et al. 2018).




Figure 3. Sequence of steps (e.g., S1) for Pacific Herring spawn index calculations for spawn survey observation type $x = \{\text{surface, Macrocytis, understory}\}$ (Table 1). Legend: rounded rectangles indicate start, rectangles indicate observations and calculations, parallelograms indicate output, arrows show order of operation, ‘§’ indicates section, and ‘Eq’ indicates equation.

Essentially, the **R** package is a step towards ‘good enough’ practices in scientific computing (Wilson et al. 2016) for Pacific Herring spawn index calculations.

2 STATISTICAL FRAMEWORK

Each spawn survey observation type has a different statistical framework and sampling protocol (section 3): surface observations, and two types of dive observations (Macrocystis and understory).

2.1 SURFACE SPAWN FRAMEWORK

Historical and recent surface spawn surveys are often opportunistic given particulars of access, weather, and tides, as well as available sampling tools such as boats, rakes, and bathyscopes. Data are analysed assuming simple random sampling.

2.2 DIVE SPAWN FRAMEWORK

Underwater dive surveys using SCUBA gear were instituted in 1988 and follow a two-stage systematic sampling design where transects are the first sampling stage and individual quadrats within transects are the second sampling stage (Jessen 1978). First we describe the understory spawn framework, then we describe the Macrocystis spawn framework which is related to understory, but simpler.

2.2.1 UNDERSTORY SPAWN FRAMEWORK

Two steps are required to calculate mean understory egg density in each surveyed spawn s (Table 1). First, mean egg density for transect t is

$$\bar{\rho}_{txsnry} = \frac{1}{Q_{txsnry}} \sum_{q=1}^{Q_{txsnry}} \rho_{qtxsnry}, \quad (1)$$

where Q_{txsnry} is the number of quadrats, $\rho_{qtxsnry}$ is egg density in thousands of eggs per square metre ($10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$; Table 2), and $\bar{\rho}_{txsnry}$ is in $10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$. Throughout this report, ‘ $\bar{-}$ ’ indicates the mean of the variable that it covers (e.g., $\bar{\rho}$ is the mean of ρ in Equation 1). Note that we report eggs in thousands (i.e., $10^3 \cdot \text{eggs}$) in this document and in the **R** package (section 13). Before we calculate mean egg density for spawn s , we determine the mean number of potential quadrats for survey type x in spawn s

$$\bar{Q}'_{xsnry} = \frac{1}{T_{xsnry}} \sum_{t=1}^{T_{xsnry}} Q'_{txsnry}, \quad (2)$$

Table 1. Index notation for Pacific Herring spawn index calculations. Legend: stock assessment region (SAR) and spawn-on-kelp (SOK). Note that ‘‘ indicates related variables (e.g., T' is related to T).

Name	Description	Range
y	Year	y_1, y_2, y_3, \dots, Y
Y	Last year of time series	
r	SAR	$1, 2, 3, \dots, R$
R	Number of SARs	
n	Location	$1, 2, 3, \dots, N_{ry}$
N_{ry}	Number of locations	
s	Spawn	$1, 2, 3, \dots, S_{nry}$
S_{nry}	Number of spawns	
x	Spawn survey observation type	$1, 2, 3, \dots, X_{snry}$
X_{snry}	Number of spawn survey observation types	
j	Sample	$1, 2, 3, \dots, J_{xsnry}$
J_{xsnry}	Number of samples	
i	Substrate type (surface)	$1, 2, 3, \dots, I_{jxsnry}$
I_{jxsnry}	Number of substrate types (surface)	
t	Transect	$1, 2, 3, \dots, T_{xsnry}$
T_{xsnry}	Number of transects	
T'_{xsnry}	Number of potential transects	
p	Plant	$1, 2, 3, \dots, P_{txsnry}$
P_{txsnry}	Number of plants	
k	Stalk	$1, 2, 3, \dots, K_{ptxsnry}$
$K_{ptxsnry}$	Number of stalks	
q	Quadrat	$1, 2, 3, \dots, Q_{txsnry}$
Q_{txsnry}	Number of quadrats	
Q'_{txsnry}	Number of potential quadrats	
i'	Substrate type (understory)	$1, 2, 3, \dots, I'_{qtxsnry}$
$I'_{qtxsnry}$	Number of substrate types (understory)	
v	Algae (i.e., vegetation) type	$1, 2, 3, \dots, V_{qtxsnry}$
$V_{qtxsnry}$	Number of algae types	
g	SOK fishery	$1, 2, 3, \dots, G_{ry}$
G_{ry}	Number of SOK fisheries	

where T_{xsnry} is the number of transects, Q'_{txsnry} is the number of potential quadrats (i.e., based on spawn width), and \bar{Q}'_{xsnry} is in number of quadrats. Then, we calculate mean egg density in eggs per square metre for survey type x in spawn s

$$\bar{\rho}_{xsnry} = \frac{1}{T_{xsnry} \bar{Q}'_{xsnry}} \sum_{t=1}^{T_{xsnry}} Q'_{txsnry} \bar{\rho}_{txsnry}, \quad (3)$$

where T_{xsnry} is the number of transects, \bar{Q}'_{xsnry} is the mean number of potential quadrats from Equation 2, Q'_{txsnry} is the number of potential quadrats, $\bar{\rho}_{txsnry}$ is the mean transect egg density from Equation 1, and $\bar{\rho}_{xsnry}$ is in $10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$. The egg density estimator from Equation 3 is unbiased, and the variance is

$$\begin{aligned} \sigma_{xsnry}^2 &= \frac{T'_{xsnry} - T_{xsnry}}{T_{xsnry} T'_{xsnry}} \sum_{t=1}^{T_{xsnry}} \frac{(Q'_{txsnry} \bar{\rho}_{txsnry} - \bar{Q}'_{xsnry} \bar{\rho}_{xsnry})^2}{\bar{Q}'_{xsnry}^2 (T_{xsnry} - 1)} \\ &\quad + \frac{f'_{xsnry}}{T_{xsnry}^2} \sum_{t=1}^{T_{xsnry}} \left(\frac{Q'_{txsnry}}{\bar{Q}'_{xsnry}} \right)^2 \frac{(1 - f''_{txsnry}) \sigma_{txsnry}^2}{Q_{txsnry}}, \end{aligned} \quad (4)$$

where T'_{xsnry} is the number of potential transects (i.e., based on spawn length), T_{xsnry} is the number of transects, Q'_{txsnry} is the number of potential quadrats, $\bar{\rho}_{txsnry}$ is the mean egg density from Equation 1, \bar{Q}'_{xsnry} is the mean number of potential quadrats from Equation 2, $\bar{\rho}_{xsnry}$ is the mean spawn egg density from Equation 3, f'_{xsnry} is the transect sampling fraction for survey type x in spawn s

$$f'_{xsnry} = \frac{T_{xsnry}}{T'_{xsnry}}, \quad (5)$$

f''_{txsnry} is the quadrat sampling fraction for transect t

$$f''_{txsnry} = \frac{Q_{txsnry}}{Q'_{txsnry}}, \quad (6)$$

where Q_{txsnry} is the number of quadrats and σ_{txsnry}^2 is the within transect egg density variance

$$\sigma_{txsnry}^2 = \frac{1}{Q_{txsnry} - 1} \sum_{q=1}^{Q_{txsnry}} (\rho_{qtxsnry} - \bar{\rho}_{txsnry})^2, \quad (7)$$

where $\rho_{qtxsnry}$ is egg density and σ_{txsnry}^2 is in units of ρ^2 .

Table 2. Notation for Pacific Herring spawn survey statistical framework. Legend: metres (m). Note that “ \prime ” indicates related variables (e.g., f' is related to f'').

Name	Description	Value or unit
ρ	Egg density	$10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$
f'	Transect sampling fraction	$0 < f' \leq 1$
f''	Quadrat sampling fraction	$0 < f'' \leq 1$
σ^2	Egg density variance	ρ^2

The calculation of the mean egg density for each spawn requires estimates of total spawn length, mean spawn width, length of each transect sampled, and estimated egg density in each sampling quadrat. The sampling protocol was determined through a series of studies conducted in the Strait of Georgia in 1981 and 1983 (Schweigert et al. 1985, 1990), and on the West Coast of Vancouver Island in 1982 (Schweigert et al. 1990). In the 1981 study, the location of transects and sampling quadrats along transects was determined using random allocation (Schweigert et al. 1985). However, this proved to be logically difficult because neither the spawn length or width is known a priori and divers had difficulty making the necessary calculations underwater. Nevertheless, data from these studies were used to determine a sampling protocol to estimate mean egg density with standard error $\text{SE} \leq 25\%$. The results indicated that the sampling required to achieve this level of precision included surveying three transects per kilometre of spawn length and sampling at least five quadrats per transect (i.e., spawn width). The sampling design was tested during a 1983 survey in the Strait of Georgia that applied a systematic rather than a random sampling protocol to simplify the logistics; variance estimates were similar to those from the 1981 study. This sampling protocol was further re-evaluated after additional surveys occurred in all areas of the coast during 1984 and 1985; the protocol was found to be robust and has been in routine use since 1988 (Schweigert et al. 1990). Although samples are collected systematically within each spawn, we assume that transects and quadrats are located randomly with respect to the underlying spawn distribution, and so these estimators are applicable (Schaeffer et al. 2012).

2.2.2 MACROCYSTIS SPAWN FRAMEWORK

Giant kelp, *Macrocystis* (*Macrocystis pyrifera*), requires a different framework than the understory spawn framework using quadrats. *Macrocystis* plants are unsuitable for quadrat sampling because *Macrocystis* plants routinely reach heights of 15 m,

and once weighed down with Pacific Herring eggs the plants can sink to lay flat on the substrate. After sampling Pacific Herring eggs on *Macrocystis* plants, Haegele and Schweigert (1990) determined that plant height, number of fronds per plant, and number egg layers per plant were key counts required to estimate the number of eggs per plant. The survey design employed to capture these data relies on determining the average plant height, number of fronds in each plant holdfast, and number of *Macrocystis* plants occurring within a 1 m swath on both sides of the transect line in a given spawn. These data are used to determine the total egg deposition on *Macrocystis* plants for each spawn (section 6.3).

3 SAMPLING PROTOCOL

The following is a brief summary of the spawn survey sampling protocol from the [Pacific Herring spawn survey manual](#). Pacific Herring in BC primarily spawn in sheltered bays and inlets, depositing eggs on rocks and algae between depths of 1.5 m above and 18 m below the 0-tide level (Humphreys and Hourston 1978; Haegele and Schweigert 1985). We identify distinct spawns (both spatially and temporally) by a unique combination of year, Location, and ‘spawn number’ $s = 1, 2, 3, \dots, S_{nry}$ (Table 1). Spawn numbers identify distinct spawns, which we define as a continuous stretch of shoreline with no detectable break in egg deposition. For a given survey type x , a spawn s is the finest scale at which we calculate the spawn index. A break in egg deposition is determined by the absence of Pacific Herring spawn on two consecutive transects, or by a temporal gap in spawning. Most spawns are also characterized by longitude and latitude, as well as start and end dates of spawning. Surveyors usually collect longitude and latitude at the start and end of each transect; for surface spawn surveys that are not along transects (section 3.1), surveyors collect longitude and latitude at the start and end of the spawn (i.e., overall length and width).

Pacific Herring spawns typically extend along the shore; from above, spawns are identified by a milky or turquoise discolouration of the ocean caused by the release of milt, and often appear as bands running parallel to the shore (Figure 4). Thus, spawn ‘length’ refers to distance parallel to the shore and spawn ‘width’ refers to distance perpendicular to the shore. Similarly, *Macrocystis* bed length L' and algae (i.e., vegetation) bed length L'' refer to distances that *Macrocystis* and algae beds extend parallel to the shore, respectively.

Most areas of the BC coast have ‘permanent transect’ locations recorded on charts which enable surveyors to place transects in the same place each year. When permanent transects are unavailable for a given spawn, surveyors set new transects




Figure 4. Aerial view of Pacific Herring spawn taken during a spawn reconnaissance flight in the Strait of Georgia. Spawn is identified by the band of discoloured water parallel to the shore.

perpendicular to the shore, beginning 200 m in from one end of the spawn, and spaced 350 m apart along the length of the spawn. The end of the spawn is determined by the absence of eggs. We digitize new transects to make them available as permanent transects in subsequent surveys. Transects generally go from the deep edge of the spawn towards shore until divers reach the near-shore edge of the spawn, which can be on the beach depending on the tide height.

Pacific Herring spawn surveyors first determine the spatial extent of the spawn in terms of length of shoreline to survey (Figure 3, steps S1, M1, and U1); this is done by raking (section 3.1) or brief dives to determine the presence or absence of spawn. Surveyors place the first transect perpendicular to the shore, in from one end (i.e., at the first permanent transect, or 200 m if there are no permanent transects) to avoid surveying areas with patchy and sparse egg layers. Within the spawn area, surveyors use transects to determine spawn width, quadrat placement, and which *Macrocystis* plants to survey. In some cases, we adjust observed spawn width to increase the accuracy of spawn index estimates (Appendix A).

After determining the spatial extent, surveyors determine the number of egg layers on substrate and algae according to sampling protocols described in section 3.1, section 3.2, and section 3.3. For eggs on substrate, one egg layer is a layer of eggs one

egg thick over the entire spawned surface (Figure 5a). For eggs on algae, surveyors count egg layers one of two ways depending on whether the algae is flat or round in cross-section. Egg layers on flat algae are counted on both sides of the algae (Figure 5b), while egg layers on round algae are counted across the diameter of the algae (Figure 5c).

3.1 SURFACE SPAWN PROTOCOL

Surface spawn surveyors use the aforementioned transect interval when possible, but the sampling interval relies on surveyor judgement and available resources. If the spawn area is sufficiently large, surface surveyors usually sample along permanent transects. Small spawns can still be mapped as they were historically, with surveyors deciding how to sample the spawn. To determine spawn coverage and intensity, surveyors deploy specialized rakes throughout the spawn to determine algae (i.e., vegetation) type, number of egg layers, and percent cover. Surveyors may deploy a bathyscope in shallow water, and at low tide a portion of the spawn may be visible for direct observation. We refer to each of these surface spawn observations as ‘samples.’ Recall that there are two cases of surface spawn surveys: all surveys prior to 1988 and surveys since 1988 when dive surveys are not possible.




Figure 5. Cross-sections showing the number of Pacific Herring egg layers on substrate (a), flat algae (b), and round algae (c). Diagrams copied with permission from the [Pacific Herring spawn survey manual](#).

3.1.1 SPAWN INTENSITY CATEGORIES

From 1928 to 1978, surface spawn surveyors categorized spawn by subjective ‘intensity’ categories instead of direct egg layer estimates (Table 3). From 1928 to 1968 there were five intensity categories described as very light, light, medium, heavy, and very heavy (numbered 1 to 5, respectively). Starting in 1969 there were nine intensity categories; the change from five to nine categories was probably to accommodate the practice of reporting intermediate categories such as 3.5 (Hay and Kronlund 1987). Starting in 1979, spawn surveyors estimated the number of egg layers directly and continued to record intensity categories until 1981 to provide overlap between the two methods. In addition to recording the number of egg layers, surveyors sometimes recorded intensity after it was officially discontinued in 1981. We have converted spawn intensity observations in the Pacific Herring spawn survey database from five to nine categories for spawns that used the five category scale between 1951 and 1968. Thus, spawn data used for stock assessments are represented either by a nine-category intensity scale or a direct estimate of the number of egg layers.

Table 3. Spawn intensity categories, description, and number of egg layers for Pacific Herring surface spawn surveys for the periods 1928 to 1968, and 1969 to 1978 (Hay and Kronlund 1987; Schweigert and Stocker 1988). Uncertainty in the number of egg layers is not available.

Intensity category		Description	Number of egg layers
1928–1968	1969–1978		
1	1	Very light	0.5529
	2		0.9444
2	3	Light	1.3360
	4		2.1496
3	5	Medium	2.9633
	6		4.1318
4	7	Heavy	5.3002
	8		6.5647
5	9	Very heavy	7.8291

3.2 MACROCYSTIS SPAWN PROTOCOL

Macrocystis spawn surveyors take a census of Macrocystis plants within 1 m of the transect line, on both the left- and right-hand sides. We refer to the swath of substrate along Macrocystis transects as the transect swath $\chi_{txsnry} = 2$ m. Divers categorize Macrocystis plants as either ‘mature’ or ‘immature’ based on stipe height; mature plants have stipes ≥ 1 m high, and are the only plants used for Macrocystis spawn index calculations. Immature plants are excluded because Pacific Herring spawn on Macrocystis fronds, not stipes; immature plants have limited fronds and slimy stipes that prevent egg adhesion. In addition, Pacific Herring typically deposit spawn higher up Macrocystis plants. For each mature plant, divers record the number of stalks. For each transect, divers record average number of egg layers and average plant height. Haegele and Schweigert (1990) provide a description of the sampling technique and the basis for estimating the total number of eggs per plant.

3.3 UNDERSTORY SPAWN PROTOCOL

Understory spawn surveyors place quadrats along transects, with a target frequency of ≥ 5 quadrats per transect, given a minimum spacing of 2 m and a maximum spacing of 40 m. Similar to how the first transect is moved in from one end of the spawn, the first quadrat is moved in from the edge of the spawn to the first 5 m mark on the transect line, to avoid surveying areas with patchy and sparse egg layers. Note that quadrat position along permanent transects varies among years due to the location and extent (i.e., width) of spawn with respect to the shoreline: spawn location causes quadrats to shift seaward or shoreward and spawn width causes transects to be shorter or longer. Understory spawn surveys use 0.5 m² quadrats; other sizes (e.g., 0.25 and 1.0 m²) have been used for research, but are not used to calculate the spawn index (Schweigert 1993). Within each quadrat, divers record the dominant (i.e., most heavily spawned) substrate type, proportion of the quadrat covered by spawn, and number of egg layers. In addition, divers identify the three most abundant algae types that have spawn. For each of these algae types, divers record the proportion of the quadrat covered by algae and the number of egg layers.

4 CONVERT EGGS TO BIOMASS

After estimating the number of eggs by survey type x in spawn s , we use a conversion factor to estimate the biomass of Pacific Herring that spawned to produce those eggs. Female Pacific Herring produce an average of approximately 200 000 eggs per kilogram (kg) of total body weight (Hay 1985); we refer to this as fecundity. Average

fecundity is derived from studies of BC Pacific Herring in 1974 and 1980 (Prince Rupert District, Strait of Georgia, and West Coast of Vancouver Island; Hay 1985), and California Pacific Herring in 1975 (Rabin and Barnhart 1975). We assume that females account for 50% of spawners, and we convert eggs to tonnes (t) of spawners using

$$\theta = \omega \varphi \frac{10^3 \text{ kg}}{t}, \quad (8)$$

where ω is female fecundity, φ is the proportion of spawners that are female, and θ is the egg conversion factor in $10^8 \cdot \text{eggs} \cdot t^{-1}$ (Table 4). Thus, we convert eggs to biomass of both sexes combined in tonnes by dividing the number of eggs by θ . Although Pacific Herring egg production is affected by environmental variability and other factors (Tanasichuk and Ware 1987; Hay and Brett 1988), we assume that bias to the spawn index from the effect of these factors on Equation 8 is negligible (Ware 1985; Schweigert 1993).

5 SURFACE SPAWN CALCULATIONS

This section describes steps S2 to S4 in Figure 3 (i.e., survey type $x = \text{surface}$). As previously mentioned, surface spawn surveyors sample along transects or using their judgement (section 3.1). Surveyors collect data at the substrate type i , sample j , and spawn s levels; we calculate metrics at the substrate type i , sample j , and spawn s levels (Table 1 and Table 5). We use the term ‘metric’ to refer to a measure of quantitative assessment. Recall that surface spawn ‘samples’ include observations collected using specialized rakes and viewing boxes (section 3.1). Occasionally, we

Table 4. Notation for converting the number of Pacific Herring eggs to spawning biomass. Legend: kilograms (kg) and tonnes (t).

Name	Description	Value or unit	References
ω	Female fecundity	$200,000 \text{ eggs} \cdot \text{kg}^{-1}$	Rabin and Barnhart (1975); Hay (1985); Hay and Brett (1988)
φ	Proportion female	0.5	
θ	Egg conversion factor	$10^8 \cdot \text{eggs} \cdot t^{-1}$	

use field data sheets to fill-in missing egg layer information for surface survey data (Appendix B).

5.1 SAMPLE OBSERVATIONS AND CALCULATIONS

Each sample j (step S2 in Figure 3) can have one or more substrate types i . The number of egg layers in substrate i is

$$E_{ijxsny} = E'_{ijxsny} \phi_{ijxsny}, \quad (9)$$

where E'_{ijxsny} is the number of egg layers, ϕ_{ijxsny} is the proportion of substrate i covered with spawn, and E_{ijxsny} is in number of egg layers. The total number of egg layers in sample j is

$$E_{jxsny} = \sum_{i=1}^{I_{jxsny}} E_{ijxsny}, \quad (10)$$

where E_{ijxsny} is the number of egg layers from Equation 9 and E_{jxsny} is in number of egg layers. For the time period when surveyors recorded spawn ‘intensity’ categories instead of direct egg layer estimates, we convert intensity to number of egg layers E_{jxsny} (Table 3). Schweigert et al. (1997) developed a model of surface egg density

Table 5. Notation for Pacific Herring surface spawn index calculations. Legend: metres (m) and tonnes (t). Note that ‘‘ indicates related variables (e.g., E' is related to E).

Name	Description	Value or unit	References
E'	Number of egg layers	$E' > 0$	
ϕ	Proportion covered in eggs	$0 < \phi \leq 1$	
E	Number of egg layers	$E > 0$	
α	Regression intercept	$14.698 \cdot 10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$	Schweigert et al. (1997)
β	Regression slope	$212.218 \cdot 10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$	Schweigert et al. (1997)
ρ	Egg density	$10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$	
L	Spawn length	m	
W	Spawn width	m	
B'	Surface spawn index (i.e., relative biomass)	t	

as a function of number of egg layers using a linear regression model¹

$$\rho_{jxsnry} = \alpha + \beta E_{jxsnry}, \quad (11)$$

where α is the regression intercept, β is the regression slope, E_{jxsnry} is the total number of egg layers from Equation 10, and ρ_{jxsnry} is in $10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$ (Figure 6). Note that $E_{jxsnry} = 0 \Rightarrow \rho_{jxsnry} = 0$ to avoid having $\rho_{jxsnry} = \alpha$ when there are no egg layers (i.e., $E_{jxsnry} = 0$).

5.2 SPAWN OBSERVATIONS AND CALCULATIONS

The mean egg density for survey type x in spawn s (step S3 in Figure 3) is

$$\bar{\rho}_{xsnry} = \frac{1}{J_{xsnry}} \sum_{j=1}^{J_{xsnry}} \rho_{jxsnry}, \quad (12)$$

where J_{xsnry} is the number of samples, ρ_{jxsnry} is egg density from Equation 11 and $\bar{\rho}_{xsnry}$ is in $10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$. Two other metrics are required at the spawn level: spawn length L_{xsnry} and spawn width W_{xsnry} , both in metres. We set W_{xsnry} to the first

¹There is an error in Schweigert et al. (1997); surface egg density is in thousands per square metre ($10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$).




Figure 6. Egg density in thousands of eggs per square metre (m; line) as a function of number of egg layers for Pacific Herring surface spawn surveys; based on Equation 11 (Schweigert et al. 1997). Note that the number of egg layers can exceed those shown in this figure.

non-missing value of median pool width, median Section width, median SAR width, or observed width W'_{xsnry} (in that order; section A.1). A pool is a group of Locations within a Section that are often adjacent, contain similar algae and substrate, and can be treated as a group with likely similar widths.

The surface spawn index in spawn s is

$$B'_{xsnry} = \frac{\bar{\rho}_{xsnry} L_{xsnry} W_{xsnry} 10^3}{\theta}, \quad (13)$$

where $\bar{\rho}_{xsnry}$ is mean egg density in spawn s from Equation 12, L_{xsnry} is spawn length, W_{xsnry} is spawn width, θ is the egg conversion factor from Equation 8, and B'_{xsnry} is in tonnes (step S4 in Figure 3).

6 MACROCYSTIS SPAWN CALCULATIONS

This section describes steps M2 to M5 in Figure 3 (i.e., survey type $x = \text{Macrocystis}$). Macrocytis spawn surveyors use SCUBA gear to collect underwater data for individual plants p , transects t , and spawns s (section 3.2), where divers enumerate every Macrocytis plant within the transect swath χ . We calculate metrics at the transect t and spawn s levels (Table 1 and Table 6).

6.1 PLANT OBSERVATIONS

For each mature plant p (step M2 in Figure 3), surveyors count the number of stalks $K_{ptxsnry}$.

6.2 TRANSECT OBSERVATIONS AND CALCULATIONS

At the transect t level (step M3 in Figure 3), spawn width is W_{txsnry} and transect swath is $\chi_{txsnry} = 2\text{ m}$, both in metres. We calculate the area in transect t as

$$A_{txsnry} = W_{txsnry} \chi_{txsnry} \quad (14)$$

where W_{txsnry} is spawn width in metres, χ_{txsnry} is transect swath in metres, and A_{txsnry} is in square metres. In addition, divers estimate summary statistics for mature Macrocytis plants p along transect t : mean height \bar{H}_{txsnry} in metres and mean number of egg layers \bar{E}_{txsnry} . We calculate the total number of stalks K in transect t as

$$K_{txsnry} = \sum_{p=1}^{P_{txsnry}} K_{ptxsnry}, \quad (15)$$

where $K_{ptxsnry}$ is the number of stalks and K_{txsnry} is in number of stalks.

Table 6. Notation for Pacific Herring *Macrocystis* spawn index calculations. Legend: metres (m) and tonnes (t). Note that ‘‘ indicates related variables (e.g., L' is related to L).

Name	Description	Value or unit	References
W	Spawn width	m	
χ	Transect swath	2 m	
A	Area	m^2	
H	Plant height	m	
E	Number of egg layers	$\overline{E} > 0$	
L'	Macrocystis bed length	m	
L	Spawn length	m	
κ	Number of stalks per plant	$\bar{\kappa} > 0$	
ξ	Regression slope	$0.073 \text{ eggs} \cdot 10^3 \cdot \text{plant}^{-1}$	Haegele and Schweigert (1990)
γ	Regression exponent on E	0.673	Haegele and Schweigert (1990)
δ	Regression exponent on H	0.932	Haegele and Schweigert (1990)
ϵ	Regression exponent on $\bar{\kappa}$	0.703	Haegele and Schweigert (1990)
ψ	Number of eggs per plant	$\text{eggs} \cdot 10^3 \cdot \text{plant}^{-1}$	
ρ	Egg density	$10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$	
B'	Macrocystis spawn index (i.e., relative biomass)	t	

6.3 SPAWN OBSERVATIONS AND CALCULATIONS

At the spawn s level (step M4 in Figure 3), the *Macrocystis* bed length is L'_{xsnry} in metres. If L'_{xsnry} is inadvertently not recorded, we set L'_{xsnry} to the spawn length L_{xsnry} . The mean width for survey type x in spawn s is

$$\overline{W}_{xsnry} = \frac{1}{T_{xsnry}} \sum_{t=1}^{T_{xsnry}} W_{txsnry}, \quad (16)$$

where T_{xsnry} is the number of transects, W_{txsnry} is the spawn width, and \bar{W}_{xsnry} is in metres. The total area of transects for survey type x in spawn s is

$$A_{xsnry} = \sum_{t=1}^{T_{xsnry}} A_{txsnry}, \quad (17)$$

where A_{txsnry} is the transect area from Equation 14, and A_{xsnry} is in square metres.

Three metrics are required to calculate the number of eggs on *Macrocystis* plants (Haegele and Schweigert 1990): number of egg layers, plant height, and number of stalks per plant. First, the mean number of egg layers for survey type x in spawn s is

$$\bar{E}_{xsnry} = \frac{1}{T_{xsnry}} \sum_{t=1}^{T_{xsnry}} \bar{E}_{txsnry}, \quad (18)$$

where T_{xsnry} is the number of transects in spawn s , \bar{E}_{txsnry} is the mean number of egg layers t , and \bar{E}_{xsnry} is in number of egg layers. Second, the mean plant height in spawn s is

$$\bar{H}_{xsnry} = \frac{1}{T_{xsnry}} \sum_{t=1}^{T_{xsnry}} \bar{H}_{txsnry}, \quad (19)$$

where T_{xsnry} is the number of transects, \bar{H}_{txsnry} is the mean plant height, and \bar{H}_{xsnry} is in metres. The third metric is the number of stalks per plant, which we calculate in three steps. The total number of observed stalks for survey type x in spawn s is

$$K_{xsnry} = \sum_{t=1}^{T_{xsnry}} K_{txsnry}, \quad (20)$$

where K_{txsnry} is the number of stalks from Equation 15 and K_{xsnry} is in number of stalks. The total number of observed plants for survey type x in spawn s is

$$P_{xsnry} = \sum_{t=1}^{T_{xsnry}} P_{txsnry}, \quad (21)$$

where P_{txsnry} is the number of plants and P_{xsnry} is in number of plants. Thus, the mean number of stalks per plant for survey type x in spawn s is

$$\bar{\kappa}_{xsnry} = \frac{K_{xsnry}}{P_{xsnry}}, \quad (22)$$

where K_{xsnry} is the number of stalks from Equation 20, P_{xsnry} is the number of plants from Equation 21, and $\bar{\kappa}_{xsnry}$ is in number of stalks per plant.

Haegele and Schweigert (1990) developed a model of the number of eggs per plant as a function of the three aforementioned metrics (number of egg layers, plant height, and number of stalks per plant) using a nonlinear multiple regression model

$$\bar{\psi}_{xsnry} = \xi \bar{E}_{xsnry}^{\gamma} \bar{H}_{xsnry}^{\delta} \bar{\kappa}_{xsnry}^{\epsilon} 10^3, \quad (23)$$

where ξ is the regression slope, \bar{E}_{xsnry} is the mean number of egg layers from Equation 18, γ is the regression exponent on \bar{E}_{xsnry} , \bar{H}_{xsnry} is the mean plant height from Equation 19, δ is the regression exponent on \bar{H}_{xsnry} , $\bar{\kappa}_{xsnry}$ is the mean number of stalks per plant from Equation 22, ϵ is the regression exponent on $\bar{\kappa}_{xsnry}$, and $\bar{\psi}_{xsnry}$ is in eggs $\cdot 10^3 \cdot \text{plant}^{-1}$ (Figure 7). Mean Macrocystis egg density in spawn s is

$$\bar{\rho}_{xsnry} = \frac{\bar{\psi}_{xsnry} P_{xsnry}}{A_{xsnry}}, \quad (24)$$

where $\bar{\psi}_{xsnry}$ is the mean number of eggs per plant from Equation 23, P_{xsnry} is the number of plants from Equation 21, A_{xsnry} is the total area of transects from Equation 17, and $\bar{\rho}_{xsnry}$ is in $10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$.

The Macrocystis spawn index in spawn s is

$$B'_{xsnry} = \frac{\bar{\rho}_{xsnry} L'_{xsnry} \bar{W}_{xsnry} 10^3}{\theta}, \quad (25)$$

where $\bar{\rho}_{xsnry}$ is the mean egg density from Equation 24, L'_{xsnry} is the Macrocystis bed length, \bar{W}_{xsnry} is the mean spawn width from Equation 16, θ is the egg conversion factor from Equation 8, and B'_{xsnry} is in tonnes (step M5 in Figure 3).

7 UNDERSTORY SPAWN CALCULATIONS

This section describes steps U2 to U5 in Figure 3 (i.e., survey type x = understory). Understory spawn surveyors use SCUBA gear to collect underwater data for substrate i' , algae (i.e., vegetation) types v , quadrats q , transects t , and spawns s (section 3.3); we calculate metrics at the substrate i' , algae type v , quadrat q , transect t , and spawn s levels (Table 1 and Table 7).

7.1 QUADRAT OBSERVATIONS AND CALCULATIONS

We calculate two separate estimates of egg density at the quadrat level (step U2 in Figure 3): spawn on substrate i' and spawn on algae v . Haegele et al. (1979) developed a model of substrate i' egg density from egg layers using a linear model

$$\rho_{i'qtxsnry} = \varphi E_{i'qtxsnry} \phi_{i'qtxsnry}, \quad (26)$$




Figure 7. Number of eggs in thousands per *Macrocystis* plant (lines) as a function of number of egg layers on plants, plant height in metres (m), and number of stalks per plant for Pacific Herring *Macrocystis* spawn surveys; based on Equation 23 (Haegle and Schweigert 1990). Note that the number of egg layers, plant height, and number of stalks per plant can exceed those shown in this figure.

where φ is the regression slope, $E_{i'qtxsnry}$ is the number of egg layers, $\phi_{i'qtxsnry}$ is the proportion of substrate covered by spawn, and $\rho_{i'qtxsnry}$ is substrate egg density in $10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$ (Figure 8).

Although quadrats have only one substrate type, they can have up to three algae

Table 7. Notation for Pacific Herring understory spawn index calculations. Legend: metres (m) and tonnes (t). Note that ‘‘’ indicates related variables (e.g., L'' is related to L).

Name	Description	Value or unit	References
φ	Regression slope for substrate	$340 \cdot 10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$	Haegele et al. (1979)
E	Number of egg layers	$E > 0$	
ϕ	Proportion covered in eggs	$0 < \phi \leq 1$	
ρ	Egg density	$10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$	
ϑ	Regression slope for algae	$600.567 \cdot 10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$	Schweigert (2005)
ϱ	Regression exponent on E	0.6355	Schweigert (2005)
ς	Regression exponent on ϕ	1.413	Schweigert (2005)
C	Algae coefficient	See Table 8	
W	Spawn width	m	
ρ	Egg density	$10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$	
L''	Algae bed length	m	
L	Spawn length	m	
B'	Understory spawn index (i.e., relative biomass)	t	

types v (section 3.3). Schweigert (2005) developed a model of algae egg density from egg layers, proportion of the quadrat covered by algae, and algae coefficients using a generalized linear model. Algae coefficients account for the effect of algae morphology on Pacific Herring egg density (Table 8). Egg density on algae v (Schweigert 2005) is

$$\rho_{vqtxsnry} = \vartheta E_{vqtxsnry}^\varrho \phi_{vqtxsnry}^\varsigma C_v, \quad (27)$$

where ϑ is the regression slope, $E_{vqtxsnry}$ is the number of egg layers, ϱ is the regression exponent on $E_{vqtxsnry}$, $\phi_{vqtxsnry}$ is the proportion covered by algae, ς is the regression exponent on $\phi_{vqtxsnry}$, C_v is the coefficient for algae v , and $\rho_{vqtxsnry}$ is in $10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$ (Figure 9).




Figure 8. Substrate egg density in thousands of eggs per square metre (m; lines) as a function of number of egg layers on substrate and proportion of substrate covered in spawn for Pacific Herring underwater spawn surveys; based on Equation 26 (Haegle et al. 1979). Note that the number of egg layers can exceed those shown in this figure.

Table 8. Algae (i.e., vegetation) types v and coefficients C for Pacific Herring understory spawn surveys (Schweigert 2005). Uncertainty in algae coefficients is not available. Algae types are described in the [Pacific Herring spawn survey manual](#).

Algae type v	Coefficient C
Grasses	0.9715
Grunge	1.0000
Kelp (flat)	0.9119
Kelp (standing)	1.1766
Leafy algae	0.6553
Rockweed	0.7793
Sargassum	1.1766
Stringy algae	1.0000




Figure 9. Algae egg density in thousands of eggs per square metre (m; lines) as a function of number of egg layers on algae, proportion of substrate covered in algae, and algae coefficients (Table 8) for Pacific Herring underwater spawn surveys; based on Equation 27 (Schweigert 2005). Note that the number of egg layers can exceed those shown in this figure.

The total understory egg density for quadrat q is

$$\rho_{qtxsnry} = \rho_{i'qtxsnry} + \sum_{v=1}^{V_{qtxsnry}} \rho_{vqtxsnry}, \quad (28)$$

where $\rho_{i'qtxsnry}$ is substrate egg density from Equation 26, $\rho_{vqtxsnry}$ is algae egg density from Equation 27, and $\rho_{qtxsnry}$ is in $10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$. Thus, we assume that eggs on substrate and algae are independent, and can be combined.

7.2 TRANSECT OBSERVATIONS AND CALCULATIONS

At the transect level (step U3 in Figure 3), the mean understory egg density in transect t is

$$\bar{\rho}_{txsnry} = \frac{1}{Q_{txsnry}} \sum_{q=1}^{Q_{txsnry}} \rho_{qtxsnry}, \quad (29)$$

where Q_{txsnry} is the number of quadrats, $\rho_{qtxsnry}$ is total understory egg density from Equation 28, and $\bar{\rho}_{txsnry}$ is in $10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$. Note that we update spawn width to correct for errors regarding the assumed accuracy of transect lines used to measure spawn width for understory surveys between 2003 and 2014 (section A.2).

7.3 SPAWN OBSERVATIONS AND CALCULATIONS

At the spawn level (step U4 in Figure 3), the mean width of survey type x in spawn s is

$$\bar{W}_{xsnry} = \frac{1}{T_{xsnry}} \sum_{t=1}^{T_{xsnry}} W_{txsnry}, \quad (30)$$

where T_{xsnry} is the number of transects, W_{txsnry} is the spawn width, and \bar{W}_{xsnry} is in metres. The algae bed length is L''_{xsnry} , also in metres. As with *Macrocystis* spawn calculations, if L''_{xsnry} is inadvertently not recorded, we set L''_{xsnry} to the spawn length L_{xsnry} . Thus, we assume that eggs on substrate and eggs on algae are represented by the same length measurement.

Next, we calculate the weighted mean egg density in spawn s , where transect egg density is weighted by spawn width at transect t , W_{txsnry} . We calculate a weighted mean because spawn width varies along the spawn length; a weighted mean ensures that transects contribute proportionally to their area. The mean egg density for survey type x in spawn s is

$$\bar{\rho}_{xsnry} = \frac{\sum_{t=1}^{T_{xsnry}} \bar{\rho}_{txsnry} W_{txsnry}}{\sum_{t=1}^{T_{xsnry}} W_{txsnry}}, \quad (31)$$

where $\bar{\rho}_{txsnry}$ is the mean understory egg density from Equation 29, W_{txsnry} is the spawn width, and $\bar{\rho}_{xsnry}$ is in $10^3 \cdot \text{eggs} \cdot \text{m}^{-2}$.

The understory spawn index in spawn s is

$$B'_{xsnry} = \frac{\bar{\rho}_{xsnry} L''_{xsnry} \bar{W}_{xsnry} 10^3}{\theta}, \quad (32)$$

where $\bar{\rho}_{xsnry}$ is the mean understory egg density from Equation 31, L''_{xsnry} is the algae bed length, \bar{W}_{xsnry} is the mean spawn width from Equation 30, θ is the egg conversion factor from Equation 8, and B'_{xsnry} is in tonnes (step U5 in Figure 3).

8 TOTAL SPAWN CALCULATIONS

This section describes step T1 in Figure 3. Recall that spawn index calculations have a nested hierarchical structure (e.g., substrate type i is nested within sample j for surface spawn calculations in section 5). However, quantities at the survey type x in spawn s level are not necessarily nested and hierarchical. For example, B'_{xsnry} is the finest scale that we calculate the spawn index (section 3), which we could aggregate by any combination of survey type x , spawn s , location n , region r , and year y . To prevent confusion, we use the notation of Montgomery (2012) in the following calculations, where ‘ \cdot ’ subscript notation indicates summation over the subscript that it replaces.

First we sum over survey type x to calculate the total spawn index in spawn s (Table 1 and Table 9)

$$B'_{\cdot snry} = \sum_{x=1}^{X_{snry}} B'_{xsnry}, \quad (33)$$

where B'_{xsnry} is the spawn index for surface, *Macrocystis*, and understory spawn survey types from Equation 13, Equation 25, and Equation 32, respectively and $B'_{\cdot snry}$ is in tonnes (step T1 in Figure 3). Then we aggregate the total spawn index by SAR r and year y

$$B'_{\cdot \cdot ry} = \sum_{n=1}^{N_{ry}} \sum_{s=1}^{S_{nry}} B'_{\cdot snry}, \quad (34)$$

where $B'_{\cdot snry}$ is the total spawn index from Equation 33 and $B'_{\cdot \cdot ry}$ is a relative index of combined sex spawning biomass in tonnes.

Table 9. Notation for Pacific Herring total spawn index calculations. Note that q represents the spawn index scaling parameter in this table, not quadrat number as in other sections of this report. Legend: tonnes (t). Note that ‘‘ indicates related variables (e.g., B' is related to B).

Name	Description	Value or unit	References
B'	Spawn index (i.e., relative biomass)	t	
q	Spawn index scaling parameter	$0.0 < q \leq 1.0$	DFO (2022)
B	Scaled biomass	t	

8.1 SCALED BIOMASS

We use $B'_{...ry}$ as an indicator of relative biomass in Pacific Herring stock assessment models (DFO 2022). We scale $B'_{...ry}$ to biomass

$$B_{...ry} = \frac{B'_{...ry}}{q_{ry}}, \quad (35)$$

where $B'_{...ry}$ is the total spawn index in tonnes from Equation 34, q_{ry} is the spawn index scaling parameter (DFO 2016), and $B_{...ry}$ is scaled biomass for SAR r and year y in tonnes. To be consistent with stock assessments, q represents the spawn index scaling parameter in this section, not quadrat number as in other sections of this report. In Pacific Herring stock assessment models, q describes the proportion of spawn observed in spawn surveys, accounting for egg loss (e.g., due to predation) and unobserved spawns (DFO 2016). Although q is uncertain and a number of assumptions are made within the assessment model, there are differences between the proportion of spawn observed in the two spawn survey periods (DFO 2022). Therefore, for each SAR we estimate one spawn index scaling parameter for the surface survey period q_1 (1951 to 1987) and a second one for the dive survey period q_2 (1988 to present).

9 SPAWN-ON-KELP CALCULATIONS

Spawn-on-kelp (SOK) fisheries collect Pacific Herring roe that adheres to algae such as *Macrocystis* after spawning. Other similar fisheries include spawn-on-bough, in which operators collect roe that adheres to submerged tree boughs; we refer to these fisheries collectively as SOK in this document. There are two types of SOK fisheries in BC: ‘open-pond’ in which operators provide algae to spawning Pacific Herring

and ‘closed-pond’ in which operators impound spawning Pacific Herring in floating nets that contain algae (Shields et al. 1985). Although SOK fisheries do not directly remove spawning Pacific Herring, they do remove eggs that could otherwise have contributed to recruitment. Note that closed-pond operations also cause incidental mortality to spawning Pacific Herring (Shields et al. 1985), but we do not address this issue here. Thus, SOK fisheries present an issue in terms of their impact to the population, and accounting in stock assessment and monitoring (Schweigert et al. 2018). For example, failing to account for SOK spawn in assessments is analogous to treating it as missed spawn via the spawn index scaling parameter q (section 8.1); conversely, accounting for SOK spawn directly in assessments may reduce uncertainty in q . Although Pacific Herring stock assessments do not account for eggs removed by SOK fisheries at this time, there are a few options to account for impacts from SOK harvest. The most direct is to estimate the quantity of eggs removed from the population and determine the biomass of Pacific Herring that produced those eggs.

Shields et al. (1985) collected information on the relationship between the number of egg layers in SOK product and the proportion of product weight that consists of eggs and kelp. They determined that *Macrocystis* kelp represents an average of 12% of the total product weight. Since SOK product is universally brined at the time of harvest, we also need to consider the uptake of salt by the eggs, which increases the overall product weight. However, there is uncertainty in the degree of brining that occurs prior to weighing the product. Nevertheless, Whyte and Englarr (1977) determined that wet product weight increases by approximately 13% due to salt uptake during a 24-hour brining period. By osmosis, brining would also draw some water from the eggs (Alderice et al. 1979); unfortunately we are unable to account for osmosis at this time. The last factor to consider is the mean fertilized egg weight, $2.38 \cdot 10^{-6}$ kg (Hay and Miller 1982).

We estimate the relative biomass of Pacific Herring that spawned and produced eggs which were removed from the population by SOK fishery g (Table 1 and Table 10)

$$B'_{gry} = \frac{H_{gry} (1 - \nu)^{\frac{1}{1+\nu}}}{w\theta}, \quad (36)$$

where H_{gry} is the weight in kilograms of Pacific Herring SOK harvest, ν is the proportion of SOK product weight that is *Macrocystis* kelp, v is the SOK product weight increase due to brining as a proportion, w is the average weight in kilograms of a fertilized egg, θ is the egg conversion factor from Equation 8, and B'_{gry} is in

tonnes. Then we aggregate the relative biomass by SAR r and year y

$$B'_{.ry} = \sum_{g=1}^{G_{ry}} B'_{gry}, \quad (37)$$

where B'_{gry} is the relative biomass from Equation 36 and $B'_{.ry}$ is in tonnes. Thus, $B'_{.ry}$ is the relative biomass of Pacific Herring that produced eggs which were removed by SOK fisheries in SAR r and year y in tonnes.

10 SOURCES OF UNCERTAINTY

Like all biological models, spawn index calculations are affected by various potential sources of uncertainty including natural variability, observation error (e.g., bias, precision), measurement error, and model structural complexity (Link et al. 2012). Some examples illustrate these sources of uncertainty:

1. Natural variability could affect Pacific Herring fecundity and the sex ratio of spawning Pacific Herring (Equation 8). Fecundity could be influenced by biological processes such as the observed non-stationarity of weight-at-age (DFO 2022) or a truncated age distribution caused by selective fishing (Brunel and Piet 2013).
2. Measurement error could affect input data (e.g., number of egg layers, spawn length and width), while model structural complexity could affect estimated

Table 10. Notation for Pacific Herring spawn-on-kelp (SOK) calculations. Legend: kilograms (kg) and tonnes (t). Note that ‘‘ indicates related variables (e.g., B' is related to scaled biomass B in Table 9).

Name	Description	Value or unit	References
H	Weight of SOK harvest	kg	
ν	Proportion of SOK product that is <i>Macrocystis</i> kelp	0.12	Shields et al. (1985)
v	SOK product weight increase due to brining (proportion)	0.132	Whyte and Englars (1977)
w	Average weight of a fertilized egg	2.38×10^{-6} kg	Hay and Miller (1982)
B'	Relative SOK spawn biomass	t	

prediction model parameters, or the form of their relationship, or both (e.g., Equation 11). In addition, spawn index prediction models are dated, and the processes could have changed in the intervening years (e.g., Equation 11, Equation 23).

3. Uncertainty in fixed parameters that are used as data without error (e.g., Equation 11); uncertainty in spawn index parameters is currently unknown.
4. Uncertainty in the number of egg layers for spawn intensity categories (Table 3) and algae coefficients (Table 8). Again, uncertainty in these values is currently unknown.

Despite these assumptions and potential sources of uncertainty, the spawn index has typically been reported without quantifying uncertainty (but see Schweigert et al. (1993) who derived a variance estimator). Reporting the spawn index without uncertainty may perpetuate the misconception that the spawn index is observed data, whereas it is derived data with assumptions and uncertainties. An additional issue for stock assessments is that the model formulation could interpret the spawn index as being more precise than it actually is, and estimates of stock status and future prognosis could be artificially precise.

There are several potential benefits to addressing spawn index uncertainty. First, quantifying uncertainty could identify parameters to target with future research. Potential analyses to quantify spawn index uncertainty include:

1. Quantify and report variability in estimated prediction model parameters and equations (e.g., Equation 11),
2. Propagate uncertainty in parameters and prediction models through spawn index calculations,
3. Incorporate the underlying data that informs prediction model equations into spawn index calculations,
4. Bootstrap observed input data (see Schweigert 1993), and
5. Conduct sensitivity analyses.

Second, acknowledging uncertainty can reduce another source of uncertainty: inadequate communication among scientists, managers, and stakeholders, which can lead to misapplication of scientific advice (Link et al. 2012). Finally, acknowledging uncertainty will increase transparency and enable users to assess potential impacts to Pacific Herring stock assessments in a management strategy evaluation (MSE) approach (e.g., DFO 2019). Addressing data and model uncertainty is a required component of MSE approaches (Punt et al. 2016).

Quantifying uncertainty may also identify options to increase survey program efficiency by targeting data that have the greatest impact on spawn index accuracy. In addition, there is a trade-off between precision of estimated quantities versus survey effort or cost. Ideally, reducing survey effort does not result in biased target variable estimates. Therefore, understanding this trade-off is important if, for example, budget reductions cause reduced survey effort. Potential strategies to improve spawn survey efficiency include:

1. Conduct underwater surveys for major spawns in core areas and surface surveys for other spawns,
2. Review transect and quadrat spacing (section 2; see Schweigert 1993), and
3. Conduct periodic versus annual surveys.

Even with a stable budget, there is a trade-off between high survey effort in some areas, versus low survey effort or no information in other areas.

11 FUTURE RESEARCH

Many of the studies to quantify parameters and prediction models used in spawn index calculations are dated (Table 11); these analyses could be reviewed with new information and updated if required. In addition, some of these studies have limited number of samples, limited spatial or temporal range, or both. In addition to work mentioned here and in section 10, potential research includes:

1. Check the assumed statistical framework for spawn index calculations,
2. Develop egg density variance estimators for surface and *Macrocystis* spawn (i.e., comparable to Equation 4 for understory spawn),
3. Check accuracy and temporal stability of fecundity and sex ratios (Equation 8),
4. Compare two methods to calculate the mean number of stalks per plant (Equation 22): ratio of means (i.e., current method) versus mean of ratios,
5. Review the assumptions that eggs on substrate and algae are:
 - (a) Independent (Equation 28) and
 - (b) Represented by the same length measurement (section 7.3),
6. Review surface spawn width adjustments (section A.1):
 - (a) Mean versus median width,
 - (b) Annual versus periodic updates, and
 - (c) Relationship between spawn width and biomass,

Table 11. Details for studies to quantify parameters and prediction models used in Pacific Herring spawn index calculations. Study details include where, when, and how many samples, when available. The major stock assessment regions are Haida Gwaii (HG), Prince Rupert District (PRD), Central Coast (CC), Strait of Georgia (SoG), and West Coast of Vancouver Island (WCVI) (Figure 2). Legend: equation (Eq), sample size (n), standard error (SE), standard deviation (SD), spawn-on-kelp (SOK), and not available (NA).

Param- eters	Description	Study details	Uncertainty	References
f', f''	Parameters for statistical framework (Eq 4).	SoG in 1981 and 1983; WCVI in 1982.	Objective is mean egg density $SE \leq 25\%$.	Schweigert et al. (1985, 1990)
Intensity	Spawn intensity categories and number of egg layers (Table 3).	NA	NA	Hay and Kronlund (1987); Schweigert and Stocker (1988)
ω	Female fecundity (Eq 8); values in eggs · kg ⁻¹ .	PRD, WCVI, and SoG in 1974 ($n = 3,293$ fish) and 1980 ($n = 1,642$); California in 1975 ($n = 37$).	$186,800 \leq \omega \leq 224,500$; $16,900 \leq SD \leq 53,500$.	Hay (1985); Hay and Brett (1988)
φ	Proportion female (Eq 8).	NA	NA	NA
α, β	Parameters for surface survey egg density model (Eq 11).	Coastwide from 1976 to 1987 ($n = 5,111$ samples).	NA	Schweigert et al. (1997)
$\xi, \gamma, \delta, \epsilon$	Parameters for number of eggs per <i>Macrocystis</i> plant model (Eq 23).	HG in 1981 and 1987 ($n = 112$ plants); PRD in 1986 ($n = 15$); CC in 1986 ($n = 5$); WCVI in 1985 and 1986 ($n = 35$).	Model accounts for 78% of variation in eggs per plant.	Haegele and Schweigert (1990)

Param- eters	Description	<i>Table 11 continued</i>		References
		Study details	Uncertainty	
φ	Parameter for substrate egg density model (Eq 26).	SoG in 1976, 1977, and 1978; PRD, CC, and WCVI in 1977.	NA	Haegele and Humphreys (1976, 1978a,b); Haegele et al. (1979)
$\vartheta, \varrho, \varsigma,$ C_v	Parameters for algae egg density model (Eq 27, Table 8).	Coastwide from 1976 to 1987 ($n = 5,111$ samples).	Model accounts for 40% of variation in egg density.	Schweigert (2005)
ν	Proportion of SOK product that is kelp (Eq 36).	HG in 1982 and 1983; PRD in 1982.	SD = 4.2.	Shields et al. (1985)
v	SOK product weight increase due to brining (proportion; Eq 36).	SoG ca. 1977.	NA	Whyte and Englart (1977)
w	Average weight of a fertilized egg (Eq 36).	SoG in 1980 ($n = 7$ samples).	SE = 3.4×10^{-7} .	Hay and Miller (1982)

7. Periodically review the accuracy of egg density models using egg layer and vegetation cover estimates collected underwater, compared to egg counts in a subset of sampled quadrats,
8. Quantify incidental mortality in SOK operations,
9. Determine the proportion of SOK product that is kelp for other algae species and tree boughs (Equation 36), and
10. Account for osmosis in SOK calculations (Equation 36).

12 CAVEATS

There are a few caveats to consider when interpreting the Pacific Herring spawn index and using spawn index data in analyses. These caveats include:

1. The spawn index is a relative index of spawning biomass,
2. The spawn survey is a presence only survey; thus the spawn index is a minimum estimate of spawning biomass,
3. There are two different spawn survey periods with substantial differences in survey effort and method (section 3.1):
 - (a) Surface period from 1951 to 1987 and
 - (b) Dive period from 1988 to present,
4. Surface spawn surveys use two different methods to estimate the number of egg layers (section 3.1.1):
 - (a) Spawn intensity categories:
 - i. Five categories from 1951 to 1968 and
 - ii. Nine categories from 1969 to 1978, and
 - (b) Direct estimates from 1979 to present,
5. The spawn index is derived from surface and dive observations of egg deposition, and includes uncertainty and assumptions (section 10), and
6. Spawn index calculations rely on dated parameters and models (section 11).

13 R PACKAGE

We developed the **SpawnIndex R** (RCT 2021) package to implement the calculations described in this document. The package is publicly accessible on the [Pacific Herring spawn index repository](#). The **SpawnIndex** package contains an example database of

Pacific Herring spawn survey observations, as well as functions to import tables from the database and calculate the spawn index. The `SpawnIndex` package is documented and includes examples to promote accessibility and transparency. There is also a vignette with an example workflow. Finally, we cross-references equations in this document with functions in the **R** package (Appendix C).

14 ACKNOWLEDGEMENTS

The authors thank Ashleen Benson for translating the calculations in the **Microsoft Access** database to an **R** script, which they referenced when developing the `SpawnIndex` package. They thank Paul Starr and Rob Kronlund for their thorough reviews that improved this document. They also thank Andy Edwards and Joe Watson for help with mathematical notation, as well as Chris Grandin for help developing the **R** package.

15 REFERENCES

- Alderice, D.F., Rosenthal, H., and Velsen, F.P.J. 1979. Influence of salinity and cadmium on the volume of Pacific herring eggs. *Helgoländer wissenschaftliche Meeresuntersuchungen* **32**: 163–178. doi:[10.1007/BF02189895](https://doi.org/10.1007/BF02189895)
- Brunel, T., and Piet, G.J. 2013. Is age structure a relevant criterion for the health of fish stocks? *ICES Journal of Marine Science* **70**: 270–283. doi:[10.1093/icesjms/fss184](https://doi.org/10.1093/icesjms/fss184)
- Cleary, J.S., Taylor, N.G., and Haist, V. 2017. Status of BC Pacific Herring (*Clupea pallasii*) in 2013 and forecasts for 2014. Research Document 2017/014, Canadian Science Advisory Secretariat, Fisheries and Oceans Canada. URL<http://cat.fsl-bsf.scitech.gc.ca/record=b4061206~S1>
- DFO (Fisheries and Oceans Canada). 2016. Stock assessment and management advice for BC Pacific herring: 2016 status and 2017 forecast. Science Response 2016/052, Canadian Science Advisory Secretariat, Fisheries and Oceans Canada. URLhttps://www.dfo-mpo.gc.ca/csas-sccs/Publications/ScR-RS/2016/2016_052-eng.html
- DFO (Fisheries and Oceans Canada). 2019. Evaluation of management procedures for Pacific Herring (*Clupea pallasii*) in the Strait of Georgia and the West Coast of Vancouver Island management areas of British Columbia. Science Advisory Report 2019/001, Canadian Science Advisory Secretariat, Fisheries and Oceans

- Canada. URL http://www.dfo-mpo.gc.ca/csas-sccs/Publications/SAR-AS/2019/2019_001-eng.html
- DFO (Fisheries and Oceans Canada). 2022. Stock status update with application of management procedures for Pacific Herring (*Clupea pallasii*) in British Columbia: Status in 2022 and forecast for 2023. Science Response 2022/046, Canadian Science Advisory Secretariat, Fisheries and Oceans Canada. URL https://www.dfo-mpo.gc.ca/csas-sccs/Publications/Scr-RS/2022/2022_046-eng.pdf
- Haegele, C.W., Hourston, A.S., Humphreys, R.D., and Miller, D.C. 1979. Eggs per unit area in British Columbia herring spawn depositions. Fisheries and Marine Service Technical Report 894, Department of Fisheries and Oceans. URL <http://cat.fsl-bsf.scitech.gc.ca/record=b3858115~S1>
- Haegele, C.W., and Humphreys, R.D. 1976. Data record for 1976 shoreline vegetation and herring spawn surveys. Data Record 15, Fisheries and Marine Service. URL <https://science-catalogue.canada.ca/record=b3848053~S6>
- Haegele, C.W., and Humphreys, R.D. 1978a. Data record for 1977 herring spawn studies. Data Report 75, Fisheries and Marine Service. URL <https://science-catalogue.canada.ca/record=b3877077~S6>
- Haegele, C.W., and Humphreys, R.D. 1978b. Data record for 1978 herring spawn studies. Data Report 107, Fisheries and Marine Service. URL <https://science-catalogue.canada.ca/record=b3864712~S6>
- Haegele, C.W., and Schweigert, J.F. 1985. Distribution and characteristics of herring spawning grounds and description of spawning behavior. *Canadian Journal of Fisheries and Aquatic Sciences* **42**(S1): 39–55. doi:[10.1139/f85-261](https://doi.org/10.1139/f85-261)
- Haegele, C.W., and Schweigert, J.F. 1990. A model which predicts Pacific Herring (*Clupea harengus pallasii*) egg deposition on giant kelp (*Macrocystis* sp.) plants from underwater observations. Canadian Manuscript Report of Fisheries and Aquatic Sciences 2056, Fisheries and Oceans Canada. URL <http://cat.fsl-bsf.scitech.gc.ca/record=b3898101~S1>
- Haist, V., and Rosenfeld, L. 1988. Definitions and codings of localities, sections, and assessment regions for British Columbia herring data. Canadian Manuscript Report of Fisheries and Aquatic Science 1994, Fisheries and Oceans Canada. URL <http://cat.cisti-icist.nrc-cnrc.gc.ca/record=b3927024~S1>
- Hart, J.L., and Tester, A.L. 1934. Quantitative studies on herring spawning. *Transactions of the American Fisheries Society* **64**(1): 307–312. doi:[10.1577/1548-8659\(1934\)64\[307:qsos\]2.0.co;2](https://doi.org/10.1577/1548-8659(1934)64[307:qsos]2.0.co;2)
- Hay, D.E. 1985. Reproductive biology of Pacific Herring (*Clupea harengus pal-*

lasi). Canadian Journal of Fisheries and Aquatic Sciences **42**(S1): 111–126. doi:[10.1139/f85-267](https://doi.org/10.1139/f85-267)

Hay, D.E., and Brett, J.R. 1988. Maturation and fecundity of Pacific Herring (*Clupea harengus pallasi*): An experimental study with comparisons to natural populations. *Canadian Journal of Fisheries and Aquatic Sciences* **45**(3): 399–406. doi:[10.1139/f88-048](https://doi.org/10.1139/f88-048)

Hay, D.E., and Kronlund, A.R. 1987. Factors affecting the distribution, abundance, and measurement of Pacific Herring (*Clupea harengus pallasi*) spawn. *Canadian Journal of Fisheries and Aquatic Sciences* **44**(6). doi:[10.1139/f87-141](https://doi.org/10.1139/f87-141)

Hay, D.E., and Miller, D.C. 1982. A quantitative assessment of herring spawn lost by storm action in French Creek, 1980. Canadian Manuscript Report of Fisheries and Aquatic Sciences 1636, Department of Fisheries and Oceans. URL<http://cat.fsl-bsf.scitech.gc.ca/record=b3849753~S1>

Humphreys, R.D., and Hourston, A.S. 1978. British Columbia herring spawn deposition manual. Miscellaneous Special Publication 38, Department of Fisheries and the Environment, Fisheris and Marine Service. URL<http://cat.fsl-bsf.scitech.gc.ca/record=b3686454~S1>

Jessen, R.J. 1978. Statistical survey techniques. John Wiley & Sons. URLhttps://openlibrary.org/books/OL4552019M/Statistical_survey_techniques

Link, J.S., Ihde, T.F., Harvey, C.J., Gaichas, S.K., Field, J.C., Brodziak, J.K.T., Townsend, H.M., and Peterman, R.M. 2012. Dealing with uncertainty in ecosystem models: The paradox of use for living marine resource management. *Progress in Oceanography* **102**: 102–114. doi:[10.1016/j.pocean.2012.03.008](https://doi.org/10.1016/j.pocean.2012.03.008)

Marwick, B., Boettiger, C., and Mullen, L. 2018. Packaging data analytical work reproducibly with R (and friends). *PeerJ Preprints (not peer reviewed)* doi:[10.7287/peerj.preprints.3192v2](https://doi.org/10.7287/peerj.preprints.3192v2)

Montgomery, D.C. 2012. Design and analysis of experiments. John Wiley & Sons, 8 ed.

Punt, A.E., Butterworth, D.S., de Moor, C.L., De Oliveira, J.A.A., and Haddon, M. 2016. Management strategy evaluation: Best practices. *Fish and Fisheries* **17**(2): 303–334. doi:[10.1111/faf.12104](https://doi.org/10.1111/faf.12104)

Rabin, D.J., and Barnhart, R.A. 1975. Fecundity of Pacific Herring, *Clupea harengus pallasi*, in Humboldt Bay. Technical Report 63(3), California Fish and Game

RCT (R Core Team). 2021. R: A language and environment for statistical computing. URL<http://www.R-project.org>. R Foundation for Statistical Computing. Vienna, Austria. Version 4.1.2 (32-bit)

- Schaeffer, R.L., Mendenhall, W., Ott, R.L., and Gerow, K. 2012. Elementary survey sampling. Duxbury advanced series. Brooks/Cole, seventh ed.
- Schweigert, J., Cleary, J., and Midgley, P. 2018. Synopsis of the Pacific Herring spawn-on-kelp fishery in British Columbia. Canadian Manuscript Report of Fisheries and Aquatic Sciences 3148, Fisheries and Oceans Canada. URL <http://cat.fsl-bsf.scitech.gc.ca/record=b4068121~S1>
- Schweigert, J.F. 1993. A review and evaluation of methodology for estimating Pacific Herring egg deposition. *Bulletin of Marine Science* **53**(2). URL <http://www.ingentaconnect.com/content/umrsmas/bullmar/1993/00000053/00000002/art00019>
- Schweigert, J.F. 2005. An assessment framework for Pacific Herring (*Clupea pallasi*) stocks in British Columbia. Research Document 2005/083, Fisheries and Oceans Canada. URL <http://cat.cisti-icist.nrc-cnrc.gc.ca/record=4025049>
- Schweigert, J.F., Fort, C., and Hamer, L. 1997. Stock assessment for British Columbia herring in 1996 and forecasts of the potential catch in 1997. Canadian Technical Report of Fisheries and Aquatic Sciences 2173, Department of Fisheries and Oceans. URL <http://cat.cisti-icist.nrc-cnrc.gc.ca/record=b4020685~S1>
- Schweigert, J.F., and Haegele, C.W. 2021. Synopsis of the development of dive surveys for Pacific Herring spawn assessment. Canadian Manuscript Report of Fisheries and Aquatic Sciences 3234, Fisheries and Oceans Canada. URL <https://cat.fsl-bsf.scitech.gc.ca/record=b4102370~S1>
- Schweigert, J.F., Haegele, C.W., and Stocker, M. 1985. Optimizing sampling design for herring spawn surveys in the Strait of Georgia, B.C. *Canadian Journal of Fisheries and Aquatic Sciences* **42**(11): 1806–1814. doi:[10.1139/f85-226](https://doi.org/10.1139/f85-226)
- Schweigert, J.F., Haegele, C.W., and Stocker, M. 1990. Evaluation of sampling strategies for scuba surveys to assess spawn deposition by Pacific Herring. *North American Journal of Fisheries Management* **10**(2): 185–195. doi:[10.1577/1548-8675\(1990\)010<0185:eoossfs>2.3.co;2](https://doi.org/10.1577/1548-8675(1990)010<0185:eoossfs>2.3.co;2)
- Schweigert, J.F., Hay, D.E., and Fort, C. 1993. Herring spawn index analysis. PSARC H93-02, Department of Fisheries and Oceans. URL <http://cat.fsl-bsf.scitech.gc.ca/record=b4018577~S1>
- Schweigert, J.F., and Stocker, M. 1988. Escapement model for estimating Pacific Herring stock size from spawn survey data and its management implications. *North American Journal of Fisheries Management* **8**. doi:[10.1577/1548-8675\(1988\)008<0063:EMFEPH>2.3.CO;2](https://doi.org/10.1577/1548-8675(1988)008<0063:EMFEPH>2.3.CO;2)

- Shields, T.L., Jamieson, G.S., and Sprout, P.E. 1985. Spawn-on-kelp fisheries in the Queen Charlotte Islands and northern British Columbia coast - 1982 and 1983. Canadian Technical Report of Fisheries and Aquatic Sciences 1372, Department of Fisheries and Oceans. URL <http://cat.fsl-bsf.scitech.gc.ca/record=b1319605~S1>
- Tanasichuk, R.W., and Ware, D.M. 1987. Influence of interannual variations in winter sea temperature on fecundity and egg size in Pacific Herring (*Clupea harengus pallasi*). *Canadian Journal of Fisheries and Aquatic Sciences* **44**(8): 1485–1495. doi:[10.1139/f87-178](https://doi.org/10.1139/f87-178)
- Ware, D.M. 1985. Life history characteristics, reproductive value, and resilience of Pacific Herring (*Clupea harengus pallasi*). *Canadian Journal of Fisheries and Aquatic Sciences* **42**: 127–137. doi:[10.1139/f85-268](https://doi.org/10.1139/f85-268)
- Whyte, J.N.C., and Englart, J.R. 1977. Aspects of the production of herring roe on *Macrocystis integrifolia* in Georgia Strait locations. Fisheries and Marine Service Technical Report 751, Fisheries and Marine Service. URL <http://cat.fsl-bsf.scitech.gc.ca/record=b1115904~S1>
- Wilson, G., Bryan, J., Cranston, K., Kitzes, J., Nederbragt, L., and Teal, T. 2016. Good Enough Practices in Scientific Computing. *PLOS* URL <https://arxiv.org/abs/1609.00037v2>
- Xie, Y. 2015. Dynamic documents with R and knitr. The R series. Chapman and Hall/CRC, Florida, USA, 2nd ed. URL <https://github.com/yihui/knitr-book>. ISBN 978-1498716963

APPENDIX A SPAWN WIDTH ADJUSTMENTS

Spawn width is a critical component in spawn index calculations. There are two cases where we adjust spawn width estimates to increase the accuracy of spawn index calculations: surface surveys in all years from 1951 to present (section A.1) and understory dive surveys from 2003 to 2014 (section A.2).

A.1 SURFACE SPAWN WIDTH

Surface surveys were the only survey type prior to 1988, while the majority of spawns since 1988 have been surveyed using SCUBA gear (section 1). Recall that we describe the spawn index as having two periods based on the predominant survey type: surface survey period from 1951 to 1987 and dive survey period from 1988 to present. One

issue with comparing these two partly overlapping protocols is that surface surveyors tend to underestimate spawn width W (Hay and Kronlund 1987).

To improve the consistency of spawn index estimates throughout the time period from 1951 to present, we adjust surface spawn width estimates using underwater estimates from dive surveys when dive data are available (Schweigert et al. 1993). Our preferred width is the median width from all dive surveys within a ‘pool.’ A pool is a group of Locations within a Section that are often adjacent, contain similar algae and substrate, and can be treated as a group with likely similar widths. We summarise spawn width by the median because widths are skewed (Schweigert et al. 1993). If there are no dive data that meet these criteria, we use the median width from all dives within the Section, or within the SAR if there are no dives within the Section. In the rare instances where no dive data meet these criteria (e.g., outside SAR boundaries), we use the observed width W'_{xsnry} from the surface survey. We update the aforementioned median width values periodically, not annually. Note that we use this process to update widths for spawns in previous years using current observations.

A.2 UNDERSTORY SPAWN WIDTH

In 2013, Fisheries and Oceans Canada (DFO) staff realized that they were inadvertently underestimating spawn width for Pacific Herring understory dive surveys (Cleary et al. 2017). The issue was caused by the assumed accuracy of transect lines used by spawn surveyors to measure spawn width. Spawn surveyors determine spawn width by placing transects perpendicular to the shore. Surveyors use weighted lead lines to ensure that lines rest on the substrate; these lines are marked in 1 m increments and are standardized to 20 m segments. Segments refer to individual sections of line, which can be linked together to make complete transects.

During the late 1990s it became apparent that the 20 m segments shrank by approximately 1 m during the first season of use and continued to shrink over time. DFO staff noticed that this issue was occurring coast wide and began re-measuring lead lines each season. They also modified the lead line marking protocol to account for shrinkage by marking 1.15 m increments; thus, segments were extended to 23 m. DFO staff derived this 15% increase by measuring and re-marking lead lines each year. Lead lines are made of a mix of polypropylene and nylon; nylon tightens up under repeated use, which we think explains the shrinkage. DFO staff re-measured lead line increments in about 2005; they found that the lines still shrank from 1.15 m to 1.0 m and they continued to use the modified protocol.

In 2013, spawn surveyors observed that some lead line increments were 1.15 m

and no longer appeared to be shrinking. Following this observation, DFO staff re-measured additional lead lines and found that lead lines were made up of a combination of 1.0 m and 1.15 m increments. The combination of observed increment lengths is explained by the lifespan of lead lines: lead lines are replaced every 5 to 10 years, with some segments being replaced more frequently. For example, inner segments are replaced more frequently than seaward segments and segments in some SARs are replaced more frequently than segments in other SARs. DFO staff think that a change in manufacturing prevents new lead lines from shrinking.

The earliest written instructions that describe the modified protocol of marking 1.15 m increments is from 2003, and this protocol was used until 2013. Note that some remote SARs continued to use old lead lines in 2014. The practice of annually re-measuring lead line increments ceased around 2005; thus we are unable to determine when lead lines ceased shrinking. Given the observations summarized above, we adjust understory spawn width estimates based on written instructions for the marking protocol in 2003. Accordingly, we estimate the years impacted by marking lead lines at 1.15 m increments (when shrinking no longer occurred) is from 2003 to 2014. However, not all SARs r and years y are impacted equally by this issue (Table A.1). That is, some SARs and years had all 1.0 m increment lengths (no correction factor needed; $\tau_{ry} = 1.000$), others had all 1.15 m increment lengths ($\tau_{ry} = 1.150$), and others had a combination of 1.0 m and 1.15 m increment lengths which we assume to be in equal proportion ($\tau_{ry} = 1.075$).

We correct understory spawn width on transect t (Table 1 and Table A.2)

$$W_{txsnry}'' = W_{txsnry}' \tau_{ry}, \quad (\text{A.1})$$

where W_{txsnry}'' is the observed spawn width, τ_{ry} is the spawn width correction factor (Table A.1), and W_{txsnry}' is in metres. Instead of updating the database permanently, we adjust spawn width in the **R** package to be transparent and to prevent mismatches between the original data sheets and databases. DFO staff now re-measure transect segments annually to ensure that this issue does not reoccur due to another change in lead line manufacturing.

APPENDIX B SURFACE SPAWN UPDATES

One record in the surface spawn database since 1951 requires an update to fill-in missing egg layer information. As with understory spawn width updates, we make this update in the **R** package:

1. Update ‘intensity’ from 0 to 1 for the records (there is 1 record) in the year 1962, Statistical Area 14, Section 142, Location code 820, and with intensity = 0. We

Table A.1. Spawn width correction factors τ_{ry} for Pacific Herring understory spawn surveys by stock assessment region (SAR, r) and year y . Legend: Haida Gwaii (HG), Prince Rupert District (PRD), Central Coast (CC), Strait of Georgia (SoG), West Coast of Vancouver Island (WCVI), Area 27 (A27), and Area 2 West (A2W).

Year	SAR							
	Major				Minor			
	HG	PRD	CC	SoG	WCVI	A27	A2W	
2003	1.000	1.075	1.075	1.075	1.075	1.075	1.075	1.000
2004	1.000	1.075	1.075	1.075	1.075	1.075	1.075	1.000
2005	1.000	1.075	1.075	1.075	1.075	1.075	1.075	1.000
2006	1.000	1.075	1.075	1.075	1.075	1.075	1.075	1.000
2007	1.000	1.075	1.075	1.075	1.075	1.075	1.075	1.000
2008	1.000	1.075	1.075	1.075	1.075	1.075	1.075	1.000
2009	1.150	1.075	1.075	1.075	1.075	1.075	1.075	1.150
2010	1.150	1.075	1.075	1.075	1.075	1.075	1.075	1.150
2011	1.150	1.075	1.075	1.075	1.075	1.075	1.075	1.150
2012	1.150	1.075	1.075	1.075	1.075	1.075	1.075	1.150
2013	1.150	1.150	1.075	1.075	1.075	1.000	1.000	1.150
2014	1.150	1.150	1.000	1.000	1.000	1.000	1.000	1.150

Table A.2. Notation for Pacific Herring understory spawn width adjustments. Legend: metres (m). Note that “ \prime ” indicates related variables (e.g., W' is related to W).

Name	Description	Value or unit
W'	Observed spawn width	m
τ	Spawn width correction factor	See Table A.1
W	Corrected spawn width	m

update intensity from 0 to 1 because spawn was surveyed but not reported. Spawn survey records prior to 1951 have additional missing or inaccurate egg layer information and are unreliable for indexing purposes. Therefore, we exclude spawn data prior to 1951 from stock assessments.

While reviewing spawn index calculations and translating them from a **Microsoft Access** database to **R**, we found several cases where index data were being overwritten with no documented reason. These updates have been omitted and affected

the following records:

1. Update E_{jxsnry} (i.e., number of egg layers; Table 5) to 2.1496 for the records (there are 15 records) in the year 1979, Statistical Area 2, and with intensity 4,
2. Update E_{jxsnry} to 0.5529 for the records (4) in the year 1981, Statistical Area 24, and with $E_{jxsnry} = 0.0$,
3. Update E_{jxsnry} to 1.3360 for the records (7) in the year 1982, Statistical Area 23, and with intensity 3,
4. Update E_{jxsnry} to 2.3300 for the records (41) in the year 1984, Statistical Area 24, and with intensity 0, and
5. Update E_{jxsnry} to 2.9800 for the records (14) in the year 1982, Statistical Area 27, and with $E_{jxsnry} = 0.0$.

In the first three cases, E_{jxsnry} was updated using intensity categories (Table 3); in the last two cases, E_{jxsnry} was updated using historical averages. These changes have negligible effects on spawn index values.

APPENDIX C R PACKAGE CROSSWALK

To facilitate pairing this document with the `SpawnIndex` **R** package ([Pacific Herring spawn index repository](#)), we cross-reference equations in this document with functions in the package (Table C.1). In addition, parameters names in this document correspond to function argument names in the **R** package. For example, theta θ in Equation 8 corresponds to `theta` in the function `eggs_to_sb()`. Similarly, egg layers E_{jxsnry} in Equation 11 corresponds to `egg_layers` in the function `dens_surf()`.

Table C.1. Crosswalk table for equations in this document (i.e., technical report) and corresponding functions in the **SpawnIndex R** package ([Pacific Herring spawn index repository](#)). Legend: ‘§’ indicates section and ‘Eq’ indicates equation.

Description	Technical report		R package
	§	Eq	Function
Egg conversion factor	4	8	<code>eggs_to_sb()</code>
Surface spawn calculations	5	9 to 13	<code>calc_surf_index()</code>
Surface egg density	5.1	11	<code>dens_surf()</code>
Macrocystis spawn calculations	6	14 to 25	<code>calc_macro_index()</code>
Number of eggs per Macrocystis plant	6.3	23	<code>eggs_macro()</code>
Understory spawn calculations	7	26 to 32	<code>calc_under_index()</code>
Understory egg density on substrate	7.1	26	<code>dens_under_sub()</code>
Understory egg density on algae	7.1	27	<code>dens_under_alg()</code>
Spawn-on-kelp calculations	9	36	<code>calc_sok_index()</code>