## Proportion of reporting SES

Study	reported	total		weight	F-T,[95% CI]
Goodman et al. 2020	30	37	<del>  ■</del>	81.1%	0.806 [0.664, 0.917]
Burnette et al. 2022	43	58	├ <del>──■</del> ─┤	74.1%	0.740 [0.619, 0.845]
Raad et al. 2008	118	206	<b>⊢</b> ■	57.3%	0.573 [0.505, 0.641]
Villarreal, 2014	94	170	<b>├──</b>	55.3%	0.553 [0.477, 0.628]
Pulli et al. 2019	37	67	<b>├──■</b>	55.2%	0.552 [0.432, 0.669]
Peter et al. 2021	6	13	<del>-</del>	46.2%	0.464 [0.212, 0.725]
Singh & Rajendra, 2024	725	1576	H <b>⊞</b> H	46.0%	0.460 [0.435, 0.485]
Waldron et al. 2018	27	69	<del>  ■</del>	39.1%	0.392 [0.280, 0.510]
Moffitt et al. 2023	211	555	<b>⊢</b> ■-1	38.0%	0.379 [0.339, 0.420]
Weisz et al. 2005	118	427	⊢ <del>≣</del> ⊣	27.6%	0.274 [0.233, 0.318]
Williamson et al. 2023	185	771	H <del>■</del> H	24.0%	0.237 [0.208, 0.268]
DeJesus et al. 2019	241	1149	H⊞H	21.0%	0.207 [0.183, 0.231]
McGorray et al. 2023	365	1762		20.7%	0.204 [0.185, 0.223]
Rad et al. 2018	34	428	H■H	7.9%	0.076 [0.052, 0.104]
Kissel & Friedman, 2023	28	1500		1.9%	0.014 [0.008, 0.021]
Sabik et al. 2021	1	100		1.0%	0.009 [0.000, 0.042]
RE Model	2263	8888			0.340 [0.213, 0.480]

0.000

0.250

0.500

Proportion

0.750

1.000