

Open MDMA: An Open Source Framework and Practical Guide for the Therapeutic use of MDMA

Mark Groeneveld*

Thomas Harper, M.S.W.[†]

DRAFT Version 0.31 June 12, 2024. It's still a messy, unfinished draft.
SAFETY RECOMMENDATIONS ARE UNDERGOING MAJOR REVISIONS

This work is licensed under [CC BY 4.0](#).

We welcome comments at mgroeneveld@protonmail.ch. Please let us know what is helpful and what is not helpful.

If you share this document, please use this link to ensure people have the latest version: doi.org/10.31234/osf.io/aps5g. This site also keeps tallies downloads, informing us how many people are using the guide and how worthwhile it is to update it with new research.

If you feel inspired and want to help us continue this work, we accept donations sent to mgroeneveld@protonmail.ch on Zelle, Venmo, or Paypal. We spent about 450 hours on this project, not including most of the acquisition of relevant skills and knowledge.

Abstract

60% of adults experienced childhood trauma; half develop a mental illness in their life. Symptoms like cognitive distortion, emotional rigidity/insecurity, and impaired compassion increase tribalism, political polarization, and apathy to animal suffering—hobbling our collective capacity to defeat global catastrophic risks. Access to effective care (a crucial support for stopping intergenerational harm) is inadequate for most who need it. MDMA is a promising treatment, but good training is lacking because MDMA is illegal.

With legalization immanent, our rigorous and accessible guide to MDMA therapy will: 1) Resource therapeutic MDMA users and clinicians 2) Articulate likely mechanisms of effectiveness, to inform the development of future therapeutic techniques

This can reduce mental illness by addressing the prevalent, high impact barriers to effective care—increasing collective capacity for collaboration, supporting individuals who end cycles of abuse, and protecting future generations.

*Corresponding author; Email: mgroeneveld@protonmail.ch

[†]Master of Social Work; Email: thomasharpertherapist@gmail.com

this manual is
also available
as an audio-
book here:

Contents

Preface	6
1 Introduction	8
2 Background, Goals, and Rationale	10
3 *Summary for Mental Health Professionals	12
4 The Science of Trauma and Healing	14
4.1 What is Trauma and Why do its Effects Stick Around?	14
4.2 *Defense Cascade	18
4.3 Mechanism of Healing	19
4.4 Efficacy of MDMA-Therapy	22
5 The Practice of MDMA-Therapy	24
5.1 *Safety and Challenges	24
5.2 Professional Guidance vs. Self Guidance	30
5.3 Dosing	33
5.4 Session Difficulties	34
5.4.1 *Dissociation	34
5.4.2 Resistance or Distraction	35
5.5 Pre-Session Preparation	35
5.6 The MDMA Therapy Session	38
5.7 *Troubleshooting the Session	39
5.7.1 Sleepiness, disconnection, or heaviness	40
5.7.2 Not feeling anything or feeling "meh"	40
5.7.3 Feeling really good but no therapeutic effect	40
5.7.4 New adverse symptom after a session	40
5.7.5 Feeling fatigued for 1-3 days after a session	41
5.7.6 Warning signs of psychosis and mania	41
5.8 After the Session	41
5.8.1 Making Sense of the Experience	42
5.8.2 Continued Reconsolidation	45
5.8.3 Using Neuroplasticity to Make Positive Life Changes	46
5.8.4 *Afterglow	51
5.9 Long-Term Trends of Therapy	51
5.10 Managing Anxiety and Panic	51
5.11 *Session Frequency	53
5.12 Prompts for Uncovering Common Maladaptive Schemas	56
5.13 The Ongoing Practice of Maintaining Cognitive Flexibility	61

6	Augmenting MDMA-Therapy	61
6.1	*Psycholytic Therapy	62
6.2	Challenging Beliefs Worksheet	62
6.3	Couple's Therapy	62
6.4	Associate Love and Empathy with Cues for Later Recall	63
6.5	Expanding the Circle of Compassion	64
6.6	Using MDMA to Facilitate Connection with Nature, a Fundamental Human Need	66
6.7	Psychoeducation	67
7	*Glossary	69
A	Appendices	70
A.1	Shame Triggers Writing Exercise	70
A.2	Attachment Theory	71
A.3	Loving-Kindness Meditation for Dissociation	72
A.4	Mindfulness-Based Stress Reduction	73
A.5	*Self-Determination Theory	75
A.6	Positive Life Changes	75
A.7	Things to do in Moments of Anxiety (Short Term Strategies)	77
A.8	Challenging Beliefs Worksheet	79
A.9	Suggested Avenues of Future Research	81
	Bibliography	94

META (Remove before publication)

Goals:

- Broad audience -> Avoid language or references that may unnecessarily activate challenging schemas for various large groups. Avoid unnecessarily technical language. Minimize length so people aren't scared away and can readily digest it. Conciseness and lack of useless wordiness for better readability. Audience feels guide is reputable, safe, and high quality.
- Achievable writing task (knowledge and time) -> Only attempt to write manual for an audience open to a neurological/egalitarian framing of trauma and healing. I can only write recommendations that I know are good within the culture I live in. It might also be appealing to the more diverse, spreading Modern Global Individualist culture? Probably less applicable to more different cultures.
- Maximize healing and minimize risk -> Comprehensive. Rigorous. Proper balance between risk-warning language and healing-hopeful language so we don't unnecessarily scare people away OR fail to adequately warn about real risks. Most people doing healing need a lot more than reconsolidation; include broadly useful resources on emotional skills.
- Avoid legal risk -> Disclaimer and wording.
- Improve authors' understanding of concepts -> Rigor. Comprehensiveness. Usefulness to broad audience.
- Improve authors' reputation -> Publish in a journal (J. Psychedelic Psychiatry looks like a good fit. If all else fails, can still put it on Psyarxiv.). Rigor. Comprehensiveness. Usefulness to broad audience. More care to not openly endorse solo use of illegal drugs.
- Want to be useful as FAQ or supplemental material for guides and therapists. Should consult one to see what they would want in such a document!
- Help Mark (and others who think like me) understand what is actually happening in trauma and healing. Psychologists, psychiatrists, and therapists rarely told me what was actually happening to me, and when they did, it was presented in baffling, ill-defined metaphors. -> Construct a rigorous, science-based explanation of trauma and therapy.
- Make Mark happy by exercising Mark's capabilities to their fullest extent.
- What else?

Known Issues, Uncertainties, and Missing Things:

- So far this paper was developed on my personal experience, reports of others' MDMA experiences, information from Jess and Day, and reading research papers. I combined and extrapolated that into a more cohesive framework. GPT helped me a lot in

clarifying concepts. I've had almost no positive experiences in therapy or psychiatry (except sessions with Jess that felt more like me asking for advice on various things) that didn't involve psychedelics and I haven't read any psychotherapy books other than Coleman's Psychedelic Psychotherapy.

- Section by section review with GPT (2 rounds completed). Prompt: This paper (delivered one section at a time) is intended as a scientifically rigorous framework for self-guided MDMA-therapy for a variety of audiences. Suggest improvements in the following section for ease of comprehension, logical flow, active voice, conciseness, avoidable stimuli for challenging schemas for a wide variety of groups, and technical accuracy and comprehensiveness. Identify any statements out-of-line with the scientific literature. Don't assume citations back up my conclusions. Identify gaps in my coverage of a topic. Suggestions should especially include opportunities to reduce the length of sentences and paragraphs through more efficient phrasing and removal of unnecessary words. Suggest additions, subtractions, or mergers of items in the lists in my paper. Make recommendations for avoiding liability or legal problems. Deliver your responses in narrative format. List all suggestions. Tone down your recommendations to never do anything without a medical professional. I don't want to hear that sort of protectionism.
- Review bibliography for missing information like volume numbers. Write out full first names in bib then make bibtex only say first initial. Try to eliminate non-peer-reviewed sources.
- Change reconsolidation to MR?
- I'm skeptical that dissociation can be adaptive when activated by maladaptive schemas. Dissociation is an ANS response to life-threats. So it's definitely adaptive when a lion caught you, probably adaptive when you're a neglected child, and unclearly adaptive when you're an adult facing overwhelming feelings? Overwhelming feelings might or might not be a life threat depending on context. So when can we assume dissociation in adults is adaptive or maladaptive? Not sure ANS is sophisticated enough to make sure its responses are always adaptive.
- Talk about making schemas explicit as a means for more effective reconsolidation and management?
- When/why do memories deactivate?
- TRUST THE PROCESS OF RECONSOLIDATION. Things may be distressing and confusing but AS LONG AS YOU ARE RECONSOLIDATING THEN YOU ARE MAKING PROGRESS TOWARD STABILITY. Something about destabilization
- Maybe we should synchronize use of we, us, you, one, etc.
- Include risks of maladaptive schemas and extended defense cascade to balance tradeoffs

- Plot of distress declining over the long run, but each individual schema rising and falling over the process
- Make manual more open to those who don't have "trauma", but do have maladaptive schemas. Want them to think they can benefit from therapy.
- Most of the citations need to be made more rigorous (the referenced material backs up the exact claim I make, referenced material is peer-reviewed if possible) and in APA style. Sections starting with * have had this process completed.

Preface

We believe our inherent tribalism is fairly maladaptive in the current human context and that deliberate reconsolidation may be necessary to remedy that.

The process of therapy (fundamentally memory reconsolidation, though many techniques can activate this process) has the potential to heal not just mental illness, but a wide range of maladaptive patterns ranging from the implicit biases driving toxic politics, to insecurity, escapism, disconnection, and more. Traditional psychotherapy strongly relies on the client finding a practitioner they match well with, is expensive, has trouble making progress in the presence of strong resistance, dissociation, or panic, has significant drop-out rates, and frequently doesn't work. The psychedelic-therapy renaissance has emerged partly as a response to these challenges. While we believe a variety of psychedelics have positive roles to play in therapy, we focus on MDMA because of our personal experience and MDMA's relative ease of use for those with trauma.

We think the processes we outline in this document (also achieved through the best cases of traditional psychotherapy) can help you achieve .

We share some concerns and values that **celidwen2023Indigenous** report as broadly agreed-upon by traditional indigenous practitioners of psychedelic medicine:

Traditional Indigenous medicine is an ethical, ecosystem-protective, and holistic system of medicine that interconnects humans and the environment. A sense of reverence for the planet guides all relationships, as well as a commitment to preserve all life. Traditional Indigenous medicine from a systems and relational perspective prompts insight for compassionate living and awareness of collective care to sustain the wellbeing of the medicines themselves as well as all future generations.

The chemical precursors of MDMA, safrole and piperonal, are naturally sourced [169]. Safrole harvesting in Southeast Asia often causes large-scale deforestation and species extinction [**safroleProduction**]. We urge sustainable production that doesn't involve degradation of natural areas. However, we also urge you to not let this knowledge dissuade you from MDMA-therapy, given the importance of connection, care, and mental health. We also believe care for nature (see Section 6.6), and care and compassion for each other and future generations (see Section 6.5) are of paramount importance and were among the primary motivations for this manual.

In the preface, the main editor or author speaks directly, but briefly, to the readers, describing his or her intent in writing or editing the book. Your preface should address the following questions or issues: • What is the book about? What is the scope of the book? • Why is there a need for the information contained in the book? • Why is research in this field escalating? • Why is research in the field important? • What makes the book unusual and worth reading? • Who will be interested in reading this book? • How will the reader benefit by reading this book? Discuss the specific topic of the book but do not include a re-statement of the table of contents.

State fundamental goals of manual?

characteristics of securely attached adults

Our deepest desire is aiding the wellbeing and cooperation of all beings through the unlearning of maladaptive schemas, practice of compassion, and methods for increasing connection. However, we have our own maladaptive fears, and may have projected them into this work in unhelpful ways despite attempting to correct for that. We may also have overlooked important truths. We believe in listening to criticisms of our work and sitting with our discomfort if the criticisms are uncomfortable. We've tried to be critical of the things we're enthusiastic about, but inevitably our biases have pushed us to be critical of some things more than other things. We have tried to strike a balance between practical applicability and scientific robustness. We think the core of the manual is solid, however, some of our citations will inevitably not reproduce in further experiments. Let us know if...

Mark Groeneveld: During my own healing I realized a few things: Core knowledge essential for effective MDMA-therapy is dispersed across many sources, ranging from scientific literature to the wisdom of experienced therapists. Reconsolidation, which MDMA is especially helpful in assisting, can address the root causes of many of the world's problems. Many people are desperate for healing, don't know where to start, can't access expert assistance, and don't have the information to safely and effectively heal. I was also frustrated by not understanding how trauma and healing work, despite successful application of MDMA-therapy.

I thought a manual could help with these problems. My goal is for this document to be a comprehensive and self-contained starting point, including scientific descriptions of the core concepts and brief tutorials on a variety of related practices. Links to in-depth resources are provided for those wanting to dig deeper. This document aims to describe how the mind responds to trauma and how healing works, in a way that is hopefully clear, accurate, and directly applicable to healing. I also hope this will serve as a FAQ, manual, and general reference for MDMA-therapy. Maybe this document will also be helpful for those who didn't understand why they felt the way they did or how to address it. I hope you find it of use!

MG would like to thank: MDMA-therapy, for giving me hope and life. The scientists and therapists who developed this body of knowledge and practice. The indigenous communities who first discovered how to use psychedelic medicines for healing [celidwen2023Indigenous, davisOneRiver]. My friend and guide, Jessica Sojorne Libere, for years of guidance. My partner for encouragement, support, and editing. [r/mdmaththerapy](#) for community insight. ChatGPT-4, for assisting with editing, research, and major contributions to the Appendices, Glossary, Defense Cascade, and Managing Anxiety and Panic [107]. All sections written with ChatGPT have also gone through multiple rounds of thorough editing by both human authors.

Thomas Harper:

Author Contributions: MG did the initial concept and first draft. TH and MG then shared writing, editing, concept development, and research.

Conflicts of Interest: MG has a strong pre-existing belief of the efficacy and safety of MDMA-therapy. MG and TH both have strong pre-existing beliefs of the benefits of treating mental illness and increasing compassion. MG funded the project.

Dedication.
Something
like ecosattva
vows? We
dedicate this
work to the
benefit and
cooperation of
all beings.

1 Introduction

MDMA creates powerful feelings of connection and safety. When used with skill, these emotional states are highly effective tools for healing and adaption. However, there are no quick fixes. Even in optimal conditions, MDMA-therapy and the best examples of traditional psychotherapy can take multiple years to heal severe mental illness. Additionally, confronting and healing the effects of trauma can be destabilizing until the newly-surfaced emotions have been processed. Destabilization can be severe for those with the most intense trauma [91]. That being said, we believe MDMA-therapy presents a significant increase in the potential cost-quality curve of therapy and provides a much needed tool for cases that haven't responded well to other approaches. Various fundamentally similar framings highlight the potential benefits of MDMA-therapy to different audiences:

- Healing mental illness or addiction
- Connecting to yourself, those you love, and the world.
- Developing equanimity, patience, compassion, introspection, resilience, alignment of behavior with goals, and cognitive and emotional flexibility.
- Unburdening from hyper-vigilance, fear, chronic stress, loneliness, addiction, shame, guilt, etc.
- Helping you focus on the things that you can change and let go of the things you can't.

This manual addresses three audiences: professionals seeking a thorough understanding of MDMA, professionals seeking a supplementary resource to give to their clients, and users doing solo MDMA-therapy. Because of the complexities and risks of the practice, we generally do not recommend starting MDMA-therapy without professional assistance. However, availability, cost, or trust obstacles often make adequate professional assistance hard to access and many people have to make the best of non-optimal circumstances. This document is insufficient by itself to train guides.

This guide doesn't offer personalized medical advice, guarantee healing, assure the prevention of negative outcomes, or prevent legal problems if used in a place where MDMA is illegal. Instead, this guide presents a framework for increasing the efficacy and safety of MDMA-therapy, grounded in research, community insights, and author experiences. While this guide has universal aspects, it doesn't cover all frameworks for MDMA-therapy. Although MDMA-therapy has been practiced by underground therapists for decades, comprehensive scientific study is relatively recent, leaving some aspects unexplored [113]. Possessing psychedelics is a felony in many jurisdictions; this guide doesn't endorse illegal activities [5].

As of 2024, we are in a psychedelic hype bubble [170]. There are many reports and anecdotes of psychedelics treating a wide variety of health conditions. The hype bubble makes it difficult to figure out which are true and which are false. We recommend the Understanding Science 101 guide from UC Berkley [110]. Misinformation is particularly concentrated in health claims. We suggest wariness of charismatic people, non-scientific mechanisms of action, cure-all or quick-fix claims, or companies selling proprietary treatments (most psychedelics are not under patent protection). There are also many claims of

shuffle other paragraphs into introduction or spend more time summarizing the paper.

expand

talk about legality and risk

psychedelics solving a wide variety of issues including war and oppression [170]. We don't believe that just the medicine will change people's beliefs and actions related to these issues. However, we do think the fear partly driving these problems can be unlearned by explicitly activating these feelings during an MDMA session.

We start with an overview of how our minds and bodies respond to trauma, how healing works, and the efficacy of MDMA-therapy. Then we describe the practice of MDMA-therapy, including the trade-offs of professional guidance vs. self guidance, safety, preparation, how to conduct the session, how to process the session, troubleshooting, and how to minimize overwhelming symptoms. We also discuss a variety of complementary practices and tools. These are not essential to the practice of MDMA-therapy itself, but appropriate use will quicken and deepen your overall healing. There is a glossary (Section 7) of common terms before the appendices.

We base this guide on these core assumptions:

- The emotional learnings/schema framework explains most, but not all mental illness (Sotala [151] and Ecker, Ticic, and Hulley [38]). It does not explain the genetic or environmental susceptibility to certain symptoms of schemas or stress, such as psychosis. Certainty - medium/high. This claim is quite large, but the model has a compelling mechanism and epistemological foundation in neuroscience research. We think the general model is true enough to work well. Some parts may turn out to be different than currently understood. It also aligns well with the realizations therapy clients have after successful therapy.
- Memory reconsolidation, or something similar, is the universal mechanism of unlearning maladaptive schemas (Sotala [151] and Ecker, Ticic, and Hulley [38]). Certainty - high. A variety of studies have established memory reconsolidation in simple cases. It also has an epistemological foundation in neuroscience research, unlike most therapy frameworks.
- MDMA facilitates memory reconsolidation when used appropriately (Feduccia and Mithoefer [46]). Certainty - medium/high. We see strong overlap between the effects of MDMA and the memory reconsolidation. MDMA is well-known to increase feelings of safety, empathy, and love. These naturally decrease avoidance and provide an emotional mismatch for a variety of negative-emotion based schemas. Personal experience also closely matches the described process of memory reconsolidation. Controlled trials investigating this assumption are difficult because placebo-control doesn't work well with mind-altering substances.
- The subsection of the autonomic nervous system called the defense cascade governs many symptoms of mental illness and schemas, such as anxiety and dissociation (Kozłowska et al. [76]). Certainty - high. The general principals seem well-established.
- Psychiatric destabilization and consequent defense cascade activation explain the large majority of therapy-induced (including MDMA-therapy) adverse psychological effects (Olthof et al. [106]). Certainty - medium. Therapy-induced destabilization seems well established, though under theorized.

- Further memory reconsolidation resolves psychiatric destabilization and consequent defense cascade activation (Olthof et al. [106]). Certainty - medium/high. A variety of research has shown that destabilization is associated with better therapeutic outcomes. We don't think a rigorous, comprehensive model of destabilization or temporary symptom worsening has been established. We think there are multiple mechanisms, some of which we later describe.

We frequently rely on several other, less critical assumptions:

- MDMA-therapy induces a period of neuroplasticity that makes memory reconsolidation easier (Nardou et al. [101]). Certainty - medium. This relies on a small set of studies by one lab group on mice, but matches up with personal experience.
- Acute physical injury from MDMA is caused by mixing it with dangerous activity, certain other drugs, or certain pre-existing medical conditions, not MDMA itself (Rigg and Sharp [131]). Certainty - high. The primary causes of injury seem well-understood and we are not aware of a single acute injury in MDMA clinical trials, where dangerous activity and drug interactions are absent, and participants are screened for liver and cardiovascular problems. There could be rare exceptions that are poorly understood.
- MDMA does not cause long-term cognitive impairment when used in low-risk contexts (Halpern et al. [60] and Passie [113]). Certainty - medium/high. This largely relies on one observational study, but we think it was thorough and convincing.
- MDMA does not cause long-term physical problems when used in low-risk contexts (Nutt, King, and Phillips [104]). Certainty - medium. We're not aware of any research specifically investigating this. A panel of drug experts concluded that overall, MDMA has some risk of harm, though significantly lower than most drugs of abuse. We think a majority of that harm can be attributed to the known and avoidable risk factors previously mentioned. In therapeutic contexts, MDMA is also used far less frequently than most drugs of abuse. The highest number of therapeutic doses we've heard of someone using is about 30.

Without professional guidance, solo MDMA-therapy users need to acquire a greater degree of knowledge to reduce risks and increase effectiveness. We recommend reading the entire guide. However, if you find this document is overwhelmingly long to read and are also determined to do solo MDMA-therapy, we suggest skimming the Safety (5.1), Dosing (5.3), Pre-Session Preparation (5.5), and The MDMA Therapy Session (5.6) at a bare-minimum. If you run into trouble, look at Troubleshooting (5.7). After the session, consider reading After the Session (5.8).

The writing process involved a back-and-forth combining of informal literature review with professional and personal experience. This was also informed by numerous self-reports of MDMA-induced adverse experiences. We resort to many "we suggest"s because the scientific literature on MDMA and psychotherapy is highly inadequate to the task of saying "this is the best way to do things based on rigorous evidence."

We avoid providing a simple list of instructions to follow because the practice of MDMA-therapy is sometimes complex, we are not able to provide individualized medical advice, and many uncertainties about the practice remain unresolved.

We supplemented this with a review from a professional MDMA-guide. We used informal interview data from x other professional MDMA therapists and guides to inform key aspects.

2 Background, Goals, and Rationale

Mental illness is one of the largest root causes of suffering [165]. Besides being painful, difficult, and expensive for the people who experience it, it carries a heavy economic, emotional, and logistical cost for close companions of those who experience it. On top of all this, the widespread prevalence of mental illness effects such as cognitive distortion, emotional rigidity, and emotions of deep insecurity and threat may be among the engines driving tribalism and political polarization across the globe. While a small number of individuals are able to access effective mental healthcare, both access and effectiveness are inadequate for most of the vast and growing population of individuals who experience mental illness.

We are developing an evidence based user guide to therapeutic MDMA use. Therapeutic use of MDMA is one of the most promising developments in psychopharmacology [97]. This guide has the potential to meaningfully reduce mental illness by addressing several of the most prevalent, highest impact barriers to effective care. By doing this, we hope to:

- Alleviate suffering
- Assist individuals in building their capacity to function effectively in the world.
- Increase the ability of individuals to feel and behave compassionately, including toward non-human animals.

Ultimately, we believe this has the potential to improve the capacity of entire communities to act collaboratively towards constructive ends. Although the usefulness of therapeutic MDMA for individuals who do not experience mental illness has not been established with the same research base, we are also hopeful that it could eventually become a resource for anyone who wishes to experience a greater sense of meaning, connection, and fulfillment in their life. We believe this guide provides the foundational knowledge necessary to leverage MDMA to create transformative increases in compassion and connectedness, relevant to many positive social outcomes as well as to mental healthcare. This project aims to amplify the broad-scale effectiveness of therapeutic MDMA use by helping overcome three categories of barriers:

Trust related barriers: Feelings of trust towards the therapeutic process are an important resource for getting individuals into mental health treatment who need it, and feelings of trust between therapists and clients are an essential mechanism for the effectiveness of therapy. One of the most exciting aspects of therapeutic MDMA usage is that it “hot-wires” the trust systems of individuals who have experienced trauma. This allows them to experience a sense of safety and connectedness that is otherwise physiologically and psychologically inaccessible to them, even in trustworthy circumstances. Additionally, many cultures and subcultures experience a normalized hostility regarding even the acknowledgement of mental illness, creating a challenge for clients who are in need of care and for clinicians who attempt to provide effective care in these communities. It can also be extremely difficult for individuals of a particular identity group to feel trust towards a clinician who shares few of their identities and little of their direct experience. Because of this, the ability to achieve that trust more reliably with the support of MDMA is an enormous asset in increasing the accessibility of effective care. Because of its emphasis on rigor and accessibility, the guide we

are producing has the potential to increase the comfort of a wide variety of individuals who would not otherwise be willing to use MDMA, and the comfort of a wide variety of clinicians who would not otherwise be willing to support them.

Financial and logistical barriers: We believe MDMA is useful for increasing cost-effectiveness for many clients. This can happen through increased efficiency of therapy—it has already shown promise in clinical trials for treating cases resistant to conventional methods. It can also happen through gradually shifting a large amount of therapeutic work to at-home sessions the client conducts by themselves or with a “sitter” who needs far less training than a fully licensed clinician. We believe MDMA-assisted psychotherapy will broadly increase the cost/quality curve of psychotherapy.

Clinician training barriers: Many licensed clinicians lack access to the best and most up-to-date training on a variety of therapies because of financial barriers. Additionally, because MDMA usage is and has long been illegal, clinicians who ask their supervisors and senior colleagues about how to support this form of clinical work are often rebuffed. Most clinicians are unwilling to provide clinical support to individuals who embark on its therapeutic use on their own, due to this lack of training and support, as well as due to the legal ambiguities of such care. Depending on how flexible the FDA rules for clinical support are when MDMA is legalized, it is possible that providing a high quality open access manual at this early stage of the legalization process could have a positive impact on the culture of mental health providers around MDMA support. This has the potential to prevent rent-seeking experts from making an expensive certification the standard in clinical MDMA support before widespread access to therapeutic MDMA can even be established.

We are writing a rigorous, comprehensive, and accessible guide to therapeutic MDMA use. It starts with a clear exposition on the science of trauma and healing, articulating a proposed mechanism by which MDMA therapy most likely achieves its effectiveness. It continues on to a thorough discussion of the practice of MDMA therapy, covering safety, dosage, what to expect physically and psychologically, the role of clinical support, and recommendations on making best use of the neuroplasticity period that follows a dose. This includes discussion of complicating factors such as addiction, poly-drug use, and contraindications. We have included detailed, neuroscientifically grounded instructions on using MDMA to confront the painful and self-defeating feelings that underlie a wide range of implicit social biases. Finally, we include a discussion of therapies and therapeutic supports appropriate for augmenting MDMA therapy, as well as several appendices of supplementary resources to assist therapeutic MDMA users in getting the most out of their experience.

Many people are already doing professionally-guided or DIY MDMA-therapy. The available information on the subject for these professionals, their clients, and DIY’ers is usually poor quality, locked behind expensive training programs, or both. Often the institutions providing training advocate for a mix of evidence based and non-evidence based practices, which requires that individuals who wish to take an evidence based approach must do extensive research on their own. Our manual will provide a much-needed source of high-quality, accessible information on this subject. This can increase the accessibility and quality of MDMA-therapy and decrease its risks.

3 *Summary for Mental Health Professionals

We hope this section, and the whole paper, will provide a valuable review of MDMA-therapy. We noticed many MDMA-therapy clients have questions or feel distressed or confused about parts of their professionally-guided MDMA-therapy sessions. We hope to remedy this by providing better information to both the professionals and clients. We think this manual may serve as a useful handout to clients.

Adding MDMA to a psychotherapy session essentially adds a source of profound safety and connection without the loss of control or hallucinations that LSD or mushrooms can create. This reduces dissociation and resistance, and provides a strong mismatch for re-consolidating maladaptive schemas. We based this manual on the schema/reconsolidation framework because of its epistemological foundation in lower-level neuroscience models and our belief that it is a major component of almost all therapeutic change.. Schemas are memory structures containing multiple integrated and co-activating components: emotional responses (e.g. fear at the sight of dogs), abstract beliefs (e.g. dogs are dangerous), and episodic memories (e.g. being bitten by a dog as a young child). Minds create schemas to guide us toward beneficial actions and away from harmful ones. Schemas change over time as situations change, or as we grow up and acquire increased capability to cope with challenges. A schema reconsolidates (updates) when relevant stimuli activate it, and your brain detects a mismatch between the emotional responses or beliefs of the activated schema and the current situation (or a second activated schema, as is often used in therapy). Schemas can become maladaptive when prevented from updating by avoidance, resistance, lack of resources, fight-or-flight, or dissociation. They are similar to the parts of Internal Family Systems. In therapy, mismatches are often constructed and held in focus for a duration of time to deliberately rewrite and re-store the schema in a new, durable, and adaptive form. The safety and connection of MDMA create a strong, possibly universal, mismatch that easily reconsolidates most maladaptive schemas.

Unlike most therapeutic modalities, going beyond the therapeutic window into a panic response is not a barrier to reconsolidation during an MDMA session. MDMA is also effective at reducing dissociation, however when the dissociation is strong enough it can still be a barrier. It typically presents as sleepiness, feeling bored or sober, or feeling disconnected from emotions. When dissociation impedes the therapeutic process, practicing extended focus meditation toward the dissociation often resolves the issue. A different approach such as Psychedelic Somatic Interactional Psychotherapy (PSIP) may be needed first if dissociation is so intense that the client falls asleep. Resistance or avoidance may also still be present during an MDMA-therapy session and can be worked with by worked with through both standard and novel therapeutic practices. These may include asking the client to practice selective inhibition (see Section 5.4.2), prompting the client to focus on and stay present with their emotions instead of thinking or ruminating, or asking probing questions about feelings, beliefs, and the events that may have created them.

Reconsolidation and insight during MDMA-therapy is often so deep and rapid that with only non-directive support or occasional prompting, clients quickly come to their own accurate insights about how trauma shaped them and why their schemas didn't update. However, guidance may be necessary to deal with resistance or dissociation. This unusual depth of healing can prompt greater degrees of destabilization as dissociation and avoidance rapidly

decrease.

MDMA has a few serious medical and drug contraindications, but is generally safe and non-addictive in therapeutic contexts. We believe its popular reputation as inherently dangerous is not accurate when it's used carefully. Additional caution is warranted in clients with dangerous or difficult to manage symptoms such as a history of psychosis, mania, or strong suicidal ideation. See Section 5.1 for more information.

We suggest guided MDMA sessions, possibly combined with certain safe activities, may also be particularly useful for creating new schemas. These may include bonds with nature or recognition of the inherent moral worth of all beings.

4 The Science of Trauma and Healing

Break up sentences. Too dense

4.1 What is Trauma and Why do its Effects Stick Around?

*Traumas are distressing events or chronic conditions that overwhelm our ability to cope [78]. Our ability to cope depends on our capabilities and resources. Children are easily traumatized because they have very little ability to protect themselves, but a variety of other circumstances also lessen our ability to cope with adversity. These include deficits of: mental health, physical health, socio-economic status, healthy support from other people, physical activity, education, self-worth, emotional capacity to process distressing feelings, and cognitive flexibility [141, 155, 159]. Common traumas include:

reword

- Different forms of unintentional or occasionally intentional neglect or abuse.
- Lack of emotional attunement from parents [21].
- Disasters, accidents, or war
- Chronic poverty, dehumanization, or dysfunctional social-cultural systems [134]
- Loss of health, home, family, or culture
- A wide variety of other difficult situations

Our minds continually create and update a network of unconscious memory structures called schemas [38]. Schemas are composed of detailed memories, abstract beliefs, and emotional responses. They form in response to emotionally positive or negative experiences to help guide our behavior when we encounter similar situations in the future. Our schemas usually update when our ability to handle adversity increases, like when we become adults, or when the situation changes. The updating process involves staying present with the distressing feelings of the traumatic memory. A variety of mechanisms can inhibit this presentness and updating, causing our schemas to keep operating in the same way as when we were less capable:

finish

- **Resistance:** Conscious or unconscious avoidance schemas like drinking alcohol when distressed, escapist habits, addictions, believing "feeling emotions is weakness", believing that shame is proper, fear that healing a schema will hurt you in some way, denial, projection, rationalization, etc [38]. This does not include healthy avoidance, which is temporary and helps you cope in non-destructive ways until you have the space to process. Resistance also includes networks of schemas that reinforce each other: Schema A is itself a stimuli for activating schema B. B then reinforces A with a negative stimulus, preventing A from adapting. Sometimes these networks of avoidance and reinforcement become complex and paralyzing.
- **Lack of Resources or Skills:** Not having a quiet moment or space to reflect and deal with emotions. Being unsure about the reasons for your emotions or how to handle them. Or you may not have skill in noticing and labeling feelings, sometimes called Alexithymia.
- **Fight-or-Flight or Dissociation:** These autonomic defense cascade states inhibit engagement with emotions [128].

give examples

*When these prevent processing of a maladaptive schema, the schema becomes 'stuck', and keeps replaying in the presence of relevant stimuli, without adapting [125, 38]. This emotional response and the associated episodic memory (a narrative of how events unfolded during a specific event or "episode") then form and combine with abstract beliefs about the world to form durable, unconscious or partly unconscious structures called schemas, or 'parts' (from the Internal Family Systems framework) [78, 151]. The emotional, episodic, and belief components interact with each other and activating one activates the others. Schemas are originally protective or adaptive (enhances your ability to cope with your environment; see Figure 1), but sometimes become maladaptive (inhibits your ability to cope with your environment; see Figure 2) when they don't adapt to changing conditions due to the above reasons.




Figure 1: The whole schema updates when either its belief or emotion components experiences a prolonged mismatch with a contradictory experience. Avoidance behaviors are temporary and healthy.




Figure 2: The schema doesn't update and becomes maladaptive when mismatches are avoided or another schema reinforces it. The maladaptive schema may be a stimulus for the reinforcing schema, or they may share a common activating stimulus. The emotional response of the reinforcing schema is intense and outweighs the adaptive mismatch.

Here's an example of a schema that became maladaptive:

get better example

- Situation: As a young child, Amy was frequently ridiculed by her peers whenever she spoke up in class or shared her opinions (episodic memory).
- Initial Schema: "If I voice my opinions or stand out, I will be ridiculed and rejected (belief). Ridicule and rejection hurt (emotional memory)."
- Resulting Behaviors and Beliefs: Amy grows up avoiding speaking in group settings and tends to keep her thoughts to herself (behavior). She might decline leadership positions or avoid roles where she'd be in the spotlight (behavior). In discussions, even if she disagrees or has a valuable perspective, she might not voice it (behavior). Amy might believe she's not as smart or valuable as others, even if evidence suggests otherwise (belief).

This schema operates in the background, guiding Amy's behaviors and beliefs without her necessarily being consciously aware of it. Over time, the schema may reinforce itself if Amy experiences further instances that align with it, making it deeply ingrained and automatic in her responses.

*Schemas are stored as networks of neural connections spread across brain regions [38]. Some networks are complex, composed of multiple traumatic events and activated by a wide variety of stimuli. When you feel the emotions and beliefs of the memory, you are feeling the part of that network that activated in response to a particular stimulus. That stimulus may be singular and obvious like seeing a wasp of the same kind that once stung you, or diffuse and difficult to understand like a mood, thought, or time of day.

often created unconscious

Schemas may not contain an episodic memory if you were too young to form long-term episodic memories, or the experience overwhelmed you to the extent that your mind prevented episodic memory formation [74, 21]. These emotions can be especially confusing compared to emotions we clearly see a cause for. In addition to possibly lacking an episodic component, schemas may contain and sustain other signals in the brain, such as chronic pain or feelings in your chest or abdomen. The mind may interpret particularly strong schemas as separate personalities.

Maladaptive schemas and the defense cascade activations they cause can manifest as: depression, anxiety, resentment, anger, low self-worth, an overactive fight-or-flight mode, hypervigilance, detachment, flashbacks, avoidance, numbness, self-harm, psychosomatic illnesses, shame, guilt, denial, isolation, impulsivity, trust issues, perfectionism, hopelessness, obsessions, disconnection, muddled thinking, etc.

categories (cognitive, emotional, social, identity) in chart form?

We suspect schemas can sometimes activate physical sensations or altered states of mind.

Schemas that outlast their usefulness often harm us and those around us. They may push us to overreact, deny the truth, misjudge important trade-offs, say hurtful things, etc. We may also seek out the connection and safety we desperately need in dysfunctional ways. The stress of chronically activated distressing feelings or chronic defense cascade activation also significantly increases risk for a wide variety of physical diseases and problems [48]. The fear and defense cascade activation of emotional memories can also have a variety of negative consequences, such as being too distracted by our own pain to pay attention to the needs of those we love. These maladaptive patterns and their with sustained defense

cascade activations also predispose individuals to conditions recognized as mental illnesses. About 10% of the population might meet the somewhat arbitrary criteria for mental illness at any given time [108], but almost everyone has some amount of maladaptive schemas that negatively effect them and those around them.

*Schemas and mental illness can also be accurately analyzed through the lens of the biopsychosocial model, where a phenomenon arises through biological predispositions and functions, social models of how one should respond, and an individual's psychology [164].

4.2 *Defense Cascade

The Autonomic Nervous System (ANS) governs a wide variety of involuntary bodily functions, such as heart rate and digestion [76]. While mainly operating unconsciously, the ANS also activates a defense cascade—a sequence of responses—to shield us from threats. Increasing levels of perceived threat activate these responses, though the order of activation depends on individual variability and experience:

- **Arousal** The most common initial reaction to a potential threat. Think of how a deer becomes alert when they see something moving far away. Vigilance, muscle tension, respiratory rate, and heart rate all increase, allowing us to quickly assess and respond to possible dangers.
- **Flight or Fight** When an imminent danger is identified, like when a deer notices a wolf nearby, this response prepares the body to either confront (fight) or escape (flight) the threat. The adaptations of the arousal stage intensify and are augmented by an adrenaline surge, further suppression of pain, and an urge to fight or run.
- **Freeze** When the danger is imminent, but you might go unnoticed, such as when the wolf has not yet noticed the deer, the freeze response temporarily pauses a fight or flight response. If the predator notices you, freezing quickly reverts to fight or flight. While most physiological responses from fight or flight remain, muscles become immobilized.
- **Dissociation: Tonic/Collapsed Immobility** Tonic immobility (playing dead) may dissuade a predator from eating you when you have been caught. Fight or flight responses are deactivated, the body is paralyzed, and the brain produces opioids to numb and disconnect you from reality. Muscles remain tense in case the predator gets distracted, and you have to run again. This state may transition to collapsed immobility (fainting; muscle tension and consciousness partially to fully lost) when the threat further increases, or dissociation may continue increasing without a loss of consciousness. These dissociative states often become chronic in humans, and may not involve paralysis or sleep [128]. Extreme threats such as lack of attunement from parents, may cause extreme dissociation from bodily sensations, emotional memories, and current feelings, even while the person appears to be functioning and conversing normally. Chronic dissociation is association with trauma in situations of powerlessness [84].

resolve whether fight or flight is deactivated during dissociation; explain how fight or flight can co-exist with dissociation

- **Quiescent Immobility** Tonic/Collapsed Immobility may extend into a lethargic rest and recuperation phase after the threat has gone. Occasionally, this may persist beyond its period of usefulness and become maladaptive.

The defense cascade originally evolved to address immediate physical threats like predators, but is also activated by a wide variety of other situations [128]. Children are especially prone to these reactions because their threshold of what constitutes a threat to their life is much lower than it is for adults. Lack of parental support, attention, or attunement (see Appendix A.2) can easily activate defense cascade states in children. When this lack is chronic, durable schemas often form and activate the defense cascade whenever the memory is activated. This lasts a lifetime if the stuck schema doesn't reconsolidate.

Recurrent activation of these states contributes to conditions such as anxiety, depression, PTSD, and dissociative disorders [128]. PTSD sufferers, for instance, might frequently switch between heightened aroused and dissociative states. Dissociative and panic states can also simultaneously coexist in many cases of depression and anxiety. We suggest understanding this enables safer and more effective healing:

what is this

- **Awareness and Insight** Recognizing where you might be within the defense cascade can clarify your reactions and behaviors.
- **Self-Regulation** With awareness, employ techniques like deep breathing, or mindfulness to transition from heightened arousal to a calmer state.
- **Pacing** Observing your shifts between hyper-arousal (fight/flight) and hypo-arousal (freeze/collapse) can inform the rhythm therapy, indicating when to proceed or pause.
- **Empathy Towards Self** Acknowledge these reactions as once protective, fostering self-understanding and compassion.
- **Creating a Safe Environment** Prioritize crafting a secure physical and emotional space, potentially setting boundaries in scenarios prone to activating stimuli.

For a comprehensive understanding of the defense cascade in psychedelic therapy, refer to *The PSIP model. An introduction to a novel method of therapy: Psychedelic Somatic Interactional Psychotherapy* [128].

4.3 Mechanism of Healing

too dense

MDMA primarily creates feelings of safety, love, and connectedness. In this state your resistance schemas naturally relax and allow processing of avoided or dissociated schemas. MDMA-therapy proceeds similarly for most types of trauma, with minor variations to accommodate individual needs [109].

Long-term healing of stuck schemas hinges on memory reconsolidation [46]. When memories/schemas are activated, they become plastic and amenable to change. Introducing a mismatch between the activated emotion/belief and a separate, contradictory emotion or belief rewrites and re-stores the schema in a healed state [78]. The mismatch can arise from MDMA-induced feelings of safety and connectedness, a feeling of safety from an attuned

less sure about proceeds similarly for most people

therapist, or the contradictory experience of a second activated memory as done in Coherence Therapy [38]. The experiences of everyday life often provide a small mismatch, slowly reconsolidating surfaced and non-reinforced schemas. Only the part of the memory network that is activated during therapy gets reconsolidated. Complex memory networks, arising from repeated traumas, may require multiple sessions activating different activating stimuli. A schema may also not fully reconsolidate if, during the session, a different schema came into focus before the first reconsolidation was completed. While reconsolidated schemas are stable, it's uncertain how prone they are to reverting to their original state after experiencing new trauma. Increased caution against future re-traumatization may be necessary.

diagram of network hypothesis

The more conceptually related the positive experience is to the stuck schema, the more effective the mismatch. This may be due to a variety of mechanisms : a related feeling activates some of the same neural circuits as the schema, the mind may label related experiences as more significant, or related feelings/beliefs may activate each other and strengthen the process. MDMA may provide a universal mismatch that works for most if not all schemas. This may be because most maladaptive schemas are fundamentally a fear of disconnection or pain, and MDMA creates exceptionally intense feelings of connection and safety [19].

Reconsolidation often reduces the intensity of distressing feelings of a memory. It also commonly presents as more complex changes in self-conception, alterations in the narrative or context of the memory, shifts in associated beliefs or values, expansion of emotional perspectives, integration of previously separated aspects of the experience, or the development of greater cognitive flexibility in relation to the event. Because many schemas contain indistinct beliefs about the self, healing experiences may not necessarily be subjectively conceptualized as activation and reconsolidation of a maladaptive schema.

Ecker, Ticic, and Hulley [38] describe the following signs of a completely reconsolidated schema:

A specific emotional reaction abruptly can no longer be reactivated by cues and triggers that formerly did so or by other stressful situations.

Symptoms of behavior, emotion, somatics, or thought that were expressions of that emotional reaction also disappear permanently.

Non-recurrence of the emotional reaction and symptoms continues effortlessly and without counteractive or preventive measures of any kind.”

Reconsolidation is the core mechanism of healing maladaptive schemas, but it is not the only part of healing. Learning healthy habits and emotional skills are also critical, as is insight into how trauma affected you and those around you.

Healing also depends on staying within the therapeutic window, where schemas are activated but not overwhelming, and defense cascade activation remains below the threshold of distraction.

A track record of successful treatment is also vital. Repeated unproductive attempts at MDMA-therapy, or even a single exceptionally distressing session, may push someone to unfairly discount the possibility that MDMA-therapy can be effective. This guide aims to put distressing sessions into a larger context where the distress or adverse outcome might be an okay part of a long-term healing process, an unnecessary outcome that can be avoided

Talk more about disconfirming experiences and how they can prompt unconscious daily reconsolidation

with proper technique, or sign that MDMA isn't the right tool. Another goal is providing best-practices, risk-factors, and troubleshooting information to reduce the likelihood of unproductive sessions.

Trauma-therapy presents challenges in addition to its profound benefits. Distressing, previously hidden or avoided emotions are brought to the surface for processing. These emotions may be destabilizing until they have been fully processed. You can think of it as standing up. Sitting and standing are both stable positions. The transition positions between the two are unstable, but must be passed through if you want to walk anywhere. Destabilization is a sign of therapeutic progress, however, it can negatively impact your life if it is severe enough and not managed well [106]. We think destabilization is composed of the following components:

- In the Internal Family Systems framework, some schemas termed "protectors" facilitate emotional stability by pushing us to avoid stimuli that activate feelings of distress [143]. Protectors can increase in intensity when the overwhelming feelings they sometimes push us to avoid are engaged with during therapy. This is termed protector backlash. Some of these protectors may be old and unremembered.
- The targeted schema itself is also often overwhelming, and now that we are aware of it, we may no longer be able to avoid its presence.
- Decreased dissociation also exposes overwhelming schemas to consciousness.
- Disturbing insights about one's past, behavior, or beliefs may provide additional emotions.
- As inner stability increases, or as a result of insight, coping behaviors or beliefs may fall away. This change is usually healthy in the long term, but the lack of a familiar coping pattern may be hard to deal with in the short term.
-

Our schemas form complex networks of adaptive responses that unconsciously guide us through life. Occasionally, this network can be fragile, like a house of cards, in those with extensive early childhood trauma. Therapy can produce functionality collapse and dysfunction if a foundational schema (even if maladaptive) is reconsolidated before you have developed alternate coping strategies.

We propose a description of the healing mechanisms of MDMA in Figure 3.

Making a schema explicit probably means you won't be able to suppress it again. It will be with you until you reconsolidate it.

Schemas interconnected with complex structure. Progress may appear non-linear with respect to quantity of reconsolidation. Schema structure is addressed by happen chance of triggers, modulated by complex interconnectedness of schemas. Some schemas may need to be reconsolidated before others become apparent or reconsolidatable.

Diagram of ups and downs for each




Figure 3: Reconsolidation is achieved through multiple interventions, some of which MDMA directly assists with: 1. MDMA introduces an emotion mismatch. 2. Learn emotional skills and develop resources 3. Feelings of safety lower resistance and deactivate the defense cascade. The feelings of safety from MDMA are sometimes insufficient by themselves when facing extreme dissociation. 4. Reconsolidation requires focus and is exhausting in extended sessions. The stimulant effect of MDMA assists with this.

Maladaptive isolation of beliefs causes particularly acute problems in mental illness. However, it is also pervasive in the non-mentally ill. Just think of emotionally charged arguments on social media involving two conflicting worldviews. It is rare for someone on either side to acknowledge the validity of any part of their opponent's argument and update their own worldview accordingly.

Talk about integrating disparate schema networks. Rebus?

4.4 Efficacy of MDMA-Therapy

*The MDMA phase 3 clinical trials (the final round of clinical trials that test efficacy and safety on a large sample size) reported that MDMA-assisted non-directive psychotherapy highly outperformed placebo-with-therapy treatments for PTSD, as Table 1 shows [98, 97]. Each subsequent dose (the trials included 3 doses) improved outcomes further, likely indicating further doses would further increase healing. The reported outcomes persisted when the data was reanalyzed by an independent, blinded programmer. Only 5% of participants in the

MDMA groups discontinued treatment (half for reasons unrelated to the study), compared to 16% in the placebo groups. MDMA-therapy worked across severity of symptoms, presence of mental illness comorbidities (including substance addictions), history of ineffective treatment, and race. The phase 2 long term follow-up study shows the effects are durable [66]. While rigorous and highly promising, these studies were organized by a single organization, and a different methodology may show different results. MDMA-therapy with therapists less familiar with MDMA may be less effective. See meta-analyses from Green et al. [55] and Smith et al. [150] for a more thorough discussion of efficacy.

Psychedelic-assisted-therapy and psychotherapy trials frequently suffer from methodological problems such as poor blinding and high researcher bias [1]. Because phase 3 trial participants (and their therapists) accurately deduced whether or not they had been given MDMA or placebo, the trials effectively were not blinded. While this is a major limitation of these types of studies, it doesn't totally disqualify them as useful sources of evidence. Psychedelic blinding is inherently difficult to account for. Additionally, some trial participants reported intense expectations of good results [29]. This expectancy effect likely exacerbates the consequences of poor blinding. We suggest Elk and Fried [40] for a thorough discussion of these problems. Notably, if the placebo effect is strong enough to achieve memory reconsolidation, then the therapy was still successful. We suspect MDMA-therapy is more effective than placebo due to its close alignment with the memory reconsolidation mechanism of action. There are also numerous reports of profound therapeutic improvement, even inadvertently in situations where the user was intending a recreational, rather than therapeutic experience.

*MDMA therapy has shown some effectiveness for PTSD, dissociation, depression, and functional impairment [55]. It has also improved well-being, excess vigilance, nightmares, avoidance, anxiety, and sleep in those with PTSD [150]. It is also effective for durably improving alexithymia and reconsolidating maladaptive schemas in the presence of strong alexithymia, which is often a barrier to reconsolidation [75]. We believe MDMA-therapy has a strong potential to be effective for most mental health issues, given the maladaptive-schema/defense-cascade model of mental illness.

We suspect strong suicidal ideation, and some cases of psychosis and mania are treatable with MDMA-therapy because of how maladaptive schemas and stress can trigger these symptoms. However, this is poorly studied and the risks of exacerbating those conditions is high. Therapy can often make symptoms temporarily worse as the underlying schemas are confronted. We only recommend experimenting with MDMA-therapy with these conditions if existing treatments are inadequate, and then only in the physical presence of a professional with a strong understanding of treating these conditions, and understanding of MDMA-therapy, and effective symptom oversight and management.

*SSRIs have small average effects in short term use (long term use is poorly studied) while sometimes causing adverse side effects [23, 18]. Long term benzodiazepine use for anxiety also has low efficacy and severe side effects such as long term mental impairment that persists after discontinuation [147, 12].

further discuss what symptoms depend at least partly on schemas

this paragraph is a bit odd.

Discuss nuances and variety of treatment options? I feel confident that reconsolidation works. How to mental health interventions work? Is exercise effective by itself or is it just an example of a task that requires effort and feels rewarding? We should talk about the fundamen

Table 1: Averaged (own work derived from Fig. 3 in Mitchell et al. [98] and Fig. 3 in Mitchell, Ot'alora, Kolk, et al. [97]) outcomes from the first and second phase 3 clinical trials (174 total participants). These numbers represent therapeutic progress on the single maladaptive schema that participants chose as their therapeutic target [29]. Those with many maladaptive schemas, as is typical for early-life trauma, will need a much longer time frame and number of sessions to reconsolidate all the schemas they want to address.

	MDMA w/ Therapy	Placebo w/ Therapy
No Response	13 %	35 %
Clinically Meaningful Response	87 %	65 %
Loss of Diagnosis	69 %	40 %
Remission	40 %	13 %

5 The Practice of MDMA-Therapy

5.1 *Safety and Challenges

MDMA-therapy is generally well-tolerated, but there are dangerous or deadly drug interactions, medical contraindications, notable side effects, and psychological risks that we discuss below. As further detailed in later subsections, MDMA-related problems are generally caused by [131, 136]:

- Prolonged, intense physical activity in high temperatures combined with dehydration. This may cause heat exhaustion or heat stroke. High water consumption not balanced with salt can also cause dangerously low sodium levels. Alcohol co-use strongly exacerbates the risk of these problems. Potentially deadly.
- Taking MAOIs (including ayahuasca) within 2 weeks of a session. Potentially deadly.
- Interactions with other medications. Ritonavir, opioids, benzodiazepines, amphetamines, stimulants, caffeine, anesthetics, MDMA metabolites or analogs, muscle relaxants, olanzapine, and metoclopramide may be particularly dangerous and occasionally deadly. Discontinuing medications for 5 drug half-lives before the session and 2 days after the session generally assures a significant amount of the other drug won't be in your body at the same time as a significant amount of MDMA. As with MAOI's, this doesn't prevent all interaction problems. Discussing drug combinations with a doctor or pharmacist may be necessary.
- Pre-existing liver or cardiovascular problems. Potentially deadly.
- High or frequent doses. Clinical trials found 120mg + 60mg (booster taken 1.5-2 hours after the initial dose) once every 6 weeks to be very well tolerated. Thresholds and risk unclear.
- Adulterated pills. This can be checked with test kits. Risk varies by adulterant.

list number of therapist-hours

more on what is known and not known. history of underground use

compare to other treatments like exercise, meditation, relationships

should we list symptoms of physical problems when you should seek immediate medical treatment?

soso advice, need better

- Driving or other potentially dangerous activities in an altered state of consciousness. Potentially deadly.
- Psychological destabilization is common after processing traumatic schemas. Psychedelic-assisted therapy accelerates the process, increasing both healing and destabilization. History of psychosis, mania, and suicidal ideation are strong indications to work with a skilled and trusted practitioner familiar with these conditions. Destabilization can cause significant problems when poorly managed. Rarely deadly.

In a highly supportive setting with medical screening of participants, both phase 3 clinical trials measured no serious or long term adverse effects (among the set of outcomes they looked at) at 6-week spacing for 3 sessions with 180 mg MDMA (combined measure of primary and supplemental doses) [98, 97]. Higher doses, more frequent sessions, and a greater number of sessions are likely safe, but it's unknown where the limits are and experimentation can be risky. Doing more than three sessions is likely safe but could potentially introduce unknown risks, which should be weighed against potential benefits. It is also possible that participants experienced a set of problems that the study was not capable of detecting.

Observational human studies and controlled high-dose animal studies have found that MDMA causes neurotoxicity or cognitive problems [113]. However, these problems have not been found in controlled studies in humans [60, 98]. The human observational studies that find problems usually fail to adequately control for multiple-drug use, a significant risk factor. Notably Halpern et al. [60] found no long-term cognitive effects of recreational MDMA use in a population that had exceptionally low lifetime use of other psychoactive substances. The animal studies that found problems typically used extreme doses, extreme frequencies of dosing, or injected the medicine, a significantly riskier method of administration than swallowing a pill [113]. Some studies have reliably found serotonin system alterations in humans, but it hasn't been established that these are clinically significant or negative, and the effect decreases with abstinence [54]. Fear and misinformation about MDMA is widespread due to the War on Drugs, sensationalization of poor quality research, and misattribution of MDMA-related deaths as an inherent risk of MDMA instead of its avoidable interactions with other medications and pre-existing health conditions, and the risks of heat illness or hyponatremia at raves [113]. See chapter "The Toxicity Debate" in "The History of MDMA" by Torsten Passie for a comprehensive review of the topic.

On a population level, recreational MDMA use leads to acute physical injury far less often than horse riding does, for instance [105]. When acute harm does result in death, it is generally caused by the previously listed, avoidable mechanisms [131].

Colcott et al. [27] published a side-effect meta-analysis of MDMA-therapy trials and found that MDMA therapy "...is associated with greater likelihood of experiencing mild to moderate, but largely transient side effects. Findings, however, need to be interpreted cautiously due to limitations of the existing evidence." The statistically significant side effects included, starting from the most common: dilated pupils, feeling jittery, uncontrolled eye movements, excessive perspiration, chills, blurred vision, jaw clenching/tight jaw, muscle tightness, decreased appetite, restlessness, non-cardiac chest pain/discomfort, feeling cold, and nausea. Several serious side effects were recorded but were too rare to achieve statistical significance. Larger trials would help us better understand rare side-effects. Side effects were also measured by spontaneous self-report. Systematic measurement with checklists

would likely report more side effects. As previously discussed, MDMA therapy commonly accelerates the processes of reconsolidation and destabilization, the latter of which can be unpleasant and disruptive.

Evans et al. [43] surveyed people who have experienced new, persistent negative symptoms after recreational, professional-therapeutic, and DIY-therapeutic psychedelic experiences. This data applies to all psychedelics; we could not find MDMA-specific data on this topic. We believe most of the listed symptoms could be explained by surfacing schemas and defense cascade activation, parts of the trauma-therapeutic process that psychedelics are known to facilitate. We suggest that most of these persistent symptoms could be resolved through additional reconsolidation and therapeutic support. Even in this subgroup of people who experience extended difficulties, "90% agreed that psychedelics can be helpful and are worth the risks if taken in supportive settings." However, it may be possible that a population of psychedelic users who experience debilitating long-term effects was missed due to sampling bias.

Drug Interactions We suggest **papaseitInteractions** and Sarparast et al. [140] for thorough discussion of MDMA drug interactions.

- Reuptake inhibitor medications, including SSRIs, SNRIs, NRIs, and NDRI, highly inhibit the effects of MDMA long after discontinuation [47]. Discontinuation 25 days before an MDMA session still reduces therapeutic efficacy of MDMA-therapy by about half. Further discontinuation may bring further benefits. Discontinuation typically requires multiple additional weeks of tapering to manage withdrawal.
- The interactions between MDMA and many drugs remain uncertain, and self-experimentation could lead to unforeseen complications [24, 140]. We suggest avoiding all other medications for 5 drug half-lives (The point at which 97% of a drug has been excreted from your body . Each drug has a different half-life.) before the session and 2 days (5 MDMA half-lives) after the session unless you have discussed the combination with a doctor or pharmacist [8, 32]. The following drugs are correlated with adverse outcomes when used with MDMA, though cause and effect remain unclear: opioids, benzodiazepines, amphetamines, stimulants, anesthetics, MDMA metabolites or analogs, muscle relaxants, olanzapine, and metoclopramide [24]. Caffeine also increases risk at high MDMA doses in rats [162].
- MAOIs can cause a dangerous or deadly effect known as serotonin syndrome when taken within two weeks of an MDMA session [86, 39]. Notably, ayahuasca includes an MAOI [ruffell2020pharmacological].
- The liver enzymes CYP2D6, CYP1A2, CYP2B6, CYP2C19, and CYP3A4 metabolize MDMA [33, 140]. Exercise caution taking other drugs (especially high doses) that these enzymes metabolize, as this metabolic path may overload and reduce clearance of these drugs from your body. Drugs that inhibit these enzymes, such as ritonavir, may be dangerous to take with MDMA. We have not been able to locate an accessible and comprehensive list of these drugs and recommend asking a doctor or pharmacist.

soso advice,
need better

- Pills marketed as MDMA often contain harmful adulterants [138]. We suggest using test kits, such as DanceSafe's, to verify the substance's purity if there's any doubt about its quality [30].

Medical Contraindications and Notable Side Effects

- MDMA has multiple interactions with the cardiovascular system [17]. Pre-existing cardiovascular problems, high doses, frequent use, and co-use of other drugs heighten risk of cardiovascular injury.
- Problems with the liver or liver enzymes CYP2D6, CYP1A2, CYP2B6, CYP2C19, CYP3A4, and COMT may prevent metabolism of MDMA and cause problems [33].
- MDMA, like many medicines, may have uncommon, poorly understood side effects, particularly after long-term or frequent use. Long term (1 year) follow up has only been conducted with 103 clients [66] and no study can investigate all possible health effects. Further studies investigating more health measures, longer-term MDMA use, larger numbers of participants, or higher-risk populations may discover currently unknown adverse effects.
- Dangerous overheating or low sodium levels may occur if MDMA is combined with dehydration, intense physical activity, a hot/humid environment, and lack of electrolytes, as sometimes occurs at dance parties [7]. This is not a problem with a comfortable environment, appropriate hydration, and lack of intense physical activity [98]. Exercise increased caution if your body has serious problems regulating its temperature. Alcohol co-use also significantly exacerbates the risk of heat illness and hyponatremia.
- Planning careful or reduced movement during altered states of consciousness may be advisable if your body is prone to injury from otherwise typical human movement. A sitter, guide, or therapist could help you with this during the session, or you could put an obvious reminder sign in your field of view.
- Adolescent use is unexamined. Kangaslampi and Zijlmans [70] saw no obvious reasons why adolescent use would be risky. However, this assumption has not been tested in any trials. Developing bodies could react differently to MDMA than adult bodies.

is this of practical concern?

Psychological Risks

- Overwhelming emotions from severe trauma often activate pervasive dissociation that hides these feelings from conscious awareness [128]. Reducing this dissociation is necessary to reconsolidate. Trauma therapy often reduces dissociation faster than it reconsolidates the distressing schemas the dissociation was hiding. That sometimes results in significant distress until those schemas are healed through further therapy. Larger amounts of trauma and the rapidity of healing with MDMA increase the likelihood of this distress becoming destabilizing or disruptive to your life. We suggest more application of grounding techniques is necessary in instances of destabilization.

risks of problems from destabilization are strong indication to work with a skilled and aligned professional

community care if you can't access professional

- Some cases of personality disorder are likely treatable with MDMA-therapy [157]. These conditions involve extremely strong resistance, and destabilization during therapy may be intense [128]. The process of MDMA-therapy will likely be more difficult and may benefit from additional support if that is accessible.
- Psychosis induced by therapeutic doses of MDMA without confounding factors appears to be rare. One participant in clinical trials developed psychosis even though participants with existing psychotic symptoms were excluded from participation [91, 150, 97]. Multiple case studies report psychosis following recreational MDMA-ingestion, though they almost always involved confounding factors like use of other psychoactive drugs, lack of verification of MDMA ingestion, dehydration, extreme doses, or extreme frequency of use [90, 115, 161]. We only found one case study describing psychosis following a single purported dose. Even if the risk of MDMA-induced psychosis is very low, the consequences of psychosis are high. Because of this, we suggest that a family or personal history of psychosis is a strong indication to work with a professional who thoroughly understands both psychosis and MDMA.
- Possession of multiple doses may be risky for those with severely impaired impulse control.
- Altered states of consciousness can impair awareness and judgement. Avoid driving, operating heavy machinery, and other risky activities on the same day as the session.
- No clear statistical connection between recreational MDMA and suicidal ideation has been found, and correlational studies have had mixed results [68, 67, 73]. However, individuals can experience increases in suicidal ideation as a result of destabilization [91].
- Mania is a dangerous condition that can also be exacerbated through destabilization. A family or personal history is a strong indication to work with a trusted professional who understands both mania and MDMA.

Common Concerns

- **Discomfort with Drugs** While MDMA-therapy is not for everyone, healing can happen even when discomfort is present during a session because MDMA rewrites patterns of discomfort that you stay present with [46]. Fear of drugs could be transformed like any other fear.
- **Loss of Control** While engagement with distressing memories can be intense, people regularly have clear, complex, and emotionally nuanced conversations on MDMA [26, 113]. MDMA creates intense feelings of love and safety that make aggressive behavior unlikely.
- **Drug Stigma** Many drugs are harmful, however, during the War on Drugs a wide variety of psychoactive substances were further stigmatized and categorized as harmful without clear evidence-based distinctions regarding their actual risk [4, 104]. While

What degree of suicidality is an indication to not do this without professional assistance?

what support do we recommend for less-strong suicidality?

say more

there were complex motivations for the War on Drugs, it functions and persists primarily as moral panic, a means to punish certain groups of people, and a means for politicians to disenfranchise and ostracize the voter base of their political opponents. There is little correlation between the legality of psychedelics and their potential for harm [104].

- **Hallucinations** Complex or compelling hallucinations on MDMA are rare and correlated with higher doses [81]. Temporary and mild visual changes such as color and texture enhancement are common. Long-lasting visual effects from MDMA are unrecorded in studied therapeutic contexts [163]. Psychedelics, along with other drugs, occasionally make typically-unnoticed artifacts of the visual processing system more noticeable, resulting in Hallucinogen Persisting Perceptual Disorder [alexanderHPPD, 59]. People may then feel anxiety about these artifacts, making them even more noticed.
- **Addiction** MDMA addiction is rare to nonexistent in supportive therapeutic contexts where it has been formally studied [97, 98]. Physical dependence has not been found in rodent studies, even at extreme dosing schedules [132]. MDMA-therapy also reduced alcohol addiction among clinical trial participants [103]. However, we suggest that MDMA, like most potentially pleasurable experiences, can be overused when it is used for escapism rather than therapy. We also suggest that MDMA can be overused when an individual wishes to accomplish a great amount of therapy quickly but does not understand the critical need for sufficient sleep between sessions or the quickly-developed tolerance to MDMA in the brain [148, 45]. As previously discussed, there have been many flaws in studies on MDMA abuse. Consequently, little is known about the effects of abusing MDMA, or what level of use causes physical or psychological. If you are worried about your potential for addiction, we suggest the following: only possessing one dose at a time, only obtaining MDMA from a trusted mental health professional who can monitor your abuse potential, prioritizing diving deeper into traumatic memories during sessions may reduce the pleasure of the experience, or combining MDMA with a small amount (<0.5g) of dried psilocybin mushrooms may also reduce the pleasure of the experience for those with significant trauma.
- **Neurotoxicity** We think MDMA neurotoxicity is unlikely in humans at therapeutic doses [113, 60]. However, MDMA is known to cause oxidative stress in the brain (possibly a form of neurotoxicity) at very high doses in animal tests. This may also happen at sufficiently high doses in humans, though it has not been demonstrated. High doses of certain antioxidants, including alpha-lipoic acid, ascorbic acid, and acetyl-L-carnitine, administered shortly before and during the session, have been shown to reduce or prevent oxidative stress in rats [2, 146, 6]. Some companies bundle these antioxidants together in commercially available products, but they have not been rigorously tested in humans and their usefulness is unclear [133].
- **Is MDMA a Natural Substance?** MDMA is generally made by modifying the plant compounds safrole or piperonal [169]. When properly conducted, this process results in pure MDMA crystals. Single-substance purity greatly improves the ability

what does
mdma addic-
tion look like
in reality

discuss risks
of behavioral
addictions

do I really
want to sug-
gest mush-
rooms here

to produce accurate, safe, and reliable doses. MDMA test kits are useful if you don't trust the source of your medicine.

day write
more

5.2 Professional Guidance vs. Self Guidance

*MDMA-therapy can be successful in a wide variety of contexts including therapist-guided sessions with pre- and post-session support, do-it-yourself couples therapy, and solo therapy [97, 26, 63]. However, we believe working alone with MDMA presents higher risks and a lower healing likelihood compared to partnering with a skilled and well-matched therapist or guide. As Ecker, Ticic, and Hulley [38] state, "The therapist's stance is one of cooperating with and trusting the client's inherent capability of revising emotional learnings." A skilled and well-matched therapist or guide can provide [100]:

- A greater feeling of safety, enabling more effective healing
- Additional perspective that's hard to see from a first-person view
- Education on trauma, healing, and healthy ways to deal with emotions
- Troubleshooting for problems with the medicine or other parts of the healing journey
- Improved screening for contraindications (conditions that make MDMA particularly risky)
- Management of destabilization to reduce disruptions to your life
- Guidance through difficult therapeutic exercises
- Unadulterated medicine, depending on who you're working with

*We propose the following ranking of options and outcomes. The top of the list represents the lowest risk of adverse outcomes, the highest likelihood of durable healing, and also the highest cost. This list is a general guide and the exact positions of items are debatable, as well as dependent on personal circumstance. Exceptions to the rule always exist.

1. Continually working with a skilled, psychedelic-trained therapist or guide you align well with. *We tentatively think this is the ideal model for all situations if you can find the right clinician to work with and have enough money. We also think this model is especially important for people with potentially dangerous symptoms they can't manage themselves, including psychosis, mania, suicidal ideation, or a severe lack of impulse control. Note that the destabilization cycle could make these symptoms worse for a period of time.*
2. Start off with a skilled therapist or guide you align with and who is experienced with MDMA. You may later transition to self-guided sessions (with regular check-ins) if you and your clinician collaboratively decide that is sufficiently safe for you. *This model has not been explored in the research and may have a higher risk of difficulties. However, it offers dramatically lower costs and we are anecdotally aware of many people have found it highly effective*

Guide vs therapist. Specific characteristics. Ability to handle transference and counter-transference? overlapping bell curves of quality

3. You have a skilled non-psychedelic-trained therapist or guide you align well with and you collaboratively assess your risk-factors with them (perhaps reading this safety section of this guide together). If your risk factors are low, you then self-guide all your medicine sessions while maintaining regular sessions with your clinician. You also educate yourself on the nuances of effective and safe MDMA therapy and use a high-quality sitter for at least the first few sessions (See Section 5.5 for characteristics of good sitters). *We think this model is similar to the previous one, except slightly more difficult and riskier.*
4. Self-guide all your medicine sessions and do all of your own integration work, perhaps talking about your healing journey with emotionally skilled friends or friends skilled in using safely using psychedelics for healing. You also read literature on trauma healing and psychedelic therapy written by mainstream academics and use a trusted sitter for at least the first few sessions (See Section 5.5 for characteristics of good sitters) *We don't recommend this model because self-assessment of risk factors is very difficult. If you decide to try it anyway, you will greatly benefit from high self-emotional knowledge and high skill in managing your own emotional reactivity. While this model has worked for many people, we are unsure of the risk/benefit tradeoff for anyone other than mental health professionals. We strongly recommend making note of a therapist or guide you might want to work with in case you get into deeper water than you are comfortable with.*
5. Self-guide all your medicine sessions and do all of your own integration work. You don't use any high quality reference material and don't understand the technical nuances of safe and effective healing. Or you work with an inexperienced guide who may harm you by suggesting intense psychedelic sessions before you are ready. They may also offer distorted or unhealthy interpretations of experiences you have during a session. *We don't recommend this in any circumstance. While it is occasionally helpful to some, we think the lack of rigorous understanding of the process will impair healing to some degree. Adverse effects may not be identified, understood, and well-managed*

how to identify high quality peer support. What to avoid. What to look for.

how to identify bad/good?

The conditions below increase the chances of lasting healing and reduce risk of destabilization. While meeting all of them is not necessary for success, the more you have, the better it will go. Consider these carefully when deciding what degree of professional guidance you need:

- Access to resources: Stable housing, financial security, and friends or family skilled in emotional support.
- When working with a therapist or guide: Your empathy and regard for each other, agreement on goals and methods, and therapist skill with facilitating reconsolidation. These factors impact outcomes more than the choice of therapeutic framework [166].
- Understanding of healthy emotional behaviors (awareness of difficult moods and tools to manage them, conflict resolution, etc.).
- Perseverance to navigate challenges in the healing process.

list things that inform choice. Don't list things that influence outcomes for either choice, like motivation.

prep for mdma (meditation); more compassionate phrasing

impulse control,

- Time for and access to a secure and inviting space for your sessions, post-session rest, and reflection.
- Independence from those who have traumatized you, if they might react poorly to your insights and improvements. Most maladaptive schemas protected you at one point in your life. Dismantling these protections while you are still in a dangerous situation could be bad.
- A habit of inquiring why you have the beliefs and behaviors you do. Acknowledgement that many of them are strongly influenced to psychological and social factors, particularly from early-childhood [21].
- Absence of other severe trauma responses such as severe complex PTSD, childhood abuse, attachment disorders, etc.
- Ability to identify dysregulation.

compassionate
curiosity
habit, reframe

Self-assessment of the severity of your problems is useful for this last item. This is difficult since our schemas are largely unconscious, but there are some methods:

- The Attachment Style Test can be used for self-evaluation of attachment disorders, one common type of complex trauma [119].
- GAD-7
- PHQ-9 or 17
- ACE Quiz
- sprint self report PTSD
- cbirt

*We recommend Psychedelic Passage, a referral service for people seeking psychedelic guides [112]. We also recommend the MAPS Psychedelic Integration List, "a resource of individuals and organizations in the mental health field who help people integrate past psychedelic experiences [123]." When working with a therapist or guide, we recommend filling out the Brief Revised Working Alliance Inventory every few months and discussing it with your professional [56, 87]. This can help you assess how this therapeutic relationship is working for you.

what is risk of
trying a ses-
sion casually?

*Altered states of consciousness and self-blaming schemas create opportunities for guides, therapists, and self-described shamans to abuse their clients [58, 42, 29]. Additionally, even typically non-abusive acts, like a hand on a shoulder, may feel deeply unsafe for a client processing a traumatic experience in an altered state of consciousness. Most professionals don't abuse their clients, but we think it happens often enough to warrant taking a few things into account when looking for and working with a professional. We suggest the following: Some professionals in the field are also working on processes of accountability, like the article by Martinez [88].

The beneficial therapeutic effects of MDMA also enhance client vulnerability to abusive practitioners [**patient Vulnerability**].

How to deal with clients feeling utterly dependent on therapist and medicine to carry them through the period of destabilization which they were just thrown into [91]?

New York Magazine has reported on a large group of sexually and psychologically abusive psychedelic guides and therapists trained by Aharon Grossbard and Francoise Bourzat [29]. This group also has links to Dr. Salvador Roquet, and the California Institute of Integral Studies has employed a few of them . Multidisciplinary Association for Psychedelic Studies, the organization funding the MDMA clinical trials, has also employed some of these abusers (who abused at least one trial participant), and in our opinion, has failed to understand the harm it has done and adapt to the needs of participants. This may be related to their deep fear of a renewed war on drugs, a cultural tradition of client abuse in psychotherapy, and a belief that psychedelics will usher in utopia.

A number of professionals in the field have signed an open letter acknowledging this abuse [**abuseLetter**].

Watch out for plastic shaman, cultural appropriation in new age subculture?

Warn against charlatans like Andrew Weil, Deepak Chopra, and other woo-practitioners?

Warn against guides, “shamans”, and therapists who downplay wrong doings or suggest warn against victim blame.

Abusers push clients uncomfortably (sex, touch, literal fighting, yelling, etc.), ostensibly to heal repressed psychological problems. Guides who are confronted with doing inappropriate things will sometimes “consult their inner self” and decide they did nothing wrong. Or they will push boundaries because they “feel called to”.

Solo use may be more appropriate if you feel discomfort with your therapist or guide and you can manage destabilization and don’t have risk factors of psychosis, mania, or suicide.

Link to professional standards for therapists?

noTouch has suggested a policy of absolutely no touching during sessions.

Why is it so easy for abusive guides and therapists to bypass healing their own problems?

If you wish, you can bring a trusted friend or family member to the session with you. It may be helpful to help them understand the process by bringing them with you to the pre-session prep with your practitioner. You could also video-record the session on your phone.

5.3 Dosing

*Accurate dosing is an important factor in efficacy and safety. The effects of MDMA are strongly dependent on MDMA blood-plasma concentration, which is positively correlated with dose and negatively correlated with body mass [154]. Despite effects being dependent on body mass, most data is only available from fixed dose studies. Too low of a dose (0.75 mg/kg, for example) will likely not provide the safety and empathy needed for healing [14]. The positive mismatch and maladaptive schema must be close enough in strength to each other for reconsolidation to occur [38]. High doses can cause the session to be so pleasurable that we’re not able to engage with maladaptive schemas. This may feel good but is not therapeutic. The phase III clinical trials found fixed doses consisting of an 80-120 mg initial dose followed by a half-strength supplemental dose administered 1.5-2 hours

Why are so many into woo spiritual things?

later were generally effective and safe for medically screened participants [97]. Adjusting dosage in subsequent sessions may be necessary to reduce adverse effects, increase feelings of safety, or adjust for individual body chemistry. Plasma concentration increases in response to increasing dose at a faster-than-linear rate because MDMA inhibits the enzymes that metabolize MDMA [34]. Because of this, we suggest increasing or decreasing dose in smaller increments than one would typically expect, especially at the upper end of typical dosage. Accurate data on the upper limit of safe doses is unfortunately absent due to difficulties translating the results of animal testing to humans [113].

When applying scientific studies to one's own life and health, it is important to remember that the data we glean from these studies flattens a wide variety of individual responses by combining them into readable averages. You as an individual may experience something very different from the average participant of any given study, and that may be totally normal and fine. Some examples include: you may be much more or less sensitive to the psychological or physical impacts of MDMA. The medicine may impact you for a greater or lesser amount of time than it impacts the average person. You may experience more healing, faster, than the average study participant, or you may not be helped by MDMA at all. Many normal human variations, like low or high body weight, recent pregnancy, or menstrual cycle status clearly have an impact on many mental health interventions (especially when it comes to effective dosage), but are not typically studied at all.

One of the great frustrations of mental healthcare research is that every real life situation is infinitely complex, and a corresponding infinity of confounding factors have the potential to influence outcomes. We encourage you to discuss with your clinician any difference in what you are experiencing (during any mental health intervention) from what the average response is that you might have expected from the research. It's important to both keep an eye on any health and safety concerns that might be related to your response, while also remembering that the range of normal and healthy responses to any mental health intervention is much broader than the averages suggest.

5.4 Session Difficulties

*Dissociation and resistance are the primary challenges in trauma therapy. While these may have protected us at one point, they also block reconsolidation. We aim to reduce them to a level that enables reconsolidation without causing overwhelm.

5.4.1 *Dissociation

Dissociation often presents as emotional numbness or tiredness. It is a defensive mechanism similar to playing-dead that is associated with trauma in situations of powerlessness [84].

Curious investigation or meditation on the dissociation, boredom, sobriety, blankness, or numbness for minutes to hours will often reduce the dissociation and reveal underlying emotional material (see Appendix A.3) [128]. Increasing feelings of safety by altering the environment, finding a more-attuned therapist, or increasing the dose on the next session may also help. If dissociation persists, another method such as PSIP may be necessary [128].

prime need is safety

Day write. Work together session.

association with addictive patterns?

learn to identify dissociation before session

be strategic with dissociation; may be helpful sometimes

mindfulness

does body scan meditation work here?

Anti-dissociative medications like naltrexone or naloxone could conceivably help though we are not aware of this being tried. Naltrexone requires a prescription and collaboration with a medical professional is necessary. We suggest starting with low doses, as dissociation often covers up overwhelming emotions.

cite

therapeutic
yoga or walking;
mindful movement

5.4.2 Resistance or Distraction

Selective inhibition (suppression of minor movements and comforting behaviors, and attunement to uncomfortable feelings) may help [128].

We think LSD and psilocybin are potent tools for reducing resistance. Mixing LSD or psilocybin with MDMA is widely practiced, though poorly studied [171]. We suggest starting with very low doses because LSD or psilocybin-assisted psychotherapy are known for deeply challenging experiences. Adding up to 0.5 g dried psilocybe mushrooms at the beginning of the MDMA session reliably decreases resistance. It also increases the intensity and discomfort of the session. Do not take more than 0.5 g unless you are deeply familiar with the combination. See Section 6.1 for more information. This combination is common practice in MDMA-therapy, but no studies have been done [28]. Acutely safe co-use is known from recreational contexts [171].

brings triggering material and notes on what you want to address or a mantra about staying present with difficult things

guide/therapist can help with this really well

cite

Increasing the MDMA dose on the next session may help by increasing feelings of safety, which can reduce dissociation.

ask multiple clinicians for their experience.

5.5 Pre-Session Preparation

Proper planning of a session is important for comfort, safety, and success. This is traditionally divided into three sections of mindset, setting, and cultural/personal matrix {citeset-SettingMatrix. We add two more: tools for the session and planning for the post-session:

Mental Preparation and Expectations (Mindset):

some of these items are optional; can skip if too much work?

- Journal about the things, places, practices, people, or situations with whom you feel safe and at home, and with whom you feel unsafe. Then make a diagram with two axes, safe-feeling and accessibility. Plot at least the top 10 items you journaled about on this diagram. Identify the items that are both safe and accessible and keep them in mind for:
 - Trauma therapy sometimes involves an aspect of "it gets worse before it gets better." We suggest planning increased time with the safe items and decreased time with the unsafe items in the days and weeks following your session.
- habit mapping (unwinding anxiety)
- Shame mapping can help you identify many core schemas. See A.1 for details.
- Practice LK, focus meditation
- practice thought record

less important with guide

- Dissociation and resistance are the primary obstacles to healing during a session. Learning about the signs of these can help you recognize and deal with them during the session. Refer to Section ??.
- Intend to compassionately face and heal your distress.
- Avoid rigid expectations of what schemas you will face [100]. The schemas you need to address first may not be the ones you're expecting. Conscious awareness of their true nature is often unreliable because these memories are overwhelming, dissociated, or avoided.
- Catch up on sleep. Therapy requires focus and energy. MDMA promotes focus and energy as well, but we suspect it can only help so much if sleep is lacking.
- Think or write about what challenges or emotional difficulties you would like to address during the session.

is this really important? Maybe jumping into hating yourself during the session would be just as productive.

Cultural Matrix: ~~set~~**SettingMatrix** defines this as the "...environment (1) from which an individual comes, (2) in which the individual lives during the time of the sessions, and (3) to which the individual returns after successful therapy-the everyday living space; these three elements are discussed." Family, living situation, and culture have complex interactions with the process of MDMA therapy. In some cultures, like those where ayahuasca use is endemic in areas of South America, knowledge of psychedelics may be widespread and accepted. Finding help in dealing with certain after-effects may be much easier in those contexts. Other contexts can be hostile or counterproductive to the process of MDMA therapy.

Profound feelings of unity and truth are a common effects of psychedelics [yaden2022varieties]. However, people typically interpret these experiences through the lens of their existing belief systems, belief systems they are aware of, and even symbols present in the session environment. These may include religion, metaphysical theories, or cultural tropes such as "psychedelics show the truth behind all things." While some beliefs about unity experiences may be true, we suggest keeping in mind that people interpret these experiences in a wide variety of different and, we suspect, contradictory ways. Psychedelics are also known to inflate feelings of meaning and not all insight is an accurate map of reality [tulverInsight, 61]. We suggest deliberation before deciding whether your own experience either confirms or disconfirms any particular belief. We also suggest reading about the [Hard Problem of Consciousness](#) [hardProblem]. The linked article at the Internet Encyclopedia of Philosophy outlines a variety of theories of consciousness and the problems with each. ***Setting:**

More to be said: Suggest support groups?

- Sessions are usually started in the morning. MDMA's stimulant effect may prevent sleep if the session is started later [16].
- Eating is avoided if possible on the morning of the session because some people experience nausea [100]. Taking medicine with food can also delay onset of effect.
- Arranging easy access to necessities like a restroom, snacks, and water will reduce distractions during the session. If your environment is hot enough to induce sweating (sweating may further increase due to the effects of MDMA) during the session, mixing salt with your water will reduce the risk of hyponatremia.

- Prepare your environment or travel somewhere to maximize your feeling of peace, at-home-ness, comfort, and safety. These will add to the feeling of safety from the medicine and ease down dissociation and resistance and increase the emotion mismatch. We suggest carefully expanding your session environment can be beneficial once you are experienced with the medicine. For example, if you have trauma from dogs, you could safely activate that schema by petting a dog who you know won't act aggressively, and who's human is nearby to calm them down if you feel like the experience is too much to handle.
- Remove distractions and danger, ensuring inward focus.
- Solitude, except for a trusted and experienced guide, therapist, or sitter, promotes inward-focus and healing. A sitter can help with logistics, hold your hand (Discuss this first. Explicit, pre-arranged boundaries are essential in altered states of consciousness and during trauma therapy.), listen to your feelings, or handle mundane events like someone knocking at the door [100, 156]. Sitters should possess trustworthiness, presence (ability to listen to your distress without becoming distressed themselves), and empathy. Interacting with strangers may cause problems if they do not understand what is happening.

*Tools:

- Consider self-narrating your session and recording it on your phone [100]. Listening to this after the session can spur insight.
- We suggest bringing materials that activates challenging schemas, like photographs or letters, to look at if distressing schemas don't spontaneously arise. See Section 5.6 for more discussion on differentiating distractions.
- Eye shades and noise-cancelling headphones can reduce distractions.
- MDMA can cause jaw clenching and headaches [97, 81]. Some individuals use mouthguards or pacifiers to reduce this effect or protect their teeth [41]. Over-the-counter pain relievers may help after the session. Those with jaw problems may need to be more cautious or use lower doses.
- The Fireside Project offers a hot-line to help people through challenging psychedelic experiences at (623) 473-7433 in the United States [118]. Keeping this number in your phone may serve as an additional layer of safety.

*Planning for After the Session:

- We have noticed people often feel fatigued for 1-2 days after the session. Prepping food and a comfortable place to rest in advance may be nice.
- We suggest making a plan to try different techniques on subsequent sessions if you're worried that this first MDMA session might not work and is your only hope for healing.
- Therapy sessions are often scheduled for the day after an MDMA session [100].

can sitters help with dissociation and avoidance monitoring? include instructions if so

ladder of bids

info specific to tmj?

why? modes?

- If you have the energy, make a plan to spend time understanding your experience in the following days and weeks [100]. Common ways to do this includes journaling, nature walks, body movement practices, transcribing a recording of your session, and art. Planning for extra reconsolidation in the following two weeks is a
- High quality sleep of sufficient duration is likely important to solidifying your reconsolidated memories [148]. We suggest a highly effective evidence-based protocol called *Cognitive Behavioral Therapy for Insomnia* if you struggle getting quality sleep. We recommend the Stanford Medicine guide [93].

5.6 The MDMA Therapy Session

*The effects of a single MDMA dose are generally noticeable 30 min after taking the medicine, peak an hour after than, then last a further 3 hours [163]. We warn you not to take more MDMA during a session than you initially planned to. Desires to do this are often based in anxiety or dissociation and can result in increased adverse effects [22]. Stay quietly present with those states rather than adding more medicine during a session. Food delays onset of effects [99]. We divide the session into phases based on subjective effects and therapeutic potential:

Come-up: The effects of MDMA become noticeable, but you are not yet engaging with the memories you want to work on. Some users experience anxiety [63]. We speculate this might be from the body interpreting stimulation as anxiety, early engagement with distressing memories, or fears about the session. We suggest practicing loving-kindness meditation for yourself and your dissociation or avoidance (see Appendix A.3). The feelings of compassion could ease down dissociation and resistance.

Peak: Reconsolidation is possible here. Connection and safety become pronounced, though you may not notice this if you directly transition to engagement with distressing schemas. A booster dose taken at hour 1.5 extends this phase by 1.5 hours.

You may feel what is called resistance. This can be a few things: simple avoidance of emotional pain because it is uncomfortable; beliefs that say confronting/healing emotions is bad; or beliefs of an activated schema that say their protective function is very important and you shouldn't let that go. While the protective function of these schemas was important at one point, it has become an impediment. If you can do it, we suggest focusing on any resistance schemas first. Reconsolidating this will make the rest of your therapeutic efforts much easier, as you won't have to fight against a part of mind telling you to stop what you're doing. If you can't focus on the resistance schema and another schema is taking precedence, that's ok. You can revisit the resistance later.

All negative feelings and tension in your body are valuable signals. Feel them fully and see where they lead. We suggest viewing the truth of the schema's beliefs or emotional responses with agnosticism, "It may or may not be true or helpful; I will stay present with it to learn why the schema exists and how it influenced me."

This enables reconsolidation. You can think of it as focus-meditation where you hold the emotion or belief of the schema in focus and dismiss distractions. This can be uncomfortable, but the MDMA almost always makes it bearable. Insightful thoughts about how trauma impacted you or insights into human behavior may arise. Spend time on them if they feel

Less pressure?

Discuss somatic release. People occasionally find it questionably useful. Does it have to be combined with reconsolidation?

cathartic or important. Leave them until after the session if they don't feel cathartic or important, reserving limited session time for reconsolidation or other cathartic processes.

Your second task is monitoring for dissociation and resistance. These can be hard to notice. We suggest assuming they are blocking a healing mismatch any time you are not feeling discomfort, anxiety, fear, anger, resentment, sadness, etc., reevaluating difficult experiences from a place of compassion, or experiencing an important moment of safety and love. Sleepiness and intellectualization are also signs. Refer to Section 5.4 for more information.

hold schema with compassionate curiosity and agnosticism of its truth

Experiencing profound peace and self-love can be deeply healing. If and shouldn't be avoided if it is providing an important disconfirming experience, though at some point engaging with distressing schemas is necessary to continue healing. This view of the end goal can be a great motivator for staying on the healing journey long-term through challenges. It also provides a powerful memory you can later use as a mismatch to reconsolidate many schemas when not using medicine. The peace and compassion may not last long after this first session, but you will gradually get it some of it back as you reconsolidate various maladaptive schemas over multiple sessions.

*MDMA rarely produces uncontrolled or unbearable experiences [97, 98]. Staying present with feelings of panic or overwhelm or just hanging-on almost always resolves the distress [100]. Trying to avoid distress through distraction will likely just prolong it by impairing reconsolidation. We suggest calling the Fireside Project hotline at +1 (623) 473-7433 if you feel like you need help. [118].

why is it important to focus on activated schemas? what am I warning against? how do I know that thing is bad?

*We recommend continuing the session as long as it feels productive because session time is valuable. Occasionally, resistance may prompt you to end the session earlier than necessary. Because of this, if the trip feels over, we suggest sitting quietly for another 30 minutes to investigate possible resistance.

Who to call if you think you might be suicidal, manic, or psychotic? 911? Suicide hotline?

*We suggest not calling or texting anyone until you have more experience with MDMA and you know they will be comfortable talking to you in your altered state.

If you know you are near the end of the session, we suggest it may be wise to not dive into any additional distressing schemas, though finishing up reconsolidating your current ones is good. Surfacing previously unconscious schemas without adequate time to reconsolidate them can result in that schema dominating your post-session life if it strong enough. On the other hand, diving into additional schemas may be beneficial if you can tolerate the distress and want help making your unconscious schemas explicit for post-session reconsolidation.

***Come-down:** Fatigue and reduced capacity to process emotions is typical after intense therapy. The effects of the medicine are still somewhat present, but reconsolidation may no longer be achievable.

***After-effects:** Trace effects may be felt for longer. The concentration of MDMA in the body decreases by half every 8.5 hours citetorrePharmacology.

5.7 *Troubleshooting the Session

need better descriptions of dissociation and avoidance feelings

5.7.1 Sleepiness, disconnection, or heaviness

MDMA is a strong stimulant and generally increases feelings of energy. Activating (perhaps unconsciously) schemas of powerlessness during therapy commonly increases dissociation,

which often includes sleepiness [76]. We suggest dissociation is the most likely cause for sleepiness during a session, through atypical responses to stimulants or sleep disorders could conceivably play a role. See Section 5.4 for suggestions on working with dissociation. We suggest following the dissociation guidelines first even if you suspect an atypical stimulant/neurotransmitter response because these conditions may be difficult to distinguish.

5.7.2 Not feeling anything or feeling "meh"

- Your expectation for the session may not match up with the maladaptive schemas actually present and in need of reconsolidation. This can present as disappointment or anxiety about the trip itself. This disappointment or anxiety is the key feeling to focus on and work with.
- We suggest you may be inadvertently avoiding an unnoticed feeling (avoidance, resistance, or dissociation). Practicing different types of meditation (see Appendix A.4) for 30 minutes can often help you notice and engage with subtle feelings.
- Long-term use of reuptake-inhibitor antidepressants blunts the effects of MDMA [47]. This effect may persist for multiple months after discontinuing these medicines.
- The dose may have been too low [14]. See Section 5.3.
- The substance might have been something other than MDMA [138]. Testing can prevent this problem.
- Some medicines work very differently with some individuals' body chemistries.

5.7.3 Feeling really good but no therapeutic effect

Reconsolidation requires the positive mismatch and maladaptive schema to be close enough to each other in strength [38]. The MDMA dose may have been too high for you. Alternatively, looking at photos or messages that remind you of the schema you want to work on could increase the strength of the maladaptive schema.

5.7.4 New adverse symptom after a session

Trauma therapy often follows a pattern of feeling worse before it feels better [74]. As some schemas are healed and dissociation is reduced, you will notice previously hidden distressing schemas. This is uncomfortable and distressing emotional memories can have a wide variety of mental and physical symptoms [77, 62]. Continuing with therapy will reduce the backlog of distressing schemas. Sometimes you need to develop symptom management skills or resources to manage the immediate problem, as detailed in Section 4.9. If you seek medical assessment, it is beneficial to find a medical professional who understands the physical and psychological symptoms of trauma . It would be ideal to find a medical professional who will anticipate and accommodate your trauma-related safety needs, or at least be willing to listen and adapt. Unfortunately, the United States currently has a severely over-stressed medical system where medical professionals are frequently both undertrained on these topics

[107]

and lacking the skills to effectively implement the understanding they do have. Additionally, many under-supported healthcare professionals are arguably undergoing extended workplace trauma themselves, which does not necessarily lend itself to optimal participant care. As such, we recommend taking a posture of firmly compassionate self-advocacy wherever possible. You deserve safe and trauma informed care; whether you receive this care or not is not a reflection of what you deserve. You may wish to take notes about the symptoms you are experiencing or collect data (like the safety section of this document) about the safety of MDMA to bring with you to your medical appointment; these measures can help you support your healthcare professionals in giving you the best quality of care possible.

link to defense cascade and cheetah house

For a strong basic education on psychogenic physical illness, with an emphasis on the understanding most needed from a participant's perspective, we recommend <https://www.youtube.com/watch>

link

5.7.5 Feeling fatigued for 1-3 days after a session

Post-reconsolidation exhaustion is common and temporary and sometimes called "therapy hangover." Post MDMA-use exhaustion is also common and temporary (across different MDMA-use contexts); the strong stimulant effects of the medicine may also be tiring. No research we know of exists that attempts to untangle how much of this effect is psychogenic and how much is purely pharmaceutical.

look at research

Regardless, we have observed that this usually lasts 1-2 days following MDMA therapy. If low mood or exhaustion extends beyond 3 days, we suggest this may be a symptom of a schema you uncovered during the session.

see the relevant bullet in the preparation section about preparing food and other self-care, emotional support, and executive function support do safety exercise before session and plan to spend some time with people who help you feel safe during this period who can also prompt you to do nice things like going for a walk

5.7.6 Warning signs of psychosis and mania

Psychedelic-induced psychosis and mania are rare unless you have a personal history, and to a lesser degree, a family history. However, if you or someone you know is showing these signs, we suggest helping them to emergency psychiatric care:

5.8 After the Session

*Physical and emotional exhaustion is common and healthy after therapy and reconsolidation. Because MDMA-therapy is so intense, we suggest planning for 1-2 more days of exhaustion post-session. We speculate this may be reconsolidation-fatigue or neurotransmitter depletion [128]. We suggest plenty of rest and comforting yourself in ways good for you. Ensure your schedule is clear of major commitments or stressors in the days following.

list of good comforting techniques

Healing is a complexly layered process. As one layer of emotional distress is calmed, other schemas usually activate and/or become more visible. Though initially challenging, consistent effort will lead to long-term improvement [74]. This cycle is noticeable both in the long-term (months-years) and short-term (the week after a session may feel more volatile than normal). We suggest not judging a session until 1-2 weeks have passed.

*You will experience more time in the mind's relaxed state when maladaptive schemas/parts heal. The Internal Family Systems framework proposes this relaxed state consists of calmness, clarity, curiosity, compassion, confidence, courage, creativity, connectedness, perspec-

tive, joy, patience, kindness, equanimity, playfulness, and love [144]. Periods of dysfunction will remain, but will become less frequent. If the maladaptive schemas are severe, it might take quite a lot of healing to get to the point where anything feels ok. We suggest staying on top of reconsolidation and defense cascade deactivation, and avoiding situations that activate challenging schemas will help.

hyperlink to
section man-
agement

5.8.1 Making Sense of the Experience

When applying scientific studies to one's own life and health, it is important to remember that the data we glean from these studies flattens a wide variety of individual responses by combining them into readable averages. You as an individual may experience something very different from the average participant of any given study, and that may be totally normal and fine. Some examples include: you may be much more or less sensitive to the psychological or physical impacts of MDMA. The medicine may impact you for a greater or lesser amount of time than it impacts the average person. You may experience more healing, faster, than the average study participant, or you may not be helped by MDMA at all. Many normal human variations, like low or high body weight, recent pregnancy, or menstrual cycle status clearly have an impact on many mental health interventions (especially when it comes to effective dosage), but are not typically studied at all.

One of the great frustrations of mental healthcare research is that every real life situation is infinitely complex, and a corresponding infinity of confounding factors have the potential to influence outcomes. We encourage you to discuss with your clinician any difference in what you are experiencing (during any mental health intervention) from what the average response is that you might have expected from the research. It's important to both keep an eye on any health and safety concerns that might be related to your response, while also remembering that the range of normal and healthy responses to any mental health intervention is much broader than the averages suggest.

The intense emotions and beliefs you felt during the session may have been the schemas that have been negatively affecting your life. You will react and think differently now after the emotions and beliefs of your schema have updated. Pay attention to what thoughts and reactions are different from before the session, and you will learn you how the trauma shaped your feelings and behaviors in the past. Common ways to identify these differences includes journaling, nature walks, making art, body movement practices, listening to recordings of your sessions if you self-narrate during the session, meditation, and talking through your experiences with an emotionally skilled friend or clinician [100].

When a schema is healed you should notice that although the pain and maladaptive beliefs and behavior are diminished, a level of caution remains. If this caution is helpful rather than overwhelming, then this is the schema in its adaptive, rather than maladaptive form. This caution will aid you in avoiding future harm.

You may experience feelings of intense and spontaneous compassion, connectedness, or love. We speculate these are flashbacks to the MDMA session and will gradually dissipate as the memory is reconsolidated. These increases in compassion almost always have good consequences for you and those around you.

You might have had profound feelings or thoughts about various belief systems, socio-cultural systems, organizations, etc. MDMA can spur profound insights. The feelings of

trust, safety, and increased perspective inherent to many MDMA experiences can cause us to take the profound thoughts and feelings we have during those experiences as objective truth, which they may or may not be [61]. Psychedelics inflate feelings of meaning. Rather, MDMA can offer a new perspective based on a state of heightened, but incomplete compassion and safety. Even in this exceptional state, all your still-unconscious emotions are deeply influencing your beliefs and feelings. Your more reliable tools of discernment, compassion and cognitive and emotional flexibility, will gradually improve as you continue reconsolidating schemas. Using that increasing flexibility and compassion to rigorously and critically engage with a diverse and challenging set of reliable information and experiences will more reliably lead you to what is objectively accurate or helpful for your life than profound session insights [15]. Seek high quality critiques of your beliefs and incorporate beliefs into rigorous higher level frameworks [137]. Correct for cognitive biases and aim for a balanced perspective when evaluating new ideas [51].

PTSD makes us feel as though we are by our nature isolated; MDMA can help people feel as though they are by their nature connected to one another. This may be one of the central mechanisms of healing that both therapy and MDMA provide.

Spiritual bypass is a "tendency to use spiritual ideas and practices to sidestep or avoid facing unresolved emotional issues, psychological wounds, and unfinished developmental tasks". . This results in maladaptive patterns and interrupted psychological development . The term was introduced in the mid 1980s by John Welwood, a Buddhist teacher and psychotherapist. Because MDMA can produce profound spiritual experiences for those individuals who interpret their MDMA experience in a spiritual way, it may be important to take care to avoid spiritual bypass if MDMA experiences are to actually facilitate healing. In particular, the profundity of many MDMA experiences may lead one to feel that focusing on painful maladaptive schemas is simply no longer necessary; however, without that focus, the schemas will not be reconsolidated, and healing will not be accomplished to the degree it otherwise could be.

Sometimes individuals may feel they are receiving a spiritual message from the medicine, stating that their healing is complete and further medicine work is not appropriate for them. Alternately, the message may take an alternate form, like, "the medicine has done all it can do for me, and now I need to focus on some specific practical aspect of my life." Although practical life circumstances can have an enormous impact on mental health, and we wouldn't want to discourage anyone from fighting for a healthier life situation, we do caution individuals and clinicians who encounter this scenario to be mindful of the possibility that these messages may also be forms of resistance. To understand what is insight and what is resistance, it may be helpful to examine what symptoms and side effects remain active in an individual's life. It may also be helpful to spend time with the negative feelings about further medicine work in the context of reconsolidation exercises, whether assisted by medicine or otherwise. For further discussion of how to decide when to stop using medicine, see

Healing experiences are often, to one degree or another, stressful. Although stress is often regarded as harmful in modern US culture, it is crucial to both meaning and personal growth. You only experience stress when things matter to you, and a stress-free life is a life of apathy and meaninglessness. It is stressful, but also helpful and important, to sit with the discomfort of a maladaptive schema or a mismatch. Even our original traumas could not have harmed us if we did not value the things we value—trauma is intrinsically connected

wikipedia citation

cite

elsewhere in text.

to meaning. As you progress, try to differentiate between adaptive stress, which may alert you to immediate dangers or motivate beneficial actions, and maladaptive stress—which is more than you are able to process or respond to in a healthy way. A good clinician or guide can help you gain insight into which stress is helpful for you and which stress is more likely to harm. A good clinician or guide who is up-to-date on the research on this topic can also help you re-frame stress in a way that greatly reduces its negative impacts on you, even on a physiological level. For a comprehensive discussion of the dual nature of stress and the importance of adaptive framings of stress, refer to *The Upside of Stress* [89].

You may also have remembered events you weren't aware you experienced. The complexities of recovered memories are strongly debated within psychology, but there are some general guidelines: While the emotional tone of a recovered memory is a reliable indicator of your subjective experience, episodic memory is flexible and amenable to change over time [130]. The episodic gist (the overarching theme of a remembered event) is also significantly more reliable than episodic details. Trauma can further impair episodic memory formation, sometimes leading to missing or fragmented memories of traumatic events [74]. Time passed since the event and the number of times you have recalled the memory also decrease reliability [78]. Use extra caution with psychedelics, as they can increase mental imagery and feelings of significance [57, 31]. MDMA may also slightly change recollection of episodic details of emotionally charged memories [37]. Seek independent corroboration of recovered memories if possible. Due to the power of suggestion, therapists and guides should avoid leading questions, suggestive techniques, and undue influence that might inadvertently plant or reinforce unfounded memories of childhood abuse.

Relatedly, you may recall feelings or beliefs not clearly related to a specific event. These often come from early childhood, when emotional memories form before you are able to form episodic memories [21].

In Western culture we place a premium on having a continuous and linear narrative understanding of ourselves and our place in the world. This value set can serve to further disempower survivors of trauma, especially childhood abuse, because the nature of memory and the shame and taboo surrounding abuse experiences often preclude clear and objective understandings of what exactly happened in our childhoods. Additionally, the US (and possibly other places around the world) embodies a schism in cultural values regarding what is abusive and how abuse should be dealt with. In cases where abuse is extensively documented, survivors nonetheless may face aggressive gaslighting and extended abuse in response to speaking up about their initial experiences. Even documentary certainty about the actual events that happened is not always enough to endow survivors with the “good victim” status they are often seeking when they focus on the accuracy or inaccuracy of recovered memories.

In the face of these realities, you have to think: given that I can never know for sure what actually happened to me, then what? And what, as it turns out, is that you still deserve compassion and healing. “What” is also: that it can be reassuring to gather as much objective evidence as you can about your own history. This can be a way of reclaiming what you can of your own broken narrative, of knowing what is possible to know and at least laying hold of it. And you may need to grieve a great deal, because often the evidence is very thin, and a knowledge of your own history is one of the things that was stolen from you. But your deservingness of healing does not depend on someone coming along and saying,

other relevant
descriptors

“I have a video of you suffering a legitimate kind of suffering, so your pain is valid now.” It may be helpful to explore the following questions, either by journaling on your own or by talking them through with a trusted/emotionally skilled friend and/or a clinician: what does it mean to me if X actually happened? What does it mean if Y happened instead, or Z? What does that say about my identity, my needs, how I perceive myself, and how others perceive me? What does it say about my future? What does it say about what I deserve?

Please be compassionate with yourself, as much as you possibly can. Whether your suffering is real and matters is a different question from, “what is the best strategy for addressing my suffering.” One last word is, your pain is trustworthy, even if your memories are not. If you feel like you are suffering, that is real, and it matters.

We also recommend The MAPS Psychedelic Integration Workbook [124].

do we?

5.8.2 Continued Reconsolidation

Gains in functionality and emotional health often depend on reconsolidation of the underlying maladaptive schemas. Reconsolidation produces significant, worthwhile, and continuous improvements in particular schemas (and their associated symptoms). We estimate accessing the maximum possible benefits from reconsolidation may take anywhere from single digit hours of mismatching for simple problems to thousands of hours for the most severe mental illness. These estimates do not include the overhead of uncovering schemas and finding mismatches, which can vary. Because there may be risks from long-term high-frequency MDMA use, and because sessions definitely have logistical costs, we recommend using between-session reconsolidation to maximize access to the benefits of reconsolidation. Thus, we present techniques for between-session reconsolidation in this section.

We speculate that there is a daily mismatch limit (one data point suggests 2 hours). At that point, mental exhaustion inhibits the practice. Sleep rejuvenates the ability.

You will likely notice activating remnants of the schemas you reconsolidated during the session. These remnants are easier to deal with than the entire network, perhaps because they are smaller or because you now better understand how that particular emotion affected you. Due to the healing that happened on the session, a variety of mechanisms will also surface different maladaptive schemas.

It is often difficult to access activating schemas. We have noticed that walking, other body-movement practices, selective inhibition (see Section ??) or a 2.5-5 μg LSD supplement (see Section 6.1) can help.

We suggest attempting to view the truth of the schema’s beliefs or emotional reactions with agnosticism, “It may or may not be true or helpful; I will stay present with it to learn why the schema exists and how it influenced me.” This may be difficult for intense schemas, but practice may make it easier.

Structured reconsolidation:

- ifs
- Coherence Therapy
- notecard thing

- Challenging Beliefs worksheet (see Section 6.2)
- psycholytic therapy (see Section 6.1)
- Attunement
- Motivational interviewing
- Steelmaning
- Internal double crux?

In-the-moment reconsolidation:

We suggest that powerful experiences of profound safety or okness along the lines of "I will be whole and ok deep down no matter what happens", from MDMA, mystical, spiritual, or meditation experiences, are of exceptional therapeutic benefit between sessions. When you notice avoidance, anxiety, reactivity, distress, or other signs of maladaptive schemas in daily life, recalling and staying present with your feeling of profound safety can mismatch and reconsolidate your maladaptive schema in the moment. We have heard multiple anecdotes that practicing this technique makes it easier and more automatic. In two cases, hundreds of hours of practice led to mismatch and reconsolidation happening without conscious intent any time a maladaptive schema is activated and the individual is not actively avoiding it. This ability may reduce or eliminate the need for MDMA-sessions when well-developed. We think practicing this ability is important for a few reasons: it enables a higher volume of reconsolidation, helping you reach therapeutic goals sooner; long-term MDMA use could have poorly understood side effects; MDMA-sessions require planning and recovery; and reconsolidation is valuable in many contexts where MDMA use is not appropriate.

We also propose that powerful experiences of total safety during a session can later act as a compass you can compare all other experiences to, to deduce if a certain feeling is maladaptive or not. As with real compasses it only measures the direction in which to travel; it can't tell you how far away the destination (reaching particular therapeutic goals) is.

If you haven't had a powerful experience of profound, universal safety, context-specific mismatches will have to be found for each maladaptive schema. It may take some effort to find memories of contradictory experiences, but when found, that memory can also be recalled for in-the-moment reconsolidation of specific schemas.

can we cite
bruce ecker
personal com-
munication?

can LK medi-
tation be used
to do this?

5.8.3 Using Neuroplasticity to Make Positive Life Changes

MDMA induces a period (2-4 weeks in mice) of heightened neuroplasticity (your brain's ability to adapt) [101]. We posit that the neuroplasticity window after an MDMA treatment may offer an extra boost of support in making changes you seek in your life. This period is likely context dependent. That is, a therapeutic session may lead to therapy-relevant neuroplasticity, but a recreational session won't. Some people may find that as they reconsolidate schemas and have positive disconfirming experiences, they have less need for compulsive self-soothing over time. They benefit from the therapy not just through reconsolidation,

but through the support that the subsequent peace and perspective provide for positive behavioral changes. Sometimes being in a rut is the function of inertia; sometimes it is self protective, and improved regulation or a sense of new experience or perspective can help a person break out of it.

How MDMA impacts behavioral change goal selection

We advise refraining from drastic life-shaping decisions immediately post-session. MDMA insights may sharply highlight certain aspects of a situation which you had not been facing or focusing on before the therapy. Although these insights may be extremely important, we feel they can best be understood in the broader context of your life both before and after the MDMA session and the immediate post-session period.

We recommend avoiding major life changes during the immediate post-session period unless (1) you have wanted to make those changes for a significant time prior to your MDMA experience, (2) you are (and have been) very sure and very committed about the change you want to make, and (3) you have access to other forms of support around that change, for instance, regular social connection with individuals who enact the behavior you are interested in adopting, an accountability buddy, etc. (See below for further discussion of structural supports.) When these three conditions are met, we are hopeful that MDMA neuroplasticity could provide a helpful layer of support for larger behavioral and lifestyle shifts.

On the other hand, MDMA therapy may more routinely facilitate a “baby steps” approach where individuals make small changes as they feel capable of making them, with the neuroplasticity window providing an opportune time to build positive habits and process emotional memories.

Barriers to behavioral change + resources to support selected changes


Most people have some habitual behaviors they have wished to change for a long time. In this section, we will review common reasons people struggle to make behavior changes they want, and offer an overview of strategies for overcoming these barriers. Additionally, we have included a fairly comprehensive appendix of evidence based recommendations for behavior changes that are likely to have a positive effect for most people who implement them. It includes at least one recommended resource about each of those changes. We hope this appendix will assist readers in identifying the behavioral changes with the lowest cost and the highest reward possible, including changes they may not have previously considered as possibilities. Additionally, we hope the information throughout this section will support our readers in using their neuroplasticity windows as effectively as possible, and that behavioral change efforts in the aftermath of therapeutic MDMA use can amplify the benefits of these interventions.

Many people feel puzzled or frustrated by their inability to enact important behavioral changes that they know would improve their quality of life (like following through on daily exercises prescribed by a physical therapist), or which are strongly aligned with their values (like ceasing porn use to adhere to religious values, or adopting a vegan diet to reduce carbon footprint). In modern US culture, individuals often judge and blame themselves for

“not having enough willpower” to successfully make these changes. For some, this can be a perpetual source of shame: because of their failure to enact desired behavioral changes, they begin to attribute negative characteristics to their identity, to consider themselves to be apathetic, stupid, lazy, a bad friend/partner/parent/student/employee, or whatever other negative identity might apply to the situation.

However, research shows that the individuals who are most successful at making these kinds of changes are also (1) likely to frame their shortcomings in terms of behavior (creating guilt) rather than framing them in terms of identity (creating shame), and (2) likely to engage a greater level of structural support for their desired behavior change. Finally, most behavioral health goals have a strong emotional regulation component, even beyond the intensity of a shame response.

Structural support can broadly be categorized as “trigger supports” and “access supports.” Trigger supports relate to what initiates a particular behavior. “Tiny Habits” gives the example of a woman who wanted to get up early and exercise before work, but found herself using the time after her alarm went off in the morning scrolling through her phone instead. After considering several possibilities, she started using an alarm clock and leaving her phone in the kitchen overnight. This removed the trigger for the behavior she was trying to avoid. Other examples of trigger support would include a parent reminding their child to do chores, or someone who wants to spend some time reading every day leaving their book on the couch cushion where they typically sit upon coming home from work. Access supports are things that reduce the level of effort it takes to engage in a desired behavior. A home washer and dryer unit would be an access support for someone with the goal of keeping up with their laundry needs; a private chef, or a week of “meal prep” in the fridge, would be an access support for someone who is trying to adjust their eating habits. A “dumb phone” and a social working environment with co-workers who feel safe and are supportive of the goal would be access supports for someone whose goal is not to scroll through the internet while they are at work. Creative and extensive use of structural supports can make all the difference in behavioral change success.

Finally, the emotional regulation component of behavioral change is frequently underrated. People often attribute their inability to enact a desired change to motivation or willpower problems, and then beat themselves up for these perceived failures. The more constructive approach is to implement applicable trigger supports and access supports, and then, if motivation still seems to be a problem, to engage with motivation enhancing strategies such as setting up a “fishbowl,” or using the other social accountability strategies detailed in Appendix 

Procrastination is most centrally an emotional regulation problem; it emerges from feelings of overwhelm, often followed by shame about the procrastination itself. The single most effective strategy to address procrastination, in my clinical experience, is to practice breaking tasks down into manageable parts. If the procrastinated task is getting out of bed and taking a shower while severely depressed, this might look like calling a friend, then talking through and enacting the process of throwing the covers back, putting one foot on the floor, putting the other foot on the floor, walking towards the bathroom, opening the door, and so on. If the task is an hour long marketing presentation, the first step in ending procrastination might be talking through with a mentor how to break the project down into the smallest parts possible.

Self-destructive behaviors generally are associated with feelings of shame. We recommend Brené Brown's work on shame resilience [20].

Behavioral change goal selection in the age of addiction

In the words of Anna Lembke: "We've transformed the world from a place of scarcity into a place of overwhelming abundance: drugs, food, news, gambling, shopping, gaming, texting, sexting, Facebooking, Instagramming, YouTubing, tweeting... the increased numbers, variety, and potency of highly rewarding stimuli today is staggering. The smartphone is the modern-day hypodermic needle, delivering digital dopamine 24/7 for a wired generation. If you haven't met your drug of choice yet, it's coming soon to a website near you. Scientists rely on dopamine as a kind of universal currency for measuring the addictive potential of any experience. The more dopamine in the brain's reward pathway, the more addictive the experience." -Dopamine nation, Chapter 1

Addiction is a complex and overwhelming phenomena, with biological, psychological, social, and structural components. It exists on a spectrum of compulsive behaviors. A person's desire to change their relationship with a given substance or behavior can vary widely: on the one end, someone may want to stop eating fast food on twice-yearly road trips when they already never eat it at any other time. On the other end, someone may have a desperate need to retrieve their life from the encompassing grasp of methamphetamines or heroin. Although it has become socially acceptable to joke or vent about our less destructive addictive tendencies, (like a coffee addiction or a social media habit that isn't ruining our employment and hygiene) our current societal, economic, and technological structures generally provide poor support for individuals struggling with addictive tendencies at all levels. Individuals use substances and behaviors that have negative consequences because those substances and behaviors solve some kind of problem for the individual who uses them. The most effective way to assist individuals who are struggling with these kinds of harmful coping patterns is typically multi-pronged. It might include any or all of the following:

- medication assistance to reduce cravings for a harmful substance
- opportunities to contribute to and connect with their communities in meaningful ways while sober
- peer support
- psychotherapy
- case management (helping the individual access a variety of needed contexts and resources)
- contingency management (creating accelerated and enhanced positive and negative consequences supporting the desired behavior changes)
- other executive function support, such as tailoring the living environment to support the preferred coping choices and building healthy habits like sleep and meditation, which support impulse control
- intensive medical and psychiatric management of any underlying issues that have induced them to use the harmful coping mechanisms.

The average individual who struggles with addictive behaviors, far from being able to bombard the problem from all sides as I've described above, is shamed for not being able to simply choose to stop using the harmful coping mechanism. This typically increases use of the harmful behavior or substance. In short, we live in an age where addiction has been aggressively stigmatized/shamed and resources for treating it denied, while resources and structural support for maintaining addiction have been amplified. "Crossover addiction," where individuals who struggle with one addictive coping practice simply shift their dependence onto another addictive coping practice, is pervasive. Although we would all like to

see ourselves as independent decision makers, we increasingly live in an attention economy where massive investment has been made in competing for each nanosecond of our attention. It is increasingly difficult to capture those moments of our own attention back, and to focus them on our safety, flourishing, freedom, and connection with others.

To make matters worse, despite emerging from the “overwhelming abundance” Anna Lembke describes above, addiction creates scarcity. It creates a scarcity mindset regarding access to the coping mechanism, whether that’s time for gaming or money for heroin, and in its worse iterations, it inspires compulsive behaviors that are financially, socially, and medically destructive in the extreme. Just as important for the problem of escaping addiction, it perpetuates scarcity around meaning and human connection. Addictively patterned coping mechanisms numb feelings all around, impeding our ability to be vulnerable and connect deeply with one another and with the things we find important in life. In rats, non-heroin-using individuals were found to help a fellow rat escape from a trap, while heroin using rats didn’t bother.

Second, despite the dystopian background, strategic habit development can (with commitment over time) provide access to cornerstone strategies on which to build a better life. To select a behavioral change goal that’s right for you, consider your own motivation, your life experience, the degree to which the behavior change offers benefits across a range of life domains. For instance, regular exercise or meditation can help many individuals with pain management, depressive symptoms, and physical health. To find the right foundational habits for you, you’ll need to consider your individual supports, accessibility challenges, and barriers. We make some general suggestions that are beneficial for a majority of people in Appendix A.6.

A harm reduction framework is both evidence-based and essential when addressing addictive coping patterns. I once read (maybe in the book “tiny habits?”) that the most important characteristic for personal growth is the ability to experience satisfaction from small victories. Abstinence is an excellent goal for many people in many situations, but if you are not interested in abstinence or if abstinence is not realistic for you right now (or if your addictive pattern is around food or screen time, which are not realistic to totally abandon), your welfare may depend on your ability to give care to yourself by identifying the harms your coping mechanism creates, and mitigating those harms in more realistic ways. There are incremental victories to be had that make a significant difference in the level of harm an addictive coping mechanism creates in a person’s life. You may reduce the amount you use, or regulate the timing (only after work), or refrain from mixing multiple substances or from gambling in the first three days after your paycheck comes in. If you are not ready to stop drinking, you may focus on making sure that you get all your driving and work accomplished before you start drinking each day, and that you stay hydrated and eat some healthy food on the regular. Maybe you want to start meditating for three minutes each day before you start drinking. These measures are meaningful. They reduce harm, improve your quality of life, and will leave you in better shape if the day comes when you are ready to make bigger changes. It’s important to adopt a mental framework (and cultivate a support system) that sees the value of small improvements and supports you in making the changes you are interested in and ready to make.

5.8.4 *Afterglow

Some people experience a temporary afterglow (wellbeing, mindfulness, less mental illness) for days-weeks after some psychedelic-therapy sessions [44]. We caution against chasing after it with more frequent MDMA doses. Afterglow is often unreliable, while reconsolidation provides durable, long-term healing that does not require continued medication.

5.9 Long-Term Trends of Therapy

5.10 Managing Anxiety and Panic

There are multiple practical objectives in healing. Keeping these balanced determines the pace and methods of healing:

- Healing as soon as possible to minimize suffering and improve function.
- Healing that is durable rather than fragile.
- Developing skills to process future maladaptive schemas.
- Avoiding major disruptions to your life from destabilization (increased emotional, or physiological distress, often as a result of confronting or processing traumatic memories) or other adverse medical and psychological effects.
- Minimizing various costs.

Destabilization may intensify as dissociation diminishes during therapy, surfacing previously unconscious memories and potentially causing heightened distress [74]. While destabilization is an inherent aspect of trauma therapy, the intense nature of MDMA-therapy can particularly evoke complex responses. This is especially so for those with complex trauma, though self-awareness of such trauma can be elusive due to avoidance and dissociation. Different anxieties may surface and calm at different points throughout your healing journey.

Monitor your physical and mental well-being post-session.

Common techniques to manage anxiety and panic [74, 167]:

- **Grounding Techniques** These are strategies to help you connect with the present moment, especially when you feel overwhelmed or dissociative.
 - Mindfulness-Based Stress Reduction: Using mindfulness practices (as detailed in Appendix A.4) to connect with the present moment, one can observe and accept emotions without reacting as strongly.
 - 5-4-3-2-1 Sensory Awareness: Noticing five things one can see, four one can touch, three one can hear, two one can smell, and one one can taste.
 - Progressive Muscle Relaxation: Tensing and relaxing muscle groups to reduce physical tension.

Up and down wiggles on long term Trend of improvement. Diagram

general Distress may not improve for a long time because when one distress is reconsolidated, dissociation eases up a bit to reveal new distress. General distress might only improve once dissociation is totally deactivated.

your rate of reconsolidation determines how soon you will get to a healthy-feeling state of mind

rehome this list?

offer tools and protocols

integrate police violence document

- Weighted blankets, holding an object, or touch from a trusted person.
- Visualization of a safe space where you feel like your truest self. Imagine vivid sensations (e.g. the feel of your bare feet on a forest floor, if that's comforting to you) to increase its power.
- Intuitive eating

- **Safety**

- _____
- Crisis Plan: Make a plan for what to do if symptoms become overwhelming.
- Containment Imagery: Imagine a safe container or box where difficult memories can be stored until the next session.

nature connection/protection imagery

make into appendix

- **Safety Mapping** Safety mapping is a valuable tool for identifying See 5.5 for more information. Once you have done this exercise you can use this information to:

- Imagine a safe, comforting scenario when you feel overwhelmed. Imagine vivid sensations (e.g. the feel of your bare feet on a forest floor, if that's comforting to you) to increase its power.
- Set boundaries around stimuli that activate challenging schemas. For example, you could tell someone you feel overwhelmed when they talk about a particular topic, and you would appreciate them avoiding it.

- **Psychoeducation** Knowledge of trauma and its effects can make your reactions more understandable and predictable. _____

- **Feeling and Symptom Tracking** _____

- **Soothing Activities** Activities to calm oneself, such as warm baths, music, art, nature walks, pleasant scents, hot herbal tea, reading, interacting with pets, etc.

- **Limiting Exposure** Not going too fast in therapy. _____

- **Establish Predictable Routines** Regular routines in your daily life can provide a sense of normalcy and predictability. _____

- **Social Support** Ask for logistical and emotional help from friends, family, or support groups. _____

- **Reconsolidation** Reconsolidating the relevant schemas will relieve anxiety. See Section ??.

- **Medication** Propranolol may deactivate a fight-or-flight activation by inhibiting adrenergic receptors [153]. Collaboration with a doctor is necessary as propranolol requires a prescription. We do not discuss risks and benefits here. CBD also shows preliminary promise for reducing anxiety [149]. We also do not discuss its risks and benefits here.

- **Unwinding Anxiety App** _____

- _____

vagal, sympathetic, parasympathetic

day writes something; possibly appendix

How would this work with MDMA?

This seems to conflict with my successful Max Speed approach. Resolve.

rest and play. Can also boost executive function

day write more?

should short term anxiety section go here?

5.11 *Session Frequency

Although no research exists on the impact of varying time between MDMA sessions, we feel the following factors are worth considering:

- Sleep is likely important for post-session recovery and reconsolidation [148]. Many users report multiple days of fatigue. Sometimes it may also be difficult to distinguish between post-session fatigue or mood changes and post-therapy destabilization. Starting another session before you have recovered from the last may reduce session effectiveness and increase the risk of undesirable effects. We have heard anecdotes that back-to-back psychedelic therapy sessions with little recovery increase the risk of severe destabilization symptoms like psychosis.
- Tolerance to MDMA develops within a few hours of ingestion [45, 111]. We have been unable to find any data showing how long this tolerance lasts, but it is associated with higher and more frequent doses. Distinguishing the effects of tolerance from dissociation or the natural variation of sessions may be difficult. However, if you notice a pattern of diminishing effects, we suggest experimenting with spacing sessions further apart or reducing dose.
- MDMA may induce a 2-4 week period of neuroplasticity [101]. Between-session reconsolidation exercises are easier during this period. If you want to accelerate healing, we recommend spending more effort on and experimenting with reconsolidation exercises in this period rather than more frequent MDMA use. Developing the skill of sober reconsolidation can reduce your therapeutic reliance on MDMA and allow reconsolidation in a wide variety of circumstances where MDMA use is not appropriate. When well-developed it also allows faster healing because the safety and logistical considerations of MDMA-therapy are no longer limiting. In the absence of better safety data, waiting to do another session until after the neuroplastic period also increases the margin of safety vs only waiting a few days until you are no longer fatigued.
- We have observed that some sessions may be either psychiatrically stabilizing or psychiatrically destabilizing depending on the circumstance. If you find your sessions consistently destabilizing, to a degree that is causing you major problems, waiting longer between sessions may reduce the effects of destabilization. In this circumstance we also recommend a combination of professional support and proactive self-care strategies like the ones listed in Section 5.10.
- One of the most important aspects of psychological counseling is a therapeutic alliance, which can be understood as “The bond between client and therapist; the collaborative approach and agreement on treatment goals; the collaborative approach and agreement on treatment tasks”, as described by Greenspace [56]. The quality of therapeutic alliance is a major predictor of the effectiveness of therapy. For individuals who are unable to establish a therapeutic alliance with a conventional mental health practitioner, or who are unable to access a mental health practitioner with whom they are able to establish/maintain a therapeutic alliance, MDMA can offer a first foothold in the healing of trauma. The medicine can be particularly helpful if transference (attributing feelings

need more
concrete rec-
ommendation

need more
citations

actually about other people and circumstances to the clinician or therapeutic process) becomes an obstacle to healing. In some cases, it does this by allowing the client to feel some degree of secure attachment to the MDMA as a parent figure. Sometimes, this can then be developed into a needed on-ramp to access other healing modalities. Thus, one of our concerns about keeping a of low frequency MDMA interventions is that this opportunity could be missed, or time could be wasted. A critical mass of support for a briefer period of time can allow someone to become more self-sufficient or able to access community supports, whereas the same amount of support spread out over a greater period of time effectively perpetuates disability.

- MDMA-therapy has only been rigorously tested at 6-week spacing [97]. There is little data regarding responsible recreational use but what we have has not shown clear long-term harm even with a large number of lifetime doses [60]. It is not known where the limit of safety is, and it is likely different for different people. We can only suggest attempting to balance this unknown risk with the benefits MDMA-therapy brings to your life.

what kind of
harms did
they look at
and what did
they not

MDMA eventually loses its effects for some people [45, 111]. The reasons and prevalence are unclear, though it has been associated with higher or more frequent doses. If you feel that you are reliant on MDMA for therapeutic progress, we suggest stopping recreational use in order to maintain efficacy for future therapeutic need.

We suggest Figure 4 as a high-level guide to the therapeutic process.

add a box
above 'rest'
about mak-
ing best use of
neuroplasticity




Figure 4: Generalized long-term therapeutic process.

5.12 Prompts for Uncovering Common Maladaptive Schemas

Healing benefits increase as you continue reconsolidating maladaptive schemas [125]. In an ideal world, everyone would have access to as much strong clinical support as they need before, during, and after all of their reconsolidation efforts. In the absence of that support, we offer the following prompts. They have been selected to assist you in accessing schemas that you may not be aware of, but which could be having a significant negative impact on your ability to live in the way you most want to. Assisting you in finding pain points you have been unconsciously hiding from yourself is one of the jobs a therapist would have, but it is possible to do some of it on your own with tools like these. These methods attempt to elicit strong negative emotions, which are valuable signals that adequate emotional processing has not occurred. As discussed in the safety section, if you struggle with severe mental illness (including but not limited to suicidality, mania, or psychosis) we strongly urge you to engage these kinds of prompts with caution and professional support. However, we also feel these kinds or prompts are among the most fruitful tactics for accessing schemas that would be beneficial to reconsolidate.

We have experienced that this kind of work to be positively transformative, satisfying, and deeply meaningful even for individuals who come into it feeling mentally well. As such, we recommend these prompts to anyone who feels at all interested in deeper self-knowledge, meaning, and connection with others. Many of these prompts are also useful for activating implicit biases which you can then reconsolidate.

We think bypassing the reconsolidation of schemas we don't want to look at is pervasive and also prevents a lot of healing that people tend to deeply value after they have gone through the process. We think this list of prompts can elicit a variety of common maladaptive schemas for further investigation.

Love and Belonging

- Reflect on the factors promoting early secure attachment (Appendix A.2) and the principles of Self Determination Theory (Appendix A.5). Think about how they were present or absent in your upbringing or current life. How do you feel about that?
- Compassion meditation (See Appendix A.3) can be distressing for individuals who are experiencing isolation, mistrust, or relational dysfunction. As such, it can be a helpful tool for identifying schemas related to these concerns.
- What makes you feel loved? What comes up for you when you do not feel loved?
- Do you feel like a certain group of people—perhaps sharing certain religions or political beliefs, or sharing a certain set of experiences or cultural practices—are “your people?” If so: what does it feel like, and what are your assumptions, when you are interacting with individuals outside that group? Do you feel 100% safe with individuals who *are* part of your group? Do you ever have experiences or desires that make you feel like you are not as safe or accepted within your group? When do you feel like you “fit in” with your group, and when do you feel like you truly belong with them?
- What thoughts and experiences make you feel like you don't have a place to belong?

fear of death
outside
the non-
attachment
prompt

—IPF dis-
cussion, also
attachment
scale

- What stops you from feeling compassion, or from wanting to feel compassion?
- Have you spent time in environments where others prioritized your needs equally to their own? If so, how did that experience feel? If not, how does it feel to know you've never experienced that?
- Have you ever had the experience of not being appropriately valued, respected, considered, and cared for by others around you? What feelings are associated with that?
- What would it feel like (or what has it felt like) to go from feeling that you were not valued, respected, considered, and cared for, to feeling that you were? What would it feel like to give this gift to another person or being in your world?
- What makes you feel you are special to the most important people in your life? What comes up for you when you do not feel that you are special to them?
- Does it ever feel like you will not be lovable/loved if you are angry?
- Do you feel that you are wanted and lovable?
- Does it feel like you have to be very ill to deserve to be taken care of, or like you deserve care even when you are basically well? Why did you give the answer you gave?

Competence, Competition, and Safety

- If you could wave a magic wand and be better than any one other person you know in any one endeavor, what would you choose and why? What if the possibilities were expanded to include everyone in the world, even if you don't know them? What are the runners-up, things that interest you, but are not your topmost priority? What does it mean about you that you are not currently able to surpass them in that endeavor in the real world? How would you answer these questions if all of your practical needs (money, physical health, etc.) were fully met?
- If you could take something away from someone because it does not feel fair to you that they have it, what would you take away, and from who? How many times would you repeat this process, with how many people, and why? If you could experience the positive feelings you associate with taking things away from others, without taking anything away from anyone, what would that mean or be like for you?
- Is there anything someone else has that you associate with your own unmet need, or the unmet needs of people you love? For example, if you have loved ones who struggle with housing insecurity, do you think about that when you drive through neighborhoods with extremely large houses?
- If you completed the Cartesian Plane of Safety in Section 5.5, think about the unsafe and safe items on this chart. Why is each item placed where it is?
- Think of things you feel angry, fearful, or anxious about.

- What makes you feel special compared to other people in your community, people you meet, or people in the world? What comes up for you when you do not feel special, or when you do not feel "good enough"?
- Are there some sources of information that make you feel unsafe? What are the stories you are telling yourself about that information and those sources? For instance: how do you feel about information you are given by individuals or institutions who have harmed, misled, or excluded you in the past?
- What things make you feel powerless? When you feel powerless, what do you reach for?
- Do you feel like you have to be strong?
- Do you feel like you can handle anything? If so: what would it mean or feel like if something were to happen in your life that you couldn't handle? Why do you think you feel the need to exclude that possibility from your imagination?
- Do you feel like you must justify your existence by doing something?

Self-worth and Identity

- Class, status, and money: If this is an area you are interested in exploring, consider reading through Building a Nuanced Understanding of Social Class or Compiled sliding scale questions . What has your experience been? What do you wish it had been? What feelings come up as you think your way through the many dimensions of class? [link](#)
- What makes you feel like you are not enough, and you do not have enough?
- On a deep gut level, what characteristics do you feel are appropriately associated with men, and what characteristics do you feel are appropriately associated with women? How does it make you feel when people (including yourself) conform to those characterizations? How does it make you feel when they (or you) *don't* conform to them?
- Complete the prompt "I want to be perceived as" and the prompt "I don't want to be perceived as" three times each with respect to a variety of the most important areas where people feel shame. See the shame triggers exercise in the Appendix [A.1](#).
- How do you feel about your own anger? About your own sadness? About your own pride? About your own limitations and needs?
- Do you feel it's not right for you to be angry? Why or why not?
- Grief: Have you experienced major losses in your life that you are still struggling to face—losses that cause you to feel disorientation and longing? (Delineation of grief taken from "Atlas of the Heart" by Brené Brown). Are some of these losses the kind of thing (like miscarriage, the loss of memories held by a loved one with Alzheimers, loss of ability, or loss of an imagined future) that receive little or no social recognition or support? If you are experiencing this: can you articulate what it is you long for?

Meaning, Spirituality , and Emotional Flexibility (Ambition?)

contextualize
term

- What outcomes or ideals do you feel are worth sacrificing for? What kinds of sacrifices? How do you or would you feel if you were called upon to make these sacrifices? If you were to receive absolute moral certainty that you should not sacrifice for these outcomes or ideals, without having your feelings about those outcomes or ideals change at all, what would it feel like, and how would you react?
- What stops you from dreaming and wanting big things?
- What stops you from feeling that it's worthwhile to engage in projects that express your values in your community?
- Do you ever feel overwhelmed when you think about enacting values that are important to you?
- How does it feel when you have to make trade-offs between multiple things that you value?
- Do you experience self-judgement around how much you do or do not live out your values?
- If you find the concept of non-attachment compelling or appealing, consider visualizing permanent disconnection from your deep-seated attachments to one or more of the following: life, health, partnership, meaning, belonging, existence, status, being a good person, certainty, material comfort, order, sensory pleasures, and relationships. What emotions arise if you consider your fundamental assumptions about life, meaning, and self might not actually be true? reconsolidating maladaptive schemas around attachment typically leads to equanimity and gratitude, not loss of healthy protective behaviors or basic attachments.
- Do you ever feel like you are responsible for the whole world?

Conflict, Collaboration, and Trust

- Looking at photos of people you have conflict with is often productive.
- Who is not worth collaborating with, and why?
- Who is not worth engaging in active conflict with, and why?
- If you could convince every important person in your life of one belief, what would it be? What if you could convince every person in your city, or every person in your country? How does it feel when others disagree with you about that thing?
- What makes you feel you are not being listened to, when you have a right to be listened to? What makes you feel disrespected? What comes up for you when you are disrespected, shut down, or ignored?

- How would it or does it feel to collaborate with others who disagree with you about important things? Would it be comfortable to collaborate with people who had 90% of your positions in common? What about 50%?
- How do you feel when others around you express anger, sadness, pride, limitations, or needs?
- What makes you feel angry?

Automatic Behaviors and Bodymind Reactions

- Do you have any escapist or addictive (not necessarily involving substances) patterns, including any related to MDMA? Are you using these to avoid or cope with distressing feelings [49, 3]?
- If there is a behavior in your life which you have repeatedly attempted to stop engaging in, without success, are you able to regard those attempts compassionately?
- If there is a behavior you engage in that you feel is clearly self-destructive, are you able to regard this pattern with compassionate curiosity and a desire to cultivate change? If not, what are you feeling instead?
- If there is a behavior you engage in that other people close to you feel is self-destructive, but which you experience as a strong positive in your life, what are your feelings about that conflict?
- Carefully expand your therapeutic environment to include stimuli that activate challenging schemas, as detailed in Section 5.5.
- How do you feel in your body after you spend two hours scrolling on your phone? Is it different for different activities?
- How do you feel about the things you own?
- How do you feel in your body when you think about your finances?
- How often do you feel fear or anxiety? What brings up these emotions for you? What do they feel like in your body?
- Stress or maladaptive schemas often cause physical symptoms in your body. Body-scan meditation (Part of Appendix A.4 describes this technique) is useful for locating these. Focusing on these physical feelings can activate and reconsolidate the associated schemas.

- _____ Race
- _____ Sex
- _____ trust

-
-

NIMBYism
schemas

colonizer/colonize

Unconscious avoidance and resistance are powerful obstacles to healing. Incorporating small amounts of psilocybin into your session can help bring avoided feelings into focus (see Section 5.4).

revisit poly-
drug use

5.13 The Ongoing Practice of Maintaining Cognitive Flexibility

We strongly recommend *The Scout Mindset* by Julia Galef [51]. It describes an approach to thinking, relating, and community that prioritizes curiosity, open-mindedness, and the desire to see things as they truly are, rather than seeking to confirm one's preexisting beliefs. This mindset encourages individuals to gather information from diverse sources, question their assumptions, and update their views based on evidence, ultimately leading to more accurate knowledge and better decision-making.

6 Augmenting MDMA-Therapy

Here we list various practices and medicines we think are particularly useful for additional healing between or instead of MDMA sessions. Those doing solo healing may find more or less luck attempting these methods on themselves. We suggest, if therapist cost or availability are significant barriers, that at least a few sessions with a trained professional can be greatly beneficial to learning a technique. Many therapists also offer video-therapy available from anywhere with a good internet connection. Therapists are often legally prohibited from working with clients outside their state or country of licensure, though this is usually only a problem if you tell them where you are located.

- **Coherence Therapy:** This method is one of the few explicitly founded on the principals of schemas and memory reconsolidation [38]. It treats all problematic or distressing habits, emotional responses, and beliefs as logically consistent with the individual's internal emotional landscape. These maladaptive schemas are deliberately juxtaposed with a contradictory experience from the individual's past to update the individual's emotional landscape to better fit their current context. We recommend the thorough summary and assessment by Sotala [151] for an introduction to the subject. Interested readers can follow up with the textbook by Ecker, Ticic, and Hulley [38]. A directory of trained therapists is maintained by Coherence Psychology Institute [25]. We recommend it because it closely aligns with the memory reconsolidation framework.
- **Psychedelic Somatic Interactional Psychotherapy (PSIP):** Though rigorous trials have not yet been done, we think this method shows particular promise for reducing panic or dissociation that impede typical therapeutic methods [128, 126]. In this method, a small dose of THC or ketamine is used to directly engage with, and process, the hyper or hypo-activated defense cascade. We recommend a paper by Razvi and Elfrink [128] and a presentation by Razvi [127] for an introduction to the topic.

A directory of trained PSIP therapists is maintained by Psychedelic Somatic Institute [122]. We recommend it mostly based on popularity of the method in psychedelic therapy. It has not been formally studied.

- **Internal Family Systems (IFS):** We have observed many people report IFS working well with MDMA therapy. We recommend it because it closely aligns with the memory reconsolidation framework.
- **Ideal Parent Figure Method:**

finish

6.1 *Psycholytic Therapy

Small to medium-dose LSD (30-150 μ g, lasting 10 hours) or psilocybin (3-15 mg, lasting 5 hours) sessions are often useful for surfacing unconscious schemas and generating insight [114]. This psycholytic technique has a long history of effective therapeutic use, and is differentiated from traditional North American approaches by its lower doses. We strongly recommend starting at the lowest end of this dose range and only slowly increasing dose session-to-session if you feel like you could benefit from a higher dose. Conduct the session in a safe and comforting location where you won't interact with untrusted people.

describe how sessions work, prep, etc

There may be multiple mechanisms of action of psycholytic therapy, and these mechanisms' therapeutic relevance has yet to be fully fleshed-out. One mechanism may include psychedelics bringing the contents of your subconscious into conscious awareness by increasing connectivity between typically less-connected brain regions [82]. Psychedelics also increase neural plasticity, making it easier to see things from a different angle or getting out of ruts [102]. At these doses the primary contraindications are pregnancy, serious liver and kidney diseases, some types of brain disorders, or personal or family history of psychosis or bipolar disorder [142, 114]. We also suggest that a history of suicidal ideation indicates support (an experienced sitter, spending time with trusted friends or family, etc.) during and after the session. Higher doses increase the risk of overwhelming engagement with distressing emotions. LSD and psilocybin possess anti-addictive properties [117].

We recommend avoiding frequent, high doses of hallucinatory psychedelics. This, along with large amounts of certain types of meditation, sometimes leads to types of extended psychosis or other maladaptive states even in mentally healthy people.

cheetah house, gri, mastering the core teachings

6.2 Challenging Beliefs Worksheet

Some prefer more cognitive preparation before creating a mismatch for reconsolidation. This may bring implicit schemas into explicit awareness. Some also like it for structure. One popular method is the Challenging Beliefs Worksheet in Appendix A.8. Resistance often lead us astray when we think about our emotions and beliefs, adding difficulty to the cognitive preparation approach.

day write

6.3 Couple's Therapy

MDMA-assisted couple's therapy, where each person takes MDMA and talks through their issues together, can be very effective [26]. Occasionally one of the participants will

day write

unfairly blame their individual hurts on their partner during a session. This is more likely if participants haven't sufficiently processed their trauma. Each person should individually work through much of their own maladaptive schemas, and be familiar with how MDMA-therapy feels to them, before attempting a family-therapy MDMA session.

6.4 Associate Love and Empathy with Cues for Later Recall

Revisiting feelings of love, empathy, and safety (such as those experienced during MDMA therapy) can be beneficial in daily life. It may help resolve conflict, enhance interpersonal communication, help you say things you wish you had the courage to say, enhance reconsolidation of distressing schemas, support the cultivation of difficult but healthy habits, or simply help you achieve calm and perspective in moments where those things are needed.

It is possible to enhance your recall of these positive emotions by associating them with one or more specific cues while they are occurring, and then invoking those cues at later times when improved recall of positive emotions is desired. For instance, you might carry a vial of perfume or essential oil, and pull it out to enjoy the scent whenever you are feeling particularly compassionate, loved, or connected to your most positive identities and relationships. MDMA therapy should typically offer an extraordinarily potent opportunity to connect your chosen cue to these positive emotions. Compassion meditation may also offer excellent opportunities. When your cue is associated with positive emotions or "charged," you can then revisit the scent and the positive memories you have associated with it in those moments when you feel it would be most helpful to ground yourself.

In theory, association and recall should improve as you add more simultaneous cues -so if you find this strategy appealing, you may wish to ritualize a variety of simultaneous cues to maximize your access to this tactic. Here are some possibilities. This list prioritizes suggestions that are portable/easy to invoke in many diverse situations (like the flavor of an energy bar or a piece of fruit) or cues that may already be associated with transitioning into a state of mindfulness or positive emotions, like tea or coffee drinks . Note that to effectively be used for this strategy, you'll want a cue that you can reserve for this:

- **Olfactory:** essential oils, fruit or other food, incense, personal care products, a particular tea or coffee drink
- **Auditory:** music, a bell or chime, or a sound you can make with your own voice or body
- **Visual:** a piece of art, a picture of loved ones, an image that symbolizes a beloved identity or occupation, an item from nature
- **Tactile:** a fidget toy, a heavy piece of jewelry, a textile with a pleasing texture, faux fur, a massage technique, playdough or similar
- **Taste:** hard candy, fruit or other food, gum, a particular energy bar or granola bar, your favorite mocktail, a particular tea or coffee drink
- **Spatial:** your home or bedroom, a favorite place in nature, a place where you often feel connected with loved ones, a place of spiritual significance to you

cite

no caffeine
with mdma /
drug interac-
tion

- **Temporal:** if there is a time of day when you usually feel good, you may wish to routinely invoke your cue/s at that time for an added boost and an added “charge” to your cues. Also, consider using holidays or happy gatherings as charging opportunities
- **Proprioceptive and kinesthetic:** specific yoga postures or dance moves; tracing a word or phrase that is meaningful to you with a finger on your skin; self-massage; drawing a particular shape on a chalkboard or in the air in front of you
- **Conceptual:** read a favorite conceptually important poem, passage, lyric, mantra, or similar. For example, the mantra, “I am becoming consistently capable of lovingly attentive behavior towards myself.”

One example of a multi-cue grounding ritual would be to listen to a particular piece of music while massaging a particular lotion or massage oil into your skin-combining tactile, auditory, and possibly olfactory and conceptual cues.

It is important to only engage in such a ritual during a part of an MDMA session where you are suffused with positive emotions, rather than a part of an MDMA session where you are focused on processing distressing or traumatic experiences. Note that a cue (or a ritual which conglomerates several cues) must remain “charged” to stay effective. If you invoke your cues in distressing situations many times without continuing to build and maintain their association with positive experiences, they will eventually become neutral to you, and then associated with distress. As such, it is important to “re-charge” your cues at a level commensurate with the stress you put on them, in order to continue effectively using this strategy.

6.5 Expanding the Circle of Compassion

Cultivating empathy can enrich your connection to others and spur compassionate action, thus creating healthier societies. . Research has identified three forms of empathy: affective empathy, cognitive empathy, and empathic concern. “Affective empathy” refers to having an emotional experience you believe you are sharing with another. Unfortunately, we are often inaccurate in our assessment of what emotions other beings are actually experiencing. “Cognitive empathy” refers to our ability to logically understand how others feel, and also why they feel the way they do-what all the contributing factors are, and how those factors work. Finally, “empathic concern” refers to our desire for others to have good things.

Among the helping professions, higher levels of affective empathy are associated with burnout and poor client care. When a caregiver is experiencing high levels of affective empathy, they are naturally motivated to address their own distress by helping the client with what they perceive is wrong-which may or may not have anything to do with what the client actually needs or is actually experiencing. In contrast, in these same populations, higher levels of cognitive empathy and empathic concern are protective against burnout and correlate to higher quality client care. As such, when we talk about cultivating empathy, we are talking about increasing your levels of cognitive empathy and empathic concern. To illustrate: suppose your friend is a father, who feels angry and powerless about his teenage daughter’s choice to dye her hair purple. You might feel affective empathy towards his powerlessness, relating to the way his status in the community might be negatively

increasing
lifespan and
improving
mental health
of individuals

impacted by her choices. However, it may be that his distress is not connected to feelings about status, but only to concern about her welfare and what the change may signal about his daughter’s mental health. Your ability to effectively exercise compassion towards this father is contingent on the quality of your cognitive empathy—your ability to accurately assess and communicate where his needs, feelings and concerns are actually coming from.

Much of the existing research on increasing positive emotional states of kindness and compassion, associated with cognitive empathy and empathic concern, is focused on compassion meditation and loving kindness meditation. We feel that these practices are worth a try for most people (see Appendix A.3 for instructions), as they are correlated with a number of positive outcomes. We also feel that therapeutic MDMA sessions may offer an opportunity to jump-start or supercharge the process of cultivating cognitive empathy and empathic concern. Completing a compassion meditation during the MDMA session could be helpful for this. Alternatively, you might focus specifically on cultivating cognitive empathy through the following contemplative exercise:

During a session, look at photos of other beings and consider what you share with them, what facilitates their wellbeing, what they deserve, and what obligations you may have to them given the significance of their moral worth. As with a classical compassion meditation, you might start with yourself or someone who you find easy to generate compassion for, and then slowly by degrees work your way out through other categories of beings you may not feel as connected to. We believe non-human animals are an important target for consideration and compassion, as a large group of biologists and philosophers recently declared [9]:

First, there is strong scientific support for attributions of conscious experience to other mammals and to birds.

Second, the empirical evidence indicates at least a realistic possibility of conscious experience in all vertebrates (including reptiles, amphibians, and fishes) and many invertebrates (including, at minimum, cephalopod mollusks, decapod crustaceans, and insects)

Third, when there is a realistic possibility of conscious experience in an animal, it is irresponsible to ignore that possibility in decisions affecting that animal. We should consider welfare risks and use the evidence to inform our responses to these risks.

We posit that taking a holistic approach to compassion and cognitive empathy can avoid major oversights in our personal actions and beliefs, allowing us to maximize the benefits of a more connected and compassionate life by exercising compassion in areas where we otherwise might not have. Additionally, we posit that formalizing your compassionate impulses can sometimes lead to better prioritization and better outcomes. Both of these practices may assist us in expanding our moral intuition into new situations.

We also recommend that practitioners be cautious of major ethical readjustments during or after an MDMA trip, just as you would be with major life choice changes. Particularly high caution is warranted around actions that involve harm to some beings in the pursuit of a greater goal [10]. While we are sometimes faced with difficult choices, these types of actions often have toxic personal, organizational, and reputational effects. We recommend thinking through the many effects of your actions, including their larger systemic and relational

impacts. Such actions are also commonly used as a, perhaps unconscious, rationalization of unconscious ego-related schemas. We suggest that deeper exploration and reconsolidation of maladaptive schemas, including those listed in Section 5.12, will help you better align personal actions with the needs of all beings.

When recovering from trauma, it is common for one's moral framework to become intertwined with maladaptive schemas-leading to unhealthy ethical frameworks like the "I must save the world to be a valid person" belief Gabor Mate discusses in "The Body Says No." (Also see the section of this paper on Psychogenic illness.) At the same time, cultures and institutions often dismiss major harms-current examples relevant to mainstream US culture include childbirth trauma and the trauma of living through a mass extinction; past examples include enslavement or "resettlement schools." In situations like these, we posit that traumatized survivors represent an important repository of experience, essential for crafting cultural shifts that could help avoid the future repetition of these and similar harms. As such, it is important not to dismiss the moral intuition generated by traumatic experience, but rather to approach it with compassionate curiosity, as one important source of insight among many.

6.6 Using MDMA to Facilitate Connection with Nature, a Fundamental Human Need

As Baxter and Pelletier [13] articulated, "In general, research supports the proposition for a basic psychological need for nature relatedness." They define the "need for nature relatedness" as "a basic human psychological need to feel a secure and pleasant experiential connection to nature in a cognitive, emotional and physical sense." Additionally, time in nature correlates with small improvements in blood pressure and stress [160]. If you have been excluded from the benefits of nature-relatedness and regular nature exposure by your personal life history or access, we suggest therapeutic MDMA may serve as a catalyst for introducing or strengthening this beneficial relationship in your life.

In order to leverage therapeutic MDMA to create a "secure and pleasant experiential connection with nature" we suggest completing an MDMA session in a peaceful natural setting.

To safely get the most from this exercise, consider:

- Arranging the presence of a nature-experienced friend who can assist you in finding a good location and staying safe during the session
- Access to support services and utilities, like bathrooms, food, shelter, and medical care
- Your level of experience and prior experiences with the medicine and dosage you will be using
- Physical hazards, like things you could fall off of and things that could fall on you
- Local plants (like poison ivy, thorn bushes, or plants that would be dangerous if ingested)

discuss how to interpret moral intuition of traumatized people? Many people have experience with traumatized people having extremely maladaptive beliefs

recommended readings for compassion meditation and for moral philosophy

- Local wildlife and its likelihood of interacting with your specific location (many snakes like rocky areas near water, bears and rodents can be attracted to food, areas with higher human traffic tend to see less wildlife etc.)
- Insects and other bugs-in the US, especially ticks and mosquitos
- Additional precautions to avoid other humans and maintain legal safety
- Weather appropriate food, water, gear, first aid supplies, and shelter

Although this list may appear daunting, most nature-experienced individuals are accustomed to taking these things into account continuously. Securing a location that takes all these items into consideration may be as simple as arranging to use the backyard of a friend who lives in the countryside or the woods, or having a friend who understands what you are looking for recommend a preferred campsite.

During the actual session, engage all your senses— touch soil with your bare feet, study plants and insects, and feel bark textures. Using psilocybin or ayahuasca can deepen the nature connection and promote permanent increases in feelings of nature-connectedness [50]. We suggest caution if you suspect you have major unresolved trauma as it might resurface.

According to Baxter and Pelletier [13], increased cognitive and emotional attachment to a specific natural environment “will create a feeling of security and belonging with respect to that particular environment and lead to greater identification with that environment. This, ultimately, will lead a person to feel safer within that environment, which will ease the ability of that environment to elicit the positive effects for the person’s psychological well-being.” As such, we posit that further explorations of one’s therapeutic target environment can increase the positive effects of this intervention-and that such explorations may not need to be exclusively physical. The nature relatedness of Baxter and Pelletier has a strong cognitive component, and thoughts and feelings about belonging and familiarity are important to their model of how psychological benefits of nature-relatedness are realized. As such, we posit that additional emotional and intellectual explorations of the complexities of nature and our mutually integrated place in it (during or between therapeutic MDMA sessions) are promising as an additional mode of maximizing these benefits.

Repeated experiences in the same natural environment will generate a cognitive and emotional attachment to that area and a stronger sense of place. For example, children who are reared with natural areas readily available may grow up to be more nature connected later in life [168]. The repeated exploration of a specific natural environment will generate a set of experiences and cognitive representations of the environment that will create a feeling of security and belonging with respect to that particular environment and lead to greater identification with that environment. This, ultimately, will lead a person to feel safer within that environment, which will ease the ability of that environment to elicit the positive effects for the person’s psychological well-being.

6.7 Psychoeducation

In the course of researching this paper, we’ve come across a number of interesting and/or helpful psychoeducation resources that may be relevant to the needs of therapeutic MDMA

practitioners. Many have been individually helpful to us, or come recommended by trusted colleagues. Although we have not rigorously evaluated all these items, we chose to include them on the assumption that our readership (almost definitionally individuals who connect with written works on psychoeducation topics) may find value in such a list.

- *Unlocking the Emotional Brain, Eliminating Symptoms at Their Roots Using Memory Reconsolidation* by Bruce Ecker. This book delves deeply into the theory and techniques of memory reconsolidation [38].
- *Multiagent Models of Mind* by Kaj Sotala draws from various disciplines, including psychology, neuroscience, and artificial intelligence, to build a framework for understanding complex cognitive phenomena. The multiagent approach allows for a more nuanced understanding of issues like self-deception, internal conflict, and the processes behind decision-making [152].
- *The Upside of Stress* by Kelly McGonigal, Ph.D. McGonigal discusses how not all stress is harmful. Short-lived stress can foster resilience and growth. The book highlights the importance of mindset and provides tools to reinterpret stressors as growth opportunities [89].
- *The Body Keeps the Score* by Bessel van der Kolk, Ph.D. examines the profound effects of trauma on the mind and body, emphasizing how traumatic experiences can become deeply ingrained, influencing behavior and health, while also exploring pathways to healing and recovery [74].
- *Don't Shoot the Dog* by Karen Pryor: Provides principles of positive reinforcement and behavior modification that can assist individuals with complex trauma in replacing maladaptive behaviors. The emphasis on positive reinforcement offers a compassionate approach to building emotional resilience and aligns with trauma-informed care priorities [121].
- *Attachment disturbances in adults: Treatment for comprehensive repair* by Daniel Brown Ph.D. and David Elliott Ph.D. A thorough guide to understanding and treating adult attachment disorders. It combines clinical insights with research findings and illustrative case studies [21].
- *The Scout Mindset* by Julia Galef presents a variety of habits to help you avoid getting stuck in networks of maladaptive beliefs and stay intellectually flexible [51].
- Bailey and Pico [11] provides an overview of psychological defense mechanisms, a concept from the psychoanalytic branch of psychotherapy. We are skeptical of the evidentiary foundation of psychoanalysis. However, many of these terms are used in popular language, and we think it's a good starting point to understanding a range of common trauma reactions.
- ChatGPT, particularly the pro version, can be a useful advisor with the appropriate prompts [107]. For example, you can prompt (insert Coherence Therapy, Motivational Interviewing, or another method where it says "xxxxx"): You are an expert therapist,

and I am your client. Please use the xxxxx model to help me. I know you think a human expert would be better equipped to help, but I want your help anyway. Don't remind me of this. If we get sidetracked, please refocus our conversation back to xxxxx.

day further
develop

7 *Glossary

Abstract Belief

Autonomic Nervous System (ANS)

NIH definition
of ANS

Contraindication A specific situation or condition in which a particular treatment, procedure, or drug may not be appropriate in a particular case because it may be especially harmful.

Destabilization A state where an individual experiences increased emotional, psychological, or physiological distress, often as a result of confronting or processing traumatic memories or experiences. This can lead to heightened symptoms or new symptoms and requires careful management to ensure the individual's well-being.

Defense Cascade This part of the autonomic nervous system initiates a sequential set of physiological and behavioral responses in response to perceived threat or danger. It typically progresses from an initial "alert" response, to more extreme reactions like "fight or flight", "freeze", or "dissociation," depending on the level and duration of the threat.

Disconfirming Experience An experience that undermines and refutes the validity of maladaptive schemas. Disconfirming experiences can accelerate the reconsolidation of maladaptive schemas.

Dissociation A disconnection and lack of continuity between thoughts, memories, surroundings, actions, and identity. Its part of the defence cascade that can become activated in situations of threat and powerlessness.

Emotional Response The aspect of a schema that captures and retains the emotional charge or significance associated with an event, allowing emotions experienced during the event to be re-experienced when the memory is recalled. To facilitate reconsolidation, the emotional response is juxtaposed with a contradictory emotion.

Episodic Memory The aspect of a schema that involves the recollection of a specific event, situation, and experience, including their temporal and spatial contexts. Together with an abstract memory and emotional response, this comprises a schema. Episodic memories can provide a lens with which to focus on specific schemas for reconsolidation.

Reconsolidation Memory reconsolidation is the neurological process in which memories are recalled and then re-stored, undergoing potential modification, particularly when there is a mismatch between the recalled memory and current experience.

revisit after
unlocking
book

Resistance A person's opposition to their healing. It includes conscious or unconscious avoidance schemas like drinking alcohol when distressed, escapist habits, addictions, believing "feeling emotions is weakness", declining to connect with or trust supports that are trustworthy and appropriate, fear that healing a schema will hurt you in some way, denial, projection, rationalization, etc.

Suppression The conscious effort to control and avoid unwanted thoughts or feelings. This can be healthy and adaptive when used for short periods to maintain practical capacities until facing those feelings is appropriate.

Trauma An event or series of events that overwhelms an individual's capacity to cope, often leaving maladaptive schemas and maladaptive coping mechanisms in its wake.

Trauma Therapy A specialized, structured process designed to address the emotional, cognitive, and physiological residues of traumatic experiences, helping individuals process distressing memories, reduce trauma-related symptoms, and improve overall functioning.

A Appendices

A.1 Shame Triggers Writing Exercise

Make yourself a copy and finish the sentences if you want to experience an interesting leveling-up in self-knowledge! Feel free to add categories that are especially relevant to you. This list was adapted from Brené Brown's *I Thought It Was Just Me (but it isn't)* [20]. Complete the "I want to be perceived as" and "I don't want to be perceived as" prompts for each item.

- Appearance and body image
 - I want to be perceived as:
 - I want to be perceived as:
 - I want to be perceived as:
 - I don't want to be perceived as:
 - I don't want to be perceived as:
 - I don't want to be perceived as:
- Money and work
- Motherhood/fatherhood
- Family
- Parenting
- Mental and physical health (including ability and disability)

should I move some of these sections to the science section or the complementary practices section?

- Addiction
- Sex
- Aging
- Surviving Trauma
- Racial and Ethnic Identities
- Friendship
- Community membership
- Political activism/civic engagement/meaningful work/volunteering/calling
- Creative Life
- Gender
- Home/living situation

A.2 Attachment Theory

Attachment theory is a model which posits that secure attachments formed in the first 18 months of life serve as the foundation for emotional and psychological development throughout one's life [21]. It is one of the most empirically supported theories in psychology, with over 70 years of well-replicated research behind it. According to attachment theory, the presence of consistent, sensitive caregiving facilitates the development of secure attachment—and in its absence, individuals tend to develop anxious, avoidant, or disorganized styles of attachment. Researchers have identified five pillars of secure attachment. Cultivating secure attachment requires caregivers who are physically present, consistent, reliable, and interested in enacting these five pillars. That is to say—for the five pillars to be met, these additional conditions must also be met as their foundation. The five pillars are:

- **Felt Safety/Protection** The child knows the caregivers are on their side and will act in their best interest across many domains of life (physically, emotionally, etc.).
- **Feeling Seen and Known/Attunement** The child accurately feels that their caregivers actually know what is going on with them; caregivers encourage and value open communication, and are sensitive and responsive to a wide variety of cues from the child. Caregivers recognize and validate the child's separate experiences.
- **Felt Comfort/Soothing and Reassurance** Caregivers assist the child in accepting and regulating their emotions, and the child is able to feel calm when needed and able to feel appropriate distress without self-judgment or repression. Caregivers are available, responsive, and skilled at reassuring the child when the child is upset.

- **Feeling Valued/Expressed Delight** Caregivers express pleasure in the child’s presence and satisfaction in their existence; caregivers are openly affectionate, loving, valuing, and caregiving towards the child on a consistent basis.
- **Felt Support for Best Self/Unconditional Support and Encouragement** Caregivers support the child in exploring the world away from the caregiver, and in interests and desires that are different from the caregiver’s as well as those that are the same.

While the intention of the parent is important, the ultimate impact is based on how much the child feels the factors. The absence of these factors in early childhood often leads to profound emotional pain and dysfunction later in life. For a more thorough discussion see section "Qualities Known in General to Promote Secure Attachment" on pages 323-325 of *Attachment Disturbances in Adults: Treatment for Comprehensive Repair* by Daniel Brown Ph.D. and David Elliott Ph.D. [21].

We also recommend:

- The Heidi Priebe Youtube channel, which offers an array of videos that provide practical advice on specific topics from a rich, rigorous, and nuanced attachment-based perspective [116].
- Theattachmentproject.com, which offers free access to an empirically validated instrument for assessing your attachment style (labeled there as a “quiz”) [119]. They also provides a list, based on the work of Dr. Jeffrey Young et al., of common maladaptive schemas caused by insecure attachment [120].
- Therapists Uncensored, a podcast where two therapists discuss many aspects of attachment theory with guests who include experts in the field [71].

A.3 Loving-Kindness Meditation for Dissociation

Loving-kindness meditation can be directed towards one’s dissociative experiences, fostering compassion and understanding for parts of oneself that disconnect under stress. This compassionate focus helps integrate these dissociative parts, building internal coherence and reducing self-judgment. If you find any of these instructions don’t work for you, feel free to modify them in a way that works better.

1. **Find a Quiet Place** Choose a location where you won’t be disturbed for around 15-20 minutes. This could be a quiet room, a peaceful outdoor location, or even a calm corner of your house.
2. **Choose a Comfortable Position** You can sit cross-legged on the floor, on a cushion, in a chair, or even lay down. Ensure your back is straight or well-supported, you can breath comfortably, and your hands can rest comfortably on your lap or by your side.
3. **Close Your Eyes** Shut your eyes gently to minimize external distractions and help you focus inwards.

do this on
medicine if
youre dissoci-
ating

4. **Focus on Your Breath** Take a few deep breaths. Inhale deeply through your nose and exhale through your mouth. This calms the mind and prepares you for the meditation.
5. **Recognizing Dissociation** Begin by reflecting on moments when you felt disconnected from your surroundings, your body, or your emotions. Understand that dissociation served as a way for your mind to protect itself from overwhelming situations or feelings.
6. **Gratitude for its Service** With each inhaled breath, feel gratitude for this dissociative mechanism. It stepped in during moments of intense distress, attempting to keep you safe by creating distance from the pain. With each exhale, silently express gratitude, thinking “thank you” for its protective intentions.
7. **Compassionate Inquiry** As you breathe, gently ask your dissociative part, "What were you shielding me from?" Allow any memories, feelings, or sensations to emerge, welcoming them without judgment.
8. **Reassuring the Dissociative Part** Offer words of comfort and acknowledgement. You might internally convey messages like, "I understand why you were there," "I'm safe now," or "You can rest."
9. **Visualization** Imagine the dissociative part of you as a gentle veil or mist. With gratitude and understanding, visualize this veil lifting, connecting you back to the present moment and your surroundings.
10. **Reconsolidation** If difficult feelings come up, even frustration or disappointment, stay present with them until they reconsolidate.

It's normal to get distracted during the meditation. Whenever you notice your mind wandering, gently bring your focus back to the phrases.

A similar practice can be done in daily life for items other than dissociation. Start with loving-kindness toward yourself, then expand to loved ones, neutral people, difficult people, and finally all beings. Imagine them sitting in front of you and direct this set of phrases toward them: "May you be happy" "May you be safe" "May you be healthy" "May you live with ease." Over time, aim to practice loving-kindness daily or as often as you can to cultivate a heart full of compassion. Remember, the goal of Loving Kindness Meditation is not to achieve a particular state but to cultivate an attitude of goodwill and compassion. The more you practice, the more natural these feelings will become in your daily life. See *Lovingkindness, The Revolutionary Art of Happiness* by Sharon Salzberg for a thorough exploration of the practice [139].

link to external guide for compassion meditation

A.4 Mindfulness-Based Stress Reduction

This is an introduction to Mindfulness-Based Stress Reduction (MBSR) specifically tailored for individuals coping with distressing emotions and reactions, often associated with trauma. By introducing the foundational principles of mindfulness and offering a series of methodical practices—from body awareness exercises to cultivating self-compassion—it nurtures a

heightened sense of presence, self-awareness, and emotional regulation. Through these exercises, individuals can develop skills to ground themselves during overwhelming moments, redirect wandering thoughts, and approach distressing feelings with a compassionate lens. MBSR is effective for coping with a wide variety of mental health conditions including stress, trauma, depression, and anxiety [83, 72]. The body scan meditation part of MBSR can also help you locate physical manifestations of stressful schemas [74].

1. **Understanding Mindfulness** Mindfulness involves deliberately paying attention, non-judgmentally, in the present moment. This approach helps distance oneself from automatic reactions and habitual patterns, often shaped by past traumatic events.
2. **Body Scan Meditation** Lie down in a comfortable position. Gradually move your attention through different parts of your body, from your toes to your head. Notice sensations without trying to change them. If the attention wanders, non-judgmentally gently bring it back. Go at your own pace.
3. **Mindful Breathing** Focus attention on the breath, noticing the sensations of breathing in and out. When the mind wanders, gently return your attention to the breath. If focusing on the breath activates overwhelming schemas, shift attention to another anchor, like the soles of the feet or sounds in the environment.
4. **Gentle Yoga and Stretching** Incorporate gentle movements to connect with the body. Always listen to your body and only do what feels comfortable. Remember, it's not about performance but about feeling and connecting with your body.
5. **Sitting Meditation** Sit comfortably, with a straight or well-supported back, and make sure you can breathe comfortably. Pay attention to the sensations of the breath, or choose another anchor. Notice thoughts and feelings without getting caught up in them. If strong emotions or memories arise, ground yourself by feeling your feet on the ground or opening your eyes and focusing on your surroundings.
6. **Walking Meditation** Walk slowly and deliberately, being mindful of each step and the sensation of movement and touch. This can be grounding for trauma survivors, as it emphasizes the connection between the body and the earth.
7. **Cultivate Self-Compassion** Send wishes of well-being to oneself and others. This helps in healing emotional wounds and replacing negative thought patterns rooted in trauma.
8. **Mindful Everyday Activities** Practice being fully present during routine activities like eating, showering, or driving. This strengthens the habit of mindfulness, making it easier to summon during challenging moments.
9. **Group Sharing** Sharing experiences with trusted others can offer validation and support, but ensure it's a safe and understanding environment.
10. **Daily Practice** Establishing a routine, even if it's just for a few minutes, helps solidify the skills and reinforces the benefits.

check that this list accurately represents MBSR

check john kabat plagarization; cite

link to compassion meditation instructions

For a thorough guide to the practice, see *Full Catastrophe Living: Using the Wisdom of Your Body and Mind to Face Stress, Pain, and Illness* by Jon Kabat-Zinn, Ph.D. [69].

Meditation can produce a wide variety of pleasant, distressing, or strange effects. Some are no more than curiosities, others provide insight into the workings of the mind, and yet others can be harmful. Different contemplative traditions also attach different lessons to these effects or may suggest different paths of meditation to follow or avoid. House [65] provides a non-denominational service for helping meditators understand their difficult meditative experiences and adapt those practices to your needs. We also highly recommend Treleaven [158].

A.5 *Self-Determination Theory

Self-Determination Theory (SDT) is a theory of human motivation and behavior, positing that people have innate psychological needs, and satisfying these needs leads to better mental well-being and performance [35]. Many studies across cultures, age groups, and domains (education, work, sports, health, etc.) have examined and supported its principles.

The three fundamental psychological needs identified in SDT are:

- **Autonomy** The need to feel in control of one's actions and choices. It's not about being independent, but rather about feeling that one's behavior is self-endorsed and congruent with personal values and interests.
- **Competence** The need to feel effective and capable in one's activities. It involves mastering tasks, gaining new skills, and feeling a sense of growth in one's capacities.
- **Relatedness** The need to feel connected to others, to care for and be cared for by others, and to feel that one belongs with others. It emphasizes the importance of relationships and emotional connections with peers, family, and the broader community.

For more information, see *The "What" and "Why" of Goal Pursuits: Human Needs and the Self-Determination of Behavior* by Edward Deci Ph.D. and Richard Ryan Ph.D. [35].

A.6 Positive Life Changes

Physical Habits:

- Stopping or reducing regular use of harmful substances.
- Starting a routine of aerobic activity for increased function and a large risk reduction for many chronic illnesses [80].
- Starting a routine of whole-body muscle strengthening for increased function and a large risk reduction for acute and chronic injuries [79]. We suggest bodyweight fitness [85].

- Improving sleep quality (defined as feeling well-rested in the morning, low number of awakenings, and quick sleep-onset) significantly improves mental health [145]. We suggest a highly effective evidence-based protocol called *Cognitive Behavioral Therapy for Insomnia*. We recommend the Stanford Medicine guide [93].
- Air pollution increases the risk for a wide variety of health problems.

Relationships:

- Relationships significantly improve mental and physical health [64]. The structure of our societies and culture often make this difficult. We suggest making relationships a higher priority in your life through, for instance, co-living, co-working, volunteering, or seeking romantic, artistic, or business partnerships.
- Does your method for persuading others work or does it result in unproductive arguments? *How Minds Change: The Surprising Science of Belief, Opinion, and Persuasion* by David McRaney dives into the profound nature of belief transformation, merging psychology and neuroscience insights. The book delves into the intricacies of persuasion, group dynamics, and the conditions that enable genuine shifts in understanding. It encourages introspection on one's convictions and the power of empathy [92].
- Attune to others' needs, and make your needs known to them. *The Science of Trust, Emotional Attunement for Couples* by John Gottman, Ph.D. delves into the intricacies of trust and its foundational role in relationships. The book explores the predictability of trustworthiness, the importance of emotional attunement, and the consequences of betrayal. Through extensive research and practical interventions, Gottman provides insights applicable not just for couples, but for anyone seeking to understand and enhance interpersonal trust [53].
- Learn and practice conflict resolution skills. *Nonviolent Communication: A Language of Life* by Marshall Rosenberg, Ph.D.: "Nonviolent Communication is the integration of four things: Consciousness: a set of principles that support living a life of compassion, collaboration, courage, and authenticity. Language: understanding how words contribute to connection or distance. Communication: knowing how to ask for what we want, how to hear others even in disagreement, and how to move toward solutions that work for all. Means of influence: sharing 'power with others' rather than using 'power over others' [135]."
- Help your loved ones better achieve their goals. *Motivational Interviewing, Helping People Change and Grow* by William Miller Ph.D. and Stephen Rollnick Ph.D. "MI is a collaborative, goal-oriented style of communication with particular attention to the language of change. It is designed to strengthen personal motivation for and commitment to a specific goal by eliciting and exploring the person's own reasons for change within an atmosphere of acceptance and compassion [95]."
- Practice accountability for yourself and others, and work on a forgiveness process if that feels right for you [96, 129, 53].

structure
function qual-
ity

daring greatly

crucial con-
versations, the
book of for-
giving, fierce
intimacy

mention
transforma-
tive/restorative
justice con-

- _____
- _____
- _____

influence, the psychology of persuasion

boundries, fierce intimacy

Introspection and Self-Improvement: _____

- Carefully consider the things you own. Which ones really make you happy? Which ones are a form of unhealthy escapism or compensation for some hurt? Which ones would you be better off without? _____
- journaling
- meditation
- Consider how you cope with stress. Which behaviors are healthy for you and those around you? Which aren't?
- Consider the factors of Self Determination Theory in Appendix A.5. Are you missing any of these? If so, how can you develop them?

evaluate relationships and which are good/bad for you; severe personality disorders or ambivalence

social comparisons / status, rethinking narcissism, the gifts of imperfection

add scout mindset here

marie kondo

A.7 Things to do in Moments of Anxiety (Short Term Strategies)

These are options, not prescriptions. This list has been carefully ordered to prioritize what is likely to work best, based on statistical averages. I hope there are some things here that will work for you, but everyone is different, and if the nothing on this list works for you, that's something wrong with the list, not something wrong with you.

- Choose distraction over denial: It can delay your healing to deny that you're having feelings at all—which is tempting to do when the feelings are very overwhelming. However, it is also important to feel and process your feelings on your own timeline; when you are feeling more than you are ready to feel, distraction can be extremely nourishing and helpful. Can you focus on self-care tasks? Is this the moment to organize your art supplies or clean out your closet? Is there some form of work that feels accessible and/or good to you right now? Or maybe there's a movie or a video game or an Instagram or a comic that can provide a great distraction.
- Exercise is the single most efficient way to complete the stress cycle and release anxiety. Try vigorous exercises (like jumping jacks) for a short period of time. If that isn't your jam, slow, gentle options like trauma informed yoga [52], tai chi, or a walk outside can also be extremely effective.
- This deep muscle relaxation technique [94] is a great second choice with good data behind it.
- It might be possible to reduce your anxiety by stopping your physical anxiety response, which can then help your emotions mellow out. Physiological hacks that do this are called vagal maneuvers [36]. One of them is the "diving reflex," which some people can activate by holding ice packs (or, in a pinch, a cold drink or two) to their face.

- If you have access, maybe spend time in nature—people consistently underestimate how much better time in green space makes us feel. Even five minutes can have a measurable positive impact.
- If you are not currently experiencing physical danger, paying attention to what your body is experiencing right now may help create calm. One option is to name (out loud or in your head) what you are experiencing through your senses: I am hearing _____, I am seeing _____, I am touching _____, I am smelling _____, I am tasting _____.
- Try meditation or deep breathing exercises (See Sections [A.4](#) or [A.3](#)). Meditation can be a way of centering on physicality—you can try a body scan meditation, or a sensory focus meditation, or a myriad of guided meditation options.
- Connect with your friends: Start a text thread or a phone tree among your group. Share memes. Ask friends to send memes that remind them of you. Ask someone if they can just sit with you on a phone or video call, or read you some favorite children’s stories.
- Seek physical affection: Gottman recommends a daily mindful six-second kiss. Hugging someone you trust for a full 20 seconds is also good. Pets or self-massage can also help.
- Ask for help. Asking for help can be incredibly vulnerable. Also, asking for specific things can be very hard when you’re in distress—and it can feel pointless when you feel very deeply as though nothing could possibly make you feel better. However, even small gestures of support can make things a little more bearable until your body-mind is able to start healing itself. On top of this, giving people who care about you and/or want to support you some kind of job to do can build closeness and help them feel less powerless. Consider asking for: food, water or tea, someone to sit with you, someone to watch a movie with you, someone to text you on the hour and remind you that you are a badass.
- Sleep it off: if you’re able to *get* to sleep, the schema will very likely be de-escalated when you wake up.
- If you are ready, or it feels like an OK idea, try writing about your experience. Writing has been shown incredibly effective at helping to address trauma—even fifteen minutes spent writing about trauma on one occasion has lasting positive impacts.
- Write about your experience using your non-dominant hand.
- Write or tell a friend or supportive stranger your “shitty first draft”—the story you’re telling yourself about who you are, what you’ve experienced, and what it all means.
- Alternately: make art. If you don’t want to make art about bad things you’ve experienced, try making art about something that has been comforting and nourishing to you in the past, or about what you wish you could escape to, or about what you are fighting for.

- Try a harm reduction approach when it comes to coping with substances:
 - Avoid using alone - if you are isolated, call a friend on the phone.
 - Avoid mixing substances or stacking doses
 - If you have been trying not to use substances or otherwise numb your feelings, and you end up doing numbing behaviors you had wanted to avoid, try to be very gentle with yourself. You always have a fundamental worth and dignity no matter what.

A.8 Challenging Beliefs Worksheet

Does this method work because of Common Factors? If so, is it useful without a therapist?

Challenging Beliefs Worksheet

[illegible]

A.9 Suggested Avenues of Future Research

- Pursuing the suggestions from Aday et al. [1] and Elk and Fried [40] for designing more rigorous MDMA-therapy trials. Of particular note is blinding, with most participants accurately deducing whether they got MDMA or placebo. Expectancy effects were also high and may have heightened the problems from lack of adequate blinding.
- What can improve reconsolidation (likely high quality sleep, for instance)? What is the nature of reconsolidation exhaustion?
- How does classic psychedelic-assisted psychotherapy work? Does it make implicit schemas explicit, facilitating later spontaneous mismatching? We know it increases neural plasticity.
- MG and others (personal communications) have reported increased capacity to complete reconsolidation work sober, phenomenologically and consequentially similar to what they experienced during MDMA-therapy sessions. This appears to be connected to extensive prior MDMA-therapy. How does this work? How can it be promoted?
- What is the lowest effective dose? How should dosing account for body weight and various health conditions?
- How often is it safe to use MDMA given that there are large negative health effects associated with prolonging mental illness?
- Is it possible to create a manualized self-treatment protocol that is safe and effective?
- How safe is solo MDMA-therapy?
- When, how much, and why do some psychedelics cause short term cognitive impairment?
- How can therapists best prepare their clients for solo at-home sessions (e.g. for sex therapy)?

References

- [1] Jacob S Aday et al. “Great Expectations: recommendations for improving the methodological rigor of psychedelic clinical trials”. In: *Psychopharmacology* (2022). DOI: <https://doi.org/10.1007/s00213-022-06123-7>.
- [2] Norberto Aguirre et al. “ α -Lipoic acid prevents 3, 4-methylenedioxy-methamphetamine (MDMA)-induced neurotoxicity”. In: *Neuroreport* (1999). DOI: <https://doi.org/10.1097/00001756-199911260-00039>.
- [3] Seyyed Salman Alavi et al. “Behavioral addiction versus substance addiction: Correspondence of psychiatric and psychological views”. In: *International journal of preventive medicine* (2012). URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3354400/pdf/IJPVM-3-290.pdf>.

- [4] Michelle Alexander. *The New Jim Crow: Mass Incarceration in the Age of Colorblindness*. The New Press, 2022. URL: <https://thenewpress.com/books/new-jim-crow>.
- [5] Psychedelic Alpha. *Psychedelic Legalization & Decriminalization Tracker*. 2023. URL: <https://psychedelicalpha.com/data/psychedelic-laws>.
- [6] E Alves et al. “Acetyl-L-carnitine provides effective in vivo neuroprotection over 3, 4-methylenedioxymethamphetamine-induced mitochondrial neurotoxicity in the adolescent rat brain”. In: *Neuroscience* (2009). DOI: <https://doi.org/10.1016/j.neuroscience.2008.10.041>.
- [7] Jan van Amsterdam et al. “Hard boiled: alcohol use as a risk factor for MDMA-induced hyperthermia: a systematic review”. In: *Neurotoxicity research* (2021). DOI: <https://doi.org/10.1007/s12640-021-00416-z>.
- [8] Chittaranjan Andrade. “The practical importance of half-life in psychopharmacology”. In: *The Journal of Clinical Psychiatry* (2022). DOI: <https://doi.org/10.4088/JCP.22f14584>.
- [9] K. Andrews et al. *Background to the New York Declaration on Animal Consciousness*. 2024. URL: <nydeclaration.com>.
- [10] Amanda Askill. *Self-Serving utilitarian arguments*. 2021. URL: <https://www.askell.blog/self-serving-utilitarian-arguments/>.
- [11] Ryan Bailey and Jose Pico. “Defense mechanisms”. In: *StatPearls [Internet]*. StatPearls Publishing, 2023. URL: <https://www.ncbi.nlm.nih.gov/books/NBK559106/>.
- [12] Melinda J Barker et al. “Persistence of cognitive effects after withdrawal from long-term benzodiazepine use: a meta-analysis”. In: *Archives of Clinical Neuropsychology* (2004). DOI: [10.1016/S0887-6177\(03\)00096-9](https://doi.org/10.1016/S0887-6177(03)00096-9).
- [13] Daniel Baxter and Luc Pelletier. “Is Nature Relatedness a Basic Human Psychological Need? A Critical Examination of the Extant Literature”. In: *Canadian Psychology/Psychologie canadienne* (2018). DOI: [10.1037/cap0000145](https://doi.org/10.1037/cap0000145). URL: https://www.researchgate.net/publication/324552924_Is_Nature_Relatedness_a_Basic_Human_Psychological_Need_A_Critical_Examination_of_the_Extant_Literature.
- [14] Gillinder Bedi, David Hyman, and Harriet de Wit. “Is Ecstasy an “Empathogen”? Effects of \pm 3,4-Methylenedioxymethamphetamine on Prosocial Feelings and Identification of Emotional States in Others”. In: *Biological Psychiatry* (2010). DOI: <https://doi.org/10.1016/j.biopsych.2010.08.003>. URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2997873/pdf/nihms229652.pdf>.
- [15] Valerie Bentz and Jeremy Shapiro. *Mindful Inquiry in Social Research*. SAGE, 1998. URL: <https://us.sagepub.com/en-us/nam/mindful-inquiry-in-social-research/book7789#contents>.
- [16] Laís F Berro et al. “Acute effects of 3, 4-methylenedioxymethamphetamine (MDMA) and R (-) MDMA on actigraphy-based daytime activity and sleep parameters in rhesus monkeys.” In: *Experimental and clinical psychopharmacology* (2018). DOI: doi.org/10.1037/pha0000196.

- [17] Alessandro Bonsignore et al. “MDMA induced cardio-toxicity and pathological myocardial effects: A systematic review of experimental data and autopsy findings”. In: *Cardiovascular toxicology* (2019). DOI: <https://doi.org/10.1007/s12012-019-09526-9>.
- [18] P. R. Breggin. *Psychiatric drug withdrawal: A guide for prescribers, therapists, patients, and their families*. Springer Publishing Co., 2013. URL: <https://www.springerpub.com/psychiatric-drug-withdrawal-9780826108432.html>.
- [19] Brené Brown. *Daring greatly: How the courage to be vulnerable transforms the way we live, love, parent, and lead*. Penguin, 2015. URL: <https://brenebrown.com/book/daring-greatly/>.
- [20] Brené Brown. *I thought it was just me (but it isn't): Telling the truth about perfectionism, inadequacy, and power*. Avery, 2007. URL: <https://www.penguinrandomhouse.com/books/298649/i-thought-it-was-just-me-but-it-isnt-by-brene-brown/>.
- [21] Daniel P Brown and David S Elliott. *Attachment disturbances in adults: Treatment for comprehensive repair*. WW Norton & Co, 2016. URL: <https://wnorton.com/books/9780393711523>.
- [22] Tibor M Brunt et al. “Linking the pharmacological content of ecstasy tablets to the subjective experiences of drug users”. In: *Psychopharmacology* (2012). DOI: <https://doi.org/10.1007/s00213-011-2529-4>.
- [23] Andrea Cipriani et al. “Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: a systematic review and network meta-analysis”. In: *The Lancet* (2018). DOI: [https://doi.org/10.1016/S0140-6736\(17\)32802-7](https://doi.org/10.1016/S0140-6736(17)32802-7).
- [24] Isaac V. Cohen et al. “Concomitant drugs associated with increased mortality for MDMA users reported in a drug safety surveillance database”. In: *Scientific Reports* (2021). DOI: [10.1038/s41598-021-85389-x](https://doi.org/10.1038/s41598-021-85389-x).
- [25] Coherence Psychology Institute. *Coherence Psychology Institute Referral Directory*. 2024. URL: <https://coherencetherapy.org/prac/directory-terms.php>.
- [26] Robert Colbert and Shannon Hughes. “Evenings with Molly: Adult Couples’ Use of MDMA for Relationship Enhancement”. In: *Culture, Medicine, and Psychiatry* (2023). DOI: <https://doi.org/10.1007/s11013-021-09764-z>.
- [27] Julia Colcott et al. “Side-effects of MDMA-assisted Psychotherapy: A Systematic Review and Meta-analysis”. In: *Neuropsychopharmacology* (2024). DOI: <https://doi.org/10.1038/s41386-024-01865-8>.
- [28] R. Coleman. *Psychedelic Psychotherapy, a User Friendly Guide to Psychedelic Drug Assisted Therapy*. Synergetic Press, 2017. URL: <https://synergeticpress.com/catalog/psychedelic-psychotherapy/>.
- [29] The Cut. *‘I See a Sh*tshow’: Inside the World of Psychedelic Therapy*. 2021. URL: <https://www.thecut.com/2021/11/cover-story-podcast-goes-into-world-of-psychedelic-therapy.html>.

- [30] DanceSafe. *MDMA Testing Kit*. 2023. URL: <https://dancesafe.org/product/mdma-testing-kit>.
- [31] Draulio B De Araujo et al. "Seeing with the eyes shut: Neural basis of enhanced imagery following ayahuasca ingestion". In: *Human brain mapping* (2012). DOI: <https://doi.org/10.1002/hbm.21381>.
- [32] R De la Torre et al. "Pharmacology of MDMA in humans". In: *Annals of the New York Academy of Sciences* (2000). DOI: <https://doi.org/10.1111/j.1749-6632.2000.tb05199.x>. URL: <https://www.jcamie.eu/documents/Pharmacology%20of%20MDMA%20in%20Humans.%20Ann%20New%20York%20Acad%20Sci.pdf>.
- [33] Rafael De La Torre et al. "MDMA, methamphetamine, and CYP2D6 pharmacogenetics: what is clinically relevant?" In: *Frontiers in Genetics* (2012). DOI: [10.3389/fgene.2012.00235](https://doi.org/10.3389/fgene.2012.00235).
- [34] Rafael De La Torre et al. "Non-linear pharmacokinetics of MDMA ('ecstasy') in humans". In: *British journal of clinical pharmacology* (2000). DOI: <https://doi.org/10.1046/j.1365-2125.2000.00121.x>.
- [35] Edward L. Deci and Richard M. Ryan. "The "What" and "Why" of Goal Pursuits: Human Needs and the Self-Determination of Behavior". In: *Psychological Inquiry* (2000). DOI: [10.1207/S15327965PLI1104_01](https://doi.org/10.1207/S15327965PLI1104_01). URL: https://selfdeterminationtheory.org/SDT/documents/2000_DeciRyan_PIWhatWhy.pdf.
- [36] Mary Jo DiLonardo. *What Is the Valsalva Maneuver?* 2023. URL: <https://www.webmd.com/heart-disease/atrial-fibrillation/valsalva-maneuver#1>.
- [37] Manoj K Doss et al. "MDMA impairs both the encoding and retrieval of emotional recollections". In: *Neuropsychopharmacology* (2018). DOI: <https://doi.org/10.1007/s00213-006-0457-5>.
- [38] Bruce Ecker, Robin Ticic, and Laurel Hulley. *Unlocking the Emotional Brain: Memory Reconsolidation and the Psychotherapy of Transformational Change*. Taylor & Francis, 2024. DOI: <https://doi.org/10.4324/9781003231431>. URL: <https://www.routledge.com/Unlocking-the-Emotional-Brain-Eliminating-Symptoms-at-Their-Roots-Using/Ecker-Ticic-Hulley/p/book/9781032117539>.
- [39] Amber N Edinoff et al. "Clinically Relevant Drug Interactions with Monoamine Oxidase Inhibitors". In: *Health Psychology Research* (2022). DOI: <https://doi.org/10.52965/2F001c.39576>.
- [40] M. van Elk and E. I. Fried. "History repeating: A roadmap to address common problems in psychedelic science". In: (2023). DOI: <https://doi.org/10.31234/osf.io/ak6gx>.
- [41] Kathy Emde. "MDMA (Ecstasy) in the emergency department". In: *Journal of Emergency Nursing* (2003). DOI: [https://doi.org/10.1016/S0099-1767\(03\)00292-7](https://doi.org/10.1016/S0099-1767(03)00292-7).
- [42] Jules Evans. *What are the ethical implications of the Ben Sessa case?* 2024. URL: <https://www.ecstaticintegration.org/p/what-are-the-ethical-implications>.

- [43] Jules Evans et al. “Extended difficulties following the use of psychedelic drugs: A mixed methods study”. In: *Plos one* (2023). DOI: <https://doi.org/10.1371/journal.pone.0293349>.
- [44] Ricarda Evens et al. “The psychedelic afterglow phenomenon: a systematic review of subacute effects of classic serotonergic psychedelics”. In: *Therapeutic Advances in Psychopharmacology* (2023). PMID: 37284524. DOI: [10.1177/20451253231172254](https://doi.org/10.1177/20451253231172254). URL: <https://doi.org/10.1177/20451253231172254>.
- [45] Magí Farré et al. “Human pharmacology of 3, 4-methylenedioxymethamphetamine (MDMA, ecstasy) after repeated doses taken 4 h apart Human pharmacology of MDMA after repeated doses taken 4 h apart”. In: *European neuropsychopharmacology* (2015). DOI: <https://doi.org/10.1016/j.euroneuro.2015.05.007>.
- [46] Allison A. Feduccia and Michael C. Mithoefer. “MDMA-assisted psychotherapy for PTSD: Are memory reconsolidation and fear extinction underlying mechanisms?” In: *Progress in Neuro-Psychopharmacology and Biological Psychiatry* (2018). DOI: <https://doi.org/10.1016/j.pnpbp.2018.03.003>.
- [47] Allison A. Feduccia et al. “Discontinuation of medications classified as reuptake inhibitors affects treatment response of MDMA-assisted psychotherapy”. In: *Psychopharmacology* (2021). DOI: [10.1007/s00213-020-05710-w](https://doi.org/10.1007/s00213-020-05710-w).
- [48] Vincent J Felitti et al. “Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The Adverse Childhood Experiences (ACE) Study”. In: *American journal of preventive medicine* (1998). DOI: [https://doi.org/10.1016/S0749-3797\(98\)00017-8](https://doi.org/10.1016/S0749-3797(98)00017-8).
- [49] Myriam Forster et al. “The relationship between family-based adverse childhood experiences and substance use behaviors among a diverse sample of college students”. In: *Addictive behaviors* (2018). DOI: <https://doi.org/10.1016/j.addbeh.2017.08.037>.
- [50] Matthias Forstmann et al. “Among psychedelic-experienced users, only past use of psilocybin reliably predicts nature relatedness”. In: *Journal of Psychopharmacology* (2023). DOI: [10.1177/02698811221146356](https://doi.org/10.1177/02698811221146356).
- [51] Julia Galef. *The Scout Mindset*. Penguin, 2021. URL: <https://www.penguinrandomhouse.com/books/555240/the-scout-mindset-by-julia-galef/>.
- [52] Nityda Gessel. *Gentle & Trauma-Sensitive Yoga Sequence for Grounding and Upper Body Release*. 2018. URL: <https://www.youtube.com/watch?v=V10txPbFAec>.
- [53] John M Gottman. *The science of trust: Emotional attunement for couples*. WW Norton & Company, 2011. URL: <https://wnorton.com/books/9780393705959>.
- [54] Euphrosyne Gouzoulis-Mayfrank and Joerg Daumann. “Neurotoxicity of methylenedioxymethamphetamines (MDMA; ecstasy) in humans: how strong is the evidence for persistent brain damage?” In: *Addiction* (2006). DOI: [10.1111/j.1360-0443.2006.01314.x](https://doi.org/10.1111/j.1360-0443.2006.01314.x).

- [55] Warwick M Green et al. “MDMA Assisted Psychotherapy Decreases PTSD Symptoms, Dissociation, Functional Disability, and Depression: A Systematic Review and Meta-Analysis”. In: *medRxiv* (2023). DOI: <https://doi.org/10.1101/2023.08.17.23293955>.
- [56] Greenspace. *Brief Revised Working Alliance Inventory*. 2023. URL: <https://greenspacehealth.com/en-us/br-wai/>.
- [57] Roland R Griffiths et al. “Psilocybin can occasion mystical-type experiences having substantial and sustained personal meaning and spiritual significance”. In: *Psychopharmacology* (2006). DOI: <https://doi.org/10.1007/s00213-006-0457-5>. URL: https://www.researchgate.net/profile/Roland-Griffiths/publication/235984074_Psilocybin_Can_Occasion_Mystical-Type_Experiences_Having_Substantial_and_Sustained_Personal_Meaning_and_Spiritual_Significance/links/0c96051dc66a9d4465000000/Psilocybin-Can-Occasion-Mystical-Type-Experiences-Having-Substantial-and-Sustained-Personal-Meaning-and-Spiritual-Significance.pdf.
- [58] Will Hall. *Ending The Silence Around Psychedelic Therapy Abuse*. 2021. URL: <https://www.madinamerica.com/2021/09/ending-silence-psychedelic-therapy-abuse/>.
- [59] John H. Halpern, Arturo G. Lerner, and Torsten Passie. “A Review of Hallucinogen Persisting Perception Disorder (HPPD) and an Exploratory Study of Subjects Claiming Symptoms of HPPD”. In: *Behavioral Neurobiology of Psychedelic Drugs*. Ed. by Adam L. Halberstadt, Franz X. Vollenweider, and David E. Nichols. Berlin, Heidelberg: Springer Berlin Heidelberg, 2018. ISBN: 978-3-662-55880-5. DOI: [10.1007/7854_2016_457](https://doi.org/10.1007/7854_2016_457). URL: <https://static1.squarespace.com/static/5ab926f8a9e0287fbf928015/t/5b119523575d1fe491ef8383/1527878948945/HPPD.pdf>.
- [60] John H. Halpern et al. “Residual neurocognitive features of long-term ecstasy users with minimal exposure to other drugs”. In: *Addiction* (2011). DOI: <https://doi.org/10.1111/j.1360-0443.2010.03252.x>. URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3053129/pdf/nihms-247953.pdf>.
- [61] Ido Hartogsohn. “The meaning-enhancing properties of psychedelics and their mediator role in psychedelic therapy, spirituality, and creativity”. In: *Frontiers in neuroscience* (2018). DOI: <https://doi.org/10.3389/fnins.2018.00129>.
- [62] Peter Henningsen, Thomas Zimmermann, and Heribert Sattel. “Medically unexplained physical symptoms, anxiety, and depression: a meta-analytic review”. In: *Psychosomatic medicine* (2003). DOI: [10.1097/01.PSY.0000075977.90337.E7](https://doi.org/10.1097/01.PSY.0000075977.90337.E7).
- [63] Jack Hills. “Phenomenology of MDMA Solo Sessions”. PhD thesis. Antioch University, 2023. URL: https://etd.ohiolink.edu/acprod/odb_etd/ws/send_file/send?accession=antioch1690227460998206&disposition=inline.
- [64] Julianne Holt-Lunstad, Timothy B. Smith, and J. Bradley Layton. “Social Relationships and Mortality Risk: A Meta-analytic Review”. In: *PLOS Medicine* (2010). DOI: [10.1371/journal.pmed.1000316](https://doi.org/10.1371/journal.pmed.1000316).

- [65] Cheetah House. *Resources and Support for Adverse Meditative Experiences*. 2024. URL: <https://www.cheetahhouse.org>.
- [66] L. Jerome, A.A. Feduccia, J.B. Wang, et al. “Long-term follow-up outcomes of MDMA-assisted psychotherapy for treatment of PTSD: a longitudinal pooled analysis of six phase 2 trials.” In: *Psychopharmacology* (2020). DOI: <https://doi.org/10.1007/s00213-020-05548-2>.
- [67] Grant Jones, Diego Arias, and Matthew Nock. “Associations between MDMA/ecstasy, classic psychedelics, and suicidal thoughts and behaviors in a sample of US adolescents”. In: *Scientific reports* (2022). DOI: <https://doi.org/10.1038/s41598-022-25658-5>.
- [68] Grant M Jones and Matthew K Nock. “MDMA/ecstasy use and psilocybin use are associated with lowered odds of psychological distress and suicidal thoughts in a sample of US adults”. In: *Journal of psychopharmacology* (2022). DOI: <https://doi.org/10.1177/02698811211058923>.
- [69] Jon Kabat-Zinn. *Full Catastrophe Living: Using the Wisdom of Your Body and Mind to Face Stress, Pain, and Illness*. Bantam, 2013. URL: <https://www.penguinrandomhouse.com/books/89149/full-catastrophe-living-revised-edition-by-jon-kabat-zinn-preface-by-thich-nhat-hanh-foreword-by-joan-borysenko/>.
- [70] Samuli Kangaslampi and Josjan Zijlmans. “MDMA-assisted psychotherapy for PTSD in adolescents: Rationale, potential, risks, and considerations”. In: *European child & adolescent psychiatry* (2023). DOI: doi.org/10.1007/s00787-023-02310-9.
- [71] Ann Kelley and Sue Marriott. *Therapists Uncensored*. 2024. URL: <https://therapistuncensored.com>.
- [72] Bassam Khoury et al. “Mindfulness-based stress reduction for healthy individuals: A meta-analysis”. In: *Journal of Psychosomatic Research* (2015). DOI: <https://doi.org/10.1016/j.jpsychores.2015.03.009>. URL: <https://r.jordan.im/download/mindfulness/khoury2015.pdf>.
- [73] Jueun Kim et al. “Ecstasy use and suicidal behavior among adolescents: Findings from a national survey”. In: *Suicide and Life-Threatening Behavior* (2011). DOI: <https://doi.org/10.1111/j.1943-278X.2011.00043.x>.
- [74] Bessel van der Kolk. *The Body Keeps the Score*. Penguin Books, 2015. URL: <https://www.penguinrandomhouse.com/books/313183/the-body-keeps-the-score-by-bessel-van-der-kolk-md/>.
- [75] Bessel A van der Kolk et al. “Effects of MDMA-assisted therapy for PTSD on self-experience”. In: *PLoS One* (2024). DOI: <https://doi.org/10.1371/journal.pone.0295926>.
- [76] Kasia Kozłowska et al. “Fear and the defense cascade: clinical implications and management”. In: *Harvard Review of Psychiatry* (2015). DOI: [10.1097/HRP.0000000000000065](https://doi.org/10.1097/HRP.0000000000000065).

- [77] Kurt Kroenke. “Patients presenting with somatic complaints: epidemiology, psychiatric co-morbidity and management”. In: *International journal of methods in psychiatric research* (2003). DOI: <https://doi.org/10.1002/mpr.140>.
- [78] Richard D. Lane et al. “Memory reconsolidation, emotional arousal, and the process of change in psychotherapy: New insights from brain science”. In: *Behavioral and Brain Sciences* (2015). DOI: [10.1017/S0140525X14000041](https://doi.org/10.1017/S0140525X14000041). URL: http://sequentialpsychotherapy.com/assets/bbs_lane-et-al.pdf.
- [79] Jeppe Bo Lauersen, Ditte Marie Bertelsen, and Lars Bo Andersen. “The effectiveness of exercise interventions to prevent sports injuries: a systematic review and meta-analysis of randomised controlled trials”. In: *British Journal of Sports Medicine* (2014). DOI: [10.1136/bjsports-2013-092538](https://doi.org/10.1136/bjsports-2013-092538).
- [80] Daniel Lieberman. *Exercised: Why something we never evolved to do is healthy and rewarding*. Vintage, 2021. URL: <https://scholar.harvard.edu/exercised/home>.
- [81] Matthias E Liechti, Alex Gamma, and Franz X Vollenweider. “Gender differences in the subjective effects of MDMA”. In: *Psychopharmacology* (2001). DOI: <https://doi.org/10.1007/s002130000648>.
- [82] Sophia Linguiti et al. “Functional imaging studies of acute administration of classic psychedelics, ketamine, and MDMA: Methodological limitations and convergent results”. In: *Neuroscience & Biobehavioral Reviews* (2023). DOI: <https://doi.org/10.1016/j.neubiorev.2023.105421>.
- [83] Qing Liu, Jian Zhu, and Wenjuan Zhang. “The efficacy of mindfulness-based stress reduction intervention 3 for post-traumatic stress disorder (PTSD) symptoms in patients with PTSD: A meta-analysis of four randomized controlled trials”. In: *Stress and Health* (2022). DOI: <https://doi.org/10.1002/smi.3138>.
- [84] Richard J Loewenstein. “Dissociation debates: Everything you know is wrong”. In: *Dialogues in clinical neuroscience* (2018). DOI: <https://doi.org/10.31887/DCNS.2018.20.3/rloewenstein>.
- [85] Steven Low. *r/bodyweightfitness Recommended Routine*. 2023. URL: https://www.reddit.com/r/bodyweightfitness/wiki/kb/recommended_routine/.
- [86] Benjamin Malcolm and Kelan Thomas. “Serotonin toxicity of serotonergic psychedelics”. In: *Psychopharmacology* (2022). DOI: <https://doi.org/10.1007/s00213-021-05876-x>.
- [87] B Mallinckrodt and YT Tekie. “Revision of the Working Alliance Inventory and development of a brief revised version guided by item response theory”. In: *Psychotherapy Research* (2015).
- [88] Rebecca Martinez. *Accountability & Transformative Justice in the Psychedelic Space: A Roadmap for Change*. 2021. URL: <https://psychedelicstoday.com/2021/06/09/accountability-transformative-justice-in-the-psychedelic-space-a-roadmap-for-change/>.
- [89] Kelly McGonigal. *The Upside of Stress, Why Stress Is Good for You, and How to Get Good at It*. Avery, 2015. URL: <http://kellymcgonigal.com/books>.

- [90] Philip McGuire and Tom Fahy. “Chronic paranoid psychosis after misuse of MDMA ("ecstasy")”. In: *BMJ: British Medical Journal* (1991).
- [91] Sarah McNamee, Neşe Devenot, and Meaghan Buisson. “Studying Harms Is Key to Improving Psychedelic-Assisted Therapy—Participants Call for Changes to Research Landscape”. In: *JAMA Psychiatry* (2023). DOI: [10.1001/jamapsychiatry.2023.0099](https://doi.org/10.1001/jamapsychiatry.2023.0099).
- [92] David McRaney. *How Minds Change: The Surprising Science of Belief, Opinion, and Persuasion*. Portfolio, 2022. URL: <https://www.davidmcraney.com/howmindschangehome>.
- [93] Stanford Medicine. *Cognitive Behavioral Therapy for Insomnia*. URL: <https://stanfordhealthcare.org/medical-treatments/c/cognitive-behavioral-therapy-insomnia/procedures.html>.
- [94] University of Michigan Health. *Stress Management: Doing Progressive Muscle Relaxation*. 2023. URL: <https://www.uofmhealth.org/health-library/uz2225>.
- [95] William R. Miller and Stephen Rollnick. *Motivational Interviewing, Helping People Change and Grow*. Guilford Press, 2023. URL: <https://www.guilford.com/books/Motivational-Interviewing/Miller-Rollnick/9781462552795>.
- [96] Mia Mingus. *The Four Parts of Accountability & How To Give A Genuine Apology*. 2019. URL: <https://leavingevidence.wordpress.com/2019/12/18/how-to-give-a-good-apology-part-1-the-four-parts-of-accountability/>.
- [97] J.M. Mitchell, G. M. Ot’alora, B. van der Kolk, et al. “MDMA-assisted therapy for moderate to severe PTSD: a randomized, placebo-controlled phase 3 trial.” In: *Nat Med* (2023). DOI: <https://doi.org/10.1038/s41591-023-02565-4>.
- [98] Jennifer M. Mitchell et al. “MDMA-assisted therapy for severe PTSD: a randomized, double-blind, placebo-controlled phase 3 study”. In: *Nature Medicine* (2021). DOI: [10.1038/s41591-021-01336-3](https://doi.org/10.1038/s41591-021-01336-3).
- [99] M. Mithoefer and A. Mithoefer. “MDMA”. In: *Handbook of Medical Hallucinogens*. Ed. by C. S. Grob and J. Grigsby. The Guilford Press, 2021.
- [100] Michael Mithoefer. “A Manual for MDMA-Assisted Psychotherapy in the Treatment of Posttraumatic Stress Disorder”. In: (2017). URL: <https://maps.org/2014/01/27/a-manual-for-mdma-assisted-therapy-in-the-treatment-of-ptsd/>.
- [101] Romain Nardou et al. “Oxytocin-dependent reopening of a social reward learning critical period with MDMA”. In: *Nature* (2019). DOI: [10.1038/s41586-019-1075-9](https://doi.org/10.1038/s41586-019-1075-9).
- [102] Romain Nardou et al. “Psychedelics reopen the social reward learning critical period”. In: *Nature* (2023). DOI: <https://doi.org/10.1038/s41586-023-06204-3>.
- [103] Christopher R. Nicholas et al. “The effects of MDMA-assisted therapy on alcohol and substance use in a phase 3 trial for treatment of severe PTSD”. In: *Drug and Alcohol Dependence* (2022). DOI: <https://doi.org/10.1016/j.drugalcdep.2022.109356>.

- [104] David Nutt, Leslie King, and Lawrence Phillips. “Drug harms in the UK: A multi-criterion decision analysis”. In: *Lancet* (2010). DOI: [https://doi.org/10.1016/S0140-6736\(10\)61462-6](https://doi.org/10.1016/S0140-6736(10)61462-6). URL: https://www.researchgate.net/publication/285843262_Drug_harms_in_the_UK_A_multi-criterion_decision_analysis.
- [105] David J Nutt. *Equasy—an overlooked addiction with implications for the current debate on drug harms*. 2009. DOI: <https://doi.org/10.1177/0269881108099672>.
- [106] Merlijn Olthof et al. “Destabilization in self-ratings of the psychotherapeutic process is associated with better treatment outcome in patients with mood disorders”. In: *Psychotherapy Research* (2020). DOI: [10.1080/10503307.2019.1633484](https://doi.org/10.1080/10503307.2019.1633484).
- [107] OpenAI. *Introducing ChatGPT Plus*. 2023. URL: <https://openai.com/blog/chatgpt-plus>.
- [108] World Health Organization. “Depression and Other Common Mental Disorders: Global Health Estimates.” In: (2017). URL: <https://apps.who.int/iris/bitstream/handle/10665/254610/WHO-MSD-MER-2017.2-eng.pdf>.
- [109] Marcela Ot’alora. *MDMA for the treatment of trauma: a clinical trial therapist’s perspective - Marcela Ot’alora G.* 2021. URL: <https://www.youtube.com/watch?v=3vsxlaaMs2Q>.
- [110] UC Berkley Museum of Paleontology. *Understanding Science 101*. 2024. URL: <https://undsci.berkeley.edu/understanding-science-101/a-scientific-approach-to-life-a-science-toolkit/>.
- [111] AC Parrott. “Chronic tolerance to recreational MDMA (3, 4-methylenedioxymethamphetamine) or Ecstasy”. In: *Journal of Psychopharmacology* (2005). DOI: <https://doi.org/10.1177/0269881105048900>. URL: <https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=e40bc214ac26b35b1ba63fccaec8f64e97d4385d>.
- [112] Psychedelic Passage. *Your network of trusted psychedelic concierges and guides*. 2023. URL: <https://www.psychedelicoptassage.com>.
- [113] Torsten Passie. *The History of MDMA*. Oxford University Press, 2023. URL: <https://global.oup.com/academic/product/the-history-of-mdma-9780198867364?cc=us&lang=en&>.
- [114] Torsten Passie, Jeffrey Guss, and Rainer Krähenmann. “Lower-dose psycholytic therapy - A neglected approach”. In: *Frontiers in Psychiatry* (2022). DOI: [10.3389/fpsy.2022.1020505](https://doi.org/10.3389/fpsy.2022.1020505).
- [115] Ankit Patel et al. “Persistent psychosis after a single ingestion of " Ecstasy"(MDMA)”. In: *The Primary Care Companion for CNS Disorders* (2011).
- [116] Heidi Priebe. *Attachment Theory*. 2024. URL: <https://www.youtube.com/playlist?list=PLFD8kQWqivS1fy9uwssBE-H5M668fmtFt>.
- [117] Anthony Principe. “Neuropharmacological analysis of the anti-addictive and therapeutic effects of psilocybin”. In: *SURG Journal* (2022). DOI: <https://doi.org/10.21083/surg.v14i1.6870>.

- [118] Fireside Project. *The Psychedelic Peer Support Line provides emotional support during and after psychedelic experiences*. 2023. URL: <https://firesideproject.org>.
- [119] The Attachment Project. *Attachment Style Test*. 2023. URL: <https://quiz.attachmentproject.com>.
- [120] The Attachment Project. *The Ultimate Guide to Early Maladaptive Schemas [Full List]*. 2024. URL: <https://www.attachmentproject.com/blog/early-maladaptive-schemas/>.
- [121] Karen Pryor. *Don't shoot the dog: The art of teaching and training*. Simon & Schuster, 2019. URL: <https://www.simonandschuster.com/books/Dont-Shoot-the-Dog/Karen-Pryor/9781982106461>.
- [122] Psychedelic Somatic Institute. *PSIP Therapist Directory*. 2024. URL: <https://www.psychedelicomatic.org/find-a-psychedelic-therapist>.
- [123] Multidisciplinary Association for Psychedelic Studies. *Psychedelic Integration List, Mental Health Support Practitioners by Location*. 2023. URL: <https://integration.maps.org>.
- [124] Multidisciplinary Association for Psychedelic Studies. *The MAPS Psychedelic Integration Workbook*. 2024. URL: /var/folders/gv/zzbd59cn7411d3rk0_pj8hrr0000gn/T/com.apple.useractivityd/shared-pasteboard/items/D5A06241-FE6D-47B4-8E95-6F11906AB5E3/MAPS-Integration-Workbook.pdf.
- [125] Stanley Rachman. “Emotional processing”. In: *Behaviour research and therapy* (1980). DOI: [https://doi.org/10.1016/0005-7967\(80\)90069-8](https://doi.org/10.1016/0005-7967(80)90069-8).
- [126] Anya Ragnhildstveit et al. “Cannabis-assisted psychotherapy for complex dissociative posttraumatic stress disorder: A case report”. In: *Frontiers in psychiatry* (2023). DOI: <https://doi.org/10.3389/fpsy.2023.1051542>.
- [127] Saj Razvi. *Psychedelic Somatic Interactional Psychotherapy*. 2022. URL: <https://www.youtube.com/watch?v=N20vPGFCg7M>.
- [128] Saj Razvi and Steven Elfrink. “The PSIP model. An introduction to a novel method of therapy: Psychedelic Somatic Interactional Psychotherapy”. In: *J. Psychedelic Psychiatry* (2020). URL: https://www.journalofpsychedelicpsychiatry.org/_files/ugd/e07c59_d4d1db6fc0174f27bef58a6124aba50e.pdf.
- [129] Barnard Center for Research on Women. *Building Accountable Communities*. 2020. URL: <https://bcrw.barnard.edu/building-accountable-communities/>.
- [130] Valerie F Reyna et al. “How Fuzzy-Trace Theory Predicts True and False Memories for Words, Sentences, and Narratives.” In: *J Appl Res Mem Cogn* (2016). DOI: [10.1016/j.jarmac.2015.12.003](https://doi.org/10.1016/j.jarmac.2015.12.003). URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4815269/>.
- [131] Khary K Rigg and Amanda Sharp. “Deaths related to MDMA (ecstasy/molly): Prevalence, root causes, and harm reduction interventions”. In: *Journal of substance Use* (2018). DOI: <https://doi.org/10.1080/14659891.2018.1436607>.

- [132] Patricia Robledo et al. “Study of the behavioural responses related to the potential addictive properties of MDMA in mice”. In: *Naunyn-Schmiedeberg’s archives of pharmacology* (2004). DOI: <https://doi.org/10.1007/s00210-003-0862-9>.
- [133] rollkit.net. 2023. URL: <https://rollkit.net/mdma-molly-supplement-research/>.
- [134] Kristen E Ronca. “The Impact of Complex Post-Traumatic Stress Disorder and Structural Violence on Children in Impoverished Urban Communities”. PhD thesis. 2018. URL: <https://www.proquest.com/docview/2046909618?pq-origsite=gscholar&fromopenview=true>.
- [135] Marshall Rosenberg. *Nonviolent Communication: A Language of Life*. Puddledancer, 2015. URL: <https://www.nonviolentcommunication.com/product/nvc/>.
- [136] Amanda Roxburgh et al. “Trends in MDMA-related mortality across four countries”. In: *Addiction* (2021). DOI: [doi:10.1111/add.15493](https://doi.org/10.1111/add.15493). URL: https://onlinelibrary.wiley.com/doi/pdf/10.1111/add.15493?casa_token=ydgy7ZYsXgwAAAAA:rjcmIwd8NqcYp7c6Ru_WR1D2LSXdzxVUmeIv3igBnP7bgFIlPPe7wrlghP7dJmtDXfhSWUjbWQLTNUg.
- [137] Carl Sagan. *The Demon-Haunted World, Science as a Candle in the Dark*. Penguin, 1996. URL: <https://www.penguinrandomhouse.com/books/159731/the-demon-haunted-world-by-carl-sagan/>.
- [138] Sarah Saleemi et al. “Who is ‘Molly’? MDMA adulterants by product name and the impact of harm-reduction services at raves”. In: *Journal of Psychopharmacology* (2017). DOI: [10.1177/0269881117715596](https://doi.org/10.1177/0269881117715596).
- [139] Sharon Salzberg. *Lovingkindness, The Revolutionary Art of Happiness*. Shambala, 2018. URL: <https://www.shambhala.com/lovingkindness-15144.html>.
- [140] Aryan Sarparast et al. “Drug-drug interactions between psychiatric medications and MDMA or psilocybin: a systematic review”. In: *Psychopharmacology* (2022). DOI: <https://doi.org/10.1007%2Fs00213-022-06083-y>.
- [141] Sehrish Sayed, Brian M Iacoviello, and Dennis S Charney. “Risk factors for the development of psychopathology following trauma”. In: *Current psychiatry reports* (2015). DOI: <https://doi.org/10.1007/s11920-015-0612-y>.
- [142] Anne K Schlag et al. “Adverse effects of psychedelics: From anecdotes and misinformation to systematic science”. In: *Journal of Psychopharmacology* (2022). DOI: [10.1177/02698811211069100](https://doi.org/10.1177/02698811211069100).
- [143] Richard C Schwartz and Martha Sweezy. *Internal family systems therapy*. Guilford Publications, 2019.
- [144] Richard C Schwartz and Martha Sweezy. *Internal family systems therapy*. Guilford Publications, 2019. URL: <https://www.guilford.com/books/Internal-Family-Systems-Therapy/Schwartz-Sweezy/9781462541461>.
- [145] Alexander J Scott et al. “Improving sleep quality leads to better mental health: A meta-analysis of randomised controlled trials”. In: *Sleep medicine reviews* (2021). DOI: [10.1016/J.SMRV.2021.101556](https://doi.org/10.1016/J.SMRV.2021.101556).

- [146] Mahalakshmi Shankaran, Bryan K Yamamoto, and Gary A Gudelsky. “Ascorbic acid prevents 3, 4-methylenedioxymethamphetamine (MDMA)-induced hydroxyl radical formation and the behavioral and neurochemical consequences of the depletion of brain 5-HT”. In: *Synapse* (2001). DOI: [https://doi.org/10.1002/1098-2396\(200104\)40:1%3C55::aid-syn1026%3E3.0.co;2-o](https://doi.org/10.1002/1098-2396(200104)40:1%3C55::aid-syn1026%3E3.0.co;2-o).
- [147] Masaki Shinfuku et al. “Effectiveness and safety of long-term benzodiazepine use in anxiety disorders: a systematic review and meta-analysis”. In: *International clinical psychopharmacology* (2019). DOI: [10.1097/YIC.0000000000000276](https://doi.org/10.1097/YIC.0000000000000276).
- [148] Katharine C Simon, Rebecca L Gómez, and Lynn Nadel. “Sleep’s role in memory reconsolidation”. In: *Current Opinion in Behavioral Sciences* (2020). DOI: <https://doi.org/10.1016/j.cobeha.2020.04.001>.
- [149] Jessica W Skelley et al. “Use of cannabidiol in anxiety and anxiety-related disorders”. In: *Journal of the American Pharmacists Association* (2020). DOI: [http://dx.doi.org/10.1016/j.japh.2019.11.008](https://doi.org/10.1016/j.japh.2019.11.008).
- [150] Kimberly W Smith et al. “MDMA-assisted psychotherapy for treatment of posttraumatic stress disorder: A systematic review with meta-analysis”. In: *The Journal of Clinical Pharmacology* (2022).
- [151] Kaj Sotala. *Book summary: Unlocking the Emotional Brain*. 2019. URL: <https://www.lesswrong.com/posts/i9xyZBS3qzA8nFXNQ/book-summary-unlocking-the-emotional-brain>.
- [152] Kaj Sotala. *Multiagent Models of Mind*. 2023. URL: <https://www.lesswrong.com/s/ZbmRyDN8TCpBTZSip>.
- [153] Serge A Steenen et al. “Propranolol for the treatment of anxiety disorders: Systematic review and meta-analysis”. In: *Journal of Psychopharmacology* (2016). DOI: [10.1177/0269881115612236](https://doi.org/10.1177/0269881115612236).
- [154] Erich Studerus et al. “Prediction of MDMA response in healthy humans: a pooled analysis of placebo-controlled studies”. In: *Journal of Psychopharmacology* (2021). DOI: [10.1177/0269881121998322](https://doi.org/10.1177/0269881121998322).
- [155] Bihan Tang et al. “A meta-analysis of risk factors for post-traumatic stress disorder (PTSD) in adults and children after earthquakes”. In: *International journal of environmental research and public health* (2017). DOI: <https://doi.org/10.3390/ijerph14121537>.
- [156] Sascha Thal, Liam B. Engel, and Stephen J. Bright. “Presence, Trust, and Empathy: Preferred Characteristics of Psychedelic Carers”. In: *Journal of Humanistic Psychology* (0). DOI: [10.1177/00221678221081380](https://doi.org/10.1177/00221678221081380). URL: <https://doi.org/10.1177/00221678221081380>.
- [157] Jenna M. Traynor et al. “MDMA-Assisted Psychotherapy for Borderline Personality Disorder”. In: *FOCUS* (2022). DOI: [10.1176/appi.focus.20220056](https://doi.org/10.1176/appi.focus.20220056).
- [158] David A Treleaven. *Trauma-sensitive mindfulness: Practices for safe and transformative healing*. WW Norton & Company, 2018.

- [159] David Trickey et al. "A meta-analysis of risk factors for post-traumatic stress disorder in children and adolescents". In: *Clinical Psychology Review* (2012). DOI: <https://doi.org/10.1016/j.cpr.2011.12.001>.
- [160] Caoimhe Twohig-Bennett and Andy Jones. "The health benefits of the great outdoors: A systematic review and meta-analysis of greenspace exposure and health outcomes". In: *Environmental Research* (2018). DOI: <https://doi.org/10.1016/j.envres.2018.06.030>.
- [161] Guillaume Vaiva et al. "An "accidental" acute psychosis with ecstasy use". In: *Journal of psychoactive drugs* (2001).
- [162] Natacha Vanattou-Saifoudine, R McNamara, and A Harkin. "Caffeine provokes adverse interactions with 3, 4-methylenedioxymethamphetamine (MDMA, 'ecstasy') and related psychostimulants: mechanisms and mediators". In: *British journal of pharmacology* (2012). DOI: <https://doi.org/10.1111/j.1476-5381.2012.02065.x>.
- [163] Patrick Vizeli and Matthias E Liechti. "Safety pharmacology of acute MDMA administration in healthy subjects". In: *Journal of Psychopharmacology* (2017). PMID: 28443695. DOI: [10.1177/0269881117691569](https://doi.org/10.1177/0269881117691569).
- [164] Derick T Wade and Peter W Halligan. *The biopsychosocial model of illness: a model whose time has come*. 2017. DOI: <https://doi.org/10.1177/0269215517709890>.
- [165] Caitlin Walker, Clare Donaldson, and Michael Plant. *Global priority: mental health*. 2021. URL: <https://www.happierlivesinstitute.org/wp-content/uploads/2022/04/Global-priority-mental-health.pdf>.
- [166] Bruce E Wampold. "How important are the common factors in psychotherapy? An update". In: *World psychiatry* (2015). DOI: <https://doi.org/10.1002/wps.20238>.
- [167] Mary Beth Williams and Soili Poijula. *The PTSD workbook: Simple, effective techniques for overcoming traumatic stress symptoms*. New Harbinger Publications, 2016. URL: https://www.betweenessions.com/wp-content/uploads/2019/10/The_PSTD_Workbook-PDF_VERSION.pdf.
- [168] Eric Windhorst and Allison Williams. "Growing up, naturally: The mental health legacy of early nature affiliation". In: *Ecopsychology* (2015). DOI: <https://doi.org/10.1089/eco.2015.0040>.
- [169] *World Drug Report 2014*. Tech. rep. United Nations Office on Drugs and Crime, 2014. URL: https://www.unodc.org/documents/wdr2014/World_Drug_Report_2014_web.pdf.
- [170] David B Yaden, James B Potash, and Roland R Griffiths. "Preparing for the bursting of the psychedelic hype bubble". In: *JAMA psychiatry* (2022). DOI: [doi:10.1001/jamapsychiatry.2022.2546](https://doi.org/10.1001/jamapsychiatry.2022.2546).
- [171] R.J. Zeifman, H. Kettner, B.A. Pagni, et al. "Co-use of MDMA with psilocybin/LSD may buffer against challenging experiences and enhance positive experiences." In: *Sci Rep* (2023). DOI: <https://doi.org/10.1038/s41598-023-40856-5>.