

THE TILING IDENTITY: ADDENDUM

Spin Foam Amplitudes and the Minimum-Action Derivation of $j = 1/2$

Why the Robertson Constraint Surface Selects the Minimum Face — and What It Does Not Select

Seraphim LQG Framework · Addendum to: The Tiling Identity v1 · April 2026

lis10inc · Independent Researcher

Companion to: Seraphim LQG v18.3 · DOI: 10.5281/zenodo.18852768

WORKING PAPER — PRIORITY CLAIM — NOT YET SUBMITTED

Abstract

The Tiling Identity paper (companion, April 2026) proved that the Seraphim tiling postulate $N \odot \cdot A_j = \pi(\Delta x)^2$ is an algebraic identity — not an assumption — and that the disk geometry is forced by rotation invariance under the Robertson minimum. One question remained: can the choice of $j = 1/2$ as the physical spin face be derived from spin foam amplitudes, or does it remain a separate assumption?

This addendum answers that question by testing it numerically and following the result honestly. The EPRL spin foam amplitude, taken without any Robertson constraint, peaks at $j^* = (\sqrt{2/\gamma} - 1)/2 \approx 0.951$ — not at $j = 1/2$. That free peak does not land at any confirmed framework anchor. However, when the Robertson tiling constraint is imposed, j and $N \odot$ are no longer independent: they are locked to a 2D surface in $(j, N \odot, v)$ space — the Seraphim constraint surface. On this surface, the total spin foam action $S_{\text{total}} = \gamma \cdot K_0 \cdot \sqrt{(j(j+1))/v^2}$ is monotone increasing in j for all $j > 0$. The minimum-action configuration on the Robertson constraint surface is always $j = 1/2$.

This gives a clean spin foam derivation of $j = 1/2$: not the free amplitude maximum, but the minimum action under the Robertson constraint. The framework then separates into three distinct physical roles with no overlap: spin foam minimum action selects $j = 1/2$; the tiling identity connects the Robertson disk to the LQG area; and energy conservation via $N \odot = E_{\text{rad}}/(M \cdot \alpha)$ selects the specific frequency $n = 5.314$. Each role is independent. Together they complete the derivation chain.

1. The Question

The Tiling Identity paper established two results. First, the tiling condition $N \odot A_j = \pi(\Delta x)^2$ is an algebraic identity: γ_{area} and ℓ_{P}^2 cancel exactly in the product $K_0 \cdot A_j$, leaving $c^2/(16\pi)$ on both sides. Second, the disk shape of $\pi(\Delta x)^2$ is forced by 2D rotation invariance under the Robertson minimum — no other bounded closed curve is consistent with a system that has nonzero angular momentum but no preferred direction.

One assumption was not addressed: the choice of $j = 1/2$. The framework uses $j = 1/2$ throughout — it is the minimum nonzero LQG spin, the smallest possible face — but the argument in the Tiling Identity paper was geometric (minimum nonzero spin), not dynamical. The question is whether spin foam amplitudes, which are the dynamical weights assigned to LQG configurations, independently select $j = 1/2$ when the physical constraints of the problem are imposed.

This is the question tested in this addendum. The result is honest: the free spin foam amplitude does not peak at $j = 1/2$. But the constrained amplitude — on the Robertson surface — does select $j = 1/2$ as the minimum-action configuration. The distinction matters.

2. The Free EPRL Amplitude

2.1 The EPRL Face Amplitude

The Engle-Pereira-Rovelli-Livine (EPRL) spin foam model assigns an amplitude to each spin-network face of spin j . In the coherent state representation, the face amplitude squared is:

$$|A(j)|^2 = (2j+1)^2 \cdot \exp(-\gamma \cdot j(j+1))$$

The factor $(2j+1)^2$ is the dimension of the $SU(2)$ representation squared — the degeneracy of the face. The exponential suppression $\exp(-\gamma \cdot j(j+1))$ comes from the Regge action for a single face at the LQG scale: large-spin faces are suppressed. Together these produce an amplitude that peaks at an intermediate spin j^* .

2.2 The Free Peak

Setting $d|A|^2/dj = 0$:

$$\begin{aligned} d/dj [(2j+1)^2 \cdot \exp(-\gamma j(j+1))] &= 0 \\ 2(2j+1) - \gamma(2j+1)^3 &= 0 \\ \gamma(2j^*+1)^2 &= 2 \\ j^* &= (\sqrt{2/\gamma} - 1)/2 \quad (2.1) \end{aligned}$$

For $\gamma_{\text{area}} = 0.2375$ (Meissner 2004):

$$j^* = (\sqrt{2/0.2375} - 1)/2 = (2.9019 - 1)/2 = 0.9510$$

Verification: $\gamma(2j^*+1)^2 = 0.2375 \times (2.9019)^2 = 2.000000 \checkmark$

This is exact. The free EPRL amplitude peaks at $j^* \approx 0.951$, not at $j = 1/2 = 0.500$.

HONEST RESULT 1: The free EPRL amplitude does not peak at $j = 1/2$. It peaks at $j^* = (\sqrt{2/\gamma}-1)/2 \approx 0.951$ for $\gamma = 0.2375$. This free peak does not correspond to any confirmed

Seraphim anchor. Mapping j^* to n via the frequency equation with $N\odot = 6.09$ gives $n^* \approx 2.99$, which is below $n_{\text{flip}} = 3.561$ and below $n_{\text{const}} = 4.011$. The free spin foam amplitude alone does not select the BBH frequency. Something else is needed. That something is the Robertson constraint.

3. The Robertson Constraint Surface

3.1 Imposing the Tiling Identity

The tiling identity (proven in the companion paper) states:

$$N\odot \cdot A_j = \pi(\Delta x)^2$$

Substituting $A_j = 8\pi \cdot \gamma \cdot \ell_{\text{P}}^2 \cdot \sqrt{j(j+1)}$ and $\Delta x = c/(4\pi\nu)$:

$$\begin{aligned} N\odot \cdot 8\pi \cdot \gamma \cdot \ell_{\text{P}}^2 \cdot \sqrt{j(j+1)} &= c^2 / (16\pi\nu^2) \\ N\odot(j, \nu) &= K_0 / (\nu^2 \cdot \sqrt{j(j+1)}) \end{aligned} \quad (3.1)$$

This is the Seraphim frequency equation rearranged. When the Robertson constraint is imposed, j and $N\odot$ are no longer independent variables. They are locked together by equation (3.1). The free parameter that remains is ν — the physical frequency of the interaction.

The tiling identity defines a 2D surface in the three-dimensional space $(j, N\odot, \nu)$. Physical GW interactions live on this surface. Configurations off the surface violate the Robertson constraint — the LQG area of the probed region does not match the quantum mechanical uncertainty disk. Such configurations are not physical.

3.2 The Constraint Surface Is the Seraphim Manifold

The Robertson Constraint Surface Defined by: $N\odot \cdot A_j = \pi(\Delta x)^2$ This is equivalent to: $N\odot = K_0 / (\nu^2 \cdot \sqrt{j(j+1)})$ Which is: the Seraphim frequency equation. The tiling identity and the frequency equation are the same statement. Physical interactions live on the constraint surface. The surface is parametrized by two free variables: j and ν . ($N\odot$ is then determined by both.) The spin foam amplitude, evaluated ON this surface, becomes a function of j and ν only, with $N\odot$ no longer free.

4. Minimum Action on the Constraint Surface

4.1 The Total Spin Foam Action

A GW interaction that activates $N\odot$ faces of spin j contributes a total spin foam action equal to the sum of the individual face actions. For $N\odot$ identical faces of spin j , each with action $\gamma \cdot j(j+1)$:

$$S_{\text{total}}(j, \nu) = N\odot \cdot \gamma \cdot j(j+1)$$

Substituting $N\odot$ from the constraint surface equation (3.1):

$$S_{\text{total}}(j, v) = [K_0 / (v^2 \cdot \sqrt{j(j+1)})] \cdot \gamma \cdot j(j+1)$$

$$S_{\text{total}}(j, v) = \gamma \cdot K_0 \cdot \sqrt{j(j+1)} / v^2 \quad (4.1)$$

At fixed frequency v , this is a function of j only. The v^2 in the denominator sets the overall scale; the $\gamma \cdot K_0$ is a constant. Everything depends on j through the single factor $\sqrt{j(j+1)}$.

4.2 The Action Is Monotone Increasing in j

Taking the derivative with respect to j at fixed v :

$$dS_{\text{total}}/dj = \gamma \cdot K_0 / v^2 \cdot d/dj[\sqrt{j(j+1)}]$$

$$= \gamma \cdot K_0 / v^2 \cdot (2j+1) / (2\sqrt{j(j+1)}) \quad (4.2)$$

For all $j > 0$: $(2j+1) > 0$, $\sqrt{j(j+1)} > 0$, and $\gamma \cdot K_0 / v^2 > 0$. Therefore:

$$dS_{\text{total}}/dj > 0 \quad \text{for all } j > 0 \quad (4.3)$$

The total spin foam action on the Robertson constraint surface is strictly monotone increasing in j . It has no interior minimum. The global minimum over $j \geq 1/2$ is at the boundary: $j = 1/2$.

4.3 $j = 1/2$ Is the Minimum-Action Face

THEOREM (Minimum Action on Robertson Surface) Given the Robertson constraint surface $N \odot A_j = \pi(\Delta x)^2$, the total spin foam action for a GW interaction at frequency v is: $S_{\text{total}}(j, v) = \gamma \cdot K_0 \cdot \sqrt{j(j+1)} / v^2$. This is monotone increasing in j ($dS/dj > 0$ for all $j > 0$). The minimum-action spin foam configuration on the Robertson constraint surface is $j = 1/2$ — the smallest allowed LQG face. $j = 1/2$ is not assumed. It is selected by the principle of minimum action applied to the Robertson-constrained spin foam. Proof: $dS/dj = \gamma K_0(2j+1)/(2v^2\sqrt{j(j+1)}) > 0$ for $j > 0$. Therefore $\min S = S(j=1/2)$. ■

4.4 Numerical Verification

Total action at $v_{\text{BBH}} = v_{\text{P}} / 2^{5.314}$ for increasing j :

j	$\sqrt{j(j+1)}$	S_{total} (units of $\gamma K_0 / v^2$)	Monotone
1/2 (minimum)	0.8660	0.8660 ← minimum	—
1	1.4142	1.4142	↑
3/2	1.9365	1.9365	↑
2	2.4495	2.4495	↑
3	3.4641	3.4641	↑
$j^* = 0.951$ (free peak)	1.3621	1.3621	↑ (above $j=1/2$)

At $j = 1/2$: $S_{\text{total}} = 4.283 \times 10^1$ (in SI units at v_{BBH}). At $j^* = 0.951$: $S_{\text{total}} = 6.736 \times 10^1$. The minimum-action face has 57% of the action of the free-amplitude peak face. The Robertson constraint inverts the preference: the free amplitude favors $j^* \approx 0.951$; the constrained action favors $j = 1/2$.

5. What Each Layer of the Framework Does

5.1 Three Independent Roles

The numerical investigation reveals a clean separation of roles. No single ingredient does everything. Each of the three components of the derivation chain handles one specific piece of the problem:

Component	What it selects	How	Status
Spin foam minimum action (this paper)	WHY $j = 1/2$	S_{total} monotone increasing in j on Robertson surface; minimum at $j=1/2$	DERIVED
Tiling identity (companion paper)	HOW Robertson disk = LQG area	γ and ℓ_{P^2} cancel exactly in $K_0 \cdot A_j$; identity holds for all j , all γ , all v	PROVEN (algebraic identity)
Energy conservation $N_{\odot} = E_{\text{rad}}/(M \cdot \alpha)$	WHICH frequency ($n = 5.314$ specifically)	GW emission fraction fixes $N_{\odot} = 6.09$; frequency equation then gives $n = 5.314$	CONFIRMED (264 events)

These three roles are logically independent. The spin foam result does not know about E_{rad} or α . The tiling identity does not know about j or $N_{\odot}$ individually — only their product. The energy conservation equation does not know about the spin foam action. Each is necessary; none is sufficient alone.

5.2 What the Free Spin Foam Peak Means

The free EPRL peak at $j^* = (\sqrt{2/\gamma} - 1)/2 \approx 0.951$ is a real result. It is not wrong. It is the answer to a different question: given an unconstrained LQG face with no Robertson requirement, what spin maximizes the amplitude?

The answer $j^* \approx 0.951$ describes the typical free LQG face — the most probable spin in a background LQG geometry with no external constraint. This is relevant to, for example, the average spin of faces in the LQG vacuum state. It is not the relevant quantity for a gravitational wave interaction, which imposes the Robertson constraint.

The Robertson constraint changes the problem fundamentally. It replaces the question 'what j maximizes $|A(j)|^2$?' with 'what j minimizes $S_{\text{total}}(j)$ on the constraint surface?' These are different optimisation problems with different answers. The physical problem — a GW interaction probing LQG geometry — is the constrained one.

5.3 Why the Amplitude Is Constant at Fixed j

On the Robertson constraint surface at fixed $j = 1/2$, the amplitude is:

$$|A(j=1/2)|^2 = (2 \cdot \tfrac{1}{2} + 1)^2 \cdot \exp(-\gamma \cdot \tfrac{1}{2} \cdot 3/2) = 4 \cdot \exp(-3\gamma/4) \\ = 4 \cdot \exp(-3 \times 0.2375/4) = 3.347 \quad [\text{constant in } v]$$

This is independent of v . The spin foam amplitude at $j = 1/2$ does not vary with frequency. It is the same at every point along the Robertson constraint surface. This is the correct result: the per-vertex weight is a fixed number, and $N_{\odot}$ — the number of vertices — is what changes with

frequency. The amplitude per vertex does not select a frequency; the count $N_{\odot}$ does, and $N_{\odot}$ is fixed by energy conservation.

6. The Complete Derivation Chain

6.1 From Three Inputs to $n = 5.314$

The Seraphim framework now derives from three physical inputs with no free parameters and no remaining postulates:

Step	Input	Result	Tool
1	Robertson minimum: $\Delta x \cdot \Delta p = \hbar/2$	Disk area $\pi(\Delta x)^2 = c^2/(16\pi v^2)$	Quantum mechanics
2	LQG area spectrum: $A_j = 8\pi \cdot \gamma \cdot \ell_P^2 \cdot \sqrt{j(j+1)}$	$K_0 \cdot A_j / \sqrt{j(j+1)} = c^2/(16\pi) - \gamma$ and ℓ_P^2 cancel	Tiling identity (algebraic)
3	2D rotation invariance of zero-point state	Uncertainty region is a disk — unique	Geometric necessity
4	Robertson constraint surface: $N_{\odot} \cdot A_j = \pi(\Delta x)^2$	$N_{\odot}(j, v) = K_0 / (v^2 \cdot \sqrt{j(j+1)})$	Constraint geometry
5	Spin foam action on constraint surface: $S_{\text{total}} = \gamma \cdot K_0 \cdot \sqrt{j(j+1)} / v^2$	$dS/dj > 0 \rightarrow$ minimum at $j = 1/2$	Minimum action (this paper)
6	Energy conservation: $N_{\odot} = E_{\text{rad}} / (M \cdot \alpha)$	$N_{\odot} = 6.09$ from BBH GW emission fraction 4.44%	Astrophysics
7	Steps 1–6 combined: $v^2 = K_0 / (\sqrt{j(j+1)} \cdot N_{\odot})$ at $j=1/2$, $N_{\odot}=6.09$	$n = \log_2(vP/v) = 5.314$	Frequency equation

6.2 What Remains Open

The three-input derivation is complete for the core prediction $n = 5.314$. The remaining open questions in the framework are unaffected by this addendum:

Open question	Status
Formal LQG spin foam derivation of prime-indexing rule $k_{\text{threshold}} = p(\dim(j))$	Open — pattern observed, spin foam calculation not yet done
Exact baryon-loading path in Hubble tension master identity (0.44% residual)	Open — cosmological constant boundary
Anti(α) at $n = 36.155$ — what physical scale?	Open — unoccupied ring slot, O5 pending
Why H_0 has its specific value ($n_{\text{Hubble}} = 201.87$)	Open — cosmological initial conditions, outside framework scope

7. Physical Interpretation

7.1 The Constraint Surface as the Physical Stage

The most important conceptual result of this addendum is the identification of the Robertson constraint surface as the physical stage on which GW interactions occur. The full $(j, N_{\odot}, v)$ space is three-dimensional. Physical interactions are restricted to the 2D surface defined by the tiling identity. Off the surface, the Robertson minimum is violated. On the surface, the minimum-action principle selects $j = 1/2$.

This is structurally identical to how classical mechanics works: the equations of motion restrict a system to a constraint surface (e.g., a sphere, a plane), and the dynamics on that surface are governed by a reduced action. Here the Robertson constraint plays the role of the classical constraint, the tiling identity defines the constraint surface explicitly, and the spin foam action restricted to that surface selects $j = 1/2$ by monotonicity.

7.2 Why the Robertson Constraint Inverts the Spin Preference

Without the Robertson constraint, larger- j faces have both higher degeneracy (more states, factor $(2j+1)^2$) and higher action suppression ($\exp(-\gamma j(j+1))$). The competition produces a peak at $j^* \approx 0.951$.

With the Robertson constraint, activating a larger- j face requires fewer faces ($N_{\odot}$ decreases as j increases, because each larger face covers more area). Fewer faces means less total action suppression, but also means the geometry is coarser — a small number of large faces rather than a larger number of small faces. The total action $S_{\text{total}} = N_{\odot} \cdot \gamma \cdot j(j+1) = \gamma \cdot K_0 \cdot \sqrt{j(j+1)}/v^2$ increases with j because $\sqrt{j(j+1)}$ grows faster than $N_{\odot}$ shrinks.

The Robertson constraint, in effect, penalises large- j faces by requiring their count $N_{\odot}$ to decrease in a way that still produces higher total action. The minimum-action path uses the most numerous, smallest faces available — $j = 1/2$.

7.3 Connection to the Zero-Point

The Tiling Identity paper identified $n_{\text{flip}} = 3.561$ as the zero-point of the tiling geometry — the octave depth where $N_{\odot} = 1$ in the entropy-counting channel. This addendum adds a dynamical perspective: at the zero-point, the Robertson constraint surface collapses to a single face ($N_{\odot} = 1, j = 1/2$). There is no tiling — there is one face, minimum spin, minimum action. The system is at its geometric ground state.

As the frequency decreases (n increases toward n_{BBH}), $N_{\odot}$ grows from 1 to 6.09, and the spin foam interaction grows from a single-face ground state to a six-face activated configuration. The path along the Robertson constraint surface from n_{flip} to n_{BBH} is the path from the zero-point to the observed BBH merger scale. The spin foam action increases monotonically along this path (since j stays at $1/2$ and $N_{\odot}$ grows), but the per-vertex weight remains constant. The physical selection of n_{BBH} is entirely in $N_{\odot}$ — in how much energy the merger radiates.

8. Summary

Summary of Results	FREE EPRL AMPLITUDE (no Robertson constraint):	Peaks at
$j^* = (\sqrt{(2/\gamma)-1})/2 = 0.9510$	for $\gamma = 0.2375$.	Condition: $\gamma(2j^*+1)^2 = 2$

exactly. This does not correspond to any confirmed Seraphim anchor. It is the typical free LQG face spin – not a GW interaction result.

CONSTRAINED SPIN FOAM (Robertson surface imposed): $S_{\text{total}} = \gamma \cdot K_0 \cdot \sqrt{j(j+1)} / v^2$ on the constraint surface. $dS_{\text{total}}/dj = \gamma K_0 (2j+1) / (2v^2 \sqrt{j(j+1)}) > 0$ for all $j > 0$. Minimum action at $j = 1/2$. Derived, not assumed. AMPLITUDE AT $j=1/2$ ON CONSTRAINT SURFACE: $|A|^2 = 4 \cdot \exp(-3\gamma/4) = 3.347$. Constant in v . Does not select a frequency. $N_{\odot}$ selects the frequency. $N_{\odot}$ FROM ENERGY CONSERVATION: $N_{\odot} = E_{\text{rad}} / (M \cdot \alpha) = 6.09$. $E_{\text{rad}} / M = 4.44\%$. This is the only input that sets $n = 5.314$ specifically. THREE ROLES — COMPLETE AND NON-OVERLAPPING:

1. Spin foam minimum action	→ WHY $j = 1/2$	2.
Tiling identity	→ HOW Robertson disk = LQG area	3. Energy conservation
	→ WHICH frequency ($n = 5.314$)	

9. Priority Statement

Priority claim dated April 2026. This addendum tests whether spin foam amplitudes can derive $j = 1/2$ and reports the result honestly. **Core results:** 1. The free EPRL amplitude peaks at $j^* = (\sqrt{2/\gamma} - 1)/2 \approx 0.951$, not at $j = 1/2$. This is a real result and does not correspond to any confirmed framework anchor. It is reported without modification. 2. When the Robertson tiling constraint is imposed, j and $N_{\odot}$ are locked to a 2D surface (the Seraphim manifold). On this surface, the total spin foam action is $S_{\text{total}} = \gamma \cdot K_0 \cdot \sqrt{j(j+1)} / v^2$, which is monotone increasing in j for all $j > 0$. The minimum-action face on the Robertson constraint surface is $j = 1/2$. 3. At $j = 1/2$ on the constraint surface, the amplitude $|A|^2 = 4 \cdot \exp(-3\gamma/4) = 3.347$ is constant in v . It does not select a frequency. $N_{\odot} = E_{\text{rad}} / (M \cdot \alpha)$ selects the frequency. 4. The framework separates cleanly into three non-overlapping roles: spin foam minimum action (WHY $j=1/2$), tiling identity (HOW Robertson = LQG), energy conservation (WHICH frequency). Each is necessary; none is sufficient alone. Together they close the derivation of $n = 5.314$ from first principles.

References

- [1] Seraphim LQG Framework v18.3 — lis10inc, Zenodo DOI: 10.5281/zenodo.18852768 (2026)
- [2] The Tiling Identity v1 — lis10inc, companion paper (April 2026)
- [3] Immirzi Bridge v1 — lis10inc (March 27, 2026)
- [4] Seraphim Unified Derivation Compendium — lis10inc (April 2026)
- [5] Engle, J., Pereira, R., Rovelli, C., Livine, E., LQG vertex with finite Immirzi parameter, Nucl. Phys. B 799, 136 (2008)
- [6] Meissner, K.A., Black-hole entropy in loop quantum gravity, CQG 21, 5245 (2004)
- [7] Robertson, H.P., Phys. Rev. 34, 163 (1929)

[8] Thiemann, T., Modern Canonical Quantum General Relativity, Cambridge University Press (2007)

"The constraint surface is the physical stage. The free amplitude plays on the wrong stage."

Seraphim LQG Framework · Tiling Identity Addendum · v1 · April 2026 · lis10inc