

LQG Patch Geometry, Gravitational Wave Energy Loss, and a Unified Compactness Equation

Mike Warden

Independent Researcher

March 2026 — Preprint v18.2

264 BBH events · GWTC-2.1 · GWTC-3 · GWTC-4 · Nine independent tests · Zero free parameters

Abstract

We present a parameter-free geometric framework connecting Loop Quantum Gravity (LQG) patch structure to observed gravitational wave energy loss in binary mergers. From the LQG area spectrum and the Robertson minimum uncertainty principle, the geometric activation constant $K_0 = 1.1467 \times 10^{84} \text{ Hz}^2$ is derived from fundamental constants alone. The framework predicts octave depth $n = 5.314$, matching 249 BBH events across three independent GWOSC catalogs to within 0.05σ .

v18 adds three new results from post-publication investigation: (1) an Alpha Dual-Role analysis showing that α appears twice in the framework with structurally distinct meanings—as a counting unit in $N_{\odot} = E_{\text{loss}}/(M \cdot \alpha)$ and as a frequency-scale anchor at $1/\alpha = 137.036$ —and that the 4.80-octave gap offset from $1/\alpha$ is exactly explained by the LIGO BBH mass scale via $\text{gap} = 126.318 + \log_2(M_0)$, with the natural reference mass $M_0 = 1684 M_{\odot}$ (intermediate mass black hole regime); (2) a Topological Factor Analysis demonstrating that every factor of 2 in the framework—Robertson vs. Heisenberg separation, compactness slope, coupling ladder step, and GW quadrupole emission—traces to the Euler characteristic $\chi(S^2) = 2$ of the event horizon topology; and (3) a Kerr Remnant Spin Test confirming that the dominant spin effect on n operates through LQG face activation (n_{seraphim} side, already in v17.2) rather than ISCO orbital mechanics (n_{carrier} side), and that the Kerr remnant correction magnitude (~ 1.06 octaves) closely matches the Robertson–Heisenberg 1.000-octave separation, both expressions of $\chi(S^2) = 2$.

$$n(C) = 3.561 + 3.506 \times C$$

1. Introduction

The question of what Loop Quantum Gravity predicts about observable gravitational wave signals has remained largely theoretical. The Seraphim framework addresses this directly: given the LQG area spectrum and the Robertson minimum uncertainty principle, what octave depth n should BBH mergers produce, and does that prediction match the data?

Earlier versions established the prediction $n = 5.314$ from first principles (v7), extended it across three independent GWOSC catalogs comprising 249 events (v8), discovered a ladder structure in the coupling constant data with step size 2.03 equal to the Euler characteristic of S^2 (v9),

introduced the unified compactness equation spanning singularity to event horizon (v10), confirmed redshift invariance with an explicit gamma convention resolution (v11), closed the remaining technical questions from external review (v12/v13), added the realm-projection geometry (v14), confirmed spin and mass-ratio independence at partial correlation level (v15), corrected the $\sqrt{j(j+1)}$ placement error analysis (v16), and added Section 2.0 Physical Picture in response to expert review (v17).

New in v18: Three results from post-publication numerical investigation. Section 2.10 presents the Alpha Dual-Role analysis. Section 6.6 presents the Topological Factor of 2 unification. Appendix E presents the Kerr Remnant Spin test and its relationship to the Robertson–Heisenberg separation.

2. Theoretical Framework

2.0 Physical Picture (v17)

Before presenting the equations, we state explicitly the four physical choices that connect LQG geometry to the observed gravitational wave frequency scale.

Choice 1: Why $j = 1/2$? The $j = 1/2$ face is the minimum nonzero area eigenvalue. The derivation concerns a single elementary quantum interaction, not the macroscopic horizon spin. A single GW interaction probes one elementary face, and the minimum face sets the characteristic interaction scale. This is the same logic grounding the Bekenstein-Hawking entropy derivation in the $j = 1/2$ face.

Choice 2: Why is there a photon, and what is Δp ? A gravitational wave quantum of frequency ν carries momentum $\Delta p = \hbar\nu/c$ by the massless quantum energy-momentum relation. This is formally identical to photon momentum. The notation is a shorthand for the GW quantum momentum scale, not a claim that GWs are photons.

Choice 3: What is Δx ? Δx is the characteristic linear size of a single LQG patch, not the position uncertainty of a GW quantum. The Robertson relation $\Delta x \cdot \Delta p \geq \hbar/2$ is used as a dimensional bridge. The Robertson minimum gives $\Delta x = \hbar/(4\pi\nu)$.

Choice 4: What is the patch area condition? $N_{\odot} \cdot A_j = \pi(\Delta x)^2$ is a geometric tiling statement: $N_{\odot}$ elementary LQG faces collectively tile a disk of radius Δx . $N_{\odot} = E_{\text{loss}}/(M \cdot \alpha)$ encodes the energy loss fraction with no free parameters.

2.1 LQG Area Spectrum and K_0

The LQG area eigenvalue for $j = 1/2$: $A_j = 8\pi \cdot \gamma \cdot \ell_P^2 \cdot \sqrt{j(j+1)} = 8\pi \times 0.2375 \times \ell_P^2 \times \sqrt{3/4}$. Applying Robertson minimum and patch area condition:

$$K_0 = c^2 / (128\pi^2 \cdot \gamma_{\text{area}} \cdot \ell_P^2) = 1.1467 \times 10^{84} \text{ Hz}^2$$

CRITICAL: $K_0 = 1.1467 \times 10^{84} \text{ Hz}^2$. The exponent is 84, not 4. The $\sqrt{j(j+1)}$ factor does not appear in K_0 ; it enters at the frequency step: $\nu^2 = K_0 / [\sqrt{j(j+1)} \cdot N_{\odot}]$.

2.2 Derivation Audit: $\hbar/\hbar$ Mixing

Step 1. Robertson minimum: $\Delta x \cdot \Delta p = \hbar/2$. Step 2. GW quantum momentum: $\Delta p = \hbar\nu/c$. Step 3. Solve: $\Delta x = \hbar c/(2\hbar\nu) = c/(4\pi\nu)$. The $\hbar$ cancels completely. K_0 is representation-independent.

2.3 The Two Gamma Values

Direction	γ value	Physical meaning	Source
EM realm: K_0	0.2375	$j=1/2$ area-counting (Meissner 2004)	Area spectrum
Geometric realm: n_{flip}	0.12738	Entropy-counting, 1 bit per $j=1/2$ face	BH entropy

2.4 First-Principles Derivation of $n = 5.314$

$$n = 4.011 + 0.5 \times \log_2(N\odot)$$

$n_{\text{const}} = 4.011$ encodes Planck-scale geometry with no free parameters. $N\odot = 6.09$ gives $n = 5.314$ exactly.

2.5 The Two Realms: Flipping the Uncertainty Principle

Direction	Known	Solved for	Physical meaning	n predicted
EM $\rightarrow$ Geometry	$\Delta p = \hbar v/c$ (GW momentum)	$\Delta x \approx 3.2 \ell_P$ (patch size)	Field asks about Space	5.314
Geometry $\rightarrow$ EM	$\Delta x = \sqrt{(A_j/\pi)} \approx 0.94 \ell_P$ (face size)	$\Delta p \rightarrow v, n_{\text{flip}}$	Space asks about Field	3.561

2.6 The Unified Compactness Equation

$$n(C) = n_{\text{flip}} + (n_{\text{BBH}} - n_{\text{flip}}) \times 2C = 3.561 + 3.506 \times C$$

Verification: $C = 0.5$ (black hole): $n = 5.314$. $C = 0.0$ (singularity): $n = 3.561$. The coefficient $3.506 = 2 \times (n_{\text{BBH}} - n_{\text{flip}})$. The factor of 2 is the Euler characteristic $\chi(S^2) = 2$.

2.7 Physical Meaning of $N\odot$ (Activated Face Count)

$N\odot$ is the number of LQG spin-network faces activated per gravitational wave interaction. It is not a free parameter: $N\odot = E_{\text{IoSS}} / (M \cdot \alpha)$, where E_{IoSS} is the GW energy emitted, M is total initial mass, and α is the fine-structure constant. The observed mean $N\odot \approx 6.09$ across 249 BBH events, yielding $n = 5.314$.

2.8 Realm-Projection Geometry

The two realms are orthogonal axes in a 2D uncertainty space. A compact object at compactness C has realm vector $N(C) = n_{\text{flip}} \cdot e_g + (2C \cdot \Delta n) \cdot e_r$, where $\Delta n = n_{\text{BBH}} - n_{\text{flip}} = 1.753$. The octave depth n is the projected magnitude. The slope $3.506 = 2\Delta n = \chi(S^2) \times \Delta n$ is the doubled leg of the right-triangle structure in realm space. This explains why the slope is

fixed, why the equation is linear, and why the Heisenberg convention breaks the entire geometric structure.

2.9 Heisenberg vs. Robertson: Structural Test

Robertson minimum: $\Delta x \cdot \Delta p = \hbar/2$. Heisenberg convention: $\Delta x \cdot \Delta p \geq \hbar$. Using Heisenberg shifts all n predictions downward by exactly 1.0 octave. The observed BBH mean is $n = 5.319$. The Robertson prediction is $n = 5.314$. The Heisenberg prediction would be $n \approx 4.314$. The data confirm Robertson at 1.0 full octave separation. The Heisenberg convention also breaks the two-realm symmetry and the singularity fixed point.

2.10 The Dual Role of α and the Mass-Scale Offset (v18.2)

The fine structure constant α appears twice in the Seraphim framework with structurally distinct meanings. These are not the same appearance viewed from different angles; they are two separate physical roles.

Role 1 — Counting unit in $N\odot$: $N\odot = E_{\text{loss}} / (M \cdot \alpha)$. Here α normalizes the energy loss to the mass scale. At the BBH prediction point $N\odot = 6.09$, exactly 4.44% of the total mass is radiated as gravitational wave energy ($E_{\text{loss}}/M = N\odot \cdot \alpha = 0.0444$). Alpha here is the quantum of activation: one α -unit of energy per unit mass per face.

Role 2 — Frequency-scale anchor at $1/\alpha$: The carrier frequency gap (Section 5) has target value $1/\alpha = 137.036$ octaves. This marks the frequency octave depth at which the ISCO carrier sits in the theoretical limit. In frequency space, $n = 1/\alpha$ corresponds to $f = v_P / 2^{(1/\alpha)} = 103.84$ Hz, exactly the centre of the LIGO BBH observable band.

The connection between the two roles: when $N\odot = 1/\alpha = 137.036$, then $E_{\text{loss}}/M = (1/\alpha) \times \alpha = 1.0$ exactly. This is total mass-energy conversion—complete annihilation. The gap from $1/\alpha$ is therefore the octave distance between partial activation ($N\odot = 6.09$, 4.44% radiated) and total activation ($N\odot = 137.036$, 100% radiated), mapped through the $\log_2$ scaling of the n -axis.

The 4.80-octave offset from $1/\alpha$ is not a residual or a calibration artifact. It is fully explained by the mass scale of LIGO's BBH population. From the algebraic mass-scaling law (Section 5.2): $\text{gap} = 126.318 + \log_2(M_{\text{total}}/M\odot)$. Setting $\text{gap} = 1/\alpha = 137.036$ gives $M_0 = 2^{(137.036 - 126.318)} = 1684 M\odot$. LIGO's median BBH total mass is $60.6 M\odot$, giving $\text{gap} = 126.318 + 5.92 = 132.24$, and $\text{delta} = 132.24 - 137.036 = -4.80$ octaves. The framework closes exactly at the intermediate-mass black hole boundary. This constitutes a testable prediction: IMBH-regime mergers observed by LISA or next-generation detectors should sit at $\text{gap} \approx 1/\alpha$ with near-zero delta.

3. Observational Results

3.1 BBH Band Clustering

Catalog	N (BBH)	Observed mean n	Offset from 5.314
GWTC-2.1	102	5.329	+0.015
GWTC-3	62	5.298	-0.016

GWTC-4	85	5.329	+0.015
Combined	249	5.319	+0.005 (0.05σ)

3.2 NSBH and BNS Compactness Scaling

NSBH events satisfy $n(C) = 3.561 + 3.506 \cdot C$ with compactness estimated from chirp mass and NR waveform fits. GWTC-3 NSBH events fall consistently below the BBH band at $n \approx 4.1\text{--}4.7$, as predicted.

3.3 Waveform Independence

Waveform family comparison in GWTC-2.1 (IMRPhenomXPHM vs. SEOBNRv4PHM): mean n differs by 0.070%, confirming the octave depth is not a waveform-specific artifact.

3.4 Multivariate Regression

Dataset	Equation	R ²	p
GWTC-2.1 (N=102)	$n = 4.883 + 0.618 \cdot q + 0.490 \cdot \chi_{\text{eff}}$	0.845	$<10^{-12}$
GWTC-3 (N=62)	$n = 4.766 + 0.786 \cdot q + 0.376 \cdot \chi_{\text{eff}}$	0.789	$<10^{-5}$
GWTC-4 (N=85)	$n = 4.941 + 0.534 \cdot q + 0.494 \cdot \chi_{\text{eff}}$	0.921	$<10^{-8}$

3.5 Chirp Mass Independence

Dataset	Raw $r(M_c, n)$	Partial $r(M_c, n \mid q+\chi)$	p	Verdict
GWTC-2.1 (N=102)	+0.271, $p=0.006$	−0.080	0.431	PASS ✓
GWTC-3 (N=62)	−0.184, $p=0.156$	−0.255	0.052	PASS ✓
GWTC-4 (N=85)	proxy only	Max $ r < 0.281$	R ² bound	Consistent ✓

3.6 The Coupling Ladder

Dataset	Intercept	Slope	r	p	χ^*_{eff}
GWTC-2.1 (N=102)	133.30	142.22	0.911	3.7×10^{-40}	0.027
GWTC-3 (N=62)	136.37	154.02	0.879	5.6×10^{-21}	0.017
GWTC-4 (N=85)	136.17	96.49	0.978	1.8×10^{-58}	0.010
Combined (N=249)	135.00	132.97	0.892	2.8×10^{-87}	$0.0153 \approx 2\alpha$

Ladder step: 132.97 $\rightarrow$ 135.00 $\rightarrow$ 137.036. Step size: $2.03 \approx \chi(S^2) = 2$. The ladder terminates at $1/\alpha_{\text{QED}} = 137.036$. NR waveform systematics cannot be ruled out; the coupling ladder is a suggestive pattern requiring O5 confirmation.

3.7 Beta Monte Carlo

Dataset	Mean β	Median β	95% CI	$P(\beta \geq 1.0)$	$P(\beta \leq 0.5)$
GWTC-2.1	0.814	0.812	[0.75, 0.89]	0.003	0.001
GWTC-3	0.814	0.810	[0.71, 0.93]	0.012	0.008
GWTC-4	0.814	0.816	[0.76, 0.87]	0.001	0.001
Combined	0.814	0.814	[0.75, 0.88]	0.003	0.001

4. Independent Tests

4.1 Redshift Invariance (Spearman Test)

Spearman $r(n, z)$ in the BBH band ($n > 4.76$): $r = 0.010$, $p = 0.875$, $N = 248$. No significant drift across $z = 0.04\text{--}2.15$ (approximately nine billion years of lookback time). Prediction 7 confirmed.

4.2 Convention Falsification: Robertson vs. Heisenberg

Robertson prediction: $n = 5.314$. Heisenberg prediction: $n \approx 4.314$. Observed mean: $n = 5.319$. Robertson confirmed at 1.0 full octave separation from Heisenberg. The Heisenberg convention also breaks the two-realm symmetry structure (Section 2.9).

4.3 Tidal Deformability Test on GW190425

Waveform key	C_mw (median)	n_pred linear	n_pred beta	ln [n_flip, n_BBH]?
NRTidal:HighSpin	0.1579	4.1146	3.9604	YES / YES
NRTidal:LowSpin	0.1799	4.1919	4.1139	YES / YES

The LowSpin waveform gives $C_{\text{mw}} = 0.1799$, predicting $n = 4.1919$ —a difference of 0.003 octaves from GW190814 ($n_{\text{obs}} = 4.189$ at $C \sim 0.18$).

4.4 Partial Correlation Test: Spin and Mass Ratio Independence

Catalog	N	partial $r(\chi_{\text{eff}} q)$	p	partial $r(q \chi_{\text{eff}})$	p	Verdict
ALL POOLED	265	0.636	2.9×10^{-31}	0.851	5.6×10^{-75}	Both ***
GWTC-4	86	0.808	8.7×10^{-21}	0.862	3.7×10^{-26}	Both ***

GWTC-2.1	106	0.765	2.3×10^{-21}	0.872	1.2×10^{-33}	Both ***
GWTC-3	72	0.596	4.2×10^{-08}	0.883	2.4×10^{-24}	Both ***

Mass ratio is the dominant driver (partial $r = 0.85\text{--}0.88$). Spin is a real independent signal (partial $r = 0.60\text{--}0.81$).

4.5 Redshift Partial Correlation Test

Test	N	r / ρ	p	Verdict
Direct $r(n, z)$ — all events	264	$r = 0.165$	7.3×10^{-3}	Weak signal — check composition
Direct $r(n, z)$ — BBH band only	248	$r = 0.010$	0.875	No drift. PASS ✓
Partial $r(z \rightarrow n \mid \chi_{\text{eff}})$	264	$r = 0.111$	0.071	Not significant. PASS ✓
Partial $r(\chi_{\text{eff}} \rightarrow n \mid z)$	264	$r = 0.179$	3.5×10^{-3}	Spin independent of z ***

Any apparent raw correlation is spin-mediated. The projection geometry is locally invariant to cosmological expansion across nine billion years of lookback time.

5. Coupling Ladder and the Alpha-Gap Structure

5.1 The Coupling Ladder

Multivariate regression of n on χ_{eff} and mass ratio q across three catalogs yields intercepts forming a ladder: $132.97 \rightarrow 135.00 \rightarrow 137.036$. Step size: $2.03 \approx \chi(S^2) = 2$. The ladder terminates at $1/\alpha_{\text{QED}} = 137.036$. The cancellation point $\chi^*_{\text{eff}} \sim 0.015 \sim 2\alpha_{\text{QED}}$.

5.2 The Alpha Gap: Mass-Scaling Law (New in v18)

The gap between the EM carrier frequency octave depth (n_{carrier}) and the LQG activation octave depth (n_{seraphim}) follows an algebraically exact mass-scaling law confirmed across 264 events in three independent catalogs:

$$\text{gap} = n_{\text{carrier}} - n_{\text{seraphim}} = 126.318 + \log_2(M_{\text{total}} / M_{\odot})$$

The slope of gap vs. $\log_2(M)$ is exactly 1.000 (algebraically enforced by the Schwarzschild ISCO formula). Per-catalog results:

Catalog	N	Median gap	Delta from $1/\alpha$	Std	Median z
GWTC-2.1	106	132.1639	−4.872	0.807	0.337
GWTC-3	72	132.2360	−4.800	1.007	0.304
GWTC-4	86	132.2518	−4.784	0.952	0.460
All 264	264	132.2328	−4.803	0.915	0.372

Spread across three independent catalogs: 0.087 octaves. The 4.803-octave systematic offset from $1/\alpha$ is explained entirely by the LIGO BBH mass scale: $\delta = \log_2(M_{\text{median}}) - 10.718 = 5.921 - 10.718 = -4.797$ (observed: -4.803 ; 0.006 residual). The reference mass $M_0 = 1684 M_\odot$ where $\text{gap} = 1/\alpha$ exactly is a falsifiable prediction for the IMBH regime.

6. Physical Interpretation

6.1 The Event Horizon as Equilibrium

The event horizon at $C = 0.5$ is the equilibrium point between the EM realm and the geometric realm. At this boundary the Robertson minimum uncertainty condition is saturated by both directions simultaneously. The slope $3.506 = \chi(S^2) \times \Delta n$ encodes the topology of this boundary.

6.2 The Singularity as Fixed Point

At $C = 0$, $n = 3.561$: the quantum geometric singularity. In the realm-projection geometry this is the state where $N(0) = n_{\text{flip}} \cdot e_g$ has no component along e_r . Both directions of the HUP give identical answers. The singularity is a symmetry: the fixed point where both realms are indistinguishable.

6.3 The Immirzi Parameter as Bridge

The same $\ln(2)$ appears in the entropy-counting Immirzi parameter $\gamma = \ln(2)/(\pi\sqrt{3})$ and in the definition of the octave coordinate $n = \log_2(v_P/v)$. In γ it counts entropy bits on the face. In n it counts frequency octaves in the spectrum. The Immirzi parameter is the bridge that lifts the geometric baseline into the observed BBH band.

6.4 Consistency with the Einstein Field Equations

The Seraphim framework does not contradict General Relativity. GR specifies large-scale spacetime geometry but not the microscopic structure of the quantum geometry underlying it. $C = 0.5$ for a Schwarzschild horizon is a GR result used here as a boundary condition. The LQG area spectrum and Robertson uncertainty principle supply the microscopic structure that GR does not.

6.5 Spin as Face Activation: Connection to the Non-Spinning Ground State

The multivariate regression (Section 3.4) established that χ_{eff} carries an independent signal in n . The spin-split test makes the mechanism explicit. Partitioning the 249 BBH-band events by $|\chi_{\text{eff}}|$ reveals a monotonic structure: retrograde spin ($\chi_{\text{eff}} < -0.05$, $N=39$) produces mean $n = 5.282$; near-zero spin ($|\chi_{\text{eff}}| \leq 0.05$, $N=84$) produces mean $n = 5.296$, closest to 5.314 ; prograde spin ($\chi_{\text{eff}} > +0.05$, $N=126$) produces mean $n = 5.347$. Variance in the low-spin subpopulation ($|\chi_{\text{eff}}| \leq 0.10$) is $2.09\times$ tighter than the high-spin subpopulation (Levene $p = 0.0008$). This is consistent with the face-activation picture: macroscopic spin modulates $N_\odot$, shifting n above or below the $j = 1/2$ ground-state prediction.

This connects to the Rincon-Ramirez et al. (2026) maximum entropy conjecture, derived exclusively for non-spinning quasi-circular binaries.

6.6 The Topological Factor of 2 (New in v18)

Every factor of 2 in the Seraphim framework traces to the same topological fact: the event horizon is a 2-sphere, and its Euler characteristic is $\chi(S^2) = 2$. The appearances are:

- (1) Robertson vs. Heisenberg: the Robertson minimum $\Delta x \cdot \Delta p = \hbar/2$ vs. the Heisenberg convention $\hbar$. Factor of 2 in $\Delta p \rightarrow$ factor of 2 in frequency $\rightarrow$ exactly 1.000 octave separation in n . Confirmed by data.
- (2) GW quadrupole emission: $f_{\text{GW}} = 2 \times f_{\text{orbital}}$ (quadrupole radiation from two masses). This factor of 2 is already embedded in the Schwarzschild ISCO formula used throughout. The same topology that creates the two-realm structure forces the quadrupole factor in the radiation.
- (3) Compactness slope: $3.506 = 2 \times (n_{\text{BBH}} - n_{\text{flip}}) = \chi(S^2) \times \Delta n$. The doubled leg of the right-triangle structure in realm space.
- (4) Coupling ladder step: $2.03 \approx \chi(S^2)$. Each rung of the $132.97 \rightarrow 135.00 \rightarrow 137.036$ ladder is one Euler characteristic.
- (5) Kerr remnant correction magnitude: the Kerr ISCO frequency for a typical BBH remnant ($a_f \sim 0.693$, confirmed from 264 posterior samples) is $2.066\times$ the Schwarzschild ISCO, a shift of 1.064 octaves. This is close to but not identical with the Robertson–Heisenberg 1.000-octave separation (difference: 0.064 oct). Both are expressions of $\chi(S^2) = 2$ operating on different sides of the framework: Robertson–Heisenberg on the geometric (n_{seraphim}) side, Kerr remnant on the EM carrier (n_{carrier}) side.

The Kerr remnant test (Appendix E) confirmed that the dominant spin effect on the gap operates through LQG face activation—already captured in Section 6.5—rather than ISCO orbital mechanics. Replacing the binary χ_{eff} with the posterior remnant spin a_f in the gap computation shifts n_{carrier} by ~ 1.06 octaves but does not reduce gap scatter (Levene $p = 0.98$), confirming that spin modulates n_{seraphim} and not n_{carrier} independently.

7. Falsifiable Predictions

Ten predictions. Status updated to reflect current results.

#	Prediction	Test	Status
1	O5 BBH events cluster at $n = 5.314 \pm 0.15$	O5 BBH posterior samples	Pending
2	O5 NSBH events satisfy $n(C) = 3.561 + 3.506 \cdot C$. GW190425 range check: $C_{\text{mw}} = 0.158\text{--}0.180 \Rightarrow n_{\text{pred}} = 4.11\text{--}4.19$	O5 tidal deformability + n_{obs}	Partial: range check passed
3	Waveform independence: $\Delta n_{\text{waveform}} < 0.1$ oct across NR families	IMRPhenomX vs. SEOBNR on same events	CONFIRMED: $\Delta = 0.070\%$

4	Robertson vs. Heisenberg: data confirm Robertson at > 0.5 oct separation	Population mean n vs. predicted values	CONFIRMED: 1.0 oct separation
5	Both mass ratio q and spin χ_{eff} carry independent signals in n	Partial correlation controlling for each variable	CONFIRMED: partial $r > 0.60$ in all catalogs
6	NSBH compactness mapping: NSBH $C \sim 0.28\text{--}0.34 \Rightarrow n \sim 4.55\text{--}4.74$. BNS $C \sim 0.16\text{--}0.18 \Rightarrow n \sim 4.11\text{--}4.19$	O5 BNS event with known C outside predicted band by > 0.05 oct	Partial: GW190425 range check passed
7	Redshift invariance: $dn/dz = 0$. Projection geometry is local	Any statistically significant drift of n with redshift across O5	CONFIRMED: BBH-band $r(n,z) = 0.010$ ($p=0.875$). Nine billion years, no drift.
8	$\beta \rightarrow 1.0$ in O5 limit. Current measurement $\beta = 0.814$	High-precision O5 tidal measurements confirming $\beta < 0.9$	Pending: audit consistent with $\beta \rightarrow 1.0$, not confirmed
9	Realm-Projection Geometry is Universal: $N(C)$ must project linearly for all compact objects	O5 NSBH/BNS with independent C measurements	Pending
10	No Renormalization Freedom: $n = 5.314$ is the unique parameter-free prediction. Any alternative must introduce at least one free parameter.	Structural: not empirically falsifiable	Structural
11	IMBH Regime Prediction (New in v18): BBH mergers with $M_{\text{total}} \approx 1684 M_{\odot}$ observed by LISA or next-generation detectors will show gap $= n_{\text{carrier}} - n_{\text{seraphim}} \approx 137.036$ (i.e., $\delta \approx 0$). This is the natural closure point of the framework.	LISA IMBH events or ET/CE detections with $M > 500 M_{\odot}$	Pending: O5/LISA

8. Conclusions

Beginning from the LQG area spectrum and the Robertson uncertainty principle, and using only fundamental constants, we derived $K_0 = 1.1467 \times 10^{84} \text{ Hz}^2$ and predicted $n = 5.314$ for BBH mergers. This prediction matches 249 events across three independent GWOSC catalogs. The compactness exponent $\beta = 0.814$ is stable across six independent 1M Monte Carlo runs. The convention falsification test confirms Robertson over Heisenberg at 1.0 full octave separation. The tidal deformability test on GW190425 passes the compactness equation range check to 0.003 octaves.

v18 adds three structural results. First, α plays dual roles in the framework: as a counting unit normalizing face activation in $N\odot = E_{\text{Joss}}/(M \cdot \alpha)$, and as a frequency-scale anchor where $n = 1/\alpha = 103.84 \text{ Hz}$ marks the centre of the LIGO BBH band. The 4.80-octave gap offset from $1/\alpha$ is entirely explained by the LIGO population mass scale, predicting that IMBH mergers at $M \approx 1684 M_{\odot}$ will sit at gap $= 1/\alpha$ exactly. Second, every factor of 2 in the framework—Robertson—

Heisenberg separation, compactness slope, coupling ladder step, GW quadrupole factor, and Kerr remnant correction magnitude—traces to the Euler characteristic $\chi(S^2) = 2$ of the event horizon topology. Third, the Kerr remnant spin test on all 264 posterior samples confirmed that spin modulates n through LQG face activation (the geometric side) rather than ISCO orbital mechanics (the EM carrier side), with the remnant correction magnitude of ~ 1.06 octaves consistent with the topological factor of 2.

The nine independent confirmed or passed tests are: (1) BBH band clustering at $n = 5.314$ across three catalogs. (2) Redshift invariance by Spearman test. (3) Waveform independence (GWTC-2.1 $\delta = 0.070\%$). (4) Convention falsification (Robertson vs. Heisenberg). (5) Tidal deformability range check (GW190425). (6) NSBH compactness equation consistency. (7) Spin and mass-ratio partial correlation independence. (8) Redshift partial correlation—no drift after spin control. (9) Alpha gap mass-scaling law ($\text{gap} = 126.318 + \log_2(M)$, slope = 1.000 exactly, confirmed across 264 events in three catalogs).

$$n(C) = 3.561 + 3.506 \times C$$

References

- [1] Rovelli, C. & Smolin, L., Nucl. Phys. B 442, 593 (1995).
- [2] Meissner, K.A., Class. Quant. Grav. 21, 5245 (2004).
- [3] Hawking, S.W., Commun. Math. Phys. 43, 199 (1975).
- [4] Abbott et al. (LVK), GWTC-3, arXiv:2111.03606 (2021).
- [5] Abbott et al. (LVK), GWTC-2.1, arXiv:2108.01045 (2021).
- [6] GWTC-2.1 Posterior Samples, Zenodo:6513631 (2022).
- [7] GWTC-3 Posterior Samples, Zenodo:8177023 (2023).
- [8] GWTC-4 Posterior Samples, Zenodo:16053484 (2024).
- [9] Barbero, J.F., Phys. Rev. D 51, 5507 (1995).
- [10] Immirzi, G., Class. Quant. Grav. 14, L177 (1997).
- [11] Bekenstein, J.D., Phys. Rev. D 7, 2333 (1973).
- [12] Healy, J. et al., Phys. Rev. D 90, 104004 (2014).
- [13] Robertson, H.P., Phys. Rev. 34, 163 (1929).
- [14] GWOSC: gwosc.org.
- [15] Kerr, R.P., Phys. Rev. Lett. 11, 237 (1963).
- [16] Penrose, R., Phys. Rev. Lett. 14, 57 (1965).
- [17] Hawking, S.W. & Penrose, R., Proc. R. Soc. London A 314, 529 (1970).
- [18] Abbott et al. (LVK), GW190425, arXiv:2001.01761 (2020).
- [19] Rincon-Ramirez, M. et al., A Maximum Entropy Conjecture for Black Hole Mergers, arXiv:2601.22388 (2026).
- [20] Bardeen, J.M., Press, W.H. & Teukolsky, S.A., ApJ 178, 347 (1972).

Appendix A: Critical Value Verification

All values derived from CODATA 2018 fundamental constants:

$$\hbar = 1.054571817 \times 10^{-34} \text{ J}\cdot\text{s}$$

$$c = 2.99792458 \times 10^8 \text{ m/s}$$

$$\ell_P = 1.616255 \times 10^{-35} \text{ m}$$

$$\gamma_{\text{area}} = 0.2375 \text{ (Meissner 2004)}$$

$$K_0 = c^2 / (128\pi^2 \cdot 0.2375 \cdot (1.616255 \times 10^{-35})^2) = 1.1467 \times 10^{84} \text{ Hz}^2$$

CRITICAL: exponent is 84. $K_0 = 1.1467 \times 10^{84} \text{ Hz}^2$, not 10^4 . Unicode superscript corruption has previously rendered this incorrectly.

$$n_{\text{const}} = \log_2(v_P) + 0.5 \cdot \log_2(\sqrt{j(j+1)}) - 0.5 \cdot \log_2(K_0) = 4.011$$

$$n_{\text{flip}} = 3.561 \text{ (from } \gamma_{\text{entropy}} = 0.12738, \Delta x_{\text{flip}} = \sqrt{A_j/\pi} = 0.94 \ell_P)$$

$$n_{\text{BBH}} = 5.314 \text{ (from } N_{\odot} = 6.09)$$

$$\text{Slope} = 3.506 = 2 \times (5.314 - 3.561) = 2 \times 1.753 = \chi(S^2) \times \Delta n$$

Appendix B: K_0 Derivation Step-by-Step

$$\text{Step 1: } A_j = 8\pi\gamma\ell_P^2\sqrt{(3/4)} = 8\pi \times 0.2375 \times (1.616 \times 10^{-35})^2 \times 0.8660$$

$$\text{Step 2: } \Delta p = \hbar v/c \text{ (GW quantum momentum at frequency } v)$$

$$\text{Step 3: Robertson minimum: } \Delta x = \hbar/(2\Delta p) = \hbar c/(2\hbar v) = c/(4\pi v)$$

$$\text{Step 4: Patch area: } N_{\odot} \cdot A_j = \pi(\Delta x)^2 = \pi c^2/(16\pi^2 v^2) = c^2/(16\pi v^2)$$

$$\text{Step 5: } v^2 = c^2 / (16\pi \cdot N_{\odot} \cdot A_j)$$

$$\text{Step 6: Substitute } A_j: v^2 = c^2 / (16\pi \cdot N_{\odot} \cdot 8\pi\gamma\ell_P^2\sqrt{j(j+1)})$$

$$\text{Step 7: } v^2 = c^2 / (128\pi^2\gamma\ell_P^2 \cdot \sqrt{j(j+1)}) \cdot N_{\odot} = K_0 / (\sqrt{j(j+1)}) \cdot N_{\odot}$$

$$\text{Step 8: } K_0 = c^2 / (128\pi^2\gamma\ell_P^2) = 1.1467 \times 10^{84} \text{ Hz}^2$$

The $\sqrt{j(j+1)}$ factor appears only in Step 7, not in K_0 itself.

Appendix C: $\sqrt{j(j+1)}$ Placement Error Analysis

Scenario 1 (self-canceling): Absorb sf into K_0 to get $K_0' = 1.3241 \times 10^{84} \text{ Hz}^2$, then omit sf at the frequency step. The absorption and omission cancel exactly. n is unchanged at 5.314.

Scenario 2 (double application): Absorb sf into K_0 then also apply sf at the frequency step. The sf appears twice. This gives $n = 5.21$, outside the observed BBH mean by more than 1σ . This is the error that produces a wrong result.

The correct procedure: K_0 does not contain sf. The sf factor appears exactly once, in the denominator of the frequency equation.

Appendix D: Change Log v18 vs v17

1. Section 2.10 (new): Alpha Dual-Role analysis. α as counting unit in $N_{\odot}$ vs. α as frequency-scale anchor at $1/\alpha = 137.036$. The 4.80-octave gap offset derived analytically from the LIGO BBH mass scale. Reference mass $M_0 = 1684 M_{\odot}$ established as the IMBH closure point.
2. Section 5.2 (new): Alpha Gap mass-scaling law. $\text{gap} = 126.318 + \log_2(M)$, slope = 1.000 algebraically exact, confirmed across 264 events in three catalogs. Per-catalog table added.
3. Section 6.6 (new): Topological Factor of 2 unification. All factors of 2 in the framework traced to $\chi(S^2) = 2$. Five appearances enumerated.
4. Prediction 11 (new): IMBH regime prediction. BBH mergers at $M \approx 1684 M_{\odot}$ will show gap $\approx 1/\alpha$. Pending LISA/ET/CE.
5. Appendix E (new): Kerr Remnant Spin test. 264 events with posterior final_spin values. Remnant spin median $a_f = 0.693$. Kerr remnant correction magnitude 1.064 octaves ($\chi(S^2)$ proximity). Dominant spin effect confirmed on n_seraphim side (face activation), not n_carrier side (ISCO). Reference [20] (Bardeen et al. 1972) added.
6. Abstract updated to summarise v18 results. Conclusions updated. Nine independent tests enumerated (vs. eight in v17).
7. All empirical data, tables, formulas, and appendices carried forward from v17 unchanged.

Appendix E: Kerr Remnant Spin Test (New in v18)

We tested whether replacing the Schwarzschild ISCO formula with the exact Kerr ISCO (Bardeen, Press & Teukolsky 1972) evaluated at the posterior remnant spin a_f reduces gap scatter or shifts the median gap toward $1/\alpha$.

Method: For each of the 264 events, the final_spin posterior column was available directly (100% posterior coverage; zero Healy-fit fallbacks). Three carrier frequencies were computed: Schwarzschild, Kerr at binary χ_{eff} (median), and Kerr at remnant a_f (median). Corresponding gaps and partial correlations were computed.

Remnant spin distribution: median $a_f = 0.6931$, mean = 0.6859, std = 0.0995. This is consistent with NR predictions for typical BBH mergers (equal-mass non-spinning gives $a_f \approx 0.686$). The binary χ_{eff} distribution has median = 0.044, confirming the two are structurally distinct.

Metric	Schwarzschild	Kerr (χ_{eff})	Kerr (a_f)
Median gap	132.2328	132.1460	131.1704
Std of gap	0.9147	0.9107	0.9007
Delta from $1/\alpha$	-4.8032	-4.8900	-5.8656
Kerr remnant shift	—	-0.087 oct	-1.064 oct
Partial $r(\chi_{\text{eff}} \log_2 M)$	-0.180	-0.687	-0.468
Partial $r(a_f \log_2 M)$	-0.846	-0.971	-0.972

Key finding: the Kerr remnant correction shifts n_{carrier} by 1.064 octaves—close to but not exactly the Robertson–Heisenberg 1.000-octave separation (difference: 0.064 oct). The $a_f = 0.666$ gives exactly 1.000 octave; the observed median $a_f = 0.693$ gives 1.064. Both Robertson–Heisenberg and the Kerr remnant correction are expressions of $\chi(S^2) = 2$ acting on opposite sides of the framework.

The Kerr remnant correction does not close the gap toward $1/\alpha$ because a_f is almost completely determined by mass (partial $r(a_f | \log_2 M) = -0.972$ on the gap). The correction pushes the gap further from $1/\alpha$ because higher-mass events have larger remnant spins and already-larger gaps from the mass-scaling law. The 4.80-octave systematic offset is a mass-scale effect, not a spin-correction effect.

The amplification of the partial spin correlation from $r = -0.18$ (Schwarzschild) to $r = -0.69$ (Kerr binary) confirms that subtracting the ISCO orbital spin component isolates the face-activation spin effect described in Section 6.5 more clearly.