

Cohomology and Spectra

Namdak Tonpa

May 2025

Abstract

This article presents formal definitions and theorems for ordinary and generalized cohomology theories, unstable and stable spectra, and spectral sequences in Abelian categories, including the Serre, Atiyah-Hirzebruch, Leray, Eilenberg-Moore, Hochschild-Serre, Filtered Complex, Chromatic, Adams, and Bockstein spectral sequences. We define slopes, sheets, coordinates, quadrants, complex filtrations, and double complexes within this framework.

Contents

1	Cohomology and Spectra	2
1.1	Ordinary Cohomology	2
1.2	Generalized Cohomology Theories	2
1.3	Unstable and Stable Spectra	2
1.4	Spectral Sequences	3
1.5	Serre Spectral Sequence	3
1.6	Atiyah-Hirzebruch Spectral Sequence	3
1.7	Leray Spectral Sequence	4
1.8	Eilenberg-Moore Spectral Sequence	4
1.9	Hochschild-Serre Spectral Sequence	4
1.10	Spectral Sequence of a Filtered Complex	4
1.11	Chromatic Spectral Sequence	4
1.12	Adams Spectral Sequence	4
1.13	Bockstein Spectral Sequence	5

1 Cohomology and Spectra

1.1 Ordinary Cohomology

Definition 1. An *ordinary cohomology theory* on the category of topological spaces and pairs is a contravariant functor $H^*(-; G) : \text{Top}^{\text{op}} \rightarrow \text{GrAb}$, assigning to each pair (X, A) a sequence of abelian groups $\{H^n(X, A; G)\}_{n \in \mathbb{Z}}$, with coefficient group G , satisfying:

1. *Homotopy:* If $f \simeq g : (X, A) \rightarrow (Y, B)$, then $f^* = g^* : H^n(Y, B; G) \rightarrow H^n(X, A; G)$.
2. *Exactness:* For (X, A) , there is a long exact sequence:
$$\cdots \rightarrow H^n(X, A; G) \rightarrow H^n(X; G) \rightarrow H^n(A; G) \xrightarrow{\delta} H^{n+1}(X, A; G) \rightarrow \cdots$$
3. *Excision:* For $U \subset A$ with $\bar{U} \subset \text{int}(A)$, the inclusion $(X \setminus U, A \setminus U) \hookrightarrow (X, A)$ induces isomorphisms $H^n(X, A; G) \cong H^n(X \setminus U, A \setminus U; G)$.
4. *Additivity:* For $X = \bigsqcup X_i$, $H^n(X; G) \cong \bigoplus H^n(X_i; G)$.
5. *Dimension:* For a point pt , $H^n(\text{pt}; G) = \begin{cases} G & n = 0 \\ 0 & n \neq 0 \end{cases}$.

1.2 Generalized Cohomology Theories

Definition 2. A *generalized cohomology theory* is a contravariant functor $h^* : \text{Top}^{\text{op}} \rightarrow \text{GrAb}$, assigning to each pair (X, A) a sequence $\{h^n(X, A)\}_{n \in \mathbb{Z}}$, satisfying:

1. *Homotopy, Exactness, Excision, and Additivity* as in Definition 1.
2. *Suspension:* There is a natural isomorphism $h^n(X, A) \cong h^{n+1}(\Sigma X, \Sigma A)$, where Σ is the reduced suspension.

The groups $h^n(\text{pt})$ form a graded ring, the coefficients of h^* .

Theorem 1. Every generalized cohomology theory h^* is representable by a spectrum $E = \{E_n, \sigma_n : \Sigma E_n \rightarrow E_{n+1}\}$, with $h^n(X) \cong [X, E_n]_*$, where $[-, -]_*$ denotes pointed homotopy classes.

1.3 Unstable and Stable Spectra

Definition 3. A *spectrum* is a sequence of pointed spaces $\{E_n\}_{n \in I}$, where $I \subseteq \mathbb{Z}$, with structure maps $\sigma_n : \Sigma E_n \rightarrow E_{n+1}$. It is:

- *Unstable* if $I \subseteq \mathbb{Z}_{\geq 0}$.
- *Stable* if $I = \mathbb{Z}$ and each σ_n is a homotopy equivalence.

Theorem 2. For an unstable spectrum E , the functor $X \mapsto [X, E_n]_*$ defines a cohomology theory on spaces of dimension $\leq n$. For a stable spectrum E , the functor $h^n(X) = [X, E_n]_*$ defines a generalized cohomology theory.

1.4 Spectral Sequences

Definition 4. A *spectral sequence* in an Abelian category \mathcal{A} is a collection of objects $\{E_r^{p,q}\}_{r \geq 1, p, q \in \mathbb{Z}}$, $E_r^{p,q} \in \mathcal{A}$, with differentials:

$$d_r^{p,q} : E_r^{p,q} \rightarrow E_r^{p+a_r, q+b_r},$$

such that:

1. $d_r \circ d_r = 0$.
2. $E_{r+1}^{p,q} = H^{p,q}(E_r, d_r) = \ker(d_r^{p,q}) / \text{im}(d_r^{p-a_r, q-b_r})$.
3. There exists a graded object $H^n \in \mathcal{A}$ with filtration $F_p H^{p+q} \subseteq H^{p+q}$, such that:

$$E_\infty^{p,q} \cong F_p H^{p+q} / F_{p-1} H^{p+q}.$$

The sequence is *first-quadrant* if $E_r^{p,q} = 0$ for $p < 0$ or $q < 0$.

Definition 5. The r -th *sheet* of a spectral sequence is the collection $\{E_r^{p,q}\}_{p,q}$. The indices (p, q) are *coordinates*, with p the filtration degree and q the complementary degree, satisfying total degree $n = p + q$. The *slope* of $d_r : E_r^{p,q} \rightarrow E_r^{p+r, q-r+1}$ is $\frac{-r+1}{r}$.

Definition 6. A *filtered complex* in $\mathcal{A} = \text{Ab}$ is a chain complex (C_*, ∂) with a filtration $\cdots \subseteq F_{p-1} C_n \subseteq F_p C_n \subseteq F_{p+1} C_n \subseteq \cdots$, compatible with ∂ . A *double complex* is a bigraded object $C_{p,q}$ with differentials $d^h : C_{p,q} \rightarrow C_{p-1,q}$, $d^v : C_{p,q} \rightarrow C_{p,q-1}$, satisfying $d^h d^h = d^v d^v = d^h d^v + d^v d^h = 0$. The total complex is $\text{Tot}(C)_n = \bigoplus_{p+q=n} C_{p,q}$.

Theorem 3. A filtered complex (C_*, F_p) induces a spectral sequence with:

$$E_0^{p,q} = F_p C_{p+q} / F_{p-1} C_{p+q}, \quad E_1^{p,q} = H_{p+q}(F_p C / F_{p-1} C) \implies H_{p+q}(C).$$

A double complex $C_{p,q}$ with filtration by p -index induces:

$$E_1^{p,q} = H_q^v(C_{p,*}), \quad d_1 = H(d^h) \implies H_{p+q}(\text{Tot}(C)).$$

1.5 Serre Spectral Sequence

Theorem 4. For a fibration $F \rightarrow E \rightarrow B$ with B path-connected, there exists a first-quadrant spectral sequence:

$$E_2^{p,q} = H^p(B; H^q(F; \mathbb{Z})) \implies H^{p+q}(E; \mathbb{Z}),$$

with $d_r : E_r^{p,q} \rightarrow E_r^{p+r, q-r+1}$.

1.6 Atiyah-Hirzebruch Spectral Sequence

Theorem 5. For a generalized cohomology theory h^* and a CW-complex X , there exists a spectral sequence:

$$E_2^{p,q} = H^p(X; h^q(pt)) \implies h^{p+q}(X),$$

with $d_r : E_r^{p,q} \rightarrow E_r^{p+r, q-r+1}$.

1.7 Leray Spectral Sequence

Theorem 6. *For a continuous map $f : X \rightarrow Y$ and a sheaf \mathcal{F} on X , there exists a spectral sequence:*

$$E_2^{p,q} = H^p(Y; R^q f_* \mathcal{F}) \implies H^{p+q}(X; \mathcal{F}),$$

with $d_r : E_r^{p,q} \rightarrow E_r^{p+r, q-r+1}$.

1.8 Eilenberg-Moore Spectral Sequence

Theorem 7. *For a pullback diagram with fibration $F \rightarrow E \rightarrow B$, there exists a spectral sequence:*

$$E_2^{p,q} = \text{Tor}_{H_*(B)}^{p,q}(H_*(F), R) \implies H_{p+q}(F; R),$$

with $d_r : E_r^{p,q} \rightarrow E_r^{p-r, q+r-1}$.

1.9 Hochschild-Serre Spectral Sequence

Theorem 8. *For a group extension $1 \rightarrow N \rightarrow G \rightarrow Q \rightarrow 1$, there exists a spectral sequence:*

$$E_2^{p,q} = H^p(Q; H^q(N; R)) \implies H^{p+q}(G; R),$$

with $d_r : E_r^{p,q} \rightarrow E_r^{p-r, q+r-1}$.

1.10 Spectral Sequence of a Filtered Complex

Theorem 9. *For a filtered complex (C_*, F_p) , there exists a spectral sequence:*

$$E_1^{p,q} = H_{p+q}(F_p C / F_{p-1} C) \implies H_{p+q}(C),$$

with $d_r : E_r^{p,q} \rightarrow E_r^{p-r, q+r-1}$.

1.11 Chromatic Spectral Sequence

Theorem 10. *For a spectrum X , there exists a spectral sequence:*

$$E_1^{n,k} = \pi_{n-k}(L_{K(k)} X) \implies \pi_{n-k}(X),$$

where $L_{K(k)} X$ is the localization at the k -th Morava K -theory, with $d_r : E_r^{n,k} \rightarrow E_r^{n+1, k-r}$.

1.12 Adams Spectral Sequence

Theorem 11. *For a spectrum X and prime p , there exists a spectral sequence:*

$$E_2^{s,t} = \text{Ext}_A^{s,t}(\text{Hom}_*(X, \mathbb{Z}/p), \mathbb{Z}/p) \implies \pi_{t-s}(X_{(p)}),$$

where A is the Steenrod algebra, with $d_r : E_r^{s,t} \rightarrow E_r^{s+r, t+r-1}$.

1.13 Bockstein Spectral Sequence

Theorem 12. *For a short exact sequence $0 \rightarrow R \rightarrow R' \rightarrow R'' \rightarrow 0$ of coefficient rings, there exists a spectral sequence:*

$$E_1^{p,q} = H^{p+q}(X; R'') \implies H^{p+q}(X; R),$$

with $d_r : E_r^{p,q} \rightarrow E_r^{p+1, q-r}$.

References

- [1] Hatcher, A., *Algebraic Topology*, Cambridge University Press, 2002.
- [2] Switzer, R., *Algebraic Topology - Homology and Homotopy*, Springer, 1975.
- [3] Kato, G., *The Heart of Cohomology*, Springer, 2006