

# Chapter 8.2

## Poisson random variable

## Review of Last Week

Variance:  $\sigma^2 = \text{Var}[X] = E[(X - \mu)^2]$  provides a measure of spread from the expected value  $\mu$ .

- Easier formula for calculating variance:

$$\text{Var}[X] = E[X^2] - \mu^2$$

Standard deviation:  $\sigma = \text{SD}[X] = \sqrt{\text{Var}[X]}$  is the typical size of the deviation from  $\mu$ .

Chebyshev's inequality: the probability that a random variable is  $k$  or more  $\sigma$  from the mean is no bigger than  $\frac{1}{k^2}$ .

$$P(|X - \mu| \geq k\sigma) \leq \frac{1}{k^2}.$$

## Review of Last Week

Geometric random variable: the number of **failures** before first success in independent trials with probability of success  $\pi$  on each trial.

$$X \sim \text{Geom}(\pi)$$

- PMF:  $f(x) = (1 - \pi)^x \pi$ ,  $x = 0, 1, 2, 3 \dots$
- For any integer  $x \geq 0$  we have the result  $P(X \geq x) = (1 - \pi)^x$ . (example 8.2)
- $E[X] = \frac{1-\pi}{\pi}$  (odds of failure)

## Poisson Experiment

Suppose some event occurs “at random times” over a fixed observation period. Let  $X$  be the random variable which counts the number of occurrences of this event over this observation period.


$X$  is called a **Poisson** random variable.

## Poisson PMF

The derivation of the PMF of  $X$  begins by approximating  $X$  with something we know, namely the binomial distribution, using the following chain of reasoning.

- Divide the time into  $n$  non-overlapping sub-intervals of equal length.
- Assume that the probability that an event occurs during a given sub-interval,  $\pi$  remains constant from sub-interval to sub-interval and is proportional to  $\frac{1}{n}$  - let's call this probability  $\lambda/n$ .
- If  $n$  is large, the probability of having two occurrences in one sub-interval is very small – we will approximate this with 0.
- The number of occurrences in one interval is independent of the number in the other sub-intervals.

# Poisson PMF



## Poisson PMF

A good approximation for  $X$  is

$$X \approx \text{Binom}(n, \frac{\lambda}{n})$$

because we have  $n$  independent sub-intervals (trials) with probability  $\pi = \lambda/n$  of occurrence in each one.

$$\begin{aligned} P(X = x) &\approx P(x \text{ of the sub-intervals contain 1 event and} \\ &\quad \text{the other } (n-x) \text{ contain 0 events}), \\ &= \binom{n}{x} \left(\frac{\lambda}{n}\right)^x \left(1 - \frac{\lambda}{n}\right)^{n-x}. \end{aligned}$$

## Poisson limit to the binomial

The binomial approximation to the Poisson experiment should get better and better as  $n \rightarrow \infty$ . In fact, when  $n$  is very large:

$$P(X = x) = \binom{n}{x} \left(\frac{\lambda}{n}\right)^x \left(1 - \frac{\lambda}{n}\right)^{n-x} \rightarrow e^{-\lambda} \frac{\lambda^x}{x!}.$$

This is referred to as the **Poisson limit** to the binomial PMF as a nod to Siméon Denis Poisson, the French mathematician who discovered it.



## Proof of the Poisson limit to binomial

$$\begin{aligned}P(X = x) &= \binom{n}{x} \left(\frac{\lambda}{n}\right)^x \left(1 - \frac{\lambda}{n}\right)^{n-x}, \\&= \frac{n!}{x!(n-x)!} \left(\frac{\lambda}{n}\right)^x \left(1 - \frac{\lambda}{n}\right)^{n-x}, \\&= \frac{n \cdot (n-1) \cdot (n-2) \dots (n-x+1)}{n^x} \cdot \frac{\lambda^x}{x!} \cdot \frac{(1 - \lambda/n)^n}{(1 - \lambda/n)^x}\end{aligned}$$

As  $n \rightarrow \infty$ , we have:

$$\left(1 - \frac{\lambda}{n}\right)^n \approx e^{-\lambda}, \quad \frac{n \cdot (n-1) \cdot (n-2) \dots (n-x+1)}{n^x} \approx 1, \quad \left(1 - \frac{\lambda}{n}\right)^x \approx 1.$$

In other words, for

$$X \sim \text{Binom}(n, \pi)$$

if  $n$  is large but  $\pi$  is small enough so that  $n\pi$  remains constant, then  $X$  is called a Poisson random variable with parameter  $\lambda = n\pi$ .

## Poisson PMF

**Definition 8.1** The PMF for a **Poisson random variable** with parameter  $\lambda (> 0)$  is

$$f(x) = e^{-\lambda} \frac{\lambda^x}{x!}, \quad x = 0, 1, 2, \dots$$

We denote  $X \sim \text{Poisson}(\lambda)$ .

Recall from calculus (Taylor series) that

$$1 + \lambda + \frac{\lambda^2}{2!} + \frac{\lambda^3}{3!} + \cdots = \sum_{x=0}^{\infty} \frac{\lambda^x}{x!} = e^{\lambda},$$

and therefore we have defined a legitimate PMF since

$$\sum_{x=0}^{\infty} f(x) = \sum_{x=0}^{\infty} e^{-\lambda} \frac{\lambda^x}{x!} = e^{-\lambda} \cdot \sum_{x=0}^{\infty} \frac{\lambda^x}{x!} = e^{-\lambda} \cdot e^{\lambda} = 1.$$

## Example 8.4

Suppose that the number of accidents occurring on a highway each day is a Poisson random variable with parameter  $\lambda = 3$ .

- a. Find the probability that 3 or more accidents occur today.

## Poisson calculations in R


```
dpois(x = 3, lambda = 3)    # $P(X = 3)$   
## [1] 0.224
```

```
ppois(q = 2, lambda = 3)    # $P(X \leq q)$   
## [1] 0.423
```


```
ppois(q = 2, lambda = 3, lower.tail = F)    # $P(X > q)$   
## [1] 0.577
```

# Probability histogram

$X \sim \text{Pois}(\lambda = 3)$



$X \sim \text{Pois}(\lambda = 10)$



## Example 8.4 contd.

Suppose that the number of accidents occurring on a highway each day is a Poisson random variable with parameter  $\lambda = 3$ .

- b. Repeat part a under the assumption that at least 1 accident occurs today.



## Expectation and variance

**Lemma 8.1** Let  $X \sim \text{Poisson}(\lambda)$ . Then

- $E[X] = \lambda$
- $\text{Var}[X] = \lambda$

The important take aways here are that if  $X \sim \text{Pois}(\lambda)$ , then

- the mean and variance of  $X$  are equal
- the parameter  $\lambda$  is the expected number of occurrences of the event during the observation period and is referred to as the **rate** parameter.

It is often the case that the number of **arrivals** at a server (ATM machine, telephone exchange, wireless network) for some specific length of time  $t$

- can be modeled by a  $Pois(\lambda t)$  distribution where  $\lambda$  is the rate per unit time
- and is such that arrivals in non-overlapping intervals are independent.

We call such a model a **Poisson process**

## Example 8.5

Customers come to a small business at an average rate of 6 per hour. Let's assume that a Poisson process is a good model for customer arrivals.

- a. Calculate the probability that there are exactly 5 customers in the next 20 minutes?

## Example 8.5

Customers come to a small business at an average rate of 6 per hour. Let's assume that a Poisson process is a good model for customer arrivals.

- b. Calculate the probability that there are exactly 5 customers in the next 20 minutes and 5 more customers in the following 10 minutes.

## Example 8.5

Customers come to a small business at an average rate of 6 per hour. Let's assume that a Poisson process is a good model for customer arrivals.

- c. Calculate the probability that the next 5 customers will arrive within 15 minutes of each other.

## Example 8.6

Is the Poisson distribution a good fit for modeling the number of fumbles in NCAA football?

```
#include packages in setup  
library(fastR2) # for the dataset Fumbles  
library(tidyverse) # for ggplot + dplyr packages
```

## Example 8.6

```
#you can type data(fumbles) in Console to load dataset in Environment  
#  
glimpse(Fumbles)
```

```
## Rows: 120  
## Columns: 7  
## $ team   <fct> Air Force, Akron, Alabama, Arizona, Arizona St, Arkansas, Arkans~  
## $ rank   <int> 53, 19, 68, 31, 94, 46, 60, 94, 18, 94, 89, 76, 4, 38, 41, 53, 4~  
## $ W      <int> 8, 1, 9, 7, 5, 9, 4, 6, 12, 4, 7, 10, 6, 2, 2, 6, 5, 8, 3, 4, 6,~  
## $ L      <int> 4, 11, 3, 4, 6, 2, 7, 5, 0, 8, 5, 1, 5, 10, 10, 5, 6, 3, 9, 6, 5~  
## $ week1  <int> 4, 2, 0, 1, 2, 0, 0, 3, 1, 2, 5, 3, 0, 1, 2, 1, 3, 3, 5, 2, 1, 0~  
## $ week2  <int> 2, 3, 3, 0, 1, 1, 0, 2, 1, 2, 2, 2, 2, 1, 3, 1, 1, 3, 5, 2, 5, 2~  
## $ week3  <int> 2, 2, 2, 2, 3, 0, 4, 0, 0, 2, 1, 2, 4, 2, 3, 3, 2, 0, 0, 2, 2, 3~
```

```
#please see STAT 311 course resources "Data verbs" slidedeck, "Data basics" lab
```

## Example 8.6

```
slice_head(Fumbles, n = 5)      #peek at first five rows
```

```
##           team rank W  L week1 week2 week3
## 1  Air Force   53 8   4    4     2     2
## 2    Akron    19 1 11    2     3     2
## 3  Alabama    68 9   3    0     3     2
## 4  Arizona    31 7   4    1     0     2
## 5 Arizona St  94 5   6    2     1     3
```



## Example 8.6

```
Fumbles %>% count(week1) #what are the values in this column and how often is each value observed?
##   week1  n
## 1     0 22
## 2     1 36
## 3     2 29
## 4     3 23
## 5     4  5
## 6     5  4
## 7     7  1

Fumbles %>% summarize(n=n(), #n() counts the number of rows
                      xbar = mean(week1), #find mean of values
                      s = sd(week1),      #find SD of values
                      min = min(week1),   #find min of values
                      max = max(week1) )  #find max of values
##      n xbar    s min max
## 1 120 1.75 1.36  0   7
```

## Example 8.6

Let  $X_i$  denote the number of fumbles made by team  $i$  in week 1. We have observed  $x_1 = 4, x_2 = 2, x_3 = 0, x_4 = 1, x_5 = 2$  and so on.


What can be said about the distribution of  $X_i$  in general?

Clearly,  $X_i$  is the number of *successes* in a given period of time, but does that automatically mean it has a Poisson distribution? Not necessarily.

## Example 8.6

Histogram of Week 1 Fumbles

Poisson probabilities with  $\lambda = 1.75$  overlaid as line segments



Source: Fumbles data from fastR2 package

## Example 8.6

### Code to make histogram with Poisson probabilities overlaid

```
# data frame containing  $P(X = x)$  assuming  $X \sim \text{Pois}(\lambda = 1.75)$ 
#
pois_fit <- tibble(
  num_fumbles = 0:7,
  f = dpois(num_fumbles, lambda = 1.75)
)

ggplot( ) +
  geom_histogram(data = Fumbles,
    mapping = aes(x = week1,
      y = after_stat(density)),
    fill = "gold",
    color = "purple",
    alpha = 0.5,
    binwidth = 1) +
  geom_segment(data = pois_fit,
    mapping = aes( x = num_fumbles,
      xend = num_fumbles,
      y = 0, yend = f),
    linewidth = 2) +
  labs(x = "number of fumbles",
    title="Histogram of Week 1 Fumbles",
    subtitle = paste("Poisson probabilities with", expression(lambda==1.75), "overlaid as line segments"),
    caption="Source: Fumbles data from fastR2 package")
```