

Volume of a simplex

Gerald Rapior

29.03.2016

1 Definition of the problem

It should be shown that the volume of a simplex tends to 0 if the dimension of simplex grows.

Say $v_n(a)$ is the volume of a simplex where n is the dimension. For convenience the volume will be determined for an arbitrary length a and later $a = 1$ will be set. Than

$$\lim_{n \rightarrow \infty} v(a = 1) = 0. \quad (1)$$

In geometry, a simplex (plural: simplexes or simplices) is a generalization of the notion of a triangle or tetrahedron to arbitrary dimensions (see Simplex in Wikipedia). For simplicity a canonical basis of unit vectors \vec{e}_i is used. The simplex is determined by a set of points:

$$C = \left\{ c_1 \vec{e}_1 + \dots + c_n \vec{e}_n \mid \forall i, c_i \geq 0, \sum_{i=1}^n c_i = 1 \right\}. \quad (2)$$

Borders of such a simplex are the unit vectors and straight lines between end points of the unit vectors respectively.

2 Determination of volume

- $n = 1$:

In figure 1 is shown a 1-dimensional simplex (a straight line). The length (volume in one dimension) is determined by:

$$v_1(a) = \int_0^a dx = \left[x \right]_0^a = a. \quad (3)$$

Later the a can be substituted by 1.




Figure 1: Volume of a 1-D simplex




Figure 2: Volume of a 2-D simplex

- $n = 2$:

In figure 2 is shown a 2-dimensional simplex (a triangle). The surface of the shaded triangle is determined by:

$$v_2(a) = \int_0^a \int_0^y dx dy = \int_0^a y dy = \left[\frac{1}{2} y^2 \right]_0^a = \frac{a^2}{2}. \quad (4)$$

- $n = 3$:

The volume of the shaded area (see fig. 3) is determined by ¹:

$$v_3(a) = \int_0^a \int_0^z \int_0^y dx dy dz = \int_0^a \int_0^z y dy dz = \int_0^a \frac{z^2}{2} dz = \left[\frac{1}{2 \cdot 3} z^3 \right]_0^a = \frac{a^3}{6}. \quad (5)$$

It seems that the volume $v_n(a)$ of dimension n for $a = 1$ is:

$$v_n(1) = \frac{1}{n!}. \quad (6)$$

3 Proof of (6)

The proof is made by means of induction. $v_n(a) = 1/n!$ for $n = 1$ is shown above (3). Say the equation is valid for n . Next it has to be shown that the equation is valid for $n + 1$.

¹The dashed triangle in fig.3 is an arbitrary similar triangle of the ‘base’. The size is direct related to the position z . At least all these triangles are ‘added together’. This shows clear the recursive way to construct volumes of higher dimensions.




Figure 3: Volume of a 3-D simplex

Proof.

$$v_{n+1}(a) = \int_0^a \underbrace{\int_0^{x_{n+1}} \cdots \int_0^{x_2} dx_1 \cdots dx_n}_{= v_n(x_{n+1})} dx_{n+1} \quad (7)$$

$$= \int_0^a \frac{1}{n!} x_{n+1}^n dx_{n+1} \quad (8)$$

$$= \left[\frac{1}{n!} \frac{1}{n+1} x_{n+1}^{n+1} \right]_0^a = \frac{a^{n+1}}{(n+1)!} \quad \square$$

In the formula above old x , y and z are replaced by x_1 , x_2 and x_3 . For $a = 1$ we get result for volume of a simplex.

At least we know that

$$\lim_{n \rightarrow \infty} v_n(1) = \lim_{n \rightarrow \infty} \frac{1}{n!} = 0. \quad (9)$$