

Assignment #3 for NMPDE

Exercise 1 Determine the variational formulation of following boundary problem:

$$\begin{cases} -u'' + g u = f, & x \in (0, 1) \\ u(0) = u(1) = 0 \end{cases}$$

and state the definition of the weak solution to the problem where $f, g \in C^\infty(0, 1)$.

Exercise 2 Determine the variational formulation of following boundary problem:

$$\begin{cases} -u'' + u = f, & x \in (0, 1) \\ u(0) = 0, u'(1) + u(1) = 0 \end{cases}$$

and state the definition of the weak solution to the problem.

Exercise 3 Which variational problem is associated to the boundary-value problem an ordinary differential equation

$$u''(x) = e^x, \quad x \in (0, 1)$$

where $u(0) = u(1) = 0$.

Exercise 4 Consider the elliptic, but not uniformly elliptic, bilinear form

$$a(u, v) = \int_0^1 x^2 u' v' \, dx$$

on $[0, 1]$. Show that the problem

$$\frac{1}{2} a(u, u) - \int_0^1 u \, dx \rightarrow \min$$

does not have a solution in $H_0^1(0, 1)$. Determine the associated differential equation.

Exercise 5 Consider a bounded, connected and regular domain $\Omega \subseteq \mathbb{R}^n$ and the

$$\begin{cases} -\Delta u = \lambda u, & \forall x \in \Omega. \\ u(x) = 0, & \forall x \in \partial\Omega. \end{cases}$$

deduce its variational formulation where λ is a constant.

Exercise 6 Justify that $u(x) = 1/|x|^\alpha \in H^1$ for $\alpha < (n-2)/2$ and $n > 3$ where $x \in \mathbb{R}^n$ and $|x| := \sqrt{x_1^2 + \cdots + x_n^2}$ is its norm.

Exercise 7 For the following equation with Robin boundary condition

$$\begin{cases} -u'' = f, \quad \forall x \in (0, 1). \\ u'(0) + \gamma_0 u(0) = \alpha_0 \\ u'(1) + \gamma_1 u(1) = \alpha_1 \end{cases}$$

show its weak formulation is given as follows: determine $u \in H^1$ satisfying

$$a(u, v) = (f, v) + (\alpha_1 - \gamma_1 u(1))v(1) - (\alpha_0 - \gamma_0 u(0))v(0), \forall v \in H^1.$$

also show that the function $w \in H^1$ that minimizes

$$J[w] = a(w, w) - 2(f, w) - 2\alpha_1 w(1) + \gamma_1 w(1)^2 + 2\alpha_0 w(0) - \gamma_0 w(0)^2$$

is u .

Exercise 8 Suppose Ω is contained in an n -dimensional cube with side length s , prove that

$$\|v\|_{L^2(\Omega)} \leq s \cdot \|\nabla v\|_{L^2(\Omega)}, \forall v \in H_0^1(\Omega).$$

Hint: establish the inequality for $v \in C_0^\infty$ and use the density of $C_0^\infty(\Omega)$ in $H_0^1(\Omega)$. Notice that $v(x_1, x_2, \dots, x_n) = \int_0^{x_1} \partial_1 v(t, x_2, \dots, x_n) dt$ and use Cauchy-Schwarz inequality.