

IMPLEMENTATION OF THE ZABR MODEL

PETER CASPERS

First Version July 17, 2012 - This Version September 7, 2013

ABSTRACT. This is mainly a repeat of [1] inserting some more intermediate steps in the calculations and a test of the numerical examples in the original paper against our own implementation in [2].

1. MODEL DESCRIPTION

We follow [1]. The general model description is given by

$$(1.1) \quad df(t) = \sigma(f(t))v(t)dW(t)$$

$$(1.2) \quad dv(t) = \epsilon(v(t))dV(t)$$

$$(1.3) \quad dW(t)dV(t) = \rho dt$$

$$(1.4) \quad v(0) = \alpha > 0$$

where the local volatility function σ only depends on the forward f (not on the time t) and is multiplied by a stochastic volatility factor v , which is driven by a correlated brownian motion V , again with a space but not time dependent local volatility function ϵ .

The SABR model is recovered as a special case by setting

$$(1.5) \quad \sigma(f) = f^\beta$$

$$(1.6) \quad \epsilon(v) = \eta v$$

with CEV parameter $\beta \in [0, 1]$ and volatility of volatility $\eta \geq 0$.

2. IMPLIED NORMAL VOLATILITY

The aim is to derive a formula for european call option prices in the general model given by the general formula

$$(2.1) \quad c(t) = E((f(T) - K)^+ | \mathcal{F}_t)$$

where T is the maturity, K is the strike and $c(t)$ is the (non discounted) call price at time t .

We define the normal implied volatility $\nu(t)$ by

$$(2.2) \quad B(f(t), \nu(t), \tau, K) = c(t)$$

where $\tau := T - t$ is the time to maturity and B is the normal Black Formula

Date: September 7, 2013.

$$(2.3) \quad B(f, \nu, \tau, K) = (f - K)\Phi\left(\frac{f - K}{\nu\sqrt{\tau}}\right) + \nu\sqrt{\tau}\phi\left(\frac{f - K}{\nu\sqrt{\tau}}\right)$$

The Ito formula applied to 2.2 yields

$$(2.4) \quad dc = -B_\tau dt + B_f df + B_\nu d\nu + \frac{1}{2}(B_{\nu\nu} d\nu d\nu + B_{ff} df df) + B_{f\nu} df d\nu$$

Setting

$$(2.5) \quad x(t) := \frac{f(t) - K}{\nu(t)}$$

the ito formula gives

$$(2.6) \quad dx = \frac{1}{\nu} df - \frac{f - K}{\nu^2} d\nu + \frac{f - K}{\nu^3} d\nu d\nu - \frac{1}{\nu^2} df d\nu = \frac{1}{\nu} (df - x d\nu) + \dots dt$$

because $dZdt = dt dt = 0$ for every brownian Z . Therefore the quadratic variation of x evaluates to

$$(2.7) \quad dxdx = \frac{1}{\nu^2} (df df - 2x df d\nu + x^2 d\nu d\nu)$$

The following equations are verified by direct computation

$$(2.8) \quad B_\nu = \nu\tau B_{ff}$$

$$(2.9) \quad B_{\nu\nu} = \left(\frac{f - K}{\nu}\right)^2 B_{ff}$$

$$(2.10) \quad B_{f\nu} = -\left(\frac{f - K}{\nu}\right) B_{ff}$$

$$(2.11) \quad 0 = -B_\tau + \frac{1}{2}\nu^2 B_{ff}$$

Since c and f are martingales, taking expectations in 2.4 and using 2.8 yields

$$(2.12) \quad 0 = -\frac{1}{2}\nu^2 B_{ff} dt + \nu\tau B_{ff} d\nu + \frac{1}{2}(x^2 B_{ff} d\nu d\nu + B_{ff} df df) - x B_{ff} df d\nu$$

Dividing by B_{ff} and using 2.7 gives

$$(2.13) \quad 0 = -\frac{1}{2}\nu^2 dt + \nu\tau E(d\nu) + \frac{1}{2}\nu^2 dxdx$$

which for $\tau \rightarrow 0$ becomes

$$(2.14) \quad 0 = -\nu^2(dt - dxdx)$$

or

$$(2.15) \quad \sigma_x := \frac{dxdx}{dt} = 1$$

The aim is to find $x(t)$, then the normal implied volatility is given by

$$(2.16) \quad \nu(t) = \frac{f - K}{x(t)}$$

Setting $x(t) = x(f(t), v(t))$, one gets $dx = x_f df + x_v dv + \dots dt$, therefore

$$(2.17) \quad 1 = \sigma_x = \frac{dxdx}{dt} = x_f^2 \sigma(f)^2 v^2 + x_v^2 \epsilon(v)^2 + 2x_f x_v \rho \sigma(f) v \epsilon(v)$$

This is a nonlinear partial differential equation in the (dummy) variables (f, v) , which can be solved using boundary condition

$$(2.18) \quad x(f = K, v) = 0$$

3. DETERMINISTIC CASE

If $\epsilon(v) = 0$, i.e. there is no stochastic volatility, the PDE 2.17 becomes

$$(3.1) \quad 1 = x_s^2 \sigma(f)^2 v^2$$

We can set $v = 1$, taking the root and get

$$(3.2) \quad x_f = \sigma(f)^{-1}$$

with solution under the boundary condition $x(K) = 0$

$$(3.3) \quad x = \int_K^f \sigma(u)^{-1} du$$

thereby yielding the normal volatility

$$(3.4) \quad \nu = \frac{f - K}{\int_K^f \sigma(u)^{-1} du}$$

Note that in the case $f = K$, $\nu = \sigma(f)$. In the lognormal case

$$(3.5) \quad \nu = \frac{\ln \frac{f}{K}}{\int_K^f \sigma(u)^{-1} du}$$

where we have to take $\nu = \sigma(f)f^{-1}$ for $f = K$. The local volatility function $\sigma(f)$ can obviously be reconstructed from x by

$$(3.6) \quad \sigma(f) = - \left(\frac{\partial x}{\partial K}(f) \right)^{-1}$$

4. THE SABR MODEL

The SABR model is recovered by setting $\sigma(f) = f^\beta$ and $\epsilon(v) = \eta v$ with $\beta \in [0, 1]$ and a volatility of volatility parameter $\eta \geq 0$. The pde 2.17 now takes the form

$$(4.1) \quad 1 = x_f^2 f^{2\beta} v^2 + x_v^2 \eta^2 v^2 + 2x_f x_v \rho f^\beta \eta v$$

We define a new variable y by setting

$$(4.2) \quad y(t) := v(t)^{-1} \int_K^{f(t)} \sigma(u)^{-1} du$$

Ito gives

$$(4.3) \quad dy = y_f df + y_v dv + \dots dt =$$

$$(4.4) \quad dW - \eta y(t) dV + \dots dt =$$

$$(4.5) \quad \sqrt{1 + \eta^2 y(t)^2 - 2\rho \eta y(t)} dB + \dots dt =:$$

$$(4.6) \quad J(y(t)) dB + \dots dt$$

It should be noted that

$$(4.7) \quad 1 + \eta^2 y(t)^2 - 2\rho \eta y(t) = (\eta y(t) - \rho)^2 + (\rho^2 - 1) > 0$$

whenever $\rho \in (-1, 1)$ what we assume in the following. Now set

$$(4.8) \quad x(t) := \int_0^{y(t)} J(u)^{-1} du$$

We aim to show that x satisfies pde 4.1. We compute

$$(4.9) \quad y_f(t) = v(t)^{-1} \sigma(f(t))^{-1}$$

$$(4.10) \quad y_v(t) = -v(t)^{-1} y(t)$$

$$(4.11) \quad x_f(t) = J(y(t))^{-1} v(t)^{-1} \sigma(f(t))^{-1}$$

$$(4.12) \quad x_v(t) = -J(y(t))^{-1} v(t)^{-1} y(t)$$

Putting this into the rhs of 4.1 we get

$$(4.13) \quad J(y(t))^{-2} + J(y(t))^{-2} v(t)^{-2} y(t)^2 \epsilon(v)^2 -$$

$$(4.14) \quad 2J(y(t))^{-2} v(t)^{-2} y(t) \rho v(t) \epsilon(v) =$$

$$(4.15) \quad J(y(t))^{-2} (1 + \eta^2 y(t)^2 - 2\rho \eta y(t)) = 1$$

which means that $x(t)$ is a solution of 4.1. We note that only the special form of $\epsilon(v) = \eta v$ was used to arrive here, whereas $\sigma(f)$ could have been arbitrary.

We explicitly compute $x(t)$ from its definition 4.8. Using the elementary identity

$$(4.16) \quad \int \frac{dh}{\sqrt{z^2 + a}} = \ln \frac{\sqrt{z^2 + a} + z}{\sqrt{a}}$$

which holds for $a > 0$ we get by a simple change of variable $\eta y(t) - \rho =: z$

$$(4.17) \quad x(t) = \frac{1}{\eta} \ln \frac{J(y(t)) + \eta y(t) - \rho}{1 - \rho}$$

where $y(t)$ is directly computed from its definition as

$$(4.18) \quad y(t) = \frac{f(t)^{1-\beta} - K^{1-\beta}}{v(t)(1-\beta)}$$

if $\beta < 1$ and

$$(4.19) \quad y(t) = \frac{1}{v(t)} \ln \frac{f(t)}{K}$$

if $\beta = 1$, where $v(0) = \alpha$. The implied normal volatility is then retrieved as above as

$$(4.20) \quad \nu(t) = \frac{f(t) - K}{x(t)}$$

This formula only holds for $K \neq f(t)$. The at the money case can be retrieved by differentiating the nominator and denominator by K and evaluating the result at $K = f(t)$. The derivative of the nominator is clearly -1 (in the lognormal case $-K^{-1}$). The denominator is handled as follows (subscripts denoting partial derivatives and omitting independent variables):

$$(4.21) \quad \frac{\partial x}{\partial K} = \frac{(1-\rho)(J_y y_K + \eta y_K)}{\eta(J + \eta y - \rho)(1-\rho)} = \frac{(J_y + \eta)y_K}{\eta(J + \eta y - \rho)}$$

$$(4.22) \quad J_y = \frac{2\eta^2 y - 2\rho\eta}{2\sqrt{1 + \eta^2 y^2 - 2\rho\eta y}}$$

$$(4.23) \quad y_K = -K^{-\beta} v^{-1}$$

Therefore

$$(4.24) \quad \frac{\partial x}{\partial K}(f(t)) = -f(t)^{-\beta} v(t)^{-1}$$

and in the normal case

$$(4.25) \quad \nu(t) = f(t)^\beta v(t)$$

where again $v(0) = \alpha$. In the lognormal case the rhs is replaced by $f(t)^{\beta-1} v(t)$. The same formulas hold for $\beta = 1$ as well.

The equivalent deterministic local volatility function 3.6 is computed using the definition of x and 4.21 as

$$(4.26) \quad -\left(\frac{\partial x}{\partial K}(f)\right)^{-1} = J(y)f^\beta v$$

We note that y has to be evaluated at $K = f$ here, while $f(t)$ in the definition of y in 4.2 stays the forward rate at time $t = 0$. Also, v means $v(t = 0)$ here.

5. THE ZABR MODEL

We generalize the SABR model by introducing a parameter $\gamma \in [0, \infty)$ and setting $\epsilon(v) = \eta v^\gamma$. For $\gamma = 1$ the original SABR model is reproduced.

An important note is that in [1] for the model formulation $v(0) = 1$ is assumed, while here we assume $v(0) = \alpha$. It is straightforward to see that the volvol parameter η' from [1] has to be transformed by

$$(5.1) \quad \eta = \eta' \alpha^{1-\gamma}$$

to get the volvol parameter η here.

As in the case of the SABR model we define a new variable y by setting

$$(5.2) \quad y(t) := v(t)^{\gamma-2} \int_K^{f(t)} \sigma(u)^{-1} du$$

and applying Ito to get

$$(5.3) \quad dy = y_f df + y_v dv + \dots dt =$$

$$(5.4) \quad v(t)^{\gamma-1} dW + (\gamma - 2)\eta v(t)^{\gamma-1} y dV + \dots dt$$

Setting $x(t) := v(t)^{1-\gamma} u(y(t))$ and again applying Ito yields

$$(5.5) \quad dx = x_y dy + x_v dv + \dots dt =$$

$$(5.6) \quad v(t)^{1-\gamma} u'(y) dy + (1 - \gamma)\eta u(y) dV + \dots dt =$$

$$(5.7) \quad u'(y) dW + ((\gamma - 2)y\eta u'(y) + (1 - \gamma)\eta u(y)) dV + \dots dt$$

Now

$$(5.8) \quad \frac{dx dx}{dt} = u'(y)^2 + ((\gamma - 2)y\eta u'(y) + (1 - \gamma)\eta u(y))^2 +$$

$$(5.9) \quad 2\rho u'(y)((\gamma - 2)y\eta u'(y) + (1 - \gamma)\eta u(y))$$

which is equal to 1 if u is a solution to the following ordinary differential equation:

$$(5.10) \quad 1 = A(y)u'(y)^2 + B(y)u'(y)u(y) + Cu(y)^2$$

$$(5.11) \quad A(y) = 1 + (\gamma - 2)^2 \eta^2 y^2 + 2\rho(\gamma - 2)\eta y$$

$$(5.12) \quad B(y) = 2\rho(1 - \gamma)\eta + 2(1 - \gamma)(\gamma - 2)\eta^2 y$$

$$(5.13) \quad C = (1 - \gamma)^2 \eta^2$$

subject to the initial condition $u(0) = 0$. This equation can be made explicit in u' :

$$(5.14) \quad u'(y) = F(y, u(y)) := \frac{-B(y)u(y) + \sqrt{B(y)^2 u(y)^2 - 4A(y)(Cu(y)^2 - 1)}}{2A(y)}$$

(... why do we discard the other solution here ...)

Suppose we have solved 5.14 then we can compute $x(t)$ directly from its definition $x(t) = v(t)^{1-\gamma} u(y(t))$ for all strikes K (note that $y(t)$ depends on K).

Also the equivalent deterministic local volatility function 3.6 is computed easily from $u(y(t))$ via

$$(5.15) \quad -\left(\frac{\partial x}{\partial K}(f)\right)^{-1}$$

with

$$(5.16) \quad \frac{\partial x}{\partial K}(f) = -v(t)^{1-\gamma} u'(y(t)) v(t)^{\gamma-2} \sigma(f)^{-1} =$$

$$(5.17) \quad -v(t)^{-1} \sigma(f)^{-1} F(y, v(t)^{\gamma-1} x(t))$$

using 5.14, which leads to

$$(5.18) \quad -\left(\frac{\partial x}{\partial K}(f)\right)^{-1} = v \sigma(f) F(y, v^{\gamma-1} x)^{-1}$$

We note that the special case formula 4.24 stays the same for the ZABR model as for the SABR model, because $F(0,0) = 1$, therefore $\partial x / \partial K = -v(t)^{-1} \sigma(f)^{-1}$. Also the same remarks concerning the evaluation of y and v as given for 4.26 apply here.

6. DUPIRE PRICING

Given a model (which in this context here will be an equivalent deterministic local volatility model generated from a SABR or ZABR model, see above) of the form

$$(6.1) \quad df(t) = \theta(f(t)) dW$$

it is possible to use the following pde derived by Dupire to price a european call option $c(T, K)$ with strike K and maturity T ,

$$(6.2) \quad \frac{\partial c}{\partial T} = \frac{\partial^2 c}{\partial K^2} \frac{\theta(K)^2}{2}$$

with initial condition $c(0, K) = (f(0) - K)^+$. For fixed maturity T this will give simultaneously call prices for all strikes K .

7. EXACT FINITE DIFFERENCE PRICES

In this section we derive the pde to price a single european call in the ZABR model. The derivation is independent in the sense that it does not make use of any of the approximations in the previous sections. The result can be used to produce benchmark prices, which are exact apart from errors introduced in the numerical procedures.

Starting with definition 1.1 consider a derivative price $c(t, f(t), v(t))$ which expands to

$$(7.1) \quad dc = c_t dt + c_f df + c_v dv + \frac{1}{2}(c_{ff} df df + c_{vv} dv dv) + c_{fv} df dv$$

$$(7.2) \quad = (c_t + \frac{1}{2}(c_{ff}\sigma(f)^2 v^2 + c_{vv}\epsilon(v)^2) + c_{fv}\sigma(f)v\epsilon(v)\rho)dt$$

$$(7.3) \quad + \dots dW + \dots dV$$

Using $dc = 0$ this yields to the pde for $c(t, f, v)$

$$(7.4) \quad c_t + \frac{1}{2}(c_{ff}\sigma(f)^2 v^2 + c_{vv}\epsilon(v)^2) + c_{fv}\sigma(f)v\epsilon(v)\rho = 0$$

with conditions

$$(7.5) \quad c(T, f, \cdot) = (f - K)^+$$

8. NUMERICAL EXAMPLES

8.1. Classic Hagan and short maturity expansions. We recompute the example from [1] (page 7, figure 1) with parameters $\alpha = 0.0873, \beta = 0.7, \nu = 0.47, \rho = -0.47$, forward rate $f = 0.0325$ (the forward is actually not mentioned in [1], but we believe this value was used there) and expiry time $t = 10.0$. The resulting log-normal implied volatilities are displayed in figure 1. The corresponding densities are plotted in figure 2





FIGURE 2. Densities, Classic Hagan formula and short maturity expansion



We reproduce figure 3 from [1]. We believe the short maturity expansion for normal volatility was used. Our result is depicted in figure 3

8.2. Equivalent deterministic local volatility. We use 4.26 or 5.18 to compute the equivalent deterministic local volatility (with the same parameters as in the previous section). The result is displayed in figure 4 which compares to [1], figure 2.

Next we use this local vol and 6.2 to produce a volatility smile. The result in terms of densities is shown in 5 together with the classic Hagan formula implied density. Clearly the local volatility approach removes the defect of the Hagan formula for strikes near zero and produces a globally arbitrage free smile.

REFERENCES

- [1] Andreasen, Jesper and Høge, Brian: ZABR – Expansions for the Masses, SSRN id 1980726, December 2011
- [2] QuantLib A free/open-source library for quantitative finance, <http://www.quantlib.org>

FIGURE 3. Normal Short Maturity Expansion ZABR model with different gammas




FIGURE 4. Equivalent Local volatility for SABR model




FIGURE 5. Densities from Dupire pricing and Classic Hagan formula

