

Nonlinear mobility continuity equations and generalized displacement convexity

J. A. Carrillo*, S. Lisini†, G. Savaré‡, D. Slepčev§

January 30, 2009

Abstract

We consider the geometry of the space of Borel measures endowed with a distance that is defined by generalizing the dynamical formulation of the Wasserstein distance to concave, nonlinear mobilities. We investigate the energy landscape of internal, potential, and interaction energies. For the internal energy, we give an explicit sufficient condition for geodesic convexity which generalizes the condition of McCann. We take an eulerian approach that does not require global information on the geodesics. As by-product, we obtain existence, stability, and contraction results for the semigroup obtained by solving the homogeneous Neumann boundary value problem for a nonlinear diffusion equation in a convex bounded domain. For the potential energy and the interaction energy, we present a non-rigorous argument indicating that they are not displacement semiconvex.

Keywords: gradient flows, displacement convexity, nonlinear diffusion equations, parabolic equations, Wasserstein distance, nonlinear mobility.

1 Introduction

Displacement convexity and Wasserstein distance. In [McC97], McCann introduced the notion of *displacement convexity* for integral functionals of the form

$$\mathcal{U}(\mu) := \int_{\Omega} U(\rho(x)) \, dx \quad \text{if } \mu = \rho \mathcal{L}^d, \quad U : [0, +\infty) \rightarrow \mathbb{R} \text{ is a convex function,}$$

*ICREA (Institució Catalana de Recerca i Estudis Avançats) and Departament de Matemàtiques, Universitat Autònoma de Barcelona, E-08193 Bellaterra, Spain. E-mail: carrillo@mat.uab.es.

†Dipartimento di Matematica “F. Casorati”, Università degli Studi di Pavia, Italy. E-mail: stefano.lisini@unipv.it.

‡Dipartimento di Matematica “F. Casorati”, Università degli Studi di Pavia, Italy. E-mail: giuseppe.savare@unipv.it.

§Department of Mathematical Sciences, Carnegie Mellon University, Pittsburgh, USA. E-mail: slepcev@math.cmu.edu.

defined on the set $\mathcal{P}_{ac}(\Omega)$ of the Borel probability measures in a convex open domain $\Omega \subset \mathbb{R}^d$, which are absolutely continuous with respect to the Lebesgue measure \mathcal{L}^d . Displacement convexity of \mathcal{U} means convexity along a particular class of curves, given by *displacement interpolation* between two given measures. These curves turned out to be the geodesics of the space $\mathcal{P}_{ac}(\Omega)$ endowed with the euclidean Wasserstein distance.

We recall that the Wasserstein distance W between two Borel probability measures μ_0 and μ_1 on Ω is defined by the following optimal transportation problem (Kantorovitch relaxed version)

$$W^2(\mu_0, \mu_1) := \min \left\{ \int_{\Omega \times \Omega} |x - y|^2 d\gamma(x, y) : \gamma \in \Gamma(\mu_0, \mu_1) \right\},$$

where $\Gamma(\mu_0, \mu_1)$ is the set of admissible plans/couplings between μ_0 and μ_1 , that is the set of all Borel probability measures on $\Omega \times \Omega$ with first marginal μ_0 and second marginal μ_1 .

We introduce the “pressure” function P , defined by

$$P(r) := rU'(r) - (U(r) - U(0)) = \int_0^r sU''(s) ds \quad \text{so that} \quad P'(r) = rU''(r), \quad P(0) = 0. \quad (1.1)$$

The main result of [McC97] states that under the assumption

$$P'(r)r \geq (1 - 1/d)P(r) \geq 0, \quad \forall r \in (0, +\infty), \quad (1.2a)$$

or, equivalently,

$$r \mapsto \frac{P(r)}{r^{1-1/d}} \quad \text{is nonnegative and nondecreasing on } (0, +\infty),$$

the functional \mathcal{U} is convex along the constant speed geodesics induced by W , i.e. for every curve $(\mu_s)_{s \in [0,1]} \subset \mathcal{P}_{ac}(\Omega)$ satisfying

$$W(\mu_{s_1}, \mu_{s_2}) = |s_1 - s_2| W(\mu_0, \mu_1) \quad \forall s_1, s_2 \in [0, 1], \quad (1.3)$$

the map $s \mapsto \mathcal{U}(\mu_s)$ is convex in $[0, 1]$. This class of curves can be, equivalently, defined by displacement interpolation, using the Brenier’s optimal transportation map pushing μ_0 onto μ_1 (see [Vil03], for example). For power-like functions U, P

$$U(\rho) = \begin{cases} \frac{1}{\beta-1} \rho^\beta & \text{if } \beta \neq 1, \\ \rho \log \rho & \text{if } \beta = 1, \end{cases} \quad P(\rho) = \rho^\beta, \quad (1.2a) \text{ is equivalent to } \beta \geq 1 - 1/d. \quad (1.4)$$

The link with a nonlinear diffusion equation. Among the various applications of this property, a remarkable one concerns a wide class of nonlinear diffusion equations. The seminal work of Otto [Ot01] contributed the key idea that a solution of the nonlinear diffusion equation

$$\partial_t \rho - \nabla \cdot (\rho \nabla U'(\rho)) = 0 \quad \text{in } (0, +\infty) \times \Omega, \quad (1.5)$$

with homogeneous Neumann boundary condition on $\partial\Omega$ can be interpreted as the trajectory of the *gradient flow* of \mathcal{U} with respect to the Wasserstein distance. This means that the equation

is formally the gradient flow of \mathcal{U} with respect to the local metric which for a tangent vector s has the form

$$\langle s, s \rangle_\rho = \int_\Omega \rho |\nabla p|^2 dx \quad \text{where} \quad \begin{cases} -\nabla \cdot (\rho \nabla p) = s & \text{in } \Omega \\ \nabla p \cdot \mathbf{n} = 0 & \text{on } \partial\Omega \end{cases}$$

where \mathbf{n} is a unit normal vector to $\partial\Omega$. Let us note here that the equation (1.5) corresponds via (1.1) to

$$\partial_t \rho - \Delta P(\rho) = 0. \quad (1.6)$$

In particular, the heat equation, for $P(\rho) = \rho$, is the gradient flow of the logarithmic entropy $\mathcal{U}(\rho) = \int_\Omega \rho \log \rho dx$. Let us also note that the metric above satisfies

$$\langle s, s \rangle_\rho = \inf \left\{ \int_\Omega \rho |\mathbf{v}|^2 dx : s + \nabla \cdot (\rho \mathbf{v}) = 0 \text{ in } \Omega \text{ and } \mathbf{v} \cdot \mathbf{n} = 0 \text{ on } \partial\Omega \right\}.$$

The key property of this metric is that the length of the minimal geodesic between given two measures is nothing but the Wasserstein distance. More precisely

$$W^2(\mu_0, \mu_1) = \inf \left\{ \int_0^1 \int_{\mathbb{R}^d} |\mathbf{v}_s(x)|^2 \rho_s(x) dx ds : \partial_s \rho + \nabla \cdot (\rho \mathbf{v}) = 0 \text{ in } (0, 1) \times \mathbb{R}^d, \right. \\ \left. \text{supp}(\rho_s) \subset \overline{\Omega}, \quad \rho_0 \mathcal{L}^d = \mu_0, \quad \rho_1 \mathcal{L}^d = \mu_1 \right\}.$$

This *dynamical formulation* of the Wasserstein distance was rigorously established by Benamou and Brenier in [BB00] and extended to more general situations in [AGS05] and [L07].

As for the classical gradient flows of convex functions in euclidean spaces, the flow associated with (1.5) is a contraction with respect to the Wasserstein distance. In [AGS05] the authors showed that one of the possible ways to rigorously express the link between the functional \mathcal{U} , the distance W , and the solution of the diffusion equation (1.5) is given by the *evolution variational inequality* satisfied by the measures $\mu_t = \rho(t, \cdot) \mathcal{L}^d$ associated with (1.5):

$$\frac{1}{2} \frac{d^+}{dt} W^2(\mu_t, \nu) \leq \mathcal{U}(\nu) - \mathcal{U}(\mu_t) \quad \forall \nu \in \mathcal{P}_{ac}(\Omega). \quad (1.7)$$

A new class of “dynamical” distances. In a number of problems from mathematical biology [H03, BFD06, BD09, DR09], mathematical physics [K93, K95, F04, F05, SSC06, CLR08, CRS08], studies of phase segregation [GL1, SI08], and studies of thin liquid films [Ber98], the mobility of “particles” depends on the density ρ itself. More precisely the local metric in the configuration space is formally given as follows: For a tangent vector s (euclidean variation)

$$\langle s, s \rangle_\rho = \int_\Omega \mathbf{m}(\rho) |\nabla p|^2 dx \quad \text{where} \quad \begin{cases} -\nabla \cdot (\mathbf{m}(\rho) \nabla p) = s & \text{in } \Omega \\ \nabla p \cdot \mathbf{n} = 0 & \text{on } \partial\Omega \end{cases}$$

where $\mathbf{m} : [0, +\infty) \rightarrow [0, +\infty)$ is the *mobility function*. The global distance generated by the local metric is given by

$$\mathcal{W}_{\mathbf{m}, \Omega}^2(\mu_0, \mu_1) := \inf \left\{ \int_0^1 \int_{\mathbb{R}^d} |\mathbf{v}_s(x)|^2 \mathbf{m}(\rho_s(x)) dx ds : \right. \\ \left. \partial_s \rho + \nabla \cdot (\mathbf{m}(\rho) \mathbf{v}) = 0 \text{ in } (0, 1) \times \mathbb{R}^d, \text{supp}(\rho_s) \subset \overline{\Omega}, \quad \rho_0 \mathcal{L}^d = \mu_0, \quad \rho_1 \mathcal{L}^d = \mu_1 \right\}. \quad (1.8)$$

This distance was recently introduced and studied in [DNS09] in the case when \mathbf{m} is concave and nondecreasing. Similarly to the case $\mathbf{m}(r) = r$, it is easy to check formally that the trajectory of the *gradient flow* of \mathcal{U} with respect to the modified distance $\mathcal{W}_{\mathbf{m},\Omega}$ solves

$$\partial_t \rho - \nabla \cdot (\mathbf{m}(\rho) \nabla U'(\rho)) = 0 \quad \text{in } (0, +\infty) \times \Omega \quad (1.9)$$

with homogeneous Neumann boundary conditions on $\partial\Omega$. Assuming that $U''\mathbf{m}$ and $U''\mathbf{m}\mathbf{m}'$ are locally integrable, we can define in this case the function P and the auxiliary function H by

$$P(r) := \int_0^r U''(z) \mathbf{m}(z) \, dz, \quad H(r) := \int_0^r U''(z) \mathbf{m}(z) \mathbf{m}'(z) \, dz = \int_0^r P'(z) \mathbf{m}'(z) \, dz,$$

so that

$$P' = \mathbf{m} U'', \quad H' = \mathbf{m}' P', \quad P(0) = H(0) = 0,$$

and, at least for smooth solutions, the problem (1.9) is equivalent to (1.6).

By means of a formal computation, detailed in Section 2, the second derivative of the internal energy functional \mathcal{U} along a geodesic curve $(\mu_s)_{s \in [0,1]}$ satisfying as in (1.3)

$$\mathcal{W}_{\mathbf{m},\Omega}(\mu_{s_1}, \mu_{s_2}) = |s_1 - s_2| \mathcal{W}_{\mathbf{m},\Omega}(\mu_0, \mu_1) \quad \forall s_1, s_2 \in [0, 1],$$

is nonnegative, i.e. $\frac{d^2}{ds^2} \mathcal{U}(\mu_s) \geq 0$, if the following generalization of McCann condition (1.2a,b) holds

$$P'(r) \mathbf{m}(r) \geq (1 - 1/d) H(r) \geq 0, \quad \forall r \in (0, +\infty). \quad (1.10a)$$

It can also be expressed by requiring that

$$r \mapsto \frac{H(r)}{\mathbf{m}^{1-1/d}(r)} = \frac{1}{\mathbf{m}^{1-1/d}(r)} \int_0^r P'(s) \mathbf{m}'(s) \, ds \quad \text{is nondecreasing in } (0, +\infty).$$

As in the case of the Wasserstein distance, in dimension $d = 1$ the condition (1.10a) reduces to the usual convexity of U . In dimension $d \geq 2$, still considering the relevant example of power-like functions U, P, \mathbf{m} as in (1.4), we get

$$U(\rho) = \begin{cases} \frac{1}{\beta-1} \rho^\beta & \text{if } \beta \neq 1 \\ \rho \log \rho & \text{if } \beta = 1 \end{cases}, \quad \mathbf{m}(\rho) = \rho^\alpha, \quad P(\rho) = \frac{\beta}{\gamma} \rho^\gamma, \quad \gamma := \alpha + \beta - 1$$

and condition (1.10a) is equivalent to

$$\alpha \in (0, 1], \quad \gamma \geq 1 - \alpha/d.$$

In this case the heat equation corresponds to $\gamma = \alpha + \beta - 1 = 1$ and it is therefore the gradient flow of the functional

$$\mathcal{U}(\rho) = \frac{1}{(2 - \alpha)(1 - \alpha)} \int_\Omega \rho^{2-\alpha} \, dx$$

with respect to the distance $\mathcal{W}_{\mathbf{m},\Omega}$ induced by the mobility function $\mathbf{m}(\rho) = \rho^\alpha$.

Another interesting example, still leading to the heat equation, is represented by the functional

$$\mathcal{U}(\rho) = \int_{\Omega} \left(\rho \log \rho + (1 - \rho) \log(1 - \rho) \right) dx, \quad 0 \leq \rho \leq 1 \quad \mathcal{L}^d\text{-a.e. in } \Omega,$$

and the distance induced by $\mathbf{m}(\rho) = \rho(1 - \rho)$, $\rho \in [0, 1]$. Notice that in this case the positivity domain of the mobility \mathbf{m} is the finite interval $[0, 1]$, a case that has not been explicitly considered in [DNS09], but that can be still covered by a careful analysis (see [LM]).

Geodesic convexity and contraction properties. Our aim is to prove rigorously the geodesic convexity of the integral functional \mathcal{U} under conditions (1.10a,b) and the metric characterization of the nonlinear diffusion equation (1.9) as the gradient flow of \mathcal{U} with respect to the distance $\mathcal{W}_{\mathbf{m},\Omega}$ (1.8). If one tries to follow the same strategy which has been developed in the more familiar Wasserstein framework, one immediately finds a serious technical difficulty, due to the lackness of an “explicit” representation of the geodesics for $\mathcal{W}_{\mathbf{m},\Omega}$. In fact, the McCann’s proof of the displacement convexity of the functionals \mathcal{U} is strictly related to the canonical representation of the Wasserstein geodesics in terms of optimal transport maps.

Existence of a minimal geodesic connecting two measures at a finite $\mathcal{W}_{\mathbf{m},\Omega}$ distance has been proved by [DNS09]. However, an explicit representation is no longer available. On the other hand in [DS08], following the eulerian approach introduced in [OW05], the authors presented a new proof of McCann’s convexity result for integral functionals defined on a compact manifold without the use of the representation of geodesics. Here, following the same approach of [DS08], we reverse the usual strategy which derives the existence and the contraction property of the gradient flow of a functional from its geodesic convexity. On the contrary, we show that under the assumption (1.10a) smooth solutions of (1.9) satisfy the following Evolution Variational Inequality analogous to (1.7)

$$\frac{1}{2} \frac{d^+}{dt} \mathcal{W}_{\mathbf{m},\Omega}^2(\mu_t, \nu) \leq \mathcal{U}(\nu) - \mathcal{U}(\mu_t), \quad \forall t \in [0, +\infty), \quad \forall \nu \in \mathcal{P}(\Omega) : \mathcal{W}_{\mathbf{m}}(\nu, \mu_0) < +\infty. \quad (1.11)$$

This is sufficient to construct a nice gradient flow generated by \mathcal{U} and metrically characterized by (1.11), as showed in [AGS05] and [AS06]. The remarkable fact proved by [DS08] is that whenever a functional \mathcal{U} admits a flow, defined at least in a dense subset of $D(\mathcal{U})$, satisfying (1.11), the functional itself is convex along the geodesics induced by the distance $\mathcal{W}_{\mathbf{m},\Omega}$. As a by-product we obtain stability, uniqueness, and regularization results for the solutions of the problem (1.9) in a suitable subspace of $\mathcal{P}(\Omega)$ metrized by $\mathcal{W}_{\mathbf{m},\Omega}$.

Concerning the assumptions on \mathbf{m} , its *concavity* is a necessary and sufficient condition to write the definition of $\mathcal{W}_{\mathbf{m},\Omega}$ with a jointly convex integrand [DNS09], which is crucial in many properties of the distance, in particular for its lower semicontinuity with respect to the usual weak convergence of measures. Since $\mathbf{m} \geq 0$ on $[0, \infty)$ the concavity implies that the mobility must be nondecreasing. This is the case considered in [DNS09]. However we are also able to treat the case when the mobility is defined on an interval $[0, M)$ where it is nonnegative and concave. In that case the configuration space is restricted to absolutely continuous measures with densities bounded from above by M . Such mobilities are of particular interest in applications as mentioned before.

Plan of the paper. In next section, we show the heuristic computations for the convexity of functionals with respect to $\mathcal{W}_{\mathbf{m},\Omega}$. Section 3 is devoted to introduce the notation and to review the needed concepts on $\mathcal{W}_{\mathbf{m},\Omega}$ from [DNS09]. Moreover, we prove a key technical regularization lemma: Lemma 3.5. Subsection 3.4 addresses the question of finiteness of $\mathcal{W}_{\mathbf{m},\Omega}(\mu_0, \mu_1)$, providing new sufficient conditions on \mathbf{m} and μ_0, μ_1 in order to ensure that $\mathcal{W}_{\mathbf{m},\Omega}(\mu_0, \mu_1) < +\infty$. After a brief review of some basic properties of the diffusion equation (1.6), in Section 4 we try to get some insight on the features of the generalized McCann condition (1.10a,b), we recall some basic facts on the metric characterization of contracting gradient flows and their relationships with geodesic convexity borrowed from [AGS05, DS08], and we state our main results Theorems 4.11 and 4.13. The core of our argument in smooth settings is collected in Section 5, whereas the last Section concludes the proofs of the main results. At the end of the paper we collect some final remarks and open problems.

2 Heuristics

We first discuss, in a formal way, the conditions for the displacement convexity of the internal, the potential and the interaction energy, with respect to the geodesics corresponding to the distance (1.8). For simplicity, we assume that $\Omega = \mathbb{R}^d$ and that densities are smooth and decaying fast enough at infinity so that all computations are justified.

2.1 Geodesics

We first obtain the optimality condition for the geodesic equations in the fluid dynamical formulation of the new distance (1.8). As in [B03], we insert the nonlinear mobility continuity equation (1.8)

$$\partial_s \rho + \nabla \cdot (\mathbf{m}(\rho) \mathbf{v}) = 0 \quad \text{in } (0, 1) \times \mathbb{R}^d. \quad (2.1)$$

inside the minimization problem as a Lagrange multiplier. As a result, we get the unconstrained minimization problem

$$\begin{aligned} \mathcal{W}_{\mathbf{m}}^2(\mu_0, \mu_1) = \inf_{(\rho, \mathbf{v})} \sup_{\psi} \Big\{ & \int_0^1 \int_{\mathbb{R}^d} \frac{1}{2} |\mathbf{v}_s(x)|^2 \mathbf{m}(\rho_s(x)) \, dx \, ds \\ & - \int_0^1 \int_{\mathbb{R}^d} [\rho_s(x) \partial_s \psi(s, x) + \mathbf{m}(\rho_s(x)) (\mathbf{v}_s(x) \cdot \nabla \psi(s, x))] \, dx \, ds \\ & + \int_{\mathbb{R}^d} \rho_1(x) \psi(1, x) \, dx - \int_{\mathbb{R}^d} \rho_0(x) \psi(0, x) \, dx \Big\}. \end{aligned}$$

Applying a formal minimax principle and thus taking first an infimum with respect to \mathbf{v} we obtain the optimality condition $\mathbf{v} = \nabla \psi$, and the following formal characterization of the distance

$$\begin{aligned} \mathcal{W}_{\mathbf{m}}^2(\mu_0, \mu_1) = \sup_{\psi} \inf_{\rho} \Big\{ & - \frac{1}{2} \int_0^1 \int_{\mathbb{R}^d} |\nabla \psi|^2 \mathbf{m}(\rho) \, dx \, ds - \int_0^1 \int_{\mathbb{R}^d} \rho \partial_s \psi \, dx \, ds \\ & + \int_{\mathbb{R}^d} \rho_1(x) \psi(1, x) \, dx - \int_{\mathbb{R}^d} \rho_0(x) \psi(0, x) \, dx \Big\}, \end{aligned}$$

which provides the further optimality condition

$$\partial_s \psi + \frac{1}{2} \mathbf{m}'(\rho_s(x)) |\nabla \psi|^2 = 0. \quad (2.2)$$

We thus end up with a coupled system of differential equations in $(0, 1) \times \mathbb{R}^d$ [DNS09, Rem. 5.19]

$$\begin{cases} \partial_s \rho + \nabla \cdot (\mathbf{m}(\rho) \nabla \psi) = 0, \\ \partial_s \psi + \frac{1}{2} \mathbf{m}'(\rho) |\nabla \psi|^2 = 0. \end{cases} \quad (2.3)$$

2.2 Internal energy

We use the formal equations (2.3) for the geodesics associated to the distance (1.8) to compute the conditions under which the internal energy functional is displacement convex. If therefore (ρ_s, ψ_s) is a smooth solution of (2.3), which decays sufficiently at infinity, we proceed as usual [CMV03, Vil03, OW05] to obtain the following formulas:

$$\frac{d}{ds} \mathcal{U}(\rho) = - \int_{\mathbb{R}^d} P(\rho) \Delta \psi \, dx,$$

and

$$\begin{aligned} \frac{d^2}{ds^2} \mathcal{U}(\rho) &= \int_{\mathbb{R}^d} (P'(\rho) \mathbf{m}(\rho) - H(\rho)) (\Delta \psi)^2 \, dx \\ &\quad + \int_{\mathbb{R}^d} H(\rho) (-\nabla \psi \cdot \nabla \Delta \psi + \frac{1}{2} \Delta |\nabla \psi|^2) \, dx \\ &\quad - \frac{1}{2} \int_{\mathbb{R}^d} P'(\rho) \mathbf{m}''(\rho) |\nabla \rho|^2 |\nabla \psi|^2 \, dx. \end{aligned}$$

As usual, the Bochner formula

$$-\nabla \psi \cdot \nabla \Delta \psi + \frac{1}{2} \Delta |\nabla \psi|^2 = |\text{Hess } \psi|^2 \geq \frac{1}{d} (\Delta \psi)^2,$$

and the fact that $H(\rho) \geq 0$, allow us to estimate it as

$$\frac{d^2}{ds^2} \mathcal{U}(\rho) \geq \int_{\mathbb{R}^d} (P'(\rho) \mathbf{m}(\rho) - (1 - 1/d) H(\rho)) (\Delta \psi)^2 \, dx - \frac{1}{2} \int_{\mathbb{R}^d} P'(\rho) \mathbf{m}''(\rho) |\nabla \rho|^2 |\nabla \psi|^2 \, dx.$$

Therefore, under conditions of concavity of the mobility $\mathbf{m}(\rho)$ and the generalized displacement McCann's condition (1.10a), the functional \mathcal{U} is convex along the geodesics of the distance $\mathcal{W}_{\mathbf{m}}$.

2.3 Potential energy

Similar heuristic formulas can be obtained for the potential and the interaction energy, as in [CMV03, Vil03]. We consider the potential energy functional

$$\mathcal{V}(\mu) := \int_{\mathbb{R}^d} V(x) \, d\mu,$$

with V a given smooth potential. As before, it is easy to check that the second derivative of \mathcal{V} along a geodesic satisfying (2.3) is

$$\begin{aligned} \frac{d^2}{ds^2} \mathcal{V}(\rho) &= \int_{\mathbb{R}^d} \mathbf{m}(\rho) \mathbf{m}'(\rho) (\text{Hess } V \nabla \psi) \cdot \nabla \psi \, dx \\ &\quad + \int_{\mathbb{R}^d} \mathbf{m}(\rho) \mathbf{m}''(\rho) \left((\nabla \rho \cdot \nabla \psi) (\nabla V \cdot \nabla \psi) - \frac{1}{2} (\nabla \rho \cdot \nabla V) |\nabla \psi|^2 \right) dx. \end{aligned}$$

This formula allows us to show that this functional cannot be convex along geodesics if \mathbf{m} is not linear. Technically, the reason is the presence of the terms linearly depending on $\nabla \rho$. We present a simple example:

Example. Let us first construct the example in one dimension. The expression for the second derivative of the functional above reduces to

$$\frac{d^2}{ds^2} \mathcal{V}(\rho) = \int_{\mathbb{R}} \mathbf{m}(\rho) \mathbf{m}'(\rho) V_{xx} \psi_x^2 \, dx + \frac{1}{2} \int_{\mathbb{R}} \mathbf{m}(\rho) \mathbf{m}''(\rho) \rho_x V_x \psi_x^2 \, dx =: I + II$$

Consider the case that V is nontrivial. Then $V_x \neq 0$ on some interval. For notational simplicity, we assume that

$$V_x > 0 \text{ on } [-2, 2].$$

Since the mobility \mathbf{m} we are considering is not a linear function of ρ there exists $z > 0$ such that $\mathbf{m}''(z) \neq 0$. Again for notational simplicity, let us assume that

$$\mathbf{m}''(z) < 0 \text{ on } \left[\frac{1}{2}, \frac{3}{2} \right].$$

The fact that we chose V_x to be positive and \mathbf{m}'' negative is irrelevant because the sign of term II can be controlled by the sign of ρ_x . Let η be a piecewise linear function on \mathbb{R} :

$$\eta(x) = \begin{cases} \frac{3}{2} & \text{if } x < -\frac{1}{2} \\ \frac{1}{2} - x & \text{if } x \in \left[-\frac{1}{2}, \frac{1}{2} \right] \\ \frac{1}{2} & \text{if } x > \frac{1}{2}. \end{cases}$$

The fact that the function is Lipschitz, but not smooth is irrelevant; smooth approximations of the given η , can also be used in the construction. Let $\eta_\varepsilon(x) = \eta\left(\frac{x}{\varepsilon}\right)$. Let $\sigma \in C_0^\infty(\mathbb{R}, [0, 1])$, supported in $[-1, 1]$, such that $\sigma = 1$ on $\left[-\frac{1}{4}, \frac{1}{4}\right]$ and $\int_{\mathbb{R}} \sigma(x) dx = 1$. Let $\rho_\varepsilon = \sigma \eta_\varepsilon$. Note that $\int_{\mathbb{R}} \rho_\varepsilon dx = 1$. A typical profile of ρ_ε is given in Figure 1.

The test velocity (tangent vector at $s = 0$) we consider also needs to be localized near zero. A simple choice is $\psi_\varepsilon(0) = \eta_\varepsilon$. Let $\rho_\varepsilon(s)$ be the corresponding geodesics given by (2.1) and (2.2). Let us observe how, at $s = 0$, the terms I and II scale with ε :

$$\begin{aligned} I_\varepsilon &\leq \max_{z \in [0, 2]} \mathbf{m}(z) \mathbf{m}'(z) \max_{x \in [-1, 1]} V_{xx}(x) \frac{1}{\varepsilon^2} \varepsilon \sim \frac{1}{\varepsilon}, \\ II_\varepsilon &\leq -\frac{1}{2} \min_{z \in [\frac{1}{2}, \frac{3}{2}]} \mathbf{m}(z) |\mathbf{m}''(z)| \frac{1}{\varepsilon} \min_{x \in [-1, 1]} V_x \frac{1}{\varepsilon^2} \varepsilon \sim -\frac{1}{\varepsilon^2}. \end{aligned}$$




Figure 1: A profile at which the potential energy is not convex.

Thus, for ε small enough, $\frac{d^2}{ds^2}\big|_{s=0} \mathcal{V}(\rho_\varepsilon(s)) < 0$. Furthermore note that the square of the length of the tangent vector $\frac{d}{dt}\rho_\varepsilon(0)$ is

$$\int_{\mathbb{R}} m(\rho_\varepsilon(0)) (\partial_x \psi_\varepsilon)^2 dx \sim \frac{1}{\varepsilon}$$

Thus for any $\lambda \in \mathbb{R}$ there exists $\varepsilon > 0$ such that

$$\frac{d^2}{ds^2}\bigg|_{s=0} \mathcal{V}(\rho_\varepsilon(s)) + \lambda \int_{\mathbb{R}} m(\rho_\varepsilon(0)) (\partial_x \psi_\varepsilon)^2 dx < 0$$

which implies that \mathcal{V} is not λ -convex for any $\lambda \in \mathbb{R}$.

Let us conclude the example by remarking that it can be extended to multidimensional domains. In particular it suffices to extend the 1-D profile to d-D to be constant in every other direction and then use a cut-off. We only sketch the elements of the construction.

We can assume that $\nabla V(0) = e_d$. Let $\tilde{\rho}_\varepsilon(x) = \rho_\varepsilon(x_d)$. Let $\hat{x} = (x_1, \dots, x_{d-1})$. To cut-off in the directions perpendicular to e_d we use the length scales $1 \gg l \gg \delta \gg \varepsilon$. Let $\theta_{l,\delta}$ be smooth cut-off function equal to 1 on $[-l, l]$ and equal to 0 outside of $[-l - \delta, l + \delta]$; with $|\nabla \theta_{l,\delta}| < \frac{C}{\delta}$ and $|D^2 \theta_{l,\delta}| < \frac{C}{\delta^2}$. Let $\rho_{l,\delta,\varepsilon}(x) = \tilde{\rho}_\varepsilon(x_d) \theta_{l,\delta}(|\hat{x}|)$. Let $\psi_{l,\delta,\varepsilon}(x) = \psi_\varepsilon(x_d) \theta_{l,\delta}(|\hat{x}|)$. Checking the scaling of appropriate terms is straightforward.

2.4 Interaction energy

Consider the interaction energy functional

$$\mathcal{W}(\rho) := \frac{1}{2} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} W(x - y) \rho(x) \rho(y) dx dy,$$

with W a given smooth potential. As before, it is easy to check that

$$\begin{aligned} \frac{d^2}{ds^2} \mathcal{W}(\rho) &= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \mathbf{m}(\rho(x)) \mathbf{m}'(\rho(x)) \rho(y) \nabla \psi(x) \cdot (\text{Hess } W(x-y) \nabla \psi(x)) dx dy \\ &\quad - \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \mathbf{m}(\rho(x)) \mathbf{m}(\rho(y)) \nabla \psi(y) \cdot (\text{Hess } W(x-y) \nabla \psi(x)) dx dy \\ &\quad + \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \mathbf{m}(\rho(x)) \mathbf{m}''(\rho(x)) \rho(y) (\nabla \rho(x) \cdot \nabla \psi(x)) (\nabla W(x-y) \cdot \nabla \psi(x)) dx dy \\ &\quad - \frac{1}{2} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \mathbf{m}(\rho(x)) \mathbf{m}''(\rho(x)) \rho(y) (\nabla \rho(x) \cdot \nabla W(x-y)) |\nabla \psi(x)|^2 dx dy. \end{aligned}$$

It can be demonstrated that if \mathbf{m} is non-linear then the interaction energy is not geodesically convex. As for the potential energy, the reason lies in the presence of derivatives of ρ in the expression above. More precisely, in one dimension the second derivative of $\mathcal{W}(\rho)$ reduces to

$$\begin{aligned} \frac{d^2}{ds^2} \mathcal{W}(\rho) &= \int_{\mathbb{R}} \int_{\mathbb{R}} \mathbf{m}(\rho(x)) \mathbf{m}'(\rho(x)) \rho(y) \psi_x^2(x) W_{xx}(x-y) dx dy \\ &\quad - \int_{\mathbb{R}} \int_{\mathbb{R}} \mathbf{m}(\rho(x)) \mathbf{m}(\rho(y)) \psi_y(y) W_{xx}(x-y) \psi_x(x) dx dy \\ &\quad + \frac{1}{2} \int_{\mathbb{R}} \int_{\mathbb{R}} \mathbf{m}(\rho(x)) \mathbf{m}''(\rho(x)) \rho(y) \rho_x(x) \psi_x^2(x) W_x(x-y) dx dy. \end{aligned}$$

It turns out that the example for the lack of (semi-)convexity provided for the potential energy is also an example (with V replaced by W) for the interaction energy. The estimates of the terms are similar, so we leave the details to the reader.

3 Notation and preliminaries

In this section, following [DNS09], we shall recall the main properties of the distance $\mathcal{W}_{\mathbf{m},\Omega}$ introduced in (1.8). For the sake of simplicity, we only consider here the case of a bounded open domain Ω , so that it is not restrictive to assume that all the measures (Radon, i.e. locally finite, in the general approach of [DNS09]) involved in the various definitions have finite total variation. Since we deal with arbitrary mobility functions \mathbf{m} , these distances do not exhibit nice homogeneity properties as in the Wasserstein case; therefore we deal with finite Borel measures without assuming that their total mass is 1.

3.1 Measures and continuity equation

We denote by $\mathcal{M}^+(\mathbb{R}^d)$ (resp. $\mathcal{M}_c^+(\mathbb{R}^d)$) the space of finite positive Borel measures on \mathbb{R}^d (resp. with compact support) and by $\mathcal{M}(\mathbb{R}^d; \mathbb{R}^d)$ the space of \mathbb{R}^d -valued Borel measures on \mathbb{R}^d with finite total variation. By Riesz representation theorem, the space $\mathcal{M}(\mathbb{R}^d; \mathbb{R}^d)$ can be identified with the dual space of $C_c^0(\mathbb{R}^d; \mathbb{R}^d)$ and it is endowed with the corresponding weak* topology. We denote by $|\boldsymbol{\nu}| \in \mathcal{M}^+(\mathbb{R}^d)$ the total variation of the vector measure $\boldsymbol{\nu} \in \mathcal{M}(\mathbb{R}^d; \mathbb{R}^d)$. $\boldsymbol{\nu}$ admits the polar decomposition $\boldsymbol{\nu} = \mathbf{w}|\boldsymbol{\nu}|$ with $\mathbf{w} \in L^1(|\boldsymbol{\nu}|; \mathbb{R}^d)$. If B is a Borel subset of \mathbb{R}^d (typically an open

or closed set) we denote by $\mathcal{M}^+(B)$ (resp. $\mathcal{M}^+(B; \mathbb{R}^d)$) the subset of $\mathcal{M}^+(\mathbb{R}^d)$ (resp. $\mathcal{M}(\mathbb{R}^d; \mathbb{R}^d)$) whose measure μ are concentrated on B , i.e. $\mu(\mathbb{R}^d \setminus B) = 0$ (resp. $|\mu|(\mathbb{R}^d \setminus B) = 0$). Notice that if B is a compact subset of \mathbb{R}^d then the convex set in $\mathcal{M}^+(B)$ of measures with a fixed total mass \mathfrak{m} is compact with respect to the weak* topology. If $\mathfrak{m} > 0$, $\mathcal{M}^+(B, \mathfrak{m})$ is the convex subset of $\mathcal{M}^+(B)$ whose measures have fixed total mass $\mu(B) = \mathfrak{m}$.

Let Ω be a bounded open subset of \mathbb{R}^d . Given $\mu_0, \mu_1 \in \mathcal{M}^+(\overline{\Omega})$ we denote by $\mathcal{CE}_\Omega(\mu_0 \rightarrow \mu_1)$ the collection of time dependent measures $(\mu_s)_{s \in [0,1]} \subset \mathcal{M}^+(\overline{\Omega})$ and $(\nu_s)_{s \in (0,1)} \in \mathcal{M}(\overline{\Omega}; \mathbb{R}^d)$ such that

1. $s \mapsto \mu_s$ is weakly* continuous in $\mathcal{M}^+(\mathbb{R}^d)$ with $\mu|_{s=0} = \mu_0$ and $\mu|_{s=1} = \mu_1$;
2. $(\nu_s)_{s \in (0,1)}$ is a Borel family with $\int_0^1 |\nu_s|(\overline{\Omega}) ds < +\infty$;
3. (μ, ν) is a distributional solution of

$$\partial_s \mu_s + \nabla \cdot \nu_s = 0 \quad \text{in } (0, 1) \times \mathbb{R}^d.$$

If $(\mu, \nu) \in \mathcal{CE}_\Omega(\mu_0 \rightarrow \mu_1)$ then it is immediate to check that the total mass $\mu_s(\mathbb{R}^d) = \mu_s(\overline{\Omega}) = \mathfrak{m}$ is a constant, independent of s . In particular, $\mu_0(\mathbb{R}^d) = \mu_1(\mathbb{R}^d)$.

3.2 Mobility and action functional

We fix a right threshold $M \in (0, +\infty]$ and a concave *mobility function* $\mathfrak{m} \in C^0[0, M)$ strictly positive in $(0, M)$. We denote by $\mathfrak{m}(M)$ the left limit of $\mathfrak{m}(r)$ as $r \uparrow M$. We can also introduce the maximal left interval of monotonicity of \mathfrak{m} whose right extreme is

$$M_\uparrow := \sup \left\{ m \in [0, M) : \mathfrak{m}|_{[0, m]} \text{ is nondecreasing} \right\}.$$

We distinguish two situation:

Case A $M = +\infty$ so that \mathfrak{m} is nondecreasing and $M_\uparrow = M = +\infty$; typically $\mathfrak{m}(0) = 0$ and the main example is provided by $\mathfrak{m}(r) = r^\alpha$, $\alpha \in [0, 1]$. This is the case considered in [DNS09]. When $\mathfrak{m}'(+\infty) := \lim_{r \uparrow +\infty} r^{-1} \mathfrak{m}(r) = \lim_{r \uparrow +\infty} \mathfrak{m}'(r) = 0$ we are in the *sublinear growth* case. A *linear growth* of \mathfrak{m} corresponds to $\mathfrak{m}'(+\infty) > 0$.

Case B $M < +\infty$, so that $0 \leq M_\uparrow \leq M$ and \mathfrak{m} is nonincreasing in the right interval $[M_\uparrow, M)$ (but we also allow \mathfrak{m} to be constant or even decreasing in $[0, M)$ with $M_\uparrow = 0$). Typically $\mathfrak{m}(0) = \mathfrak{m}(M) = 0$ (in this case $0 < M_\uparrow < M$) and the main example is $\mathfrak{m}(r) = r(M - r)$, or, more generally, $\mathfrak{m}(r) = r^{\alpha_0}(M - r)^{\alpha_1}$, $\alpha_0, \alpha_1 \in (0, 1]$.

Many properties proved in the case A can be extended to the case B, but there are important exceptions: we refer to [LM] for further details. Using the conventions

$$\begin{aligned} a/b &= 0 & \text{if } a = b = 0, \\ a/b &= +\infty & \text{if } a > 0 = b, \end{aligned} \tag{3.1}$$

the corresponding *action density function* $\phi_m : \mathbb{R} \times \mathbb{R}^d \rightarrow [0, +\infty]$ is defined by

$$\phi_m(\rho, \mathbf{w}) = \begin{cases} \frac{|\mathbf{w}|^2}{m(\rho)} & \text{if } \rho \in [0, M], \\ +\infty & \text{if } \rho < 0 \text{ or } \rho > M. \end{cases}$$

It is not difficult to check that, under the convention (3.1), the function ϕ_m is (jointly) convex and lower semi-continuous.

Given that m is concave and ϕ_m is convex, when $M = +\infty$ we can define the recession function $\varphi_m^\infty : \mathbb{R}^d \mapsto [0, +\infty]$ (recall (3.1))

$$\varphi_m^\infty(\mathbf{w}) := \lim_{r \uparrow +\infty} r\phi_m(1, \mathbf{w}/r) = \frac{|\mathbf{w}|^2}{m'(\infty)}, \quad m'(\infty) := \lim_{r \rightarrow +\infty} m'(r) = \lim_{r \rightarrow +\infty} \frac{m(r)}{r} \geq 0.$$

We introduce now the *action functional*

$$\Phi_{m,\Omega} : \mathcal{M}^+(\mathbb{R}^d) \times \mathcal{M}(\mathbb{R}^d; \mathbb{R}^d) \rightarrow [0, +\infty],$$

defined on couples of measures $\mu \in \mathcal{M}^+(\mathbb{R}^d)$, $\nu \in \mathcal{M}(\mathbb{R}^d; \mathbb{R}^d)$. In order to define it we consider the usual Lebesgue decomposition $\mu = \rho \mathcal{L}^d + \mu^\perp$, $\nu = \mathbf{w} \mathcal{L}^d + \nu^\perp$ and distinguish the following cases:

1. If the support of μ or ν is not contained in $\overline{\Omega}$ then $\Phi_{m,\Omega}(\mu, \nu) = +\infty$;
2. When $M < +\infty$ (Case B), we set

$$\Phi_{m,\Omega}(\mu, \nu) := \begin{cases} \int_{\Omega} \phi_m(\rho, \mathbf{w}) \, dx & \text{if } \mu^\perp = 0, \nu^\perp = 0 \\ +\infty & \text{otherwise;} \end{cases}$$

notice that if $\Phi_{m,\Omega}(\mu, \nu) < +\infty$ then $\rho \in L^\infty(\Omega)$ with $0 \leq \rho \leq M$ \mathcal{L}^d -a.e. in Ω and $\mathbf{w} \in L^2(\Omega; \mathbb{R}^d)$.

3. When $M = +\infty$ and $m'(\infty) = 0$ (Case A, sublinear growth) then

$$\Phi_{m,\Omega}(\mu, \nu) := \begin{cases} \int_{\Omega} \phi_m(\rho, \mathbf{w}) \, dx & \text{if } \nu^\perp = 0 \\ +\infty & \text{otherwise;} \end{cases}$$

4. Finally, when $M = +\infty$ and $m'(\infty) > 0$ (Case A, linear growth) then we set

$$\Phi_{m,\Omega}(\mu, \nu) := \begin{cases} \int_{\Omega} \phi_m(\rho, \mathbf{w}) \, dx + \int_{\Omega} \varphi_m^\infty(\mathbf{w}^\perp) \, d\mu^\perp & \text{if } \nu^\perp = \mathbf{w}^\perp \mu^\perp \ll \mu^\perp \\ +\infty & \text{otherwise.} \end{cases}$$

3.3 The modified Wasserstein distance

Let Ω be a bounded open set. Given $\mu^0, \mu^1 \in \mathcal{M}^+(\overline{\Omega})$ we define

$$\mathcal{W}_{\mathbf{m}, \Omega}(\mu^0, \mu^1) := \inf \left\{ \left(\int_0^1 \Phi_{\mathbf{m}, \Omega}(\mu_s, \nu_s) ds \right)^{1/2} : (\mu, \nu) \in \mathcal{CE}_{\Omega}(\mu^0 \rightarrow \mu^1) \right\} \quad (3.2)$$

$$= \inf \left\{ \int_0^1 \left(\Phi_{\mathbf{m}, \Omega}(\mu_s, \nu_s) \right)^{1/2} ds : (\mu, \nu) \in \mathcal{CE}_{\Omega}(\mu^0 \rightarrow \mu^1) \right\}. \quad (3.3)$$

We refer to [DNS09, Thm. 5.4] for the equivalence between (3.2) and (3.3). $\mathcal{W}_{\mathbf{m}, \Omega}(\mu^0, \mu^1) = +\infty$ if the set of connecting curves $\mathcal{CE}_{\Omega}(\mu^0 \rightarrow \mu^1)$ is empty. The following three propositions are proved in [DNS09], see Theorems 5.5, 5.6, 5.7, 5.15, and Proposition 5.14.

Proposition 3.1. *The space $\mathcal{M}^+(\overline{\Omega})$ endowed with the distance $\mathcal{W}_{\mathbf{m}, \Omega}$ is a complete pseudo-metric space (the distance can assume the value $+\infty$), inducing as strong as, or stronger topology than the weak* one.*

Given a measure $\sigma \in \mathcal{M}^+(\overline{\Omega})$, the space $\mathcal{M}_{\mathbf{m}, \Omega}^+[\sigma] := \{\mu \in \mathcal{M}^+(\overline{\Omega}) : \mathcal{W}_{\mathbf{m}, \Omega}(\mu, \sigma) < +\infty\}$ is a complete metric space whose measures have the same total mass of σ .

Moreover, for every $\mu_0, \mu_1 \in \mathcal{M}^+(\overline{\Omega})$ such that $\mathcal{W}_{\mathbf{m}, \Omega}(\mu_0, \mu_1) < +\infty$ there exists a minimizing couple (μ, ν) in (3.2) (unique, if \mathbf{m} is strictly concave and sublinear) and the curve $(\mu_s)_{s \in [0, 1]}$ is a constant speed geodesic for $\mathcal{W}_{\mathbf{m}, \Omega}$, thus satisfying

$$\mathcal{W}_{\mathbf{m}, \Omega}(\mu_t, \mu_s) = |t - s| \mathcal{W}_{\mathbf{m}, \Omega}(\mu_0, \mu_1) \quad \forall s, t \in [0, 1].$$

Proposition 3.2 (Lower semi-continuity). *If Ω_n, Ω are bounded open sets such that $\mathcal{L}^d|_{\Omega_n}$ weakly* converges to $\mathcal{L}^d|_{\Omega}$, $M_n \in (0, +\infty]$ is a nonincreasing sequence converging to M , \mathbf{m}_n is a sequence of nonnegative concave functions in the intervals $(0, M_n)$ such that*

$$\mathbf{m}_{n'}(r) \geq \mathbf{m}_{n''}(r) \quad \forall r \in (0, M_{n''}) \quad \text{if } n' \leq n'', \quad \lim_{n \rightarrow \infty} \mathbf{m}_n(r) = \mathbf{m}(r) \quad \forall r \in (0, M),$$

and μ_0^n, μ_1^n are sequences of measures weakly convergent to μ_0 and μ_1 respectively, then*

$$\liminf_{n \rightarrow +\infty} \mathcal{W}_{\mathbf{m}_n, \Omega_n}(\mu_0^n, \mu_1^n) \geq \mathcal{W}_{\mathbf{m}, \Omega}(\mu_0, \mu_1). \quad (3.4)$$

Proposition 3.3 (Monotonicity). *Let $\tilde{\Omega} \supset \Omega$, $\tilde{\mathbf{m}} \geq \mathbf{m}$, $\mu_0, \mu_1 \in \mathcal{M}^+(\overline{\Omega})$. Then the following inequality holds*

$$\mathcal{W}_{\tilde{\mathbf{m}}, \tilde{\Omega}}(\mu_0, \mu_1) \leq \mathcal{W}_{\mathbf{m}, \Omega}(\mu_0, \mu_1).$$

Proposition 3.4. *Let $k \in C_c^\infty(\mathbb{R}^d)$ be a nonnegative convolution kernel, with $\int_{\mathbb{R}^d} k(x) dx = 1$ and $\text{supp}(k) = \overline{B}_1(0)$, and let $k_\varepsilon(x) := \varepsilon^{-d} k(x/\varepsilon)$. For every $\mu, \mu_0, \mu_1 \in \mathcal{M}^+(\overline{\Omega})$ and $\nu \in \mathcal{M}(\overline{\Omega}; \mathbb{R}^d)$ we have*

$$\begin{aligned} \Phi_{\mathbf{m}, \Omega_\varepsilon}(\mu * k_\varepsilon, \nu * k_\varepsilon) &\leq \Phi_{\mathbf{m}, \Omega}(\mu, \nu) & \forall \varepsilon > 0, \\ \mathcal{W}_{\mathbf{m}, \Omega_\varepsilon}(\mu_0 * k_\varepsilon, \mu_1 * k_\varepsilon) &\leq \mathcal{W}_{\mathbf{m}, \Omega}(\mu_0, \mu_1) & \forall \varepsilon > 0, \end{aligned} \quad (3.5)$$

$$\lim_{\varepsilon \rightarrow 0} \mathcal{W}_{\mathbf{m}, \Omega_\varepsilon}(\mu_0 * k_\varepsilon, \mu_1 * k_\varepsilon) = \mathcal{W}_{\mathbf{m}, \Omega}(\mu_0, \mu_1), \quad (3.6)$$

where $\Omega_\varepsilon := \Omega + B_\varepsilon(0)$.

Proof. If $\Phi_{\mathbf{m},\Omega}(\mu, \nu) < +\infty$ then μ, ν are supported in $\overline{\Omega}$ and [DNS09, Theorem 2.3] yields

$$\Phi_{\mathbf{m},\Omega}(\mu, \nu) = \Phi_{\mathbf{m},\mathbb{R}^d}(\mu, \nu) \geq \Phi_{\mathbf{m},\mathbb{R}^d}(\mu * k_\varepsilon, \nu * k_\varepsilon) = \Phi_{\mathbf{m},\Omega_\varepsilon}(\mu * k_\varepsilon, \nu * k_\varepsilon),$$

being $\mu * k_\varepsilon, \nu * k_\varepsilon$ supported in $\overline{\Omega}_\varepsilon$. Notice that only the concavity of \mathbf{m} (and not its monotonicity) plays a role here. A similar argument and [DNS09, Theorem 5.15] yields (3.5). The limit (3.6) is an immediate consequence of (3.4) and (3.5). \square

The next technical lemma provides a crucial approximation result for curves with finite $\Phi_{\mathbf{m},\Omega}$ energy. It allows for measures to be approximated by ones with smooth, positive densities.

Lemma 3.5. *Let Ω be an open bounded convex set and let $(\mu, \nu) \in \mathcal{CE}_\Omega(\mu_0 \rightarrow \mu_1)$ with given constant mass \mathbf{m} and finite energy $\int_0^1 \Phi_{\mathbf{m},\Omega}(\mu_s, \nu_s) ds < +\infty$. For every $\varepsilon > 0, \delta \in [0, 1]$ there exist a decreasing family of smooth convex sets $\Omega^\varepsilon \downarrow \Omega$ and a family of curves $(\mu^{\varepsilon,\delta}, \nu^{\varepsilon,\delta}) \in \mathcal{CE}_{\Omega^\varepsilon}(\mu_0^{\varepsilon,\delta} \rightarrow \mu_1^{\varepsilon,\delta})$ with the following properties*

$$\mu_i^{\varepsilon,\delta} = (1 - \delta)\mu_i * k_\varepsilon + \delta\lambda^\varepsilon, \quad \lambda^\varepsilon := \frac{\mathbf{m}}{\mathcal{L}^d(\Omega^\varepsilon)} \mathcal{L}^d|_{\Omega^\varepsilon}, \quad \mu_s^{\varepsilon,\delta}(\Omega^\varepsilon) = \mathbf{m}, \quad (3.7)$$

$$\mu_s^{\varepsilon,\delta} = \rho_s^{\varepsilon,\delta} \mathcal{L}^d|_{\Omega^\varepsilon}, \quad \nu_s^{\varepsilon,\delta} = \mathbf{w}_s^{\varepsilon,\delta} \mathcal{L}^d|_{\Omega^\varepsilon}, \quad \rho^{\varepsilon,\delta}, \mathbf{w}^{\varepsilon,\delta} \in C^\infty([0, 1] \times \overline{\Omega}^\varepsilon), \quad (3.8)$$

$$\partial_s \rho_s^{\varepsilon,\delta} + \nabla \cdot \mathbf{w}_s^{\varepsilon,\delta} = 0 \quad \text{in } (0, 1) \times \Omega^\varepsilon, \quad \rho^{\varepsilon,\delta} \geq \delta \frac{\mathbf{m}}{\mathcal{L}^d(\Omega)} > 0,$$

$$\frac{1}{c_\varepsilon^2} \int_0^1 \Phi_{\mathbf{m},\Omega^\varepsilon}(\mu_s^{\varepsilon,\delta}, \nu_s^{\varepsilon,\delta}) ds \leq \int_0^1 \Phi_{\mathbf{m},\Omega}(\mu_s, \nu_s) ds = \lim_{\varepsilon, \delta \downarrow 0} \int_0^1 \Phi_{\mathbf{m},\Omega^\varepsilon}(\mu_s^{\varepsilon,\delta}, \nu_s^{\varepsilon,\delta}) ds,$$

where $c_\varepsilon := 1 + 2\varepsilon$.

Proof. Let us extend (μ_s, ν_s) outside the unit interval by setting $\nu_s \equiv 0$ and $\mu_s \equiv \mu_0$ if $s < 0$, $\mu_s \equiv \mu_1$ if $s > 1$; it is immediate to check that (μ, ν) still satisfy the continuity equation. We then consider a family of smooth and convex open sets Ω^ε satisfying $\Omega + B_{2\varepsilon}(0) \subset \Omega^\varepsilon \subset \Omega + B_{3\varepsilon}(0)$ and define $\tilde{\mu}_s^\varepsilon := \mu * k_\varepsilon, \tilde{\nu}_s^\varepsilon := \nu * k_\varepsilon$ which have smooth densities $\tilde{\rho}_s^\varepsilon, \tilde{\mathbf{w}}_s^\varepsilon$ and are concentrated in $\overline{\Omega} + B_\varepsilon(0)$. We perform a further time convolution with respect to a 1-dimensional family of nonnegative smooth mollifiers $h_\varepsilon(z) := \varepsilon^{-1}h(z/\varepsilon)$ with support in $[-\varepsilon, \varepsilon]$ and integral 1

$$\bar{\mu}_s^\varepsilon := \int_{\mathbb{R}} \tilde{\mu}_z^\varepsilon h_\varepsilon(s - z) dz, \quad \bar{\nu}_s^\varepsilon := \int_{\mathbb{R}} \tilde{\nu}_z^\varepsilon h_\varepsilon(s - z) dz,$$

with corresponding densities $\rho_s^\varepsilon, \mathbf{w}_s^\varepsilon$. Notice that $\bar{\mu}_{-\varepsilon}^\varepsilon = \mu_0^{\varepsilon,0}, \bar{\mu}_{1+\varepsilon}^\varepsilon = \mu_1^{\varepsilon,0}$ and, by the convexity of $\phi_{\mathbf{m}}$ and Jensen's inequality, we have

$$\phi_{\mathbf{m}}(\rho_s^\varepsilon, \mathbf{w}_s^\varepsilon) \leq \int_{\mathbb{R}} \phi_{\mathbf{m}}(\tilde{\rho}_z^\varepsilon, \tilde{\mathbf{w}}_z^\varepsilon) h_\varepsilon(s - z) dz, \quad \Phi_{\mathbf{m},\Omega^\varepsilon}(\bar{\mu}_s^\varepsilon, \bar{\nu}_s^\varepsilon) \leq \int_{\mathbb{R}} \Phi_{\mathbf{m},\Omega^\varepsilon}(\tilde{\mu}_z^\varepsilon, \tilde{\nu}_z^\varepsilon) h_\varepsilon(s - z) dz$$

so that, being $\bar{\nu}_s^\varepsilon = 0$ if $s < -\varepsilon$ or $s > 1 + \varepsilon$,

$$\int_{-\varepsilon}^{1+\varepsilon} \Phi_{\mathbf{m},\Omega^\varepsilon}(\bar{\mu}_s^\varepsilon, \bar{\nu}_s^\varepsilon) ds = \int_{\mathbb{R}} \Phi_{\mathbf{m},\Omega^\varepsilon}(\bar{\mu}_s^\varepsilon, \bar{\nu}_s^\varepsilon) ds \leq \int_{\mathbb{R}} \Phi_{\mathbf{m},\Omega^\varepsilon}(\tilde{\mu}_s^\varepsilon, \tilde{\nu}_s^\varepsilon) ds \leq \int_0^1 \Phi_{\mathbf{m},\Omega}(\mu_s, \nu_s) ds.$$

We eventually set

$$\mu_s^\varepsilon := \bar{\mu}_{c_\varepsilon s - \varepsilon}^\varepsilon, \quad \nu_s^\varepsilon := c_\varepsilon \bar{\nu}_{c_\varepsilon s - \varepsilon}^\varepsilon, \quad c_\varepsilon := 1 + 2\varepsilon$$

and

$$\mu_s^{\varepsilon, \delta} := (1 - \delta)\mu_s^\varepsilon + \delta\lambda^\varepsilon, \quad \nu_s^{\varepsilon, \delta} := \nu_s^\varepsilon.$$

It is then easy to check that all the requirements are satisfied. \square

3.4 Couple of measures at finite $\mathcal{W}_{\mathbf{m}, \Omega}$ distance

We discuss now some cases when it is possible to prove that the distance between two measures is finite. We already know [DNS09, Cor. 5.25] (in the case A, but the same argument can be easily adapted to cover the case $M < +\infty$) that when Ω is convex and bounded

$$\text{if } \mu_i = \rho_i \mathcal{L}^d \text{ with } \|\rho_i\|_{L^\infty(\mathbb{R}^d)} < M \text{ then } \mathcal{W}_{\mathbf{m}, \Omega}(\mu_0, \mu_1) < \infty. \quad (3.9)$$

We focus on the case A, $M = +\infty$, and exploit some ideas of [Sa08]. In order to refine the condition (3.9), we first introduce the functions

$$k_{\mathbf{m}, d}(r) := \left(r^{1+1/d} \mathbf{m}(r)\right)^{-1/2}, \quad K_{\mathbf{m}, d}(r) := \frac{1}{d} \int_r^{+\infty} k_{\mathbf{m}, d}(z) dz, \quad r > 0.$$

Observe that $K_{\mathbf{m}, d}$ is either everywhere finite or identically $+\infty$. In particular, in the case $\mathbf{m}(r) = r^\alpha$, $K_{\mathbf{m}, d}$ is finite if and only if $\alpha > 1 - 1/d$.

Theorem 3.6. *Let Ω be a bounded, open convex set of \mathbb{R}^d . Suppose that $M = +\infty$, $\mathbf{m} > 0$, and that $K_{\mathbf{m}, d}$ is finite. Then any two measures $\mu_0, \mu_1 \in \mathcal{M}^+(\bar{\Omega}, \mathbf{m})$ have finite distance $\mathcal{W}_{\mathbf{m}, \Omega}(\mu_0, \mu_1) < +\infty$ and the topology induced by $\mathcal{W}_{\mathbf{m}, \Omega}$ on the space $\mathcal{M}^+(\bar{\Omega}, \mathbf{m})$ coincides with the usual weak* topology. In particular, the metric space $(\mathcal{M}^+(\bar{\Omega}, \mathbf{m}), \mathcal{W}_{\mathbf{m}, \Omega})$ is compact and separable.*

Proof. We fix an open set B with compact closure in Ω and a reference measure $\lambda = \bar{\rho} \mathcal{L}^d|_B$ with $\lambda(\Omega) = \mathbf{m}$ and $0 < \bar{\rho}(x) \leq b$ for \mathcal{L}^d -a.e. x in B . Since $\mathcal{W}_{\mathbf{m}, \Omega}$ satisfies the triangular inequality, the first part of the theorem follows if we show that $\mathcal{W}_{\mathbf{m}, \Omega}(\lambda, \mu) < +\infty$ for every $\mu \in \mathcal{M}^+(\bar{\Omega}, \mathbf{m})$.

Let $\mathbf{r} : B \rightarrow \bar{\Omega}$ be the Brenier map pushing λ onto μ : we know that \mathbf{r} is cyclically monotone. We set $\mathbf{r}_s := (1 - s)\mathbf{i} + s\mathbf{r}$ with image $B_s \subset (1 - s)\bar{B} + s\bar{\Omega} \subset \bar{\Omega}$ and inverse $\mathbf{s}_s = \mathbf{r}_s^{-1} : B_s \rightarrow B$, and $\mathbf{v}_s := (\mathbf{r} - \mathbf{i}) \circ \mathbf{r}_s^{-1} = \mathbf{i} - \mathbf{s}_s$. It is well known that \mathbf{s}_s is a Lipschitz map with Lipschitz constant bounded by $(1 - s)^{-1}$ and that the curve $\mu_s := (\mathbf{r}_s)_\# \lambda$ belongs to $\mathcal{CE}_\Omega(\lambda \rightarrow \mu)$ with

$$\mu_s = \bar{\rho}_s \mathcal{L}^d|_{B_s}, \quad \bar{\rho}_s = \chi_{B_s} \bar{\rho}(\mathbf{s}_s) J_s, \quad J_s := \det D\mathbf{s}_s, \quad \mathbf{v}_s = \bar{\rho}_s \mathbf{v}_s \mathcal{L}^d.$$

Since the map $r \mapsto r/\mathbf{m}(r)$ is nondecreasing and $J_s \leq (1 - s)^{-d}$, it follows that

$$\Phi_{\mathbf{m}, \Omega}(\mu_s, \nu_s) = \int_{B_s} \frac{\bar{\rho}_s^2}{\mathbf{m}(\bar{\rho}_s)} |\mathbf{v}_s|^2 dx = \int_B \frac{\bar{\rho}(y) J_s(\mathbf{r}_s(y))}{\mathbf{m}(\bar{\rho}(y) J_s(\mathbf{r}_s(y)))} |\mathbf{r}(y) - y|^2 \bar{\rho}(y) dy \quad (3.10)$$

$$\leq \frac{b(1 - s)^{-d}}{\mathbf{m}(b(1 - s)^{-d})} \int_B |\mathbf{r}(y) - y|^2 \bar{\rho}(y) dy = \frac{b(1 - s)^{-d}}{\mathbf{m}(b(1 - s)^{-d})} W_2^2(\lambda, \mu). \quad (3.11)$$

Taking the square and applying (3.3), since

$$\int_0^1 \left(\frac{b(1-s)^{-d}}{\mathfrak{m}(b(1-s)^{-d})} \right)^{1/2} ds = \frac{b^{1/d}}{d} \int_b^{+\infty} \left(\frac{z}{\mathfrak{m}(z)} \right)^{1/2} z^{-1-1/d} dz = b^{1/d} K_{\mathfrak{m},d}(b)$$

we get the estimate

$$\mathcal{W}_{\mathfrak{m},\Omega}(\lambda, \mu) \leq b^{1/d} K_{\mathfrak{m},d}(b) W_2(\lambda, \mu). \quad (3.12)$$

A completely analogous calculation with $\mu := \mu_0$ (resp. $\mu := \mu_1$) and $\mu_s = \mu_{0,s}$ (resp. $\mu_s = \mu_{1,s}$) shows that

$$\mathcal{W}_{\mathfrak{m},\Omega}(\mu_{i,1-\varepsilon}, \mu_i) \leq b^{1/d} K_{\mathfrak{m},d}(b\varepsilon^{-d}) W_2(\lambda, \mu_i) \quad \forall \varepsilon > 0, \quad i = 0, 1.$$

On the other hand, taking into account that the density of $\mu_{i,1-\varepsilon}$ is bounded by $b\varepsilon^{-d}$, we can apply (3.12) with $\mu_{0,1-\varepsilon}$ instead of λ , obtaining

$$\mathcal{W}_{\mathfrak{m},\Omega}(\mu_{0,1-\varepsilon}, \mu_{1,1-\varepsilon}) \leq b^{1/d} \varepsilon^{-1} K_{\mathfrak{m},d}(b\varepsilon^{-d}) W_2(\mu_{0,1-\varepsilon}, \mu_{1,1-\varepsilon}).$$

Therefore, the triangular inequality yields

$$\mathcal{W}_{\mathfrak{m},\Omega}(\mu_0, \mu_1) \leq b^{1/d} K_{\mathfrak{m},d}(b\varepsilon^{-d}) \left(W_2(\mu_0, \lambda) + W_2(\mu_1, \lambda) + \varepsilon^{-1} W_2(\mu_{0,1-\varepsilon}, \mu_{1,1-\varepsilon}) \right).$$

Applying this estimate to a sequence μ_n weakly* converging to μ (and therefore converging also with respect to W_2), since the corresponding geodesic interpolants with λ $\mu_{n,1-\varepsilon}$ converge to $\mu_{1-\varepsilon}$ as $n \rightarrow \infty$ with respect to W_2 , we easily obtain

$$\limsup_{n \rightarrow \infty} \mathcal{W}_{\mathfrak{m},\Omega}(\mu_n, \mu) \leq 2b^{1/d} K_{\mathfrak{m},d}(b\varepsilon^{-d}) W_2(\mu, \lambda).$$

Since $\lim_{\varepsilon \downarrow 0} K_{\mathfrak{m},d}(b\varepsilon^{-d}) = 0$, taking ε arbitrarily small, we conclude. \square

In the next result we do not assume any particular condition on \mathfrak{m} , but we ask that $\mu_i \ll \mathcal{L}^d$ with densities satisfying some extra integrability assumptions.

Theorem 3.7. *Let Ω be a bounded, open convex set of \mathbb{R}^d and assume that $M = +\infty$, $\mathfrak{m} > 0$. If the measures $\mu_i = \rho_i \mathcal{L}^d|_{\Omega} \in \mathcal{M}^+(\Omega, \mathfrak{m})$, $i = 0, 1$, satisfy*

$$\int_{\Omega} \frac{\rho_i(x)^2}{\mathfrak{m}(\rho_i(x))} dx < +\infty \quad i = 0, 1, \quad (3.13)$$

then $\mathcal{W}_{\mathfrak{m},\Omega}(\mu_0, \mu_1) < +\infty$.

Proof. We argue as in the previous proof, keeping the same notation and observing that for $0 \leq s \leq 1/2$ (3.11) yields

$$\Phi_{\mathfrak{m},\Omega}(\mu_s, \nu_s) \leq \frac{b 2^d}{\mathfrak{m}(b 2^d)} W_2^2(\lambda, \mu). \quad (3.14)$$

When $1/2 \leq s \leq 1$ we invert the role of λ and $\mu = \rho \mathcal{L}^d$ in (3.10) obtaining

$$\Phi_{\mathfrak{m},\Omega}(\mu_s, \nu_s) = \int_{\Omega} \frac{\rho(y) \tilde{J}_s(\tilde{\mathbf{s}}_s(y))}{\mathfrak{m}(\rho(y) \tilde{J}_s(\tilde{\mathbf{s}}_s(y)))} |\tilde{\mathbf{s}}(y) - y|^2 \rho(y) dy \quad (3.15)$$

where $\tilde{s}_s = (1-s)s + si$ is the optimal map pushing μ onto μ_s and $\tilde{J}_s = \det D\tilde{s}_s^{-1}$ satisfies $\tilde{J}_s \leq s^{-d}$. (3.15) then yields for $1/2 \leq s \leq 1$

$$\Phi_{\mathbf{m},\Omega}(\mu_s, \nu_s) \leq 2^{d+1} \int_{\Omega} \frac{\rho(y)^2}{\mathbf{m}(\rho(y))} (|\tilde{s}(y)|^2 + |y|^2) dy. \quad (3.16)$$

Since the range of $\tilde{s}(y)$ is μ -essentially bounded, the integral in (3.16) is finite thanks to (3.13). Integrating (3.14) in $(0, 1/2)$ and (3.16) in $(1/2, 1)$ we conclude that $\mathcal{W}_{\mathbf{m},\Omega}(\lambda, \mu)$ is finite. \square

4 Geodesic convexity of integral functionals and their gradient flows

4.1 Nonlinear diffusion equations: weak and limit solutions

We consider a

$$\text{convex density function } U \in W_{\text{loc}}^{2,1}(0, M) \text{ with } \mathbf{m}U'' \in L_{\text{loc}}^1([0, M)) \quad (4.1a)$$

and a *pressure function* $P : [0, M) \rightarrow \mathbb{R}$ defined by

$$P(r) := \int_0^r \mathbf{m}(z)U''(z) dz.$$

Let us observe that $P \in W_{\text{loc}}^{1,1}([0, M))$ is nondecreasing, continuous, and $P(0) = 0$. When U has a superlinear growth at $+\infty$ the corresponding *internal energy functional* $\mathcal{U} : D(\mathcal{U}) \subset \mathcal{M}_c^+(\mathbb{R}^d) \rightarrow (-\infty, +\infty]$ is defined as

$$\mathcal{U}(\mu) := \int_{\mathbb{R}^d} U(\rho(x)) dx, \quad D(\mathcal{U}) := \left\{ \mu = \rho \mathcal{L}^d \in \mathcal{M}_c^+(\mathbb{R}^d) : U(\rho) \in L^1(\mathbb{R}^d) \right\} \quad (4.2)$$

Since U is bounded from below by a linear function and μ has compact support, the integral in (4.2) is always well defined. \mathcal{U} is lower semicontinuous with respect to weak convergence in $\mathcal{M}_c^+(\mathbb{R}^d)$ if and only if

$$U'(+\infty) := \lim_{r \uparrow +\infty} \frac{U(r)}{r} = \lim_{r \uparrow +\infty} U'(r) = +\infty.$$

When $U'(+\infty) < +\infty$ we define the functional \mathcal{U} as

$$\mathcal{U}(\mu) := \int_{\mathbb{R}^d} U(\rho) dx + U'(+\infty)\mu^\perp(\mathbb{R}^d), \quad \mu = \rho \mathcal{L}^d + \mu^\perp,$$

where μ^\perp is the singular part of μ in the usual Lebesgue decomposition.

Let $\Omega \subset \mathbb{R}^d$ be a bounded, open, and connected set with Lipschitz boundary $\partial\Omega$ and exterior unit normal \mathbf{n} . We will often suppose that Ω is convex in the sequel. We consider the homogeneous Neumann boundary value problem for the nonlinear diffusion equation

$$\partial_t \rho - \Delta P(\rho) = 0 \quad \text{in } (0, +\infty) \times \Omega, \quad \partial_{\mathbf{n}} P(\rho) = 0 \quad \text{on } (0, +\infty) \times \partial\Omega, \quad (4.3)$$

with *nonnegative* initial condition $\rho(0, \cdot) = \rho_0$. We also introduce the dissipation rate of \mathcal{U} along the flow by

$$\mathcal{D}(\rho) = \int_{\Omega} \frac{|\nabla P(\rho)|^2}{\mathfrak{m}(\rho)} dx = \int_{\Omega} \phi_{\mathfrak{m}}(\rho, \nabla P(\rho)) dx, \quad \forall 0 \leq \rho \in L^1(\Omega), P(\rho) \in W^{1,1}(\Omega). \quad (4.4)$$

We collect in the following result some well established facts [Vaz07] on weak and classical solutions to (4.3).

Theorem 4.1 (Very weak and classical solutions). *Let us suppose that Ω is bounded and $\rho_0 \in L^\infty(\Omega)$. There exists a unique solution $\rho \in L^\infty((0, +\infty) \times \Omega) \cap C^0([0, +\infty); L^1(\Omega))$ with $P(\rho) \in L^\infty((0, +\infty) \times \Omega) \cap L^2((0, +\infty); W^{1,2}(\Omega))$ to (4.3) satisfying the following weak formulation*

$$\int_0^{+\infty} \int_{\Omega} \left(\rho \partial_t \zeta - \nabla P(\rho) \cdot \nabla \zeta \right) dx dt = 0 \quad \forall \zeta \in C_c^\infty((0, +\infty) \times \mathbb{R}^d), \quad (4.5)$$

and the initial condition $\rho(0, \cdot) = \rho_0$. The energy \mathcal{U} is decreasing along the flow and satisfies the identity

$$\int_{\Omega} U(\rho(T, x)) dx + \int_0^T \int_{\Omega} \frac{|\nabla P(\rho)|^2}{\mathfrak{m}(\rho)} dx dt = \int_{\Omega} U(\rho_0(x)) dx, \quad \forall T > 0. \quad (4.6)$$

The map $\rho_0 \mapsto S_t \rho_0 := \rho(t, \cdot)$ can be extended to a C^0 contraction semigroup $S = S(P, \Omega)$ in the positive cone of $L^1(\Omega)$, whose curves $S_t \rho_0$ are also called “limit L^1 -solutions” of (4.3), and it satisfies

$$\text{ess inf}_{\Omega} \rho_0 \leq S_t \rho_0 \leq \text{ess sup}_{\Omega} \rho_0.$$

If moreover $U, \mathfrak{m} \in C^\infty(0, M)$, U is uniformly convex, Ω is smooth and $\inf_{\Omega} \rho_0 > 0$, then $\rho \in C^\infty((0, +\infty) \times \overline{\Omega})$ and is a classical solution to (4.3).

Let us briefly discuss here two useful lemma, whose proof follows from a standard variational argument.

Lemma 4.2. *If $\rho_0, U(\rho_0) \in L^1(\Omega)$ then the limit L^1 -solution $\rho = S(\rho_0)$ satisfies $P(\rho) \in L_{\text{loc}}^1([0, +\infty); W^{1,1}(\Omega))$, the weak formulation (4.5), and the energy inequality*

$$\int_{\Omega} U(\rho(T, x)) dx + \int_0^T \int_{\Omega} \frac{|\nabla P(\rho)|^2}{\mathfrak{m}(\rho)} dx dt \leq \int_{\Omega} U(\rho_0(x)) dx. \quad (4.7)$$

Proof. Let us first show that we can find a constant C depending only on $P, \omega := \mathcal{L}^d(\Omega)$, $\mathfrak{m} = \int_{\Omega} \rho dx$, and the constant c_p in the Poincaré inequality for Ω such that

$$\|P(\rho)\|_{L^1(\Omega)} \leq C \left(1 + \|\nabla P(\rho)\|_{L^1(\Omega)} \right) \quad \forall \rho \in L^1(\Omega), \int_{\Omega} \rho dx = \mathfrak{m}, P(\rho) \in W^{1,1}(\Omega). \quad (4.8)$$

In fact, setting $\mathfrak{p} := \int_{\Omega} P(\rho) dx$ and $\ell := \mathcal{L}^d(\{x \in \Omega : P(\rho) \geq \mathfrak{p}/2\})$ Poincaré and Chebyshev inequality yield

$$\frac{1}{2} \mathfrak{p}(\omega - \ell) \leq \int_{\Omega} |P(\rho) - \mathfrak{p}| dx \leq c_p \int_{\Omega} |\nabla P(\rho)| dx, \quad \frac{1}{2} \mathfrak{p} \leq P(\mathfrak{m}/\ell),$$

so that if $\ell \geq \omega/2$ we get $\mathbf{p} \leq 2P(2\mathbf{m}/\omega)$, whereas if $\ell \leq \omega/2$ we obtain $\mathbf{p} \leq 4\omega^{-1}c_p \int_{\Omega} |\nabla P(\rho)| \, dx$.

If now $\rho_t = S_t \rho_0$ is the $L^1(\Omega)$ -limit of a sequence $\rho_{n,t} = S_t \rho_{n,0}$ of bounded solutions with $U(\rho_{n,0}) \rightarrow U(\rho_0)$ in $L^1(\Omega)$ as $n \uparrow +\infty$, from the uniform bound (4.6) we obtain for every bounded Borel set $\mathcal{T} \subset (0, +\infty)$, every $B \subset \Omega$, and every nonnegative constants a, b such that $\mathbf{m}(r) \leq a + br$,

$$\begin{aligned} \int_{\mathcal{T}} \int_B |\nabla P(\rho_n)| \, dx \, dt &\leq \|\mathbf{m}(\rho_n)\|_{L^1(\mathcal{T} \times B)}^{1/2} \left(\int_{\mathcal{T} \times B} \frac{|\nabla P(\rho_n)|^2}{\mathbf{m}(\rho_n)} \, dx \, dt \right)^{1/2} \\ &\leq C \|a + b\rho_n\|_{L^1(\mathcal{T} \times B)}^{1/2}. \end{aligned}$$

Taking $\mathcal{T} = (0, T)$, $B = \Omega$ and applying (4.8), we obtain a uniform bound of the sequence $P(\rho_n)$ in $L^1(0, T; W^{1,1}(\Omega))$; since ρ_n converges to ρ in $L^1((0, T) \times \Omega)$, we obtain that $\nabla P(\rho_n)$ is uniformly integrable and therefore it converges weakly to $\nabla P(\rho)$ in $L^1((0, T) \times \Omega)$. It follows that $P(\rho) \in L^1(0, T; W^{1,1}(\Omega))$ and we can then pass to the limit in the weak formulation (4.5) written for ρ_n , obtaining the same identity for ρ . The inequality (4.7) eventually follows by the same limit procedure, recalling that the dissipation functional (4.4) is lower semicontinuous with respect to weak convergence in $L^1(\Omega)$. \square

The following stability result is used in the sequel; its proof is an easy adaption of [Vaz07, Prop. 6.10].

Proposition 4.3. *Let $\Omega^n \subset \mathbb{R}^d$ be a decreasing sequence of open, bounded, convex sets converging to Ω and let $S^n = S(P, \Omega^n)$, $S(P, \Omega)$ be the associated semigroups provided by Theorem 4.1. If (after a trivial extension to 0 outside Ω^n) $\rho_0^n \in L^1(\Omega^n)$ is converging strongly in $L^1(\mathbb{R}^d)$ to $\rho_0 \in L^1(\Omega)$, then $S_t^n(\rho_0^n) \rightarrow S_t(\rho_0)$ in the same L^1 sense, as $n \uparrow +\infty$ for every $t > 0$.*

4.2 The generalized McCann condition

We assume that $P'\mathbf{m}' \in L^1_{\text{loc}}([0, M])$ and we introduce a primitive function H of $h := P'\mathbf{m}' = U''\mathbf{m}\mathbf{m}'$,

$$H(r) := H_0 + \int_0^r P'(z)\mathbf{m}'(z) \, dz \quad \text{for some } H_0 \geq 0. \quad (4.9)$$

When the dimension d is greater than 1, we assume that $\inf_{r \in (0, M)} H(r) = 0$; this means that the locally integrability assumption on h cannot be avoided, as well as, in the case when $M < +\infty$, its integrability in $(0, M)$. In the case $M = +\infty$ we can simply choose $H_0 = 0$ in (4.9). In the case $M < +\infty$ we can choose

$$H_0 := \left(\int_0^M P'\mathbf{m}' \, dx \right)^- < +\infty. \quad (4.10)$$

Remark 4.4. Notice that in the most common case when $\mathbf{m}'(0_+) = \lim_{r \downarrow 0} r^{-1}\mathbf{m}(r) > 0$, the local integrability of h in a right neighborhood of 0 implies the local integrability of $\mathbf{m}U''$, as we already required in (4.1a), and the lower boundedness of P . When the space dimension is 1 all these restriction can be removed: we comment on this issue in the next remark 4.16. If $M < +\infty$, P is locally Lipschitz near 0 and M , and $\mathbf{m}(0) = \mathbf{m}(M) = 0$, then we get $H_0 = 0$ so that P and \mathbf{m} should satisfy the compatibility condition $\int_0^M P'\mathbf{m}' \, dr = 0$.

Definition 4.5. Let U, P, H and \mathbf{m} be defined in the interval $(0, M)$ according to (4.1a,b) and (4.9). We say that the energy density U and the corresponding pressure function P satisfy the d -dimensional generalized McCann condition for the mobility \mathbf{m} , denoted by $GMC(\mathbf{m}, d)$, if for a suitable choice of H_0

$$U''(r)\mathbf{m}^2(r) = P'(r)\mathbf{m}(r) \geq (1 - 1/d)H(r) \geq 0, \quad \forall r \in (0, M), \quad (4.11a)$$

or, equivalently,

$$r \mapsto \frac{H(r)}{\mathbf{m}^{1-1/d}(r)} = \frac{1}{\mathbf{m}^{1-1/d}(r)} \int_0^r P'(s)\mathbf{m}'(s) \, ds \quad \text{is nondecreasing in } (0, +\infty).$$

We collect in the following remarks some simple properties related to this definition.

Remark 4.6 (Elementary properties).

1. (Linear mobility) (4.11a) is consistent with the usual McCann condition (1.2a) in the linear case of $\mathbf{m}(r) = r$.
2. (Dimension $d = 1$) As in the case of McCann condition, in space dimension $d = 1$ (4.11a) is equivalent to the convexity of U or to the monotonicity of P .
3. (Local boundedness of U when $d > 1$). In dimension $d > 1$ the energy density function U is bounded in a right neighborhood of 0 (and in a left neighborhood of M , in the case $M < +\infty$). Since $U'' = P'/\mathbf{m}$ the property is immediate if $\mathbf{m}(0) > 0$. If $\mathbf{m}(0) = 0$ then $\mathbf{m}'(0) > 0$ and therefore P is bounded around 0 and the formula

$$U(r) = U(r_0) + U'(r_0)(r - r_0) + \int_0^{r_0} \frac{(z - r)^+}{\mathbf{m}(z)} P'(z) \, dz, \quad r \in (0, r_0],$$

shows that $\lim_{r \downarrow 0} U(r) < +\infty$.

4. (Constant mobility) When $\mathbf{m}(r) \equiv c > 0$ (4.11a) is still equivalent to the convexity of U .
5. (The power-like case) In the case of $P(r) = r^\gamma$ ($\gamma = \alpha + \beta - 1$ if $U(r) = r^\beta$) and $\mathbf{m}(r) = r^\alpha$, (4.11a) is satisfied if and only if

$$\gamma \geq 1 - \frac{\alpha}{d}. \quad (4.12)$$

6. (The case $P(r) = r$) It is immediate to check that the couple (r, \mathbf{m}) always satisfies (4.11a): it corresponds to the entropy function $U_{\mathbf{m}}$ whose second derivative is \mathbf{m}^{-1} . After fixing some $r_0 \in (0, M)$ (the choice $r_0 = 0$ is admissible if \mathbf{m}^{-1} is integrable in a right neighborhood of 0), we obtain

$$U_{\mathbf{m}}(r) := \int_{r_0}^r \frac{r - z}{\mathbf{m}(z)} \, dz, \quad P_{\mathbf{m}}(r) = r - r_0.$$

7. (The case of the logarithmic entropy) $U(r) = r \log r$ satisfies $GMC(r^\alpha, d)$ if and only if $\gamma = \alpha \geq d/(d + 1)$.

8. (Linearity) If P_1 and P_2 satisfy $GMC(\mathbf{m}, d)$ then also $\alpha_1 P_1 + \alpha_2 P_2$ satisfies $GMC(\mathbf{m}, d)$, for every $\alpha_1, \alpha_2 \geq 0$. Analogously, if P satisfies $GMC(\mathbf{m}_1, d)$ and $GMC(\mathbf{m}_2, d)$ then P satisfies $GMC(\alpha_1 \mathbf{m}_1 + \alpha_2 \mathbf{m}_2, d)$. In particular, if P satisfies $GMC(\mathbf{m}, d)$ then $P(r) + \alpha r$ satisfies $GMC(\mathbf{m} + \beta, d)$ for every $\alpha, \beta \geq 0$.
9. (Shift) If $M = +\infty$ and P satisfies $GMC(\mathbf{m}, d)$ then P satisfies $GMC(\mathbf{m}(\cdot + \alpha), d)$ and $P(\cdot - \alpha)$ satisfies $GMC(\mathbf{m}, d)$, for every $\alpha \geq 0$.

The next two properties are more technical and require a detailed proof.

Lemma 4.7 (Smoothing). *Let us assume that P satisfies $GMC(\mathbf{m}, d)$ and let us fix two constants $0 < M' < M'' < M$. Then there exists a family P_η, \mathbf{m}_η , $\eta > 0$, with smooth restriction to $[M', M'']$ such that $P_\eta \geq P$ is strictly increasing, $\mathbf{m}_\eta \geq \mathbf{m}$ is concave, P_η satisfies $GMC(\mathbf{m}_\eta, d)$ (in $[0, M'']$), and P_η, \mathbf{m}_η converge uniformly to P, \mathbf{m} in $[M', M'']$ as $\eta \downarrow 0$. Moreover, if P' is locally integrable in a right neighborhood of 0, then we can choose $M' = 0$.*

Proof. When $M' > 0$ it is not restrictive (up to choosing a smaller M') to assume that M' is a Lebesgue point of the derivative of P . Let H be as in (4.9) and let us set $\tilde{\mathbf{m}}_\eta(r) := \mathbf{m}(r) + \eta$, $\tilde{P}_\eta(r) = P(r) + \eta r$,

$$\tilde{H}_\eta(r) = H_0 + \eta \mathbf{m}(0) + \eta^2 + \int_0^r \tilde{P}'_\eta(s) \tilde{\mathbf{m}}'_\eta(s) ds = H(r) + \eta \mathbf{m}(r) + \eta^2 \geq \eta^2 > 0.$$

By the previous Remark (points 6 and 8) \tilde{P}_η satisfies $GMC(\tilde{\mathbf{m}}_\eta, d)$ and moreover

$$\tilde{P}'_\eta \tilde{\mathbf{m}}_\eta - (1 - 1/d) \tilde{H}_\eta = P' \mathbf{m} - (1 - 1/d) H + \frac{\eta}{d} (\eta + \mathbf{m}) \geq \frac{\eta}{d} (\mathbf{m} + \eta) \geq \eta^2/d. \quad (4.13)$$

By choosing a family of mollifiers h_δ , $\delta > 0$, with support in $[0, \delta]$, we introduce the functions

$$\begin{aligned} \tilde{P}_{\eta, \delta}(r) &:= \begin{cases} \tilde{P}_\eta(M') + \int_{M'}^r \tilde{P}'_\eta * h_\delta ds & \text{if } r \geq M', \\ \tilde{P}_\eta(r) & \text{if } r < M', \end{cases} \\ \tilde{\mathbf{m}}_{\eta, \delta}(r) &:= \begin{cases} \tilde{\mathbf{m}}_\eta(M') + \int_{M'}^r \tilde{\mathbf{m}}'_\eta * h_\delta ds & \text{if } r \geq M', \\ \tilde{\mathbf{m}}_\eta(r) & \text{if } r < M', \end{cases} \end{aligned}$$

which are smooth in $[M', M'']$ and satisfy the requested monotonicity/concavity conditions. Since $\tilde{P}'_{\eta, \delta}$ converges to \tilde{P}'_η in $L^1_{\text{loc}}(0, M']$ and $\tilde{\mathbf{m}}_{\eta, \delta}$ is uniformly bounded and converges pointwise a.e. to $\tilde{\mathbf{m}}'_\eta$ as $\delta \rightarrow 0$, we conclude that the corresponding continuous functions $\tilde{H}_{\eta, \delta}$ converge uniformly to \tilde{H}_η as $\delta \downarrow 0$. By (4.13), we can find a sufficiently small $\delta = \delta_\eta$ depending on η such that

$$\tilde{P}'_{\eta, \delta_\eta} \tilde{\mathbf{m}}'_{\eta, \delta_\eta} \geq (1 - 1/d) \tilde{H}_{\eta, \delta_\eta} \geq 0.$$

A standard diagonal argument concludes the proof. \square

Lemma 4.8 (Minimal asymptotic behaviour). *When $d > 1$ and $M = +\infty$, the function $P_{\min}(r) := \int_0^r \mathbf{m}(z)^{-1/d} dz$ satisfies $GMC(\mathbf{m}, d)$ and provides an (asymptotic) lower bound for every any other P , since for every $r_0 > 0$ there exists a constant $c_0 > 0$ such that*

$$P'(r) \geq c_0 P'_{\min}(r) = c_0 \mathbf{m}(r)^{-1/d}, \quad U''(r) \geq c_0 \mathbf{m}(r)^{-1-1/d} \quad \text{for a.e. } r \geq r_0. \quad (4.14)$$

Proof. In fact $f(r) := P' \mathbf{m}$ satisfies

$$f(r) \geq (1 - 1/d) \left(H(r_0) + \int_{r_0}^r f(r) \mathbf{m}'(r) / \mathbf{m}(r) dr \right)$$

Gronwall Lemma then yields (4.14) with $c_0 := (1 - 1/d)H(r_0)\mathbf{m}(r_0)^{1/d-1}$. \square

Notice that in the case $\mathbf{m}(r) = r^\alpha$ we obtain the functions $P_{\min}(r) = cr^{\gamma_0}$ with exponent $\gamma_0 = 1 - \alpha/d$, which is consistent with (4.12). The corresponding energy density functions are then $U_{\min}(r) = cr^{2-\alpha(1+1/d)}$: in particular, when $\alpha < d/(d+1)$, all the energy functions have a superlinear growth as $r \uparrow \infty$.

Remark 4.9 (A sufficient condition). It is possible to give a simpler sufficient condition than (4.11a), at least when $\mathbf{m}U''$ is integrable in a right neighborhood of 0 and $M = +\infty$: if

$$\text{the map } r \mapsto \mathbf{m}^{1/d}(r)P'(r) = \mathbf{m}^{1+1/d}(r)U''(r) \text{ is positive and nondecreasing in } (0, +\infty) \quad (4.15)$$

then (4.11a) is satisfied. In fact, assuming U smooth for simplicity, (4.15) is equivalent to

$$0 \leq \mathbf{m}^{1/d}P'' - (1 - 1/d)\mathbf{m}^{1/d-1}\mathbf{m}'P'.$$

Multiplying the inequality by $\mathbf{m}^{1-1/d}$ and integrating in time we get (4.11a). Condition (4.15) gives the same sharp bound (4.12) in the power case.

4.3 The metric approach to gradient flows

We recall here some basic facts about the metric notion of gradient flows, referring to [AGS05] for further details. Let (D, \mathcal{W}) be a metric space, not assumed to be complete, and let $\mathcal{V} : D(\mathcal{V}) \rightarrow (-\infty, +\infty]$ be a lower semicontinuous functional. A family of continuous maps $S_t : D \rightarrow D$, $t \geq 0$, is a C^0 -(metric) contraction gradient flow of \mathcal{V} with respect to \mathcal{W} if

$$\begin{aligned} S_{t+h}(u) &= S_h(S_t(u)), \quad \lim_{t \downarrow 0} S_t(u) = S_0(u) = u \quad \forall u \in D, \quad t, h \geq 0, \\ \frac{1}{2}\mathcal{W}^2(S_t(u), v) - \frac{1}{2}\mathcal{W}^2(u, v) &\leq t \left(\mathcal{V}(v) - \mathcal{V}(S_t(u)) \right) \quad \forall t > 0, \quad u \in D, \quad v \in D(\mathcal{V}). \end{aligned} \quad (4.16a)$$

Thanks to [DS08, Prop. 3.1], conditions (4.16a,b) imply

$S_t(D) \subset D(\mathcal{V}) \forall t > 0$ and the map $t \mapsto \mathcal{V}(S_t(u))$ is not increasing in $(0, +\infty)$,

$$\frac{1}{2} \frac{d^+}{dt} \mathcal{W}^2(S_t(u), v) + \mathcal{V}(S_t(u)) \leq \mathcal{V}(v), \quad \forall u \in D, v \in D(\mathcal{V}), t \geq 0, \quad (4.17)$$

$$\mathcal{V}(S_t(u)) \leq \mathcal{V}(v) + \frac{1}{2t} \mathcal{W}^2(u, v) \quad \forall u \in D, v \in D(\mathcal{V}), t > 0,$$

$$\mathcal{W}^2(S_{t_1}(u), S_{t_0}(u)) \leq 2(t_1 - t_0) \left(\mathcal{V}(S_{t_0}u) - \mathcal{V}_{\inf} \right) \quad \forall u \in D(\mathcal{V}), 0 \leq t_0 \leq t_1, \quad (4.18)$$

$$\mathcal{W}(S_t(u), S_t(v)) \leq \mathcal{W}(u, v) \quad \forall u, v \in D, t \geq 0.$$

In (4.17) we used the usual notation

$$\frac{d^+}{dt} \zeta(t) = \limsup_{h \rightarrow 0^+} \frac{\zeta(t+h) - \zeta(t)}{h}.$$

for every real function $\zeta : [0, +\infty) \rightarrow \mathbb{R}$.

The following approximated convexity estimate [DS08, Theorem 3.2] plays an important role in the sequel.

Theorem 4.10 (Approximated convexity). *Let us suppose that S is metric contraction gradient flow of \mathcal{V} with respect to \mathcal{W} according to (4.16a,b) and let $s \mapsto u_s \in D$, $s \in [0, 1]$, be a Lipschitz (“almost” geodesic) curve such that $u_0, u_1 \in D(\mathcal{V})$ and*

$$\mathcal{W}(u_r, u_s) \leq L|r - s| \quad \forall r, s \in [0, 1], \quad L^2 \leq \mathcal{W}^2(u_0, u_1) + \delta^2. \quad (4.19)$$

Then for every $s \in [0, 1]$ and $t > 0$, we have

$$\mathcal{V}(S_t(u_s)) \leq (1-s)\mathcal{V}(u_0) + s\mathcal{V}(u_1) + \frac{s(1-s)}{2t} \delta^2.$$

In particular, if u_s is a minimal geodesic, i.e. (4.19) holds with $\delta = 0$, then

$$\mathcal{V}(u_s) \leq (1-s)\mathcal{V}(u_0) + s\mathcal{V}(u_1) \quad \forall s \in [0, 1].$$

4.4 Main results

We state our main result about the generation of a contractive gradient flow of \mathcal{U} with respect to $\mathcal{W}_{m,\Omega}$.

Theorem 4.11 (Contractive gradient flow). *Let us assume that Ω is a bounded, convex open set, and the functions U, P, H satisfy the generalized McCann condition $\text{GMC}(\mathbf{m}, d)$. For every reference measure $\sigma \in \mathcal{M}^+(\Omega)$ with finite energy $\mathcal{U}(\sigma) < +\infty$ the functional \mathcal{U} generates a unique metric contraction gradient flow $\mathcal{S} = \mathcal{S}(\mathcal{U}, \mathbf{m}, \Omega)$ in the space*

$$D := \left\{ \mu \in \mathcal{M}^+(\Omega) : \mu \ll \mathcal{L}^d|_{\Omega}, \mathcal{W}_{m,\Omega}(\mu, \sigma) < +\infty, \mathcal{U}(\mu) < +\infty \right\}$$

endowed with the distance $\mathcal{W}_{m,\Omega}$. Moreover \mathcal{S} is characterized by the formula $S_t\mu_0 = \rho_t \mathcal{L}^d|_{\Omega}$, where $\rho_t = S_t\rho_0$ is a limit L^1 -solution of (4.3).

When \mathbf{m} satisfies the finiteness condition of Theorem 3.6 (in particular $\mathbf{m}(r) = r^\alpha$ with $\alpha > 1 - 1/d$) we obtain a much more refined result, which in particular shows the continuous dependence of \mathcal{S} on the weak* topology.

Corollary 4.12. *Under the same assumptions on Ω, U, P of the previous theorem, if moreover $M = +\infty$ and \mathbf{m} satisfies the finiteness condition of theorem 3.6, then the semigroup \mathcal{S} can be uniquely extended to a contraction semigroup on every convex set $\mathcal{M}^+(\bar{\Omega}, \mathbf{m})$, which is continuous with respect to the weak* convergence of the initial data. If U has a superlinear growth, then $\mathcal{S}_t(\mu_0) = \rho_t \mathcal{L}^d \ll \mathcal{L}^d|_\Omega$ for every $t > 0$ and ρ_t is a weak solution of (4.3) according to (4.5).*

We conclude this section with our main convexity result.

Theorem 4.13 (Convexity). *Let us assume that Ω is a bounded convex open set, and the functions U, P, H satisfy the generalized McCann condition $GMC(\mathbf{m}, d)$. For every $\mu_0, \mu_1 \in \mathcal{M}^+(\Omega) \cap D(\mathcal{W})$ with finite distance $\mathcal{W}_{\mathbf{m}, \Omega}(\mu_0, \mu_1) < +\infty$ there exists a constant speed minimizing geodesic for $\mathcal{W}_{\mathbf{m}, \Omega}$, $\mu : [0, 1] \rightarrow \mathcal{M}^+(\Omega)$ connecting μ_0 to μ_1 such that*

$$\mathcal{W}(\mu_s) \leq s\mathcal{W}(\mu_1) + (1-s)\mathcal{W}(\mu_0), \quad \forall s \in [0, 1]. \quad (4.20)$$

Remark 4.14 (Weak and strong convexity). When (4.20) holds for all the (constant speed, minimizing) geodesics, the functional \mathcal{W} is called *strongly* geodesically convex. When \mathbf{m} is strictly concave and has a sublinear growth (or $M < +\infty$) then every two measures with finite $\mathcal{W}_{\mathbf{m}, \Omega}$ -distance can be connected by a *unique* geodesic [DNS09, Theorem 5.11], so that there is no difference between strong or weak convexity and (4.20) yields that the map $s \mapsto \mathcal{W}(\mu_s)$ is convex in $[0, 1]$.

Remark 4.15 (Absolutely continuous measures). Even when geodesics are not unique, the proof of Theorem 4.13 shows in fact that (4.20) is satisfied by *any* geodesic μ_s with $\mu_s \ll \mathcal{L}^d$ for every $s \in \mathbb{R}^d$, which surely exist if U has a superlinear growth. Along this class of geodesics we still obtain that the map $s \mapsto \mathcal{W}(\mu_s)$ is convex in $[0, 1]$.

Remark 4.16 (The one-dimensional case). When the space dimension $d = 1$, then the generalized McCann condition $GMC(\mathbf{m}, 1)$ reduces to the usual convexity of U . In this case, a simple approximation argument shows that we can cover also the case of functions U which are not bounded in a right neighborhood of 0 (and in a left neighborhood of M , if $M < +\infty$) and the integrability assumptions on $U''\mathbf{m}$ of (4.1a) and on $U''\mathbf{m}\mathbf{m}'$ of (4.10) can be dropped.

5 Action inequalities in the smooth case

In this section we assume that Ω is a *smooth and bounded open set*. We consider a smooth curve

$$\mu_s := \rho_s \mathcal{L}^d|_\Omega, \quad \rho \in C^\infty([0, 1] \times \bar{\Omega}), \quad 0 < m_0 \leq \rho \leq m_1 < M, \quad \mu_s(\Omega) \equiv \mathbf{m}, \quad s \in [0, 1]. \quad (5.1)$$

We also assume that P and \mathbf{m} are of class C^∞ in $[m_0, m_1]$. We consider the semigroup $S = S(P, \Omega)$ defined by Theorem 4.1 and we set

$$\mu_{s,t} := \rho_{s,t} \mathcal{L}^d|_\Omega, \quad \rho_{s,t}(\cdot) = \rho(s, t, \cdot) := S_{st}\rho_s, \quad s \in [0, 1], \quad t \geq 0. \quad (5.2)$$

Classical theory of quasilinear parabolic equation shows that $\rho \in C^\infty([0, 1] \times [0, \infty) \times \Omega) \cap C^\infty([0, 1] \times (0, +\infty) \times \bar{\Omega})$.

Since the semigroup S_t preserve the lower and upper bounds on ρ and $\int_\Omega \partial_s \rho \, dx = 0$, for every $(s, t) \in [0, 1] \times [0, +\infty)$ we can introduce the unique solution $\zeta_{s,t} = \zeta(s, t, \cdot) \in C^\infty(\bar{\Omega})$, of the uniformly elliptic Neumann boundary value problem

$$\begin{cases} -\nabla \cdot (\mathbf{m}(\rho) \nabla \zeta) = \partial_s \rho & \text{in } \Omega, \\ \nabla \zeta \cdot \mathbf{n} = 0 & \text{on } \partial\Omega, \\ \int_\Omega \zeta(x) \, dx = 0. \end{cases} \quad (5.3)$$

It is easy to check that ζ depends smoothly on s and t . Notice that (5.3) is equivalent to

$$\int_\Omega \mathbf{m}(\rho) \nabla \zeta \cdot \nabla \eta \, dx = \int_\Omega \partial_s \rho \eta \, dx \quad \forall \eta \in C^1(\bar{\Omega}). \quad (5.4)$$

By construction, for every $t \geq 0$ the curve $s \mapsto (\mu_{s,t}, \boldsymbol{\nu}_{s,t})$ with $\boldsymbol{\nu}_{s,t} := \mathbf{m}(\rho_{s,t}) \nabla \zeta_{s,t} \mathcal{L}^d|_\Omega$ belongs to $\mathcal{CE}(\mu_0 \rightarrow \mu_{1,t})$ and its energy can be evaluated by integrating the action

$$A_{s,t} := \Phi_{\mathbf{m},\Omega}(\mu_{s,t}, \boldsymbol{\nu}_{s,t}) = \int_\Omega \mathbf{m}(\rho_{s,t}) |\nabla \zeta_{s,t}|^2 \, dx$$

with respect to s in the interval $[0, 1]$. The integral provides an upper bound of the $\mathcal{W}_{\mathbf{m},\Omega}$ -distance between μ_0 and $\mu_{1,t} = \rho_{1,t} \mathcal{L}^d$, which corresponds to the solution of the nonlinear diffusion equation (4.3) with initial datum ρ_1 . As it was shown in [DS08], evaluating the time derivative of the action $A_{s,t}$ is a crucial step to prove that (4.3) satisfies the EVI formulation (4.16a). Next lemma, which does not require any convexity assumption on Ω , collects the main calculations.

Lemma 5.1. *Let $\rho_s, \rho_{s,t}$, and $\zeta_{s,t}$ be as in (5.1), (5.2), and (5.3). Then for every $(s, t) \in [0, 1] \times (0, +\infty)$ we have*

$$\begin{aligned} \frac{1}{2} \frac{\partial}{\partial t} A_{s,t} &= \frac{\partial}{\partial t} \frac{1}{2} \int_\Omega |\nabla \zeta_{s,t}|^2 \mathbf{m}(\rho_{s,t}) \, dx = - \int_\Omega \nabla P(\rho_{s,t}) \cdot \nabla \zeta_{s,t} \, dx \\ &\quad - s \int_\Omega \left((P'(\rho_{s,t}) \mathbf{m}(\rho_{s,t}) - H(\rho_{s,t})) (\Delta \zeta_{s,t})^2 + H(\rho_{s,t}) |D^2 \zeta_{s,t}|^2 \right) \, dx \\ &\quad + s \int_\Omega P'(\rho_{s,t}) \mathbf{m}''(\rho_{s,t}) |\nabla \rho_{s,t}|^2 |\nabla \zeta_{s,t}|^2 \, dx + \frac{1}{2} s \int_{\partial\Omega} H(\rho_{s,t}) \nabla |\nabla \zeta_{s,t}|^2 \cdot \mathbf{n} \, d\mathcal{H}^{d-1}, \end{aligned} \quad (5.5)$$

where H is defined in (4.9).

Proof. For keep the notation simple, we omit the explicit dependence of ρ, ζ on s, t . By the definition of ρ we easily get

$$\partial_t \rho = s \Delta P(\rho) \quad \text{and} \quad s \int_\Omega \nabla P(\rho) \cdot \nabla \eta \, dx = - \int_\Omega \partial_t \rho \eta \, dx \quad \forall \eta \in C^1(\bar{\Omega}). \quad (5.6)$$

Further differentiation with respect to s yields

$$- \int_{\Omega} \nabla P(\rho) \cdot \nabla \eta \, dx + s \int_{\Omega} \partial_s P(\rho) \Delta \eta \, dx = \int_{\Omega} \partial_{st} \rho \eta \, dx \quad (5.7)$$

for all $\eta \in C^2(\overline{\Omega})$ with $\nabla \eta \cdot \mathbf{n} = 0$ on $\partial\Omega$. On the other hand, differentiating (5.4) with respect to t we obtain

$$\int_{\Omega} \mathbf{m}(\rho) \partial_t \nabla \zeta \cdot \nabla \eta \, dx = \int_{\Omega} \partial_{st} \rho \eta \, dx - \int_{\Omega} \partial_t \mathbf{m}(\rho) \nabla \zeta \cdot \nabla \eta \, dx \quad \forall \eta \in C^1(\overline{\Omega}). \quad (5.8)$$

The time derivative of the action functional is

$$\begin{aligned} \frac{\partial}{\partial t} \frac{1}{2} \int_{\Omega} |\nabla \zeta|^2 \mathbf{m}(\rho) \, dx &= \frac{1}{2} \int_{\Omega} \partial_t \mathbf{m}(\rho) |\nabla \zeta|^2 \, dx + \int_{\Omega} \partial_t \nabla \zeta \cdot \nabla \zeta \mathbf{m}(\rho) \, dx \\ &\stackrel{(5.8)}{=} \int_{\Omega} \partial_{st} \rho \zeta \, dx - \frac{1}{2} \int_{\Omega} \partial_t \mathbf{m}(\rho) |\nabla \zeta|^2 \, dx \\ &\stackrel{(5.7)}{=} - \int_{\Omega} \nabla P(\rho) \cdot \nabla \zeta \, dx + s \int_{\Omega} \partial_s P(\rho) \Delta \zeta \, dx - \frac{1}{2} \int_{\Omega} \partial_t \mathbf{m}(\rho) |\nabla \zeta|^2 \, dx = I + II + III. \end{aligned} \quad (5.9)$$

We evaluate separately the various contributions: concerning the second integral II we introduce the auxiliary function G

$$G(r) := P'(r) \mathbf{m}(r) - H(r), \quad \text{so that} \quad G'(r) = P''(r) \mathbf{m}(r), \quad (5.10)$$

and we get

$$\begin{aligned} II &= s \int_{\Omega} \partial_s P(\rho) \Delta \zeta \, dx = s \int_{\Omega} P'(\rho) \partial_s \rho \Delta \zeta \, dx \\ &\stackrel{(5.4)}{=} s \int_{\Omega} P'(\rho) \mathbf{m}(\rho) \nabla \Delta \zeta \cdot \nabla \zeta \, dx + s \int_{\Omega} \Delta \zeta P''(\rho) \mathbf{m}(\rho) \nabla \rho \cdot \nabla \zeta \, dx \\ &\stackrel{(5.10)}{=} s \int_{\Omega} P'(\rho) \mathbf{m}(\rho) \nabla \Delta \zeta \cdot \nabla \zeta \, dx + s \int_{\Omega} \Delta \zeta \nabla G(\rho) \cdot \nabla \zeta \, dx \\ &= s \int_{\Omega} H(\rho) \nabla \Delta \zeta \cdot \nabla \zeta \, dx - s \int_{\Omega} G(\rho) (\Delta \zeta)^2 \, dx. \end{aligned}$$

The third integral of (5.9) is

$$\begin{aligned} III &= -\frac{1}{2} \int_{\Omega} \mathbf{m}'(\rho) \partial_t \rho |\nabla \zeta|^2 \, dx = \\ &\stackrel{(5.6)}{=} \frac{s}{2} \int_{\Omega} P'(\rho) \mathbf{m}''(\rho) |\nabla \rho|^2 |\nabla \zeta|^2 \, dx + \frac{s}{2} \int_{\Omega} P'(\rho) \mathbf{m}'(\rho) \nabla |\nabla \zeta|^2 \cdot \nabla \rho \, dx \\ &\stackrel{(4.9)}{=} \frac{s}{2} \int_{\Omega} P'(\rho) \mathbf{m}''(\rho) |\nabla \rho|^2 |\nabla \zeta|^2 \, dx + \frac{s}{2} \int_{\Omega} \nabla |\nabla \zeta|^2 \cdot \nabla H(\rho) \, dx. \end{aligned} \quad (5.11)$$

A further integration by parts in the last integral and the Bochner formula

$$\nabla \zeta \cdot \nabla \Delta \zeta - \frac{1}{2} \Delta |\nabla \zeta|^2 = -|D^2 \zeta|^2$$

yield (5.5). □

Corollary 5.2. *Under the same notation and assumptions of Lemma 5.1, if Ω is convex and U satisfies the generalized McCann condition $GMC(\mathbf{m}, d)$ (4.11a,b), then*

$$\frac{1}{2} \frac{\partial}{\partial t} A_{s,t} = \frac{\partial}{\partial t} \frac{1}{2} \int_{\Omega} |\nabla \zeta_{s,t}|^2 \mathbf{m}(\rho_{s,t}) \, dx \leq -\frac{\partial}{\partial s} \mathcal{W}(\mu_{s,t}) = -\frac{\partial}{\partial s} \int_{\Omega} U(\rho_{s,t}) \, dx, \quad (5.12)$$

and

$$\frac{1}{2} \mathcal{W}_{\mathbf{m}, \Omega}^2(\mu_{1,t}, \mu_0) + t \mathcal{W}(\mu_{1,t}) \leq \frac{1}{2} \int_0^1 A_{s,t} \, ds + \int_0^t \mathcal{W}(\mu_{1,\tau}) \, d\tau \leq \frac{1}{2} \int_0^1 A_{s,0} \, ds + t \mathcal{W}(\mu_0). \quad (5.13)$$

Proof. We determine the sign of the terms in the right hand side of (5.5) thanks to (4.11a,b) and the convexity of Ω . Recalling that $|D^2 \zeta| \geq \frac{1}{d} (\Delta \zeta)^2$ and $H \geq 0$ we obtain that the second integral in the right-hand side of (5.5) is nonpositive

$$-\int_{\Omega} \left((P'(\rho) \mathbf{m}(\rho) - H(\rho)) (\Delta \zeta)^2 + H(\rho) |D^2 \zeta|^2 \right) \, dx \stackrel{(4.11a)}{\leq} 0. \quad (5.14)$$

Since P is increasing and m is concave, we have $P'(\rho) \mathbf{m}''(\rho) \leq 0$ which yields

$$\int_{\Omega} P'(\rho) \mathbf{m}''(\rho) |\nabla \rho|^2 |\nabla \zeta|^2 \, dx \leq 0. \quad (5.15)$$

Since H is nonnegative, the smoothness and convexity of Ω and the smoothness of ζ yields, see [Gri85, Ot01, GST08],

$$\nabla |\nabla \zeta|^2 \cdot \mathbf{n} \leq 0, \quad \text{on } \partial\Omega, \quad \int_{\partial\Omega} H(\rho) \nabla |\nabla \zeta|^2 \cdot \mathbf{n} \, d\mathcal{H}^{d-1} \leq 0. \quad (5.16)$$

Combining (5.14), (5.15) and (5.16), (5.5) yields the inequality

$$\frac{\partial}{\partial t} \frac{1}{2} \int_{\Omega} |\nabla \zeta|^2 \mathbf{m}(\rho) \, dx \leq -\int_{\Omega} \nabla \zeta \cdot \nabla P(\rho) \, dx. \quad (5.17)$$

On the other hand

$$\frac{\partial}{\partial s} \int_{\Omega} U(\rho) \, dx = \int_{\Omega} U'(\rho) \partial_s \rho \, dx \stackrel{(5.4)}{=} \int_{\Omega} \mathbf{m}(\rho) \nabla U'(\rho) \cdot \nabla \zeta \, dx = \int_{\Omega} \nabla P(\rho) \cdot \nabla \zeta \, dx, \quad (5.18)$$

so that (5.12) follows by (5.17) and (5.18). Integrating (5.12) with respect to s and t , we obtain the second inequality in (5.13). The first inequality in (5.13) follows from the definition of $\mathcal{W}_{\mathbf{m}, \Omega}$ and the monotonicity of $\tau \mapsto \mathcal{W}(\mu_{1,\tau})$ (see the energy identity (4.6)). \square

6 Proof of the main theorems

6.1 The generation result

Recall that $\mathcal{S}_t(\mu_0) = S_t(\rho_0) \mathcal{L}^d|_{\Omega}$ when $\mu_0 = \rho_0 \mathcal{L}^d|_{\Omega}$; Theorem 4.11 is an immediate consequence of the following result.

Theorem 6.1. *Let Ω be a bounded, convex open set of \mathbb{R}^d and let us assume that $\mu_i \in \mathcal{M}^+(\overline{\Omega})$, $i = 0, 1$, have finite distance $\mathcal{W}_{\mathbf{m}, \Omega}(\mu_0, \mu_1) < +\infty$ (and therefore the same mass $\mathbf{m} = \mu_0(\Omega) = \mu_1(\Omega)$) and satisfy $\mathcal{U}(\mu_0) < +\infty$ and $\mu_1 = \rho_1 \mathcal{L}^d|_{\Omega} \ll \mathcal{L}^d$. If U satisfies the generalized McCann condition $\text{GMC}(\mathbf{m}, d)$ (4.11a, b) then*

$$\frac{1}{2} \mathcal{W}_{\mathbf{m}, \Omega}^2(\mathcal{S}_t \mu_1, \mu_0) + t \mathcal{U}(\mathcal{S}_t \mu_1) \leq \frac{1}{2} \mathcal{W}_{\mathbf{m}, \Omega}^2(\mu_1, \mu_0) + t \mathcal{U}(\mu_0) \quad \forall t \geq 0.$$

Proof. Since $\mathcal{W}_{\mathbf{m}, \Omega}(\mu_0, \mu_1) < +\infty$ there exists a geodesic curve $(\mu, \nu) \in \mathcal{CE}_{\Omega}(\mu_0 \rightarrow \mu_1)$ such that

$$\Phi_{\mathbf{m}, \Omega}(\mu_s, \nu_s) \equiv \int_0^1 \Phi_{\mathbf{m}, \Omega}(\mu_s, \nu_s) \, ds = \mathcal{W}_{\mathbf{m}, \Omega}^2(\mu_0, \mu_1).$$

Applying Lemma 3.5, we find a family of approximating curves $(\mu^{\varepsilon, \delta}, \nu^{\varepsilon, \delta})$ and smooth convex open sets Ω_{ε} satisfying (3.7), (3.8) and

$$\int_0^1 \Phi_{\mathbf{m}, \Omega_{\varepsilon}}(\mu^{\varepsilon, \delta}, \nu^{\varepsilon, \delta}) \, ds \leq c_{\varepsilon}^2 \mathcal{W}_{\mathbf{m}, \Omega}^2(\mu_0, \mu_1).$$

Let $\mathbf{m}_{\eta}, P_{\eta}$ as in Lemma 4.7, for constants $0 < M' < M'' < M$ such that $M' \leq \rho^{\varepsilon, \delta} \leq M''$ in Ω_{ε} , and let $\zeta_s^{\varepsilon, \delta, \eta} \in C^{\infty}(\mathbb{R}^d)$ obtained by solving (5.3) with respect to \mathbf{m}_{η} and $\partial_s \rho^{\varepsilon, \delta}$ in Ω_{ε} . Since $\mathbf{m}_{\eta} \geq \mathbf{m}$, by Theorem 5.21 of [DNS09] we easily have

$$\int_{\Omega_{\varepsilon}} |\nabla \zeta_s^{\varepsilon, \delta, \eta}|^2 \mathbf{m}_{\eta}(\rho^{\varepsilon, \delta}) \, dx = \Phi_{\mathbf{m}_{\eta}, \Omega_{\varepsilon}}(\mu^{\varepsilon, \delta}, \nu^{\varepsilon, \delta}) \leq \Phi_{\mathbf{m}, \Omega_{\varepsilon}}(\mu^{\varepsilon, \delta}, \nu^{\varepsilon, \delta}) \quad \forall \eta > 0, \, s \in [0, 1].$$

If $\mathcal{S}^{\varepsilon, \eta} = \mathcal{S}(\mathcal{U}_{\eta}, \mathbf{m}_{\eta}, \Omega_{\varepsilon})$ is the semigroup associated with $S(P_{\eta}, \Omega_{\varepsilon})$ and the corresponding integral functional \mathcal{U}_{η} , (5.13) gives

$$\begin{aligned} \frac{1}{2} \mathcal{W}_{\mathbf{m}_{\eta}, \Omega_{\varepsilon}}^2(\mathcal{S}_t^{\varepsilon, \eta} \mu_1^{\varepsilon, \delta}, \mu_0^{\varepsilon, \delta}) + t \mathcal{U}_{\eta}(\mathcal{S}_t^{\varepsilon, \eta} \mu_1^{\varepsilon, \delta}) &\leq \frac{1}{2} \int_0^1 \int_{\Omega_{\varepsilon}} |\nabla \zeta_s^{\varepsilon, \delta, \eta}|^2 \mathbf{m}_{\eta}(\rho^{\varepsilon, \delta}) \, dx \, ds + t \mathcal{U}_{\eta}(\mu_0^{\varepsilon, \delta}) \\ &\leq \frac{c_{\varepsilon}^2}{2} \mathcal{W}_{\mathbf{m}, \Omega}^2(\mu_1, \mu_0) + t \mathcal{U}_{\eta}(\mu_0^{\varepsilon, \delta}). \end{aligned}$$

Passing to the limit as $\eta \downarrow 0$ (notice that the functions $\mathcal{S}_t^{\varepsilon, \eta} \rho^{\varepsilon, \delta}$ take their values in $[M', M'']$) we get

$$\frac{1}{2} \mathcal{W}_{\mathbf{m}, \Omega_{\varepsilon}}^2(\mathcal{S}_t^{\varepsilon} \mu_1^{\varepsilon, \delta}, \mu_0^{\varepsilon, \delta}) + t \mathcal{U}(\mathcal{S}_t^{\varepsilon} \mu_1^{\varepsilon, \delta}) \leq \frac{c_{\varepsilon}^2}{2} \mathcal{W}_{\mathbf{m}, \Omega}^2(\mu_1, \mu_0) + t \mathcal{U}(\mu_0^{\varepsilon, \delta})$$

where $\mathcal{S}^{\varepsilon} = \mathcal{S}(\mathcal{U}, \mathbf{m}, \Omega_{\varepsilon})$ is associated with $S(P, \Omega_{\varepsilon})$. We can then pass to the limit as $\delta \downarrow 0$: since $\rho^{\varepsilon, \delta} \rightarrow \rho^{\varepsilon}$ in $L^{\infty}(\Omega_{\varepsilon})$ we immediately have

$$\frac{1}{2} \mathcal{W}_{\mathbf{m}, \Omega_{\varepsilon}}^2(\mathcal{S}_t^{\varepsilon} \mu_1^{\varepsilon}, \mu_0^{\varepsilon}) + t \mathcal{U}(\mathcal{S}_t^{\varepsilon} \mu_1^{\varepsilon}) \leq \frac{c_{\varepsilon}^2}{2} \mathcal{W}_{\mathbf{m}, \Omega}^2(\mu_1, \mu_0) + t \mathcal{U}(\mu_0^{\varepsilon}).$$

Finally as $\varepsilon \downarrow 0$ we conclude, recalling Proposition 4.3. □

6.2 Geodesic convexity

The proof of Theorem 4.13 follows immediately from the generation result Theorem 6.1 and Theorem 4.10 if every measure μ_s of the geodesic curve is absolutely continuous with respect to $\mathcal{L}^d|_\Omega$ (see also Remark 4.15). On the other hand, this property is not known *a priori*, so we need a more refined argument.

Proof. As in Proposition 3.4 we set $\mu_s^\varepsilon := \mu_s * k_\varepsilon$, $\nu_s^\varepsilon := \nu_s * k_\varepsilon$ and we denote by $\mathcal{S}^\varepsilon = \mathcal{S}(\mathcal{W}, \mathbf{m}, \Omega_\varepsilon)$. By (3.5) and the contraction property given by Theorem 4.11 in Ω_ε we have

$$\mathcal{W}_{\mathbf{m}, \Omega_\varepsilon}(\mathcal{S}_t^\varepsilon \mu_{s_1}^\varepsilon, \mathcal{S}_t^\varepsilon \mu_{s_2}^\varepsilon) \leq \mathcal{W}_{\mathbf{m}, \Omega_\varepsilon}(\mu_{s_1}^\varepsilon, \mu_{s_2}^\varepsilon) \leq \mathcal{W}_{\mathbf{m}, \Omega}(\mu_{s_1}, \mu_{s_2}) = |s_1 - s_2| \mathcal{W}_{\mathbf{m}, \Omega}(\mu_0, \mu_1) \quad (6.1)$$

and by (3.6), Theorem 4.10 and (4.18) we have

$$\delta_\varepsilon^2 := \mathcal{W}_{\Omega, \mathbf{m}}^2(\mu_0, \mu_1) - \mathcal{W}_{\mathbf{m}, \Omega_\varepsilon}^2(\mu_0^\varepsilon, \mu_1^\varepsilon) \rightarrow 0 \quad \text{as } \varepsilon \downarrow 0,$$

$$\mathcal{W}(\mathcal{S}_t^\varepsilon \mu_s^\varepsilon) \leq (1-s)\mathcal{W}(\mu_0^\varepsilon) + s\mathcal{W}(\mu_1^\varepsilon) + \frac{\delta_\varepsilon^2}{2t}s(1-s), \quad (6.2)$$

$$\mathcal{W}_{\mathbf{m}, \Omega_\varepsilon}^2(\mathcal{S}_t^\varepsilon \mu_i^\varepsilon, \mu_i) \leq t(\mathcal{W}(\mu_i^\varepsilon) - \inf \mathcal{W}) \leq t(\mathcal{W}(\mu_i) - \inf \mathcal{W}), \quad (6.3)$$

where the second inequality in (6.3) follows from Jensen's inequality.

We choose now a countable set \mathcal{C} dense in $[0, 1]$ and containing 0 and 1, a vanishing sequence $(t_k)_{k \in \mathbb{N}}$ and another vanishing sequence $(\varepsilon_k)_{k \in \mathbb{N}}$ so that $\lim_{k \uparrow +\infty} t_k^{-1} \delta_{\varepsilon_k}^2 = 0$. By compactness and a standard diagonal argument, up to extracting a further subsequence, we can find limit points $\tilde{\mu}_s$ for $s \in \mathcal{C}$ such that

$$\mathcal{S}_{t_k}^{\varepsilon_k} \mu_s^{\varepsilon_k} \rightharpoonup \tilde{\mu}_s \quad \text{weakly as } k \uparrow +\infty.$$

By (6.1) and Proposition 3.2 we get

$$\mathcal{W}_{\mathbf{m}, \Omega}(\tilde{\mu}_{s_1}, \tilde{\mu}_{s_2}) \leq |s_1 - s_2| \mathcal{W}_{\mathbf{m}, \Omega}(\mu_0, \mu_1) \quad \forall s_1, s_2 \in \mathcal{C}. \quad (6.4)$$

(6.3) yields $\tilde{\mu}_0 = \mu_0$, $\tilde{\mu}_1 = \mu_1$ so that we can extend $\tilde{\mu}$ to a continuous curve (still denoted by $\tilde{\mu}$) connecting μ_0 and μ_1 still satisfying (6.4) for every $s_1, s_2 \in [0, 1]$. The triangular inequality shows that (6.4) is in fact an equality and the curve $\tilde{\mu}$ is a constant speed minimizing geodesic. On the other hand, the lower semicontinuity of \mathcal{W} with respect to weak convergence and (6.2) yields

$$\mathcal{W}(\tilde{\mu}_s) \leq (1-s)\mathcal{W}(\mu_0) + s\mathcal{W}(\mu_1) \quad \forall s \in \mathcal{C}. \quad (6.5)$$

A further lower semicontinuity and density argument shows that (6.5) holds for every $s \in [0, 1]$. \square

Final remarks and open problems

This paper is a first step towards the investigation of the geometry of spaces of measures metrized by $\mathcal{W}_{m,\Omega}$, the induced convexity notions for integral functionals and the corresponding generation of gradient flows with applications to various nonlinear evolutionary PDE's.

Since a sufficiently general theory is far from being developed and understood, it is in some sense surprising that one can reproduce in this setting the celebrated McCann convexity result. On the other hand, many interesting and basic problems remain open: here is just a provisional list.

- At the level of the distance $\mathcal{W}_{m,\Omega}$ only partial results on some basic properties (such as density of regular measures or necessary and sufficient conditions ensuring the finiteness of the distance), are known and a complete and accurate picture is still missing.
- The situation is even less clear in the case of unbounded domains: in this paper we restricted our attention to the bounded case only.
- The study of other integral functionals is completely open, as well as applications to different types of evolution equations, like scalar conservation laws or nonlinear fourth order equation of thin-film type, see the introduction of [DNS09].
- It would be interesting to study other metric quantities (e.g. the metric slope) and the pseudo-Riemannian structure (tangent space, Alexandrov curvature, etc) connected with the distance and the energy functionals, see [AGS05, CMV06].
- The regularization properties and asymptotic behaviour of the gradient flow \mathcal{U} and its perturbation can be studied as well: in the Wasserstein case the geodesic convexity of a functional yields many interesting estimates.
- The convergence of the so called “Minimizing movement” or JKO-scheme could be exploited in this and other situations: in the case of geodesically convex energies, further information on the Alexandrov curvature of the distance $\mathcal{W}_{m,\Omega}$ would be crucial, see [AGS05, Sa07].

Acknowledgments. J.A.C. acknowledges the support from DGI-MICINN (Spain) project MTM2008-06349-C03-03. G.S. and S.L. have been partially supported by MIUR-PRIN'06 grant for the project “Variational methods in optimal mass transportation and in geometric measure theory”. D.S. was partially supported by NSF grant DMS-0638481. D.S. would also like to thank the Center for Nonlinear Analysis (NSF grants DMS-0405343 and DMS-0635983) for its support during the preparation of this paper.

References

- [AGS05] L. Ambrosio, N. Gigli, and G. Savaré, Gradient flows in metric spaces and in the space of probability measures, Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 2005.

- [AS06] L. Ambrosio and G. Savaré, Gradient flows of probability measures, Handbook of differential equations: evolutionary equations 3, C.M. Dafermos, E. Feireisl editors, 1–136, Elsevier, 2006.
- [BB00] J.-D. Benamou and Y. Brenier, A computational fluid mechanics solution to the Monge-Kantorovich mass transfer problem, Numer. Math., 84 (2000), pp. 375–393.
- [Ber98] A. Bertozzi, The mathematics of moving contact lines in thin liquid films, Notices Amer. Math. Soc., 45 (1998), pp. 689–697.
- [B03] Y. Brenier, Extended Monge-Kantorovich theory, Optimal transportation and applications (Martina Franca, 2001), Lecture Notes in Math. 1813, 91–121, Springer-Berlin 2003.
- [BFD06] M. Burger, M. di Francesco and Y. Dolak, The Keller-Segel model for chemotaxis with prevention of overcrowding: linear vs. nonlinear diffusion, SIAM J. Math. Anal., 38 (2006), pp. 1288–1315.
- [BD09] M. Burger and M. Di Francesco, Large time behavior of nonlocal aggregation models with nonlinear diffusion, to appear in Networks and Heterogenous Media.
- [CLR08] J. A. Carrillo, P. Laurençot and J. Rosado, Fermi-Dirac-Fokker-Planck Equation: Well-posedness & Long-time Asymptotics, preprint UAB, 2008.
- [CMV03] J. A. Carrillo, R. J. McCann and C. Villani, Kinetic equilibration rates for granular media and related equations: entropy dissipation and mass transportation estimates, Rev. Matemática Iberoamericana, 19 (2003), pp. 1–48.
- [CMV06] J. A. Carrillo, R. J. McCann and C. Villani, Contractions in the 2-Wasserstein length space and thermalization of granular media, Arch. Ration. Mech. Anal., 179 (2006), pp. 217–263.
- [CRS08] J. A. Carrillo, J. Rosado and F. Salvarani, 1D nonlinear Fokker-Planck equations for fermions and bosons, Appl. Math. Lett., 21 (2008), pp. 148–154.
- [DS08] S. Daneri and G. Savaré, Eulerian calculus for the displacement convexity in the Wasserstein distance, SIAM J. Math. Anal., 3 (2008), pp. 1104–1122.
- [DR09] M. Di Francesco and J. Rosado, Fully parabolic Keller-Segel model for chemotaxis with prevention of overcrowding, to appear on Nonlinearity.
- [DNS09] J. Dolbeault, B. Nazaret, and G. Savaré, A new class of dynamic transport distances between measures, Calc. Var. & PDEs, 34 (2009), pp. 193–231.
- [F04] T. D. Frank, Classical Langevin equations for the free electron gas and blackbody radiation, J. Phys. A, 37 (2004), pp. 3561–3567.

- [F05] T. D. Frank, *Nonlinear Fokker-Planck Equations*, Springer Series in Synergetics, Springer, 2005.
- [GL1] G. Giacomini and J. Lebowitz, Phase segregation dynamics in particle systems with long range interactions. I. Macroscopic limits, *J. Statist. Phys.*, 87 (1997), pp. 37–61.
- [Gri85] P. Grisvard, *Elliptic problems in nonsmooth domains*, Monographs and Studies in Mathematics 24, Pitman (Advanced Publishing Program), Boston, MA, 1985.
- [GST08] U. Gianazza, G. Savaré and G. Toscani, The Wasserstein gradient flow of the Fisher information and the quantum drift-diffusion equation, to appear in *Arch. Ration. Mech. Anal.*
- [H03] D. Horstmann, From 1970 until present: the Keller-Segel model in chemotaxis and its consequences. I, *Jahresber. Deutsch. Math.-Verein.*, 105 (2003), pp. 103–165.
- [K95] G. Kaniadakis, Generalized Boltzmann equation describing the dynamics of bosons and fermions, *Phys. Lett. A*, 203 (1995), pp. 229–234.
- [K93] G. Kaniadakis, P. Quarati, Kinetic equation for classical particles obeying an exclusion principle, *Phys. Rev. E*, 48 (1993), pp. 4263–4270.
- [L07] S. Lisini, Characterization of absolutely continuous curves in Wasserstein spaces. *Calc. Var. Partial Differential Equations* 28, (2007), pp. 85–120.
- [LM] S. Lisini and A. Marigonda, work in preparation.
- [McC97] R. J. McCann, A convexity principle for interacting gases, *Adv. Math.*, 128 (1997), pp. 153–179.
- [Ot01] F. Otto, The geometry of dissipative evolution equation: the porous medium equation, *Comm. Partial Differential Equations*, 26 (2001), pp. 101–174.
- [OW05] F. Otto and M. Westdickenberg, Eulerian calculus for the contraction in the Wasserstein distance, *SIAM J. Math. Anal.*, 37 (2005), pp. 1227–1255.
- [Sa08] F. Santambrogio, Absolute continuity and summability of transport densities: simpler proofs and new estimates, CVGMT preprint (<http://cvgmt.sns.it/papers/san08/>), 2008.
- [Sa07] G. Savaré, Gradient flows and diffusion semigroups in metric spaces under lower curvature bounds, *C. R. Math. Acad. Sci. Paris*, 345 (2007), pp. 151–154.
- [Sl08] D. Slepčev, Coarsening in nonlocal interfacial systems, *SIAM J. Math. Anal.*, 40 (2008), pp. 1029–1048.
- [SSC06] J. Sopiak, C. Sire and P.H. Chavanis, Dynamics of the Bose-Einstein condensation: analogy with the collapse dynamics of a classical self-gravitating Brownian gas, *Phys. Rev. E*, 74 (2006), pp. 011112.

- [Vaz07] J. L. Vázquez, The porous medium equation, Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, Oxford, 2007.
- [Vil03] C. Villani, Topics in optimal transportation, Graduate Studies in Mathematics 58, American Mathematical Society, Providence, RI, 2003.
- [Vil07] C. Villani, Optimal transport, Old and New, Springer-Verlag 2008.