

Biomedical Physics & Engineering Express



PAPER

SNAIL driven by a feed forward loop motif promotes TGF β induced epithelial to mesenchymal transitionGottumukkala Sai Bhavani¹ and Anbumathi Palanisamy^{2,*} ¹ Department of Biotechnology, NIT Warangal, India² Department of Biotechnology, NIT Warangal, India

* Author to whom any correspondence should be addressed.


E-mail: anbu@nitw.ac.in**Keywords:** epithelial to mesenchymal transition (EMT), mathematical modelling, cancer metastasis, SNAIL, TGF β signaling, systems biology, coherent type 1 feed forward loop (C1FFL)Supplementary material for this article is available [online](#)RECEIVED
26 January 2022REVISED
20 May 2022ACCEPTED FOR PUBLICATION
14 June 2022PUBLISHED
24 June 2022**Abstract**

Epithelial to Mesenchymal Transition (EMT) plays an important role in tissue regeneration, embryonic development, and cancer metastasis. Several signaling pathways are known to regulate EMT, among which the modulation of TGF β (Transforming Growth Factor- β) induced EMT is crucial in several cancer types. Several mathematical models were built to explore the role of core regulatory circuit of ZEB/miR-200, SNAIL/miR-34 double negative feedback loops in modulating TGF β induced EMT. Different emergent behavior including tristability, irreversible switching, existence of hybrid EMT states were inferred through these models. Some studies have explored the role of TGF β receptor activation, SMADs nucleocytoplasmic shuttling and complex formation. Recent experiments have revealed that MDM2 along with SMAD complex regulates SNAIL expression driven EMT. Encouraged by this, in the present study we developed a mathematical model for p53/MDM2 dependent TGF β induced EMT regulation. Inclusion of p53 brings in an additional mechanistic perspective in exploring the EM transition. The network formulated comprises a C1FFL moderating SNAIL expression involving MDM2 and SMAD complex, which functions as a noise filter and persistent detector. The C1FFL was also observed to operate as a coincidence detector driving the SNAIL dependent downstream signaling into phenotypic switching decision. Systems modelling and analysis of the devised network, displayed interesting dynamic behavior, systems response to various inputs stimulus, providing a better understanding of p53/MDM2 dependent TGF β induced Epithelial to Mesenchymal Transition.

Introduction

The art of communication among cells performing various tasks in response to limitless signals around them is phenomenal. These signals play a crucial role in cell cycle control and regulation (Tyson and Novak 2014). Failure in this signal transduction to elicit appropriate biological response results in a wide array of diseases including cancer (Hanahan and Weinberg 2011; Sever and Brugge 2015). Evidence suggests that the process of carcinogenesis is a multistep process that occurs due to alterations in these signaling pathways (Hanahan and Weinberg 2000).

One of the widely observed characteristics of carcinomas is Epithelial-Mesenchymal Transition (EMT) (Thiery 2002; Thiery and Sleeman 2006; Thiery *et al* 2009; Scheel and Weinberg 2012; Nieto *et al* 2016; Lambert *et al* 2017; Moustakas and de Herreros 2017). EMT (figure 1) is a process where epithelial cell loses its cell polarity, adhesive properties and subsequently gains migratory, invasive phenotype known as mesenchymal cells (Kang and Massague 2004; Onder *et al* 2008; Lamouille *et al* 2014; Prislei *et al* 2015; Savagner 2015). Literature has illustrated the involvement of several important molecules, transcription factors and their associated pathways in cancer. Signaling pathways including EGF, EGFR, TKs, ERK,



Ras, MAPK, PI3K, mTOR, CREB, Rho-Rock, Wnt, Notch delta, Shh, $TGF\beta$ are attributed to cause the mammalian cells to undergo EMT. (Kalluri and Weinberg 2009; De Craene and Berx 2013)

Among the above-mentioned pathways, the family of $TGF\beta$ is well known for its dual role (as suppressor and promoter of tumour) and plays a remarkable part in modulating EMT (Dickson *et al* 1987; Miyazono 2000; Derynck and Zhang 2003; Brier and Moses 2006; Xu *et al* 2009; Ceppi *et al* 2010; Ikushima and Miyazono 2010; Gregory *et al* 2011; Bedi *et al* 2012; Bryant *et al* 2012; Reichl *et al* 2012; Xu *et al* 2012; Giannelli *et al* 2014). $TGF\beta$ (Transforming Growth Factor- β) and its family of cytokines are involved in countless physiological/biological processes responsible for regulating cellular differentiation, proliferation, migration, and apoptosis by activation of their serine/threonine transmembrane receptors (Attisano and Wrana 1996, 1998; Massague 1998; Attisano and Wrana 2000; Attisano and Wrana 2002; Zavadil and Bottinger 2005; Wu and Hill 2009). Activation of these receptors by phosphorylation results in a complex cascade of signaling in both SMAD dependent (Canonical) and SMAD-independent manner (Non-Canonical) (Heldin *et al* 1997; Attisano and Wrana 1998; Attisano and Wrana 2000; Johnsen *et al* 2002; Derynck and Zhang 2003; Massague *et al* 2005; Xu *et al* 2009; Lamouille *et al* 2012; Massague 2012).

SMAD dependent activation of $TGF\beta$ results in the activation of crucial transcription factors including SNAIL (SNAIL/SLUG), ZEB (ZEB1, ZEB2) family of regulators which are known to directly repress the epithelial marker E-Cadherin and upregulate the Mesenchymal marker N-cadherin (Thiery *et al* 2009; Lamouille *et al* 2013). Several miRNAs are found to be involved in this process among which families of miR-34a and miR-200 are well studied (Gregory *et al* 2008; Park *et al* 2008; Paterson *et al* 2008; Ceppi *et al* 2010; Gregory *et al* 2011; Siemens *et al* 2011). These miRNA families are well known for their double negative feedback loops with ZEB and SNAIL transcription factor families. (Bracken *et al* 2008; Burk *et al* 2008;

Wellner *et al* 2009; Gregory *et al* 2011; Siemens *et al* 2011; Tian *et al* 2013). Several mathematical models of EMT are built around these regulators to understand the dynamics of EMT.

SNAIL is a zinc finger transcription factor family comprising of SNAIL1, SNAIL2, and SNAIL3. Over-expression of SNAIL family of regulators were observed to be associated with cancer and metastasis (Hemavathy *et al* 2000; Nieto 2002; Vega *et al* 2004; Yang *et al* 2009; Franco *et al* 2010; Li *et al* 2011; Fan *et al* 2012; Zhang *et al* 2013; Yang *et al* 2017; Cai *et al* 2019). SNAIL was observed to recruit cofactors like HDAC1, CBP300, HDAC2, SIN3A, ctBP, etc to form a repressor complex to repress epithelial gene expression (Peinado *et al* 2004; Peinado *et al* 2007). It is a highly unstable protein in cytoplasm and needs to translocate into nucleus to exert its function. PAK1 is known to phosphorylate SNAIL at S246 that favours its nuclear localization (Yang *et al* 2005). The process of EMT and cancer resistance are closely linked with each other. Snail expression was also found to be associated with chemo- and radio-resistance increasing cell stemness (Smith and Bhowmick 2016). SNAIL was observed to show resistance to the drugs that have been developed promoting cancer progression and metastases. They include 5- Fluorouracil, Cisplatin, Doxorubicin in targeting Breast, HNSCC (Head and Neck Squamous Cell Carcinoma) and ovarian cancer (Hsu *et al* 2010; Li and Zhou 2011; Haslehurst *et al* 2012; Shen *et al* 2012; Zhang *et al* 2012; Mariano *et al* 2015). Considering the numerous studies that have been developed to understand the signaling molecules that drive EMT and their correlation with chemoresistance, SNAIL has become an attractive target to explore cancer resistance.

In addition to SNAIL and ZEB families, recent experimental studies have shown the evidence of MDM2 in regulating SNAIL expression by promoting $TGF\beta$ induced EMT. Studies have shown that the activation of oncogene MDM2 through $TGF\beta$ induced SMAD complex (Araki *et al* 2010). MDM2 enhances its own activity through a positive feedback loop

upregulating R-SMAD phosphorylation and thus promotes activation of SMAD complex (Araki *et al* 2010; Lu *et al* 2016; Chen *et al* 2017; Tang *et al* 2019). Previous studies also have shown complementary role of MDM2 in degrading SNAIL by ubiquitination, when in complex with wild type p53 thus inhibiting EMT (Wang *et al* 2009; Lim *et al* 2010). Wild type p53 is a well-known tumour suppressor gene notable for its regulatory functions in cell cycle (Rivlin *et al* 2011; Kim *et al* 2019). In many cancer types p53 is mutated, resulting in upregulation of MDM2 that transcriptionally activates SNAIL along with suppression of p53 (Lu *et al* 2016). p53 is also known to regulate the expressions of epithelial markers miR-34a, miR-200 which are part of the double negative feedback loops ZEB/ miR200 and SNAIL/miR34 promoting EMT (Hunziker *et al* 2010; Chang *et al* 2011; Kim *et al* 2011; Coradini *et al* 2012; Ren *et al* 2013).

Several mathematical models were built to understand the dynamics of TGF β signaling and their significance in inducing Epithelial to Mesenchymal transition (Miettinen *et al* 1994; Heldin *et al* 1997; Vilar *et al* 2006; Zi and Klipp 2007; Turner and Kohandel 2010; Vilar and Saiz 2011; Malfettone *et al* 2017; Tripathi *et al* 2020). Amongst which various models were built around the core regulatory network of TGF β induced EMT. The network mainly comprises of double negative feedback loops of miR-200/ZEB and miR-34a/SNAIL to understand the dynamics of transformation from epithelial to mesenchymal phenotype (Lu *et al* 2013; Tian *et al* 2013; Zhang *et al* 2014; Jolly Jia *et al* 2016; Jolly Tripathi *et al* 2016). Studies show the prominence of double negative feedback loops in modulating the reversible and irreversible bistable switching from epithelial to mesenchymal phenotypes along with existence of a hybrid EMT/ MET phenotypes. It was also observed that release of miR-200-mediated inhibition of TGF β is required to maintain a stable mesenchymal phenotype (Gregory *et al* 2011; Tian *et al* 2013). A model developed by Lu *et al* (2013), depicts the significance of mRNA: miRNA coupling in the process of EMT. Autoregulation of the transcription factor SNAIL was observed to act as a noise integrator in the process of EMT (Lu *et al* 2013). However, many of these models found in literature review are focused on (i) the upstream SMAD signaling and its nucleocytoplasmic shuttling, (ii) the downstream double negative feedback loops controlling the E/M, M/E (ir)reversible switch. There is no dynamic ordinary differential equation based modelling framework that explores the role of p53, MDM2 dependent alterations to the SNAIL and SNAIL dependent miRNA regulations of EMT. Therefore, motivated by the role of p53-MDM2 in TGF β induced EMT and their involvement in regulating miRNAs, a mathematical model was developed coupling SMAD regulation along with its downstream core regulatory network as shown in figure 2.

Methods

Network that regulates epithelial to mesenchymal transition

TGF β signaling pathways were observed to be the major players in the development of epithelial to mesenchymal phenotypic switch and thus migratory behavior in several cancer cells. Therefore, with the objective to explore the role of various key regulators involved in this regulatory process, a pathways network was assembled. An extensive literature survey was performed to obtain the signaling flow following the recent experiments that has revealed the regulatory roles of SMAD, p53, MDM2 in regulating SNAIL expression (Lu *et al* 2016; Chen *et al* 2017; Tang *et al* 2019). Hence, a gene regulatory network depicting TGF β induced EMT was assembled (figure 2) for a cancerous cell which includes 29 nodes (genes, receptors, miRNAs, Phenotype) connected through 45 edges (activation, repression, nucleocytoplasmic shuttling). The network consists of cancer marker genes p53, MDM2 in negative feedback loop activated through DNA Damage. ZEB, SNAIL mesenchymal markers regulated through MDM2, SMAD complex that initiates and facilitates the process of EMT through inhibition of E-Cadherin, activation of N-Cadherin, activation of latent TGF β (represented as enTGFb). Also epithelial markers miR-34 and miR-200 that are in double negative feedback loops with mesenchymal markers were considered. SNAIL activated through upstream regulatory molecules (MDM2, SMAD complex) subsequently activates the latent TGF β and its receptors TGF β R (Represented as enTGFb and TGFbR in figure 2) through a positive feedback dependent regulation, initiating the process of EMT. The network construct also considers the SMADs complex association, dissociation and nucleocytoplasmic shutting. The focus of the study was primarily on the MDM2/SNAIL axis that triggers the TGF- β /SMAD/SNAIL signaling. The network assembled thus is show in figure 2. The logical relationships among nodes was tabulated in supplementary Table S1(b). The network was further analysed and characterised using both discrete (Boolean logical modelling) and continuous dynamic modelling using ordinary differential equations.

Discrete dynamic modelling

The network constructed (figure 2) was translated into the dynamic model to explore EMT/MET and the role of various regulators in the process of transition. Discrete dynamic modelling was utilized as first step because of its computational feasibility and capacity to work with qualitative biological data. Specifically Boolean modelling approach was adopted where each network node was represented by only two known qualitative states: ON or OFF. The ON state means a regulator is in an active form or at its maximum




Table 1. Various state combinations of TCF1, TCF2 and their resulting phenotype. +/- indicates the presence or absence of hybrid phenotype.

Scenario	TCF1	TCF2	exTGF β	Hybrid	EMT (Time Steps)
1	0	0	0	+	Hybrid (Oscillations)
2	1	1	0	+	72 Time steps
3	0	1	0	+	113 Time steps
4	1	0	0	+	86 Time steps
5	0	0	1	—	Hybrid (Oscillations)
6	1	1	1	—	10 Steps
7	0	1	1	—	10 Steps
8	1	0	1	—	12 Steps

activity (represented as 1) whereas OFF state means a regulator is inactive form or is at its basal minimum activity (represented as 0). Regulatory functions for each node (Regulator) were formulated depending on their upstream regulators and are known to produce known biological outcomes (Supplementary Table S1 (available online at stacks.iop.org/BPEX/8/045012/mmedia)). The *python* library *Booleannet* downloaded from (<https://github.com/ialbert/booleannet>) (Albert *et al* 2008) was used to perform the simulations. Python 3.x was used with Ubuntu as working operating

system. Considering all the timescales involved in transcriptional and signal transduction events, asynchronous update schema was used. This accounts for different timescales in signaling networks, in which the state of each node was updated according to its own timescale (Garg *et al* 2008; Chatain *et al* 2018).

To evaluate the TGF β dependent transition of cells from epithelial to mesenchymal transition, epithelial state was assumed to be the initial state of the system responding to the signal DNA Damage. Multiple

simulations were performed with various initial conditions and varying states of certain regulators some of which are tabulated in table 1. To further assess the effect of various regulators on the reversibility of EMT, the initial epithelial conditions was over-raided after 110-time steps in each simulation when the system was assumed to be transitioned completely into mesenchymal state and attained steady state. The simulations were performed for 5000 timesteps and 500 iterations.

Continuous dynamic modelling

To further explore the role of network structure driven dynamic behaviour, continuous dynamic modelling was utilised. ODE modelling and simulations were adopted specifically to deduce the functional role of the SMAD, MDM2 on SNAIL which were observed to be interacting in a coherent type1 feed forward regulatory (C1FFL) interaction (figure 4). Two input functions S_x (TGF β) , S_y (p53) were considered to be the stimulus that activates SMAD complex and MDM2 which are required for SNAIL transcriptional activation. Various different combinations of continuous, pulsatile inputs were provided in the form of S_x and S_y to observe their influence on SNAIL transcriptional activity. Both AND gate like logic and OR gate like logic were independently explored for S_x and S_y dependent regulation of the regulator SNAIL in C1FFL. These mechanisms for the individual regulators SMAD complex, MDM2 and SNAIL were captured in terms of mass action kinetics and Hill equations-based ODE as shown below.

$$\frac{d[SMAD]}{dt} = a * S_x - Kd1 * [SMAD] \quad (1)$$

$$\frac{d[MDM2]}{dt} = b * S_y + \frac{k1 * [SMAD]^n}{km1^n + [SMAD]^n} - kd2 * [MDM2] \quad (2)$$

$$\frac{d[SNAIL]}{dt} = G(MDM2, k2; SMAD, k3) - kd2 * [SNAIL] \quad (3)$$

The regulatory function for AND gate is $G_{SNAIL}(MDM2, k2) * G_{SNAIL}(SMAD, k3)$ follows the hill functions and is given as $\frac{k2 * [MDM2]^n}{km2^n + [MDM2]^n} * \frac{k3 * [SMAD]^n}{km3^n + [SMAD]^n}$. The regulatory function for OR gate is $G_{SNAIL}(MDM2, k2) + G_{SNAIL}(SMAD, k3)$ and is given as $\frac{k2 * [MDM2]^n}{km2^n + [MDM2]^n} + \frac{k3 * [SMAD]^n}{km3^n + [SMAD]^n}$.

The ODE models were simulated using *ode23s* solver of MATLAB R2021a. The Associated parameters and their descriptions were tabulated in supplementary Figure S29 and Table S29.


RESULTS

SNAIL lies at the epicentre of p53 dependent EMT regulation

The network devised (figure 2) comprises of SMADs complex formation, SMADs nucleocytoplasmic shuttling, and the core regulatory network that accounts for the effect of p53-MDM2. We hypothesised that the network gets activated in a layered manner. The first layer comprises the activation of serine/threonine receptors by exogenous or endogenous TGF β through phosphorylation (Massague 1998; Attisano and Wrana 1996, 2002; Vander Ark *et al* 2018). The second layer comprises of SARA dependent recruitment of R-SMAD to the receptors for activation by phosphorylation. Phosphorylated R-SMAD forms a complex with Co-SMAD that shuttles into the nucleus, and thus promotes transcriptional activation of MDM2, SNAIL in presence of transcription cofactors TCF1, TCF2 (Heldin *et al* 1997; Attisano and Wrana 1998; Johnsen *et al* 2002; Derynck and Zhang 2003). Once shuttled into nucleus the complex disassociates into R-SMADP and Co-SMAD. These disassociated parts further shuttle between the nucleus and cytoplasm or gets degraded depending on the environment (Nakabayashi and Sasaki 2009). In the third layer, DNA damage dependent p53-MDM2 and their associated regulators are considered. MDM2 activates SNAIL, which directly suppresses mir-34, miR-200, E-cadherin and p53 which are required for maintaining epithelial state. By repression of these markers and by activating Zeb, N-cadherin Mesenchymal transition is promoted. These interaction have the double negative feedback loop architecture between SNAIL/miR-34a and ZEB/miR-200 (Ciliberto *et al* 2005; Burk *et al* 2008; Korpall *et al* 2008; Wang *et al* 2009; Hunziker *et al* 2010; Tryndyak *et al* 2010; Kim *et al* 2011; Siemens *et al* 2011; Liu *et al* 2013). MDM2 also enhances its own regulation through phosphorylation of R-SMADs in a positive feedback manner (Chen *et al* 2017; Tang *et al* 2019). In addition to the regulations discussed above, SNAIL also activates the endogenous TGF β that enhances the TGF β signaling, and miR-200 negatively inhibits this endogenous TGF β (Zhang *et al* 2014). We speculate a crucial role for this network in governing the downstream EMT transition. Thus, the network was assembled by reviewing the relevant literature to understand the network influenced behavior that promotes EMT.

Logical modelling and simulation of TGF β dependent EMT regulation

To understand the dynamics of signaling during the process of epithelial to mesenchymal transition, the core regulatory network shown in figure 2 was translated into Logical and Mathematical Models. Discrete dynamic modelling was used as first step because of its computational feasibility and capacity to work with qualitative biological data. Specifically, Boolean Logical modelling was used, and simulations




were conducted using the *python* library *Booleannet* downloaded from (<https://github.com/ialbert/booleannet>) (Albert *et al* 2008). Logical modelling was based on the components of the regulatory network where each regulatory gene is considered a node and the connection between nodes are considered as edges. Each network node is described by binary states: ON (active) or OFF (inactive) making the model discrete deterministic and parameter free. The regulatory functions in the network were formulated based on the influence from the edges built by logical operators AND, OR, NOT representing biological interactions found in literature (Table S1). While beginning the study we performed simulations for 5000 timesteps, 500 iterations with different input conditions (table 1) to explore the time required for complete EMT

transition. Asynchronous update scheme was preferred because it accounts for all the genes at each and every time step during the state transition (Garg *et al* 2008; Chatain *et al* 2018). The simulation captures the complete EMT transition within 100 timesteps (figure 3).

Epithelial to mesenchymal transition

Initially, for logical modelling the nodes were set according to their known epithelial cell states. Specifically, the nodes of epithelial markers p53, miR-200, miR-34a, E-Cadherin were set to ON (1) state and the mesenchymal markers SNAIL, ZEB, N-Cadherin, MDM2, SMADs, exTGF- β , enTGF β were set to OFF (0) state. The model without any stimulus or dysregulation remains at epithelial state where E-Cadherin is




at its maximum node activity with N-Cadherin at its minimum node activity as shown in Figure S2. Subsequently, to implement dysregulation in cancer cells, DNA Damage was set to ON state to observe the resulting signaling events (figure 3). When the DNA Damage is set to ON state it was observed that the presence of this dysregulation activates the p53-MDM2 signaling axis further leading to activation of SNAIL. Increased node activity of SNAIL subsequently activates the enTGF β through positive feedback that regulates SMAD signaling axis thus further enhancing transcriptional activity of MDM2 and SNAIL. This enhanced transcriptional activity of SNAIL subsequently triggers ZEB transcription, which in turn represses the epithelial markers. Turning DNA Damage to ON state ultimately led to the changes in all the nodes particularly EMT hallmarks over the time course of 72 timesteps. All the epithelial markers were observed to be at their minimum activity (0) and mesenchymal markers were observed to be at their maximum activity (1) reflecting TGF β driven EMT. If a node is reaching steady state with minimum activity (OFF state) to begin with and ends up at maximum activity (ON state) by the end of simulation, it is inferred as network dependent rule driven steady state. Existence of a hybrid EM phenotype, (both epithelial and mesenchymal states) was observed for a period of 40 timesteps before the cell transits completely into mesenchymal phenotype. Presence of epithelial markers (figure 3d) and mesenchymal markers (figure 3e) were observed from timestep 10 to timestep 40 which was interpreted as EM hybrid state for this study.

To observe the effect of external environmental stress that induces EMT, exTGF β was set to ON state. It was observed that when exTGF β is turned ON the steady state is reached within 10 timesteps signifying the role of exTGF β (any similar stress) in accelerating

the EMT switch Figure S3. Further to know the impact of the feedforward (figure 4) relationship amongst SMAD complex, MDM2 in regulating the SNAIL transcriptional activation and their role in EMT, the nodes of their transcription cofactors TCF1 and TCF2 were set to ON & OFF states by keeping (i) exTGF β at its ON state; (ii) exTGF β at its OFF state (table 1). When stimulus to MDM2, *i.e.*, TCF1 was set to OFF and exTGF β was set to OFF, lack of stimulus to MDM2 causes the p53-MDM2 axis to oscillate (Figure S4). However, there is activity of mesenchymal markers that maintains the levels of their counterpart miRNAs as shown in Figure S4. Activation of SNAIL by enTGF β and by MDM2 enhances its own activity, thus simultaneously suppressing epithelial markers and promoting mesenchymal markers leading to EMT. When the stimulus for SNAIL, *i.e.*, TCF2 along with exTGF β were set to OFF states, the activation of MDM2 occurs through DNA Damage which in turn activates SNAIL. This further regulates enTGF β through positive feedback loop which in turn activates MDM2 through SMAD axis thus further enhancing SNAIL expression which promotes EMT (Figure S5). When both the stimulus TCF1, TCF2 were set to OFF along with exTGF β set to OFF oscillations were observed (Figure S6). These oscillations were observed due to leaky expression of functioning SNAIL (driven by RSMAD-Co-SMAD complex and MDM2) that favors the maintenance of the hybrid phenotype. Table 1 summarizes various state combinations of TCF1, TCF2 and their resulting phenotype. From this it is inferred that the presence of both the signals promotes EMT.

Similar simulations were performed with the background of exTGF β set to ON state (Figures S7 – S9). EMT was observed to occur swiftly within 10 timesteps. Thus, existence of external environmental



stress speeds up the epithelial to mesenchymal transition. When SNAIL self-repression was introduced to study the effect of SNAIL auto regulation on EMT, it was observed that the modified structure exhibited nearly similar behavior as previous cases but took longer timesteps to achieve EMT Figures S10- S17. However, there was an exception, for the case when TCF2 was turned to OFF and, TCF1 was turned ON a hybrid phenotype was observed (Figure S12). We performed this set of simulations to explore the impact of self-regulatory role on EMT. SNAIL self-regulation was experimentally explored elsewhere (Peiro *et al* 2006).

Blocking SNAIL expression completely reverses EMT to MET


To assess the effects of various regulators on the reversibility of EMT (mesenchymal to epithelial transition), the simulation conditions used to generate figure 3 were updated after 110 timesteps (assuming steady state is reached after EMT transition at this time point), and a set of systematic analysis were performed. When the stimulus for MDM2 i.e., TCF1 was turned OFF mesenchymal phenotype persists because SNAIL reaches to its maximum activity (Figure S18). Similar results were observed when TCF2 was turned OFF (Figure S19). Thus, in both scenarios mesenchymal phenotype is maintained because SNAIL activity (memory) due to the initial signal that led to EMT was still persistent. Subsequently, when both TCF1 and TCF2 were turned OFF together (Figure S20) the system reaches a hybrid phenotype. To evaluate the effect of miRNAs on the phenotype switching, miR-34 and, miR-200 were expressed by setting their node to ON state. It was observed that expression of both miRNAs (E promoting) resulted in persistent activity of SNAIL and maintenance of mesenchymal markers. Although epithelial markers were activated persistent activity of SNAIL resulted in continued suppression and absence of E-cadherin thus hybrid phenotype was

observed (Figure S25). When the ZEB expression alone was knocked down N-Cadherin is downregulated, but E-cadherin is not expressed (Figure S21). When SNAIL expression alone or in combination with ZEB was knockdown altogether, (Figure S22, S23) there was a complete transition from mesenchymal to epithelial phenotype.

SNAIL driven by feed forward loop moderates epithelial to mesenchymal transition

Presence of Coherent type 1 Feed Forward Network motif architecture among MDM2, RSMAD-CoSMAD complex (referred to as SMAD hereon) and SNAIL was a surprising inference of this work during the network assembly (figures 2, 4). The FFL between the SMAD, MDM2 and SNAIL was observed to be coherent type 1 FFL where signal from $\text{TGF}\beta$ axis (S_x) activates SMAD which directly regulates MDM2. MDM2 is also activated by signal through p53 when there is DNA Damage (S_y). SMAD (direct edge) AND/OR SMAD->MDM2 (indirect edges) regulates SNAIL in a feed forward manner (figure 4). Simulation, analysis of this C1FFL may assist to further understand the specific role of SMAD and MDM2 in regulating SNAIL transcriptional activation that modulates $\text{TGF}\beta$ induced EMT switch. Hence the focus was shifted to performing a detailed analysis of the FFL using discrete (Logical) and continuous dynamics modelling approaches.

Logical modelling simulations was performed using *python* library *Booleannet* for 100 timesteps in synchronous update mode. In the first case, AND logic scenario was studied by keeping both the input signals S_x and S_y as ON. S_x activates SMAD, SMAD in turn activates MDM2 which is also activated by S_y . Together this direct and indirect edges activate SNAIL figure 5. To understand the architecture further the input signal S_y was kept as OFF and S_x was retained to be present and set as ON. For this, it was observed that with loss of the input signal (S_y) for MDM2, MDM2 is




activated only after SMAD gets activated by S_x . SNAIL was observed to be at its ON state, only when both SMAD and MDM2 were at their ON State. Therefore, a delay of 2-time steps was observed in this case as shown in (Figure S27(a)). In the second case OR logic regulation was analyzed (Figure S28).

To understand more about the dynamics of SNAIL based on this C1FFL network motif, Ordinary differential equation (ODE) based dynamic modelling was performed using *MATLAB* solver *ODE23s*. Following activation through input signals (S_x , S_y) in AND logic dependent manner a delayed activation (of ~ 2.3 time-steps) of SNAIL expression was observed (figure 6). The delay is attributed to the requirement of the necessary thresholds k_3 (SMAD on SNAIL) and k_2 (Mdm2 on SNAIL) which are simulated in AND logic manner. Different patterns of the input signals reflecting various *in vivo* scenarios were studied (shown in Figures S30). It was observed that SNAIL reaches its maximum threshold only when both the inputs stimulus S_x and S_y were provided to the network motif. Absence of S_x fails to activate SNAIL because AND logic requirement is not fulfilled (Figure S30). However, absence of S_y leads to delayed expression of SNAIL because

MDM2 accumulated due to SMAD dependent activation (k_1) promotes SNAIL expression (Figure S30).

To further study the effect of input stimulus S_x and S_y on SNAIL expression, pulse input of S_x was provided to the system in the absence of S_y . A time pulse shorter than the delay time does not lead to SNAIL expression in the C1FFL. This is because SMAD and MDM2 do not have sufficient time to reach the threshold that is essential to activate SNAIL (figure 7). Only persistent pulse longer than the delay time was observed to activate SNAIL expression. Apart from pulse detection, this motif functions like a noise filtering system with respect to S_x . When short consecutive pulses of S_x were provided there was no activation of SNAIL, and these pulses were filtered because of the AND logic like functioning of this motif.

In the subsequent scenario, input pulses of S_x and S_y were given at different intervals of time. When the inputs were provided for shorter span of time MDM2 reaches a threshold level that is sufficient to activate SNAIL figure 8. This indicates that this motif also functions like a coincident detector because of the transient memory storage capacity of MDM2 which is sustained by immediate availability of S_y even after the




removal of S_x . To assess the OFF-step dynamics for the simulation in figure 6, the stimuli (S_x and S_y) were withdrawn. Immediately after the withdrawal of S_x and S_y SNAIL expression comes to an arrest because of the AND logic dependent activation. Similarly, the SNAIL activation of C1FFL motif was also analysed for OR logic dependent regulation and various difference input stimulus combinations were also explored (Shown in Supplementary file Figure S28, S31).

Discussion

TGF β induced regulatory processes and network modelling

The regulatory process of the cell is a smooth balance between the cell proliferation and the cell death (Lopez and Tait 2015; Shrekar and Viswanathan 2021). Numerous factors including various environmental conditions result in disturbing this balance causes cancer (Selvaggio *et al* 2020). Numerous mathematical and computational formalisms are developed to analyze these regulatory processes towards understanding the functioning of the cell (Ciliberto *et al* 2005; Kolch *et al* 2005; Zi and Klipp 2007; Albert *et al* 2008; Anderson and Quaranta 2008; Schmierer *et al*

2008; Nakabayashi and Sasaki 2009; Turner and Kohandel 2010; Idikio 2011; Cohen *et al* 2015; Rodriguez *et al* 2015; Jolly and Levine 2017; Khatibi *et al* 2017; Morshed *et al* 2018; Alon 2019; Tripathi *et al* 2020). Among which logical framework proved to be one of the successful one for understanding dynamic features of complex biological signals (Beal *et al* 2018). (Kauffmann *et al* 1973, 1969) was the first to introduce the logical modelling for analysis of generic Boolean network models using synchronous update (Kauffman 1969; Glass and Kauffman 1973). Following which this framework has been used to explore the cellular phenotype/ physiological changes that occur in various cellular processes such as cell cycle, DNA repair, tumorigenesis, cancer pathways etc (Albert and Othmer 2003; Novak and Tyson 2004; Saez-Rodriguez *et al* 2009; Schlatter *et al* 2009; Saadatpour *et al* 2010; Grieco *et al* 2013; Saadatpour and Albert 2013; Tyson and Novak 2014), (Ribba *et al* 2006; Steinway *et al* 2014; Remy *et al* 2015; Steinway *et al* 2015; Rossato *et al* 2019). Such predictive parameter free models are useful to understand the state transitions and their biological implications (Beal *et al* 2018; Eduati *et al* 2020; Beal *et al* 2021; Shrekar and Viswanathan 2021).



The network assembled (figure 2) comprises of tumour suppressor genes (p53, $TGF\beta$), tumour promoting genes & transcription factors ($TGF\beta$, SNAIL, MDM2, ZEB) and were commonly observed in numerous cancer types including Breast, Ovarian, Hepatocellular, Colorectal, Lung Carcinoma (Bolós *et al* 2016; Lu *et al* 2016; Osorio *et al* 2016; Saitoh *et al* 2016; Yang *et al* 2017; Cai *et al* 2019; Jiang *et al* 2020; Lin *et al* 2020; Tripathi *et al* 2020). Presence of a feed forward network architecture was a surprise observation between the upstream $TGF\beta$ activated SMAD complex, MDM2 for the regulation of SNAIL transcriptional activation. Thus, logical modelling framework adopted for this study has demonstrated the EMT transition that promotes metastasis and the role of individual regulators guiding this process.

Implications of logical modelling and analysis on $TGF\beta$ induced phenotypic switching

Among the hallmarks of cancer (Hanahan and Weinberg 2000, 2011) EMT is one of the widely explored hallmark of the tumorigenesis. Steinway *et al* (2014) published a $TGF\beta$ induced EMT model where the concomitant activation of WNT and SHH signaling was necessary for the $TGF\beta$ dependent EMT in Hepatocellular Carcinoma (HCC) (Steinway *et al*

2014). Single and combinatorial perturbation analysis were performed on the logical model developed to observe their effect on $TGF\beta$ induced EMT. Single perturbations resulted in a hybrid phenotype and multiple interventions were observed to result in an epithelial phenotype (Steinway *et al* 2015). Analogously, combinatorial perturbations performed both theoretically and experimentally by Watanabe *et al* (2019) by manipulating the essential genes $TGF\beta$ and ZEB1 revealed the key functional role of ZEB1 in reciprocity and reversibility of EMT (Watanabe *et al* 2019). Study by rated the importance of microenvironments in controlling the cellular plasticity along EMT and suggests hybrid and mesenchymal phenotype occur via different paths (Selvaggio *et al* 2020). Several other perturbation analysis were also performed by various groups by building logical models to analyze the stability of the cellular phenotype during the transition in different cancer types (Cohen *et al* 2015; Mendez-Lopez *et al* 2017; Selvaggio *et al* 2020). Further logical models were also used to analyze the drug effects and therapeutic strategies (Flobak *et al* 2015; Traynard *et al* 2017; Zanudo *et al* 2018). Thus, logical modelling and simple perturbation analysis can be used to understand the molecular mechanism influencing various key physiological processes such

as EMT in terms of their signaling regulatory components.

The functional principles of EMT/MET was previously described by several mathematical models that revolves around the core regulatory circuit miR-34/SNAIL/miR-200/ZEB in terms of cascade switches and bistable switches (Lu *et al* 2013; Tian *et al* 2013; Jolly and Jia *et al* 2016; Jolly Tripathi *et al* 2016; Jia *et al* 2017; Jolly and Levine 2017; Tripathi *et al* 2020; Tripathi *et al* 2021). Although the response of phenotypic switching is not an all or none response, it is rather a collective response that depends on several input signals and the environment conditions (Gupta *et al* 2011; Huang *et al* 2014; Jolly *et al* 2014; Jolly *et al* 2015; Boareto *et al* 2016; Li and Balazsi 2018).

The phenotypic shift from epithelial to mesenchymal states was observed to be a stepwise process during which existence of hybrid EM state was investigated for its crucial role in tumour progression and organ fibrosis (Li *et al* 2016). Hybrid EMT was noticed to allow collective migration of tumour cells evading immune attack with higher migrating potential (Vinay *et al* 2015; Saxena *et al* 2020). Our model predicts that introducing the negative self-regulation of SNAIL, results in hybrid phenotypes with longer time span before mesenchymal transition. This opens a window for further analysis on hybrid EM phenotypes and its impact on collective migration of cancer cells. Literature shows that knockdown of ZEB (at low levels) and not SNAIL alone (Gregory *et al* 2011) or upregulation of miRNAs miR-34a or miR-200 (thus causing complete downregulation of SNAIL)(Migliore *et al* 2008; Migliore and Giordano 2009; Bader 2012; Wang *et al* 2013; Okada *et al* 2014) reverts the cell to the epithelial phenotype. Our results show that either of them do not revert the epithelial phenotype but down-regulation or suppression of SNAIL expression all together can revert the cell to epithelial phenotype. Previous studies have demonstrated the effect of various epigenetic factors / phenotype stability factors, the interconnectedness between these factors with core regulatory circuit and their effect on the stability/resistance on EMT (Padua *et al* 2008; Cieply *et al* 2012; Huang *et al* 2015; Jolly and Tripathi *et al* 2016; Jolly *et al* 2018; Bocci *et al* 2019; Jia *et al* 2019; Jia *et al* 2020; Silveira and Mombach 2020; Silveira *et al* 2020; Subbalakshmi *et al* 2020). The present work investigated the signaling network level regulation, a layer above this core regulatory circuit, which includes an additional p53/MDM2 loop that affects the same phenomenon of EMT. The present work has identified this layer involving SMAD complex, MDM2 and SNAIL to be regulated through a coherent type 1 FFL motif architecture (figure 4). p53 has been well explored for its role as a tumour suppressor, driver gene and p53/mdm2 loop in maintaining the oscillations in the presence of DNA damage (Lev Bar-Or *et al* 2000; Ciliberto *et al* 2005; Cohen *et al* 2015). However, the role of p53/MDM2 in EMT is recently demonstrated

experimentally (Wang *et al* 2009; Lim *et al* 2010; Kim *et al* 2011; 2011; Ren *et al* 2013; Wang *et al* 2013; Chen *et al* 2017; Tang *et al* 2019). This is also an attempt to analyse the effect of p53/MDM2 on the core regulatory circuit and phenotypic plasticity. MDM2 is observed to be a part of coherent type 1 FFL motif architecture above the core EMT regulatory circuit.

Our model and model analysis illustrates that for the complete transition from epithelial to mesenchymal phenotype SNAIL requires to be active which occurred within fewer time steps (10 steps). SNAIL activity is required for regulation of its downstream transcription factors. SNAIL attains maximum activity swiftly when there is external environment stress and when both TCF1&TCF2 are set to ON. Absence of any of the input signals or Presence of negative auto regulation on SNAIL results in longer durations of hybrid phenotype before attaining EMT thus providing the essential time for the system for preparations and final decision making. Knockdown of transcription cofactors alone for both SNAIL and MDM2 or over-expression of miRNAs do not reverse EMT but results in a hybrid phenotype. Whereas complete knockdown of SNAIL expression completely causes a switch from mesenchymal to epithelial phenotype. Therefore, the focus was further shifted to understand the significance of SNAIL regulated by both SMAD complex and MDM2 in a feed forward manner.

Coherent feed forward loop network architecture and guardian role of SNAIL

SNAIL was observed to be one of the prominent molecules affecting the phenotypic plasticity that favours the changes in cell shape (Vega *et al* 2004). Studies by Peiro *et al* (2006) demonstrated the self-regulatory role SNAIL1 family (Peiro *et al* 2006). SNAIL along with SMAD2/3 complex acts as a transcriptional repressor for epithelial markers (Vincent *et al* 2009;). It was also observed that elevated expression of SNAIL forces the cells to attain mesenchymal traits by repressing p53 and p53 induced miRNAs (Siemens *et al* 2011). The role of SNAIL as a noise buffering integrator and its ability to control the fate of hybrid E/M phenotype was demonstrated. (Jolly *et al* 2014; Lu *et al* 2013). Our results were observed to corroborate along with these roles discussed. SNAIL functions like noise filtering system in the C1FFL motif structure analysed and it propagates the filtered signal to the downstream core EMT regulators. SNAIL is known to drive cell proliferation in many cancer types including hepatocellular, ovarian, breast, colorectal, lung cancer and tumour progression. We postulate that perhaps the process of tumorigenesis and EMT transformation may involve the C1FFL dependent regulation characterised in this study. This proposed hypothesis can be verified *in vivo* in cancer cell lines by single cell level microscopic studies and through cell culture studies. Further,

'Build it to understand' synthetic biology strategies can be utilised to further refine the C1FFL dependent molecular regulation that lies above the core EMT regulatory switch. Such synthetic biology study may possibly provide a diagnostic tool for early assessment of E-M transitions.

FFLs involving transcription factors and their various subtypes have been analysed in the past (Schleif 2000; Shen-Orr *et al* 2002; Mangan and Alon 2003; Kalir *et al* 2005; Alon 2007; Iyengar *et al* 2017; Zhang *et al* 2018; Alon 2019; Guo and Murray 2019). It has been demonstrated that specific network motifs such as auto-negative regulation or C1FFL can filter or dampen external noise associated fluctuations (Ghosh *et al* 2005; Osella *et al* 2011; Alon 2019). Gill *et al* (2018) experimentally characterise a positive FFL that enforces YAP/TAZ engaging hippo pathway in promoting tumour (Gill *et al* 2018). Reeves *et al* (2009) studied the engineering principles of combining the FFL with a feedback loop (Reeves 2019). Work by Tej *et al* (2019) suggests that the function of small RNA FFL is critically governed by the small RNA not only in generating a speedy and strong response but also in producing a reliable response by regulating the gene expression noise (Tej *et al* 2019). Dynamic analysis of multi-input FFL by Zheng *et al* (2009) in *C. elegans*, illustrates both persistence and coincidence detection among the neuronal cells (Zheng 2009). Similarly, our analysis of the FFL in figure 4 also has demonstrated both persistence and coincidence detection which we postulate may have a crucial role in EM transition. Our study suggests that the FFL governed by two TFs causes a delay in ON(AND) and OFF step (OR) by regulating SNAIL expression. We anticipate that SNAIL functions like a noise filter reminiscent with Lu *et al* (Lu *et al* 2013). Thus, the proposed C1FFL structure through SNAIL guards the downstream miRNA network from getting activated for low range of signals.

In current framework, when considered C1FFL alone, SNAIL activity depends on the input stimulus that comes through the upstream molecules (SMAD complex, Mdm2). However, if the whole regulatory circuit is considered, the C1FFL network architecture functions like a backbone for SNAIL activation and it does not function in isolation. These varying input signals also modulate the p53 dependent regulation of EMT in presence of dysregulation such as DNA Damage, and other environmental conditions. We postulate that this signaling follows a layered manner of activation, driving the activity of elements in the EMT network. Together the logical, ODE modelling and analysis of C1 FFL revealed that both the inputs (S_x , S_y) play a significant role in activating the SMAD and MDM2 which in turn promotes SNAIL which is essential for the downstream signaling to regulate EMT switch. SNAIL do not promote the downstream signaling if there is failure to detect input signals S_x thus playing a very prominent guidance role in this

signaling cascade. We also observe that pulsatile input (analogous to noisy signal in terms of S_x) is filtered and EM transition will not be promoted by SNAIL (figure 7). However, simultaneous presence of pulsatile inputs (S_x AND S_y), is detected coincidentally (figure 8) implying the occurrence of EM transition promoted by SNAIL. Thus, SNAIL function as a decision-making element for the downstream EMT signaling regulators.

Conclusion

Diverse range of signaling molecules and pathways were observed to be involved in promoting epithelial to mesenchymal transition. Among which $TGF\beta$ and its family of signaling molecules play a prominent role. This study attempted to model and analyse the network that couples the upstream $TGF\beta$ signal level regulation of EMT which includes SMADs, SMAD-CoSMAD complexes, p53 and Mdm2.

Logical modelling and analysis revealed that for the transition from epithelial to mesenchymal phenotype to occur completely, SNAIL requires signal from both SMAD complex and MDM2. Blockage of any of the two signals causes the system to stay in hybrid EM phenotype for longer periods of time. Presence of exogenous $TGF\beta$ accelerates EMT, slimming the chances of hybrid phenotype. Negative autoregulation of SNAIL results in prolonged EM hybrid phenotype whereas complete knockdown results in MET. One of the important observations is the presence of C1FFL motif like regulation of SNAIL involving SMAD complex and MDM2. Dynamic ODE modelling and analysis of the C1FFL displayed a sign sensitive delay and persistence detection of the stimulus S_x . This motif also functioned like a coincidence detector when both stimulus (S_x and S_y) were present. These inferences led us to conclude that this signaling architecture enables SNAIL to filter the noise in terms of its input signals (S_x) to function like a 'Guardian' for the downstream networks performing the phenotypic switching. However, presence of multiple signals (S_x AND S_y), resulted in EM transition promoted by SNAIL expression. Together results from logical and dynamic modelling of the network suggests that SNAIL not only acts as a signal integrator but also plays a significant role in decision making during the phenotypic switching of EMT. We conclude that SNAIL is at the heart of the $TGF\beta$ dependent regulation of EMT bridging the C1FFL to the most frequently explored miRNA dependent core regulatory circuit of EMT. Therefore, SNAIL may be targeted for the drugs development and therapeutic interventions.

Acknowledgments

Our research is supported by grants and scholarship provided by NITW. AP is supported by NITW, RSM

(P1068) grant and GSB is supported by NITW institute scholarship. Both the authors acknowledge the facilities and support extended by NIT Warangal. We thank the anonymous reviewers for their careful reading and helpful suggestions on the earlier drafts of the manuscript.

Data availability statement


All data that support the findings of this study are included within the article (and any supplementary files).

Declarations:

The authors declare no conflict of interest. The authors declare no competing financial interests.

ORCID iDs

Gottumukkala Sai Bhavani  <https://orcid.org/0000-0002-2870-6385>

Anbumathi Palanisamy  <https://orcid.org/0000-0002-9076-5271>

References

- Albert I, Thakar J, Li S, Zhang R and Albert R 2008 Boolean network simulations for life scientists *Source Code Biol. Med.* **3** 16
- Albert R and Othmer H G 2003 The topology of the regulatory interactions predicts the expression pattern of the segment polarity genes in *Drosophila melanogaster* *J. Theor. Biol.* **223** 1–18
- Alon U 2007 Network motifs: theory and experimental approaches *Nat. Rev. Genet.* **8** 450–61
- Alon U 2019 *An introduction to systems biology: design principles of biological circuits*. 2nd edn (Boca Raton, London New York: CRC Press, Taylor & Francis Group) <https://www.routledge.com/An-Introduction-to-Systems-Biology-Design-Principles-of-Biological-Circuits/Alon/p/book/9781439837177> (<https://doi.org/10.1201/9780429283321>)
- Andersson A R and Quaranta V 2008 Integrative mathematical oncology *Nat. Rev. Cancer* **8** 227–34
- Araki S, Eitel J A, Batuello C N, Bijangi-Vishehsaraei K, Xie X J, Danielpour D, Pollok K E, Boothman D A and Mayo L D 2010 TGF-beta1-induced expression of human Mdm2 correlates with late-stage metastatic breast cancer *J. Clin. Invest.* **120** 290–302
- Vander Ark A, Cao J and Li X 2018 TGF-beta receptors: In and beyond TGF-beta signaling *Cell Signal* **52** 112–20
- Attisano L and Wrana J L 1996 Signal transduction by members of the transforming growth factor-beta superfamily *Cytokine Growth Factor Rev.* **7** 327–39
- Attisano L and Wrana J L 1998 Mads and Smads in TGF beta signalling *Curr. Opin. Cell Biol.* **10** 188–94
- Attisano L and Wrana J L 2000 Smads as transcriptional co-modulators *Curr. Opin. Cell Biol.* **12** 235–43
- Attisano L and Wrana J L 2002 Signal transduction by the TGF-beta superfamily *Science* **296** 1646–7
- Bader A G 2012 miR-34 - a microRNA replacement therapy is headed to the clinic *Front. genet* **3** 120
- Beal J, Montagud A, Traynard P, Barillot E and Calzone L 2018 Personalization of logical models with multi-omics data allows clinical stratification of patients *Front. Physiol* **9** 1965
- Beal J, Pantolini L, Noel V, Barillot E and Calzone L 2021 Personalized logical models to investigate cancer response to BRAF treatments in melanomas and colorectal cancers *PLoS Comput. Biol.* **17** e1007900
- Bedi A, Chang X, Noonan K, Pham V, Bedi R and Ravi R 2012 Inhibition of TGF-beta enhances the *in vivo* antitumor efficacy of EGF receptor-targeted therapy *Mol. Cancer. Ther.* **11** 2429–39
- Bierie B and Moses H L 2006 Tumour microenvironment: TGFbeta: the molecular Jekyll and Hyde of cancer *Nat. Rev. Cancer* **6** 506–20
- Boareto M, Jolly M K, Goldman A, Pietila M, Mani S A, Sengupta S, Ben-Jacob E, Levine H and Onuchic J N 2016 Notch-Jagged signalling can give rise to clusters of cells exhibiting a hybrid epithelial/mesenchymal phenotype *J. R. Soc. Interface* **13** 20151106
- Bocci F, Tripathi S C, Vilchez Mercedes S A, George J T, Casabar J P and Jolly M K 2019 NRF2 activates a partial epithelial-mesenchymal transition and is maximally present in a hybrid epithelial/mesenchymal phenotype *Integr. Biol. (Camb)* **11** 251–63
- Bolós V, Peinado H, Pérez-Moreno M A, Fraga M F, Esteller M and Cano A 2016 The transcription factor Slug represses E-cadherin expression and induces epithelial to mesenchymal transitions: a comparison with Snail and E47 repressors *J. Cell Sci.* **129** 1283–1283
- Bracken C P, Gregory P A, Kolesnikoff N, Bert A G, Wang J, Shannon M F and Goodall G J 2008 A double-negative feedback loop between ZEB1-SIP1 and the microRNA-200 family regulates epithelial-mesenchymal transition *Cancer Res.* **68** 7846–54
- Bryant J L, Britson J, Balko J M, Willian M, Timmons R, Frolov A and Black E P 2012 A microRNA gene expression signature predicts response to erlotinib in epithelial cancer cell lines and targets EMT *Br. J. Cancer* **106** 148–56
- Burk U, Schubert J, Wellner U, Schmalhofer O, Vincan E, Spaderna S and Brabletz T 2008 A reciprocal repression between ZEB1 and members of the miR-200 family promotes EMT and invasion in cancer cells *EMBO. Rep* **9** 582–9
- Cai J, Xia L, Li J, Ni S, Song H and Wu X 2019 Tumor-associated macrophages derived TGF-beta-induced epithelial to mesenchymal transition in colorectal cancer cells through smad2,3-4/snail signaling pathway *Cancer. Res. Treat* **51** 252–66
- Ceppi P, Mudduluru G, Kumarswamy R, Rapa I, Scagliotti G V, Papotti M and Allgayer H 2010 Loss of miR-200c expression induces an aggressive, invasive, and chemoresistant phenotype in non-small cell lung cancer *Mol. Cancer. Res* **8** 1207–16
- Chang C J, Chao C H, Xia W, Yang J Y, Xiong Y and Hung M C 2011 p53 regulates epithelial-mesenchymal transition and stem cell properties through modulating miRNAs *Nat. Cell Biol.* **13** 317–23
- Chatain T, Haar S and Paulevé L 2018 Boolean networks: beyond generalized asynchronicity *Int. Workshop on Cellular Automata and Discrete Complex Systems* (https://doi.org/10.1007/978-3-319-92675-9_3)
- Chen Y, Wang D D, Wu Y P, Su D, Zhou T Y and Yang B 2017 MDM2 promotes epithelial-mesenchymal transition and metastasis of ovarian cancer SKOV3 cells *Br. J. Cancer* **117** 1192–201
- Cieply B, Riley P T, Pifer P M, Widmeyer J, Addison J B, Ivanov A V, Denvir J and Frisch S M 2012 Suppression of the epithelial-mesenchymal transition by Grainyhead-like-2 *Cancer Res.* **72** 2440–53
- Ciliberto A, Novak B and Tyson J J 2005 Steady states and oscillations in the p53/Mdm2 network *Cell Cycle* **4** 488–93
- Cohen D P, Martignetti L, Robine S, Barillot E, Zinoviyev A and Calzone L 2015 Mathematical modelling of molecular pathways enabling tumour cell invasion and migration *PLoS Comput. Biol.* **11** e1004571

- Coradini D, Fornili M, Ambrogio F, Boracchi P and Biganzoli E 2012 TP53 mutation, epithelial-mesenchymal transition, and stemlike features in breast cancer subtypes *J. Biomed. Biotechnol.* **2012** 254085
- De Craene B and Berrux G 2013 Regulatory networks defining EMT during cancer initiation and progression *Nat. Rev. Cancer* **13** 97–110
- Derynck R and Zhang Y E 2003 Smad-dependent and Smad-independent pathways in TGF- β family signalling *Nature* **425** 577–84
- Dickson R B, Kasid A, Huff K K, Bates S E, Knabbe C, Bronzert D, Gelmann E P and Lippman M E 1987 Activation of growth factor secretion in tumorigenic states of breast cancer induced by 17 β -estradiol or v-Ha-ras oncogene *Proc. Natl. Acad. Sci. USA* **84** 837–41
- Eduati F, Jaaks P, Wappler J, Cramer T, Merten C A, Garnett M J and Saez-Rodriguez J 2020 Patient-specific logic models of signaling pathways from screenings on cancer biopsies to prioritize personalized combination therapies *Mol. Syst. Biol.* **16** e8664
- Fan F, Samuel S, Evans K W, Lu J, Xia L and Ellis L M 2012 Overexpression of snail induces epithelial-mesenchymal transition and a cancer stem cell-like phenotype in human colorectal cancer cells *Cancer. Med.* **1** 5–16
- Flobak A, Baudot A, Remy E, Thommesen L, Thieffry D, Kuiper M and Laegreid A 2015 Discovery of drug synergies in gastric cancer cells predicted by logical modeling *PLoS Comput. Biol.* **11** e1004426
- Franco D L, Mainez J, Vega S, Sancho P, Murillo M M and Nieto M A 2010 Snail suppresses TGF- β -induced apoptosis and is sufficient to trigger EMT in hepatocytes *J. Cell Sci.* **123** 3467–77
- Garg A, Di Cara A, Xenarios I, Mendoza L and De Micheli G 2008 Synchronous versus asynchronous modeling of gene regulatory networks *Bioinformatics* **24** 1917–25
- Ghosh B, Karmakar R and Bose I 2005 Noise characteristics of feed forward loops *Phys. Biol.* **2** 36–45
- Giannelli G, Villa E and Lahn M 2014 Transforming growth factor- β as a therapeutic target in hepatocellular carcinoma *Cancer Res.* **74** 1890–4
- Gill M K, Christova T, Zhang Y Y, Gregorieff A, Zhang L and Attisano L 2018 A feed forward loop enforces YAP/TAZ signaling during tumorigenesis *Nat. Commun.* **9** 3510
- Glass L and Kauffman S A 1973 The logical analysis of continuous, non-linear biochemical control networks *J. Theor. Biol.* **39** 103–29
- Gregory P A, Bert A G, Paterson E L, Barry S C, Tsykin A, Farshid G, Vadas M A, Khew-Goodall Y and Goodall G J 2008 The miR-200 family and miR-205 regulate epithelial to mesenchymal transition by targeting ZEB1 and SIP1 *Nat. Cell Biol.* **10** 593–601
- Gregory P A, Bracken C P, Smith E, Bert A G, Wright J A and Goodall G J 2011 An autocrine TGF- β /ZEB/miR-200 signaling network regulates establishment and maintenance of epithelial-mesenchymal transition *Mol. Biol. Cell* **22** 1686–98
- Grieco L, Calzone L, Bernard-Pierrot I, Radvanyi F, Kahn-Perles B and Thieffry D 2013 Integrative modelling of the influence of MAPK network on cancer cell fate decision *PLoS Comput. Biol.* **9** e1003286
- Guo S and Murray R M 2019 Construction of incoherent feedforward loop circuits in a cell-free system and in cells *ACS. Synth. Biol.* **8** 606–10
- Gupta P B, Fillmore C M, Jiang G, Shapira S D, Tao K, Kuperwasser C and Lander E S 2011 Stochastic state transitions give rise to phenotypic equilibrium in populations of cancer cells *Cell* **146** 633–44
- Hanahan D and Weinberg R A 2000 The hallmarks of cancer *Cell* **100** 57–70
- Hanahan D and Weinberg R A 2011 Hallmarks of cancer: the next generation *Cell* **144** 646–74
- Haslehurst A M, Koti M, Dharsee M, Nuin P, Evans K and Feilottter H 2012 EMT transcription factors snail and slug directly contribute to cisplatin resistance in ovarian cancer *BMC Cancer* **12** 91
- Heldin C H, Miyazono K and ten Dijke P 1997 TGF- β signalling from cell membrane to nucleus through SMAD proteins *Nature* **390** 465–71
- Hemavathy K, Ashraf S I and Ip Y T 2000 Snail/Slug family of repressors: slowly going into the fast lane of development and cancer *Gene* **257** 1–12
- Hsu D S, Lan H Y, Huang C H, Tai S K, Chang S Y and Yang M H 2010 Regulation of excision repair cross-complementation group 1 by Snail contributes to cisplatin resistance in head and neck cancer *Clin. Cancer Res.* **16** 4561–71
- Huang B, Jolly M K, Lu M, Tsarfaty I, Ben-Jacob E and Onuchic J N 2015 Modeling the transitions between collective and solitary migration phenotypes in cancer metastasis *Sci. Rep.* **5** 17379
- Huang B, Lu M, Jolly M K, Tsarfaty I, Onuchic J and Ben-Jacob E 2014 The three-way switch operation of Rac1/RhoA GTPase-based circuit controlling amoeboid-hybrid-mesenchymal transition *Sci Rep.* **4** 6449
- Hunziker A, Jensen M H and Krishna S 2010 Stress-specific response of the p53-Mdm2 feedback loop *BMC Syst. Biol.* **4** 94
- Idikio H A 2011 Human cancer classification: a systems biology-based model integrating morphology, cancer stem cells, proteomics, and genomics *J. Cancer* **2** 107–15
- Ikushima H and Miyazono K 2010 TGF β signalling: a complex web in cancer progression *Nat. Rev. Cancer* **10** 415–24
- Iyengar B R, Pillai B, Venkatesh K V and Gadgil C J 2017 Systematic comparison of the response properties of protein and RNA mediated gene regulatory motifs *Mol. Biosyst.* **13** 1235–45
- Jia D, Jolly M K, Tripathi S C, Den Hollander P, Huang B and Levine H 2017 Distinguishing mechanisms underlying EMT tristability *Cancer Converge* **1** 2
- Jia W, Deshmukh A, Mani S A, Jolly M K and Levine H 2019 A possible role for epigenetic feedback regulation in the dynamics of the epithelial-mesenchymal transition (EMT) *Phys. Biol.* **16** 066004
- Jia W, Tripathi S, Chakraborty P, Chedere A, Rangarajan A, Levine H and Jolly M K 2020 Epigenetic feedback and stochastic partitioning during cell division can drive resistance to EMT *Oncotarget* **11** 2611–24
- Jiang K, Zhao T, Shen M, Zhang F, Duan S, Lei Z and Chen Y 2020 Correction: MiR-940 inhibits TGF- β -induced epithelial-mesenchymal transition and cell invasion by targeting Snail in non-small cell lung cancer *J. Cancer* **11** 4897–8
- Johnsen S A, Subramaniam M, Janknecht R and Spelsberg T C 2002 TGF β inducible early gene enhances TGF β /Smad-dependent transcriptional responses *Oncogene* **21** 5783–90
- Jolly M K, Huang B, Lu M, Mani S A, Levine H and Ben-Jacob E 2014 Towards elucidating the connection between epithelial-mesenchymal transitions and stemness *J. R. Soc. Interface* **11** 20140962
- Jolly M K, Jia D, Boareto M, Mani S A, Pienta K J, Ben-Jacob E and Levine H 2015 Coupling the modules of EMT and stemness: A tunable ‘stemness window’ model *Oncotarget* **6** 25161–74
- Jolly M K, Jia D, Tripathi S, Hanash S, Mani S, Ben-Jacob E and Levine H 2016 Network motifs that stabilize the hybrid epithelial/mesenchymal phenotype *APS March Meeting Abstracts* **2016** 006–B039. <https://meetings.aps.org/Meeting/MAR16/Session/B39.6>
- Jolly M K and Levine H 2017 Computational systems biology of epithelial-hybrid-mesenchymal transitions *Current Opinion in Systems Biology* **3** 1–6
- Jolly M K, Preca B T, Tripathi S C, Jia D, George J T and Levine H 2018 Interconnected feedback loops among ESRP1, HAS2, and CD44 regulate epithelial-mesenchymal plasticity in cancer *APL Bioeng.* **2** 031908
- Jolly M K, Tripathi S C, Jia D, Mooney S M, Celiktas M and Levine H 2016 Stability of the hybrid epithelial/mesenchymal phenotype *Oncotarget* **7** 27067–84
- Kalir S, Mangan S and Alon U 2005 A coherent feed-forward loop with a SUM input function prolongs flagella expression in *Escherichia coli* *Mol. Syst. Biol.* **1** 2005

- Kalluri R and Weinberg R A 2009 The basics of epithelial-mesenchymal transition *J. Clin. Invest.* **119** 1420–8
- Kang Y and Massague J 2004 Epithelial-mesenchymal transitions: twist in development and metastasis *Cell* **118** 277–9
- Kauffman S A 1969 Metabolic stability and epigenesis in randomly constructed genetic nets *J. Theor. Biol.* **22** 437–67
- Khatibi S, Zhu H J, Wagner J, Tan C W, Manton J H and Burgess A W 2017 Mathematical model of TGF- β signalling: feedback coupling is consistent with signal switching *BMC Syst. Biol.* **11** 48
- Kim E, Kim J Y and Lee J Y 2019 Mathematical modeling of p53 pathways *Int. J. Mol. Sci.* **20** 5179
- Kim N H, Kim H S, Li X Y, Lee I, Choi H S and Yook J I 2011 A p53/miRNA-34 axis regulates Snail1-dependent cancer cell epithelial-mesenchymal transition *J. Cell Biol.* **195** 417–33
- Kim T, Veronese A, Pichiorri F, Lee T J, Jeon Y J and Croce C M 2011 p53 regulates epithelial-mesenchymal transition through microRNAs targeting ZEB1 and ZEB2 *J. Exp. Med.* **208** 875–83
- Kolch W, Calder M and Gilbert D 2005 When kinases meet mathematics: the systems biology of MAPK signalling *FEBS Lett.* **579** 1891–5
- Korpal M, Lee E S, Hu G and Kang Y 2008 The miR-200 family inhibits epithelial-mesenchymal transition and cancer cell migration by direct targeting of E-cadherin transcriptional repressors ZEB1 and ZEB2 *J. Biol. Chem.* **283** 14910–4
- Lambert A W, Pattabiraman D R and Weinberg R A 2017 Emerging biological principles of metastasis *Cell* **168** 670–91
- Lamouille S, Connolly E, Smyth J W, Akhurst R J and Derynck R 2012 TGF- β -induced activation of mTOR complex 2 drives epithelial-mesenchymal transition and cell invasion *J. Cell Sci.* **125** 1259–73
- Lamouille S, Subramanyam D, Belloch R and Derynck R 2013 Regulation of epithelial-mesenchymal and mesenchymal-epithelial transitions by microRNAs *Curr. Opin. Cell Biol.* **25** 200–7
- Lamouille S, Xu J and Derynck R 2014 Molecular mechanisms of epithelial-mesenchymal transition *Nat. Rev. Mol. Cell Biol.* **15** 178–96
- Lev Bar-Or R, Maya R, Segel L A, Alon U, Levine A J and Oren M 2000 Generation of oscillations by the p53-Mdm2 feedback loop: a theoretical and experimental study *Proc. Natl. Acad. Sci. USA* **97** 11250–5
- Li C and Balazsi G 2018 A landscape view on the interplay between EMT and cancer metastasis *NPJ. Syst. Biol. Appl.* **4** 34
- Li C, Hong T and Nie Q 2016 Quantifying the landscape and kinetic paths for epithelial-mesenchymal transition from a core circuit *Phys. Chem. Chem. Phys.* **18** 17949–56
- Li J and Zhou B P 2011 Activation of beta-catenin and Akt pathways by Twist are critical for the maintenance of EMT associated cancer stem cell-like characters *BMC. Cancer* **11** 49
- Li W, Liu C, Tang Y, Li H, Zhou F and Lv S 2011 Overexpression of snail accelerates adriamycin induction of multidrug resistance in breast cancer cells *Asian. Pac. J. Cancer. Prev* **12** 2575–80 <https://pubmed.ncbi.nlm.nih.gov/22320957/>
- Lim S O, Kim H and Jung G 2010 p53 inhibits tumor cell invasion via the degradation of snail protein in hepatocellular carcinoma *FEBS Lett.* **584** 2231–6
- Lin Q, Zhou C R, Bai M J, Zhu D, Chen J W and Huang M S 2020 Exosome-mediated miRNA delivery promotes liver cancer EMT and metastasis *Am. J. Transl. Res* **12** 1080–95 <https://pubmed.ncbi.nlm.nih.gov/32269736/>
- Liu Y N, Yin J J, Abou-Kheir W, Hynes P G, Casey O M and Kelly K 2013 MiR-1 and miR-200 inhibit EMT via Slug-dependent and tumorigenesis via Slug-independent mechanisms *Oncogene* **32** 296–306
- Lopez J and Tait S W 2015 Mitochondrial apoptosis: killing cancer using the enemy within *Br. J. Cancer* **112** 957–62
- Lu M, Jolly M K, Levine H, Onuchic J N and Ben-Jacob E 2013 MicroRNA-based regulation of epithelial-hybrid-mesenchymal fate determination *Proc. Natl. Acad. Sci. USA* **110** 18144–9
- Lu X, Yan C, Huang Y, Shi D, Fu Z, Qiu J and Yin Y 2016 Mouse double minute 2 (MDM2) upregulates Snail expression and induces epithelial-to-mesenchymal transition in breast cancer cells *in vitro* and *in vivo* *Oncotarget* **7** 37177–91
- Malfettone A, Soukupova J, Bertran E, Crosas-Molist E, Lastra R and Fabregat I 2017 Transforming growth factor- β -induced plasticity causes a migratory stemness phenotype in hepatocellular carcinoma *Cancer. Lett* **392** 39–50
- Mangan S and Alon U 2003 Structure and function of the feed-forward loop network motif *Proc. Natl. Acad. Sci. USA* **100** 11980–5
- Mariano G, Ricciardi M R, Trisciuglio D, Zampieri M, Ciccarone F and Reale A 2015 PARP inhibitor ABT-888 affects response of MDA-MB-231 cells to doxorubicin treatment, targeting Snail expression *Oncotarget* **6** 15008–21
- Massague J 1998 TGF- β signal transduction *Annu. Rev. Biochem.* **67** 753–91
- Massague J 2012 TGF β signalling in context *Nat. Rev. Mol. Cell Biol.* **13** 616–30
- Massague J, Seoane J and Wotton D 2005 Smad transcription factors *Genes. Dev* **19** 2783–810
- Mendez-Lopez L F, Davila-Velderrain J, Dominguez-Huttinger E, Enriquez-Olguin C, Martinez-Garcia J C and Alvarez-Buylla E R 2017 Gene regulatory network underlying the immortalization of epithelial cells *BMC Syst. Biol.* **11** 24
- Miettinen P J, Ebner R, Lopez A R and Derynck R 1994 TGF- β induced transdifferentiation of mammary epithelial cells to mesenchymal cells: involvement of type I receptors *J. Cell Biol.* **127** 2021–36
- Migliore C and Giordano S 2009 MiRNAs as new master players *Cell Cycle* **8** 2185–6
- Migliore C, Petrelli A, Ghiso E, Corso S, Capparuccia L, Eramo A, Comoglio P M and Giordano S 2008 MicroRNAs impair MET-mediated invasive growth *Cancer Res.* **68** 10128–36
- Miyazono K 2000 Positive and negative regulation of TGF- β signaling *J. Cell Sci.* **113** (Pt 7) 1101–9
- Morshed A, Dutta P and Dillon R H 2018 Mathematical modeling and numerical simulation of the TGF- β /Smad signaling pathway in tumor microenvironments *Appl. Numer. Math.* **133** 41–51
- Moustakas A and de Herreros A G 2017 Epithelial-mesenchymal transition in cancer *Mol. Oncol* **11** 715–7
- Nakabayashi J and Sasaki A 2009 A mathematical model of the stoichiometric control of Smad complex formation in TGF- β signal transduction pathway *J. Theor. Biol.* **259** 389–403
- Nieto M A 2002 The snail superfamily of zinc-finger transcription factors *Nat. Rev. Mol. Cell Biol.* **3** 155–66
- Nieto M A, Huang R Y, Jackson R A and Thiery J P 2016 EMT: 2016 *Cell* **166** 21–45
- Novak B and Tyson J J 2004 A model for restriction point control of the mammalian cell cycle *J. Theor. Biol.* **230** 563–79
- Okada N, Lin C P, Ribeiro M C, Biton A, Lai G and He L 2014 A positive feedback between p53 and miR-34 miRNAs mediates tumor suppression *Genes. Dev* **28** 438–50
- Onder T T, Gupta P B, Mani S A, Yang J, Lander E S and Weinberg R A 2008 Loss of E-cadherin promotes metastasis via multiple downstream transcriptional pathways *Cancer Res.* **68** 3645–54
- Osella M, Bosia C, Cora D and Caselle M 2011 The role of incoherent microRNA-mediated feedforward loops in noise buffering *PLoS Comput. Biol.* **7** e1001101
- Osorio L A, Farfan N M, Castellon E A and Contreras H R 2016 SNAIL transcription factor increases the motility and invasive capacity of prostate cancer cells *Mol. Med. Rep.* **13** 778–86
- Padua D, Zhang X H, Wang Q, Nadal C, Gerald W L, Gomis R R and Massague J 2008 TGF β primes breast tumors for lung metastasis seeding through angiopoietin-like 4 *Cell* **133** 66–77
- Park S M, Gaur A B, Lengyel E and Peter M E 2008 The miR-200 family determines the epithelial phenotype of cancer cells by targeting the E-cadherin repressors ZEB1 and ZEB2 *Genes. Dev* **22** 894–907

- Paterson E L, Kolesnikoff N, Gregory P A, Bert A G, Khew-Goodall Y and Goodall G J 2008 The microRNA-200 family regulates epithelial to mesenchymal transition *ScientificWorldJournal* **8** 901–4
- Peinado H, Ballestar E, Esteller M and Cano A 2004 Snail mediates E-cadherin repression by the recruitment of the Sin3A/histone deacetylase 1 (HDAC1)/HDAC2 complex *Mol. Cell. Biol.* **24** 306–19
- Peinado H, Olmeda D and Cano A 2007 Snail, Zeb and bHLH factors in tumour progression: an alliance against the epithelial phenotype? *Nat. Rev. Cancer* **7** 415–28
- Peiro S, Escriva M, Puig I, Barbera M J, Dave N, Garcia and de Herreros A 2006 Snail1 transcriptional repressor binds to its own promoter and controls its expression *Nucleic Acids Res.* **34** 2077–84
- Prisley S, Martinelli E, Zannoni G F, Petrillo M, Filippetti F and Ferlini C 2015 Role and prognostic significance of the epithelial-mesenchymal transition factor ZEB2 in ovarian cancer *Oncotarget* **6** 18966–79
- Reeves G T 2019 The engineering principles of combining a transcriptional incoherent feedforward loop with negative feedback *J. Biol. Eng.* **13** 62
- Reichl P, Haider C, Grubinger M and Mikulits W 2012 TGF-beta in epithelial to mesenchymal transition and metastasis of liver carcinoma *Curr Pharm Des* **18** 4135–47
- Remy E, Rebouissou S, Chaouiya C, Zinovyev A, Radvanyi F and Calzone L 2015 A Modeling Approach to Explain Mutually Exclusive and Co-Occurring Genetic Alterations in Bladder Tumorigenesis *Cancer Res.* **75** 4042–52
- Ren D, Wang M, Guo W, Zhao X, Tu X, Huang S, Zou X and Peng X 2013 Wild-type p53 suppresses the epithelial-mesenchymal transition and stemness in PC-3 prostate cancer cells by modulating miR145 *Int. J. Oncol.* **42** 1473–81
- Ribba B, Colin T and Schnell S 2006 A multiscale mathematical model of cancer, and its use in analyzing irradiation therapies *Theor. Biol. Med. Model* **3** 7
- Rivlin N, Brosh R, Oren M and Rotter V 2011 Mutations in the p53 Tumor Suppressor Gene: Important Milestones at the Various Steps of Tumorigenesis *Genes cancer* **2** 466–74
- Rodriguez A, Crespo I, Androsova G and del Sol A 2015 Discrete logic modelling optimization to contextualize prior knowledge networks using PRUNET *PLoS One* **10** e0127216
- Rossato V V, Silveira D A, Gupta S and Mombach J C M 2019 Towards the contribution of the p38MAPK pathway to the dual role of TGFbeta in cancer: a boolean model approach *Comput. Biol. Med.* **104** 235–40
- Saadatpour A, Albert I and Albert R 2010 Attractor analysis of asynchronous Boolean models of signal transduction networks *J. Theor. Biol.* **266** 641–56
- Saadatpour A and Albert R 2013 Boolean modeling of biological regulatory networks: a methodology tutorial *Methods* **62** 3–12
- Saez-Rodriguez J, Alexopoulos L G, Epperlein J, Samaga R, Lauffenburger D A, Klamt S and Sorger P K 2009 Discrete logic modelling as a means to link protein signalling networks with functional analysis of mammalian signal transduction *Mol. Syst. Biol* **5** 331
- Saitoh M, Endo K, Furuya S, Minami M, Fukasawa A, Imamura T and Miyazawa K 2016 STAT3 integrates cooperative Ras and TGF-beta signals that induce Snail expression *Oncogene* **35** 1049–57
- Savagner P 2015 Epithelial–mesenchymal transitions: from cell plasticity to concept elasticity *Curr Top Dev Biol.* **112** 273–300
- Saxena K, Jolly M K and Balamurugan K 2020 Hypoxia, partial EMT and collective migration: emerging culprits in metastasis *Transl. Oncol* **13** 100845
- Scheel C. and Weinberg R. A. 2012 Cancer stem cells and epithelial–mesenchymal transition: Concepts and molecular links *Seminars in Cancer Biology* **22** 396–403
- Schlatter R, Schmich K, Avalos Vizcarra I, Scheurich P, Sauter T, Borner C, Ederer M, Merfort I and Sawodny O 2009 ON/OFF and beyond—a boolean model of apoptosis *PLoS Comput. Biol.* **5** e1000595
- Schleif R 2000 Regulation of the L-arabinose operon of Escherichia coli *Trends Genet.* **16** 559–65
- Schmierer B, Tournier A L, Bates P A and Hill C S 2008 Mathematical modeling identifies Smad nucleocytoplasmic shuttling as a dynamic signal-interpreting system *Proc. Natl. Acad. Sci. U S A* **105** 6608–13
- Selvaggio G, Canato S, Pawar A, Monteiro P T, Guerreiro P S, Bras M M, Janody F and Chaouiya C 2020 Hybrid epithelial-mesenchymal phenotypes are controlled by microenvironmental factors *Cancer Res.* **80** 2407–20
- Sever R and Brugge J S 2015 Signal transduction in cancer *Cold Spring Harb Perspect Med* **5** a006098–006098
- Shen D W, Pouliot L M, Hall M D and Gottesman M M 2012 Cisplatin resistance: a cellular self-defense mechanism resulting from multiple epigenetic and genetic changes *Pharmacol Rev* **64** 706–21
- Shen-Orr S S, Milo R, Mangan S and Alon U 2002 Network motifs in the transcriptional regulation network of Escherichia coli *Nat. Genet.* **31** 64–8
- Sherekar S and Viswanathan G A 2021 Boolean dynamic modeling of cancer signaling networks: Prognosis, progression, and therapeutics *Computational and Systems Oncology* **1** e1017
- Siemens H, Jackstadt R, Hunten S, Kaller M, Menssen A, Gotz U and Hermeking H 2011 miR-34 and SNAIL form a double-negative feedback loop to regulate epithelial-mesenchymal transitions *Cell. Cycle* **10** 4256–71
- Silveira D A, Gupta S and Mombach J C M 2020 Systems biology approach suggests new miRNAs as phenotypic stability factors in the epithelial-mesenchymal transition *J. R. Soc. Interface* **17** 20200693
- Silveira D A and Mombach J C M 2020 Dynamics of the feedback loops required for the phenotypic stabilization in the epithelial-mesenchymal transition *FEBS J.* **287** 578–88
- Smith B N and Bhowmick N A 2016 Role of EMT in metastasis and therapy resistance *J. Clin. Med* **5** 17
- Steinway S N, Zanudo J G, Ding W, Rountree C B, Feith D J, Loughran T P Jr and Albert R 2014 Network modeling of TGFbeta signaling in hepatocellular carcinoma epithelial-to-mesenchymal transition reveals joint sonic hedgehog and Wnt pathway activation *Cancer Res.* **74** 5963–77
- Steinway S N, Zanudo J G T, Michel P J, Feith D J, Loughran T P and Albert R 2015 Combinatorial interventions inhibit TGFbeta-driven epithelial-to-mesenchymal transition and support hybrid cellular phenotypes *NPI. Syst. Biol. Appl* **1** 15014
- Subbalakshmi A R, Kundnani D, Biswas K, Ghosh A, Hanash S M, Tripathi S C and Jolly M K 2020 NFATc acts as a non-canonical phenotypic stability factor for a hybrid epithelial/mesenchymal phenotype *Front. Oncol* **10** 553342
- Tang Y, Xuan Y, Qiao G, Ou Z, He Z, Zhu Q, Liao M and Yin G 2019 MDM2 promotes epithelial-mesenchymal transition through activation of Smad2/3 signaling pathway in lung adenocarcinoma *Oncotargets. Ther* **12** 2247–58
- Tej S, Gaurav K and Mukherji S 2019 Small RNA driven feed-forward loop: critical role of sRNA in noise filtering *Phys. Biol.* **16** 046008
- Thiery J P 2002 Epithelial-mesenchymal transitions in tumour progression *Nat. Rev. Cancer* **2** 442–54
- Thiery J P, Acloque H, Huang R Y and Nieto M A 2009 Epithelial-mesenchymal transitions in development and disease *Cell* **139** 871–90
- Thiery J P and Sleeman J P 2006 Complex networks orchestrate epithelial-mesenchymal transitions *Nat. Rev. Mol. Cell Biol.* **7** 131–42
- Tian X J, Zhang H and Xing J 2013 Coupled reversible and irreversible bistable switches underlying TGFbeta-induced epithelial to mesenchymal transition *Biophys. J.* **105** 1079–89
- Traynard P, Tobalina L, Eduati F, Calzone L and Saez-Rodriguez J 2017 Logic Modeling in Quantitative Systems Pharmacology *CPT Pharmacometrics Syst Pharmacol* **6** 499–511
- Tripathi S, Levine H and Jolly M K 2020 The physics of cellular decision making during epithelial-mesenchymal transition *Annu. Rev. Biophys.* **49** 1–18

- Tripathi S, Xing J, Levine H and Jolly M K 2021 Mathematical modeling of plasticity and heterogeneity in EMT *Methods Mol. Biol.* **2179** 385–413
- Tryndyak V P, Beland F A and Pogribny I P 2010 E-cadherin transcriptional down-regulation by epigenetic and microRNA-200 family alterations is related to mesenchymal and drug-resistant phenotypes in human breast cancer cells *Int. J. Cancer* **126** 2575–83
- Turner C and Kohandel M 2010 Investigating the link between epithelial-mesenchymal transition and the cancer stem cell phenotype: A mathematical approach *J. Theor. Biol.* **265** 329–35
- Tyson J J and Novak B 2014 Control of cell growth, division and death: information processing in living cells *Interface Focus* **4** 20130070
- Vega S, Morales A V, Ocana O H, Valdes F, Fabregat I and Nieto M A 2004 Snail blocks the cell cycle and confers resistance to cell death *Genes. Dev.* **18** 1131–43
- Vilar J M, Jansen R and Sander C 2006 Signal processing in the TGF-beta superfamily ligand-receptor network *PLoS Comput. Biol.* **2** e3
- Vilar J M and Saiz L 2011 Trafficking coordinate description of intracellular transport control of signaling networks *Biophys. J.* **101** 2315–23
- Vinay D S, Ryan E P, Pawelec G, Talib W H, Stagg J and Kumara H S 2015 Immune evasion in cancer: Mechanistic basis and therapeutic strategies *Seminars in cancer biology* **35** S185–98
- Vincent T, Neve E P, Johnson J R, Kukalev A, Rojo F and Fuxe J 2009 A SNAI1-SMAD3/4 transcriptional repressor complex promotes TGF-beta mediated epithelial-mesenchymal transition *Nat. Cell Biol.* **11** 943–50
- Wang L G, Ni Y, Su B H, Mu X R, Shen H C and Du J J 2013 MicroRNA-34b functions as a tumor suppressor and acts as a nodal point in the feedback loop with Met *Int. J. Oncol.* **42** 957–62
- Wang S P, Wang W L, Chang Y L, Wu C T, Chao Y C and Yang P C 2009 p53 controls cancer cell invasion by inducing the MDM2-mediated degradation of Slug *Nat. Cell Biol.* **11** 694–704
- Wang Z, Jiang Y, Guan D, Li J, Yin H, Pan Y, Xie D and Chen Y 2013 Critical roles of p53 in epithelial-mesenchymal transition and metastasis of hepatocellular carcinoma cells *PLoS One* **8** e72846
- Watanabe K, Panchy N, Noguchi S, Suzuki H and Hong T 2019 Combinatorial perturbation analysis reveals divergent regulations of mesenchymal genes during epithelial-to-mesenchymal transition *NPJ Syst Biol Appl* **5** 21
- Wellner U, Schubert J, Burk U C, Schmalhofer O, Zhu F and Brabletz T 2009 The EMT-activator ZEB1 promotes tumorigenicity by repressing stemness-inhibiting microRNAs *Nat. Cell Biol.* **11** 1487–95
- Wu M Y and Hill C S 2009 Tgf-beta superfamily signaling in embryonic development and homeostasis *Dev. Cell* **16** 329–43
- Xu J, Lamouille S and Derynck R 2009 TGF-beta-induced epithelial to mesenchymal transition *Cell Res* **19** 156–72
- Xu Q, Wang L, Li H, Han Q, Li J, Qu X, Huang S and Zhao R C 2012 Mesenchymal stem cells play a potential role in regulating the establishment and maintenance of epithelial-mesenchymal transition in MCF7 human breast cancer cells by paracrine and induced autocrine TGF-beta *Int. J. Oncol.* **41** 959–68
- Yang M H, Chen C L, Chau G Y, Chiou S H, Su C W, Chou T Y, Peng W L and Wu J C 2009 Comprehensive analysis of the independent effect of twist and snail in promoting metastasis of hepatocellular carcinoma *Hepatology* **50** 1464–74
- Yang X, Han M, Han H, Wang B, Li S, Zhang Z and Zhao W 2017 Silencing Snail suppresses tumor cell proliferation and invasion by reversing epithelial-to-mesenchymal transition and arresting G2/M phase in non-small cell lung cancer *Int. J. Oncol.* **50** 1251–60
- Yang Z, Rayala S, Nguyen D, Vadlamudi R K, Chen S and Kumar R 2005 Pak1 phosphorylation of snail, a master regulator of epithelial-to-mesenchyme transition, modulates snail's subcellular localization and functions *Cancer Res.* **65** 3179–84
- Zanudo J G T, Steinway S N and Albert R 2018 Discrete dynamic network modeling of oncogenic signaling: Mechanistic insights for personalized treatment of cancer *Curr. Opin. Syst. Biol.* **9** 1–10
- Zavadil J and Bottinger E P 2005 TGF-beta and epithelial-to-mesenchymal transitions *Oncogene* **24** 5764–74
- Zhang J, Tian X J, Chen Y J, Wang W, Watkins S and Xing J 2018 Pathway crosstalk enables cells to interpret TGF-beta duration *NPJ. Syst. Biol. Appl* **4** 18
- Zhang J, Tian X J, Zhang H, Teng Y, Li R, Bai F, Elankumaran S and Xing J 2014 TGF-beta-induced epithelial-to-mesenchymal transition proceeds through stepwise activation of multiple feedback loops *Sci. Signal* **7** ra91
- Zhang L, Huang G, Li X, Zhang Y, Jiang Y and Qian C 2013 Hypoxia induces epithelial-mesenchymal transition via activation of SNAI1 by hypoxia-inducible factor -1alpha in hepatocellular carcinoma *BMC. Cancer* **13** 108
- Zhang W, Feng M, Zheng G, Chen Y, Wang X, Pen B, Yin J, Yu Y and He Z 2012 Chemoresistance to 5-fluorouracil induces epithelial-mesenchymal transition via up-regulation of Snail in MCF7 human breast cancer cells *Biochem. Biophys. Res. Commun.* **417** 679–85
- Zheng Y. 2009 *Network Motifs in the C.elegans Neural Network and Their Dynamic Analysis*. ([https://web.mit.edu/8.592/www/grades/projects/Projects\(2009\)/YannanZheng.pdf](https://web.mit.edu/8.592/www/grades/projects/Projects(2009)/YannanZheng.pdf))
- Zi Z and Klipp E 2007 Constraint-based modeling and kinetic analysis of the Smad dependent TGF-beta signaling pathway *PLoS One* **2** e936