The 262-operation $\langle 4 \times 4 \times 4 : 48 \rangle$ algorithm is also the more accurate one: error-bound exponent 9/4 and about a quarter less error

Greg Sidebottom

Draft — October 25, 2026

Abstract

Dumas, Pernet and Sedoglavic's accurate rational $\langle 4 \times 4 \times 4 : 48 \rangle$ algorithm, Table 1 of their 2026 paper, has $(\infty, 2)$-growth factor $\gamma_{\infty,2} = 18\sqrt{2} \approx 25.46$, an error-bound exponent $\log_4 \gamma_{\infty,2} \approx 2.335$. Their later 262-operation variant, in the same orbit, was published for its lower operation count. We show that it is also the more accurate algorithm. Its growth factors are smaller in all four norms they consider: $\gamma_{\infty,2} = 16\sqrt{2}$, an exponent of exactly 9/4; $\gamma_{\infty,\infty} = 132$ against 148; $\gamma_{2,\infty} = 16\sqrt{963}$ against $8\sqrt{4142}$; and $\gamma_{2,2} = 64\sqrt{2}$ against $8\sqrt{130}$. Its $\gamma_2 = 64(1 + \sqrt{2})$ is the same, and its rounding-chain factor Q_0 is 80 against 180. Measured against exact references on random matrices, in single and double precision at one to three recursion levels, its error is 0.75–0.77 of Table 1's on average (0.62–0.90 for single instances), at the same speed in a fused-pass implementation over Apple's Accelerate. In recursion down to scalars its advantage grows, to 0.72 at five levels. We rediscovered the point by minimizing $\gamma_{\infty,2}$ over the de Groote orbit and the scheme's three role assignments with an exact-gradient search. Among the orbit's 648 dyadic orthogonal points, enumerated exhaustively, it is, up to symmetry, the unique point that minimizes both $\gamma_{\infty,2}$ and $\gamma_{\infty,\infty}$. The orbit's global $\gamma_{\infty,2}$ minimum, 21.41, needs coefficients in $\mathbb{Q}(\sqrt{2})$ and measures no better than Table 1. Dyadic probe points show that neither $\gamma_{\infty,2}$ nor $\gamma_{\infty,\infty}$ alone predicts the measured error, while Q_0 and exact dyadic coefficients matter. The other known rational classes, Li, Wang and Hu's curve and Agarwal's scheme, have larger $\gamma_{\infty,2}$ minima.

1 Introduction

Two 4×4 matrices can be multiplied with 48 multiplications over any ring containing 1/2. AlphaEvolve found a rank-48 scheme with complex coefficients [14], and Dumas, Pernet and Sedoglavic (DPS) turned it into one with rational coefficients [7]. Applied recursively, such a scheme multiplies $n \times n$ matrices with $O(n^{\log_4 48}) = O(n^{2.7925})$ operations. Fast schemes trade accuracy for speed, and their error bounds are governed by a *growth factor* computed from the scheme's coefficient matrices [3, 2, 5, 12]. DPS generalized it to a (p, q)-growth factor $\gamma_{p,q}$ for any output and input norms [6, 9], and they argue that the $(\infty, 2)$-factor $\gamma_{\infty,2}$ best reflects the accuracy observed in practice.

In [8] DPS proposed an accurate variant of their rational scheme, given as Table 1 of that paper and in PLinOpt as `4x4x4_48_accurate` [11]. They found it in the de Groote isotropy orbit [4] of their first scheme: a point minimizing the smooth bound $\gamma_2 \geq \gamma_{2,2}$ along the orbit, then moved by orthogonal isotropies, which keep γ_2, to get sparse straight-line programs. Its growth factor is $\gamma_{\infty,2} \approx 25.456$, error-bound exponent $\log_4 \gamma_{\infty,2} \approx 2.335$, the smallest among the algorithms they

compare, and its programs take 284 operations. The paper's versions run from 19 March 2026 (v1) to 29 July 2026 (v3, the current one), and Table 1's scheme entered PLinOpt on 23 June 2026.

DPS then published a cheaper accurate variant with 262 operations. It entered PLinOpt on 13 October 2026 as `4x4x4_48_accurate_262` (commit `b441528`, by J.-G. Dumas), described as "accurate with 262 operations". The second version of [10], dated 2 October 2026, cites it as the accurate variant with the best leading term, $9 + 3/16$; the first version, of 10 September 2026, presents a 216-operation algorithm and gives no accurate variant in its abstract. Neither source reports the new variant's growth factors, and Table 1 of [8] remains the accurate variant on record.

We show that the 262-operation variant is also the more accurate one:

- Its growth factors are smaller than Table 1's in all four norms of [8, Table 2], with the same γ_2: $\gamma_{\infty,2} = 16\sqrt{2}$, an error-bound exponent of exactly $9/4$, against 25.456; $\gamma_{\infty,\infty} = 132$ against 148; $\gamma_{2,2} = 64\sqrt{2}$ against 91.21; and $\gamma_{2,\infty} \approx 496.5$ against 514.9. Its rounding-chain factor Q_0 is 80 against 180 (Section 3).
- Measured on random matrices against exact references, in single and double precision, its error is 0.75–0.77 of Table 1's on average at one to three recursion levels, and never above 0.90. In recursion down to scalars the advantage grows, to 0.72 at five levels. It runs at the same speed in a fused-pass implementation over Apple's Accelerate (Sections 5 and 6).

We found this point independently, before we knew of it, by minimizing $\gamma_{\infty,2}$ over the orbit and the scheme's role assignments (Section 4), and that search is the paper's other contribution. Among the orbit's dyadic orthogonal points, which we enumerate exhaustively, the points minimizing $\gamma_{\infty,\infty}$ also minimize $\gamma_{\infty,2}$, and all of them are the 262-operation variant up to symmetry. The orbit's global $\gamma_{\infty,2}$ minimum, 21.407, is reached only at points with coefficients in $\mathbb{Q}(\sqrt{2})$, and it measures no better than Table 1. The other known rational classes, members of Li, Wang and Hu's curve [13] and Agarwal's scheme [1], have larger $\gamma_{\infty,2}$ minima. So $\gamma_{\infty,2}$ alone is not a sufficient objective for an accuracy search. The rounding-chain factor Q_0 and exact dyadic coefficients matter as much, and the 262-operation variant is the best in all of them.

2 Background

2.1 Schemes, growth factors and Q_0

For an $m \times k$ matrix A, $\mathrm{Vect}(A)$ is its row-major vectorization. A bilinear algorithm with r multiplications for $\langle m \times k \times n \rangle$ is a triple of matrices $L \in \mathbb{Q}^{r \times mk}$, $R \in \mathbb{Q}^{r \times kn}$, $P \in \mathbb{Q}^{mn \times r}$ with

$$\mathrm{Vect}(AB) = P\big((L\,\mathrm{Vect}\,A) \odot (R\,\mathrm{Vect}\,B)\big)$$

for all A and B, where $\odot$ is the entrywise product. Here $m = k = n = 4$ and $r = 48$. With L_i the rows of L, R_i the rows of R, $p_{j,i}$ the entries of P and $q^\star$ the dual exponent of q, DPS's (p, q)-growth factor [9, Th. 18] is

$$\gamma_{p,q} = \Big\| \Big(\sum_{i=1}^{r} \|L_i\|_{q^\star} \|R_i\|_{q^\star} \, |p_{j,i}| \Big)_j \Big\|_p ,$$

and the forward error of ℓ recursive levels is at most $f_{p,q} \|A\|_q \|B\|_q \, \varepsilon + O(\varepsilon^2)$ with $f_{p,q} = O(\gamma_{p,q}^{\ell}) = O(n^{\log_4 \gamma_{p,q}})$. The classical algorithm has $\gamma_{\infty,\infty} = \gamma_{\infty,2} = 4$. DPS also use the smooth bound $\gamma_2 = \sum_i \|L_i\|_2 \|R_i\|_2 \|P_{*,i}\|_2 \geq \gamma_{2,2}$ [8, eq. (5)]. PLinOpt's `growthfactor` tool [11] also reports a combinatorial factor,

$$Q_0 = \max_j \Big(\max_{i:\, p_{j,i} \neq 0} \|L_i\|_0 \, \|R_i\|_0 + \|P_j\|_0 \Big),$$

where $\|\cdot\|_0$ counts nonzeros. It bounds the length of the chain of roundings that feeds one output, and the tool combines it with the growth factors in the constants of its error bounds.

2.2 Isotropies and symmetries

For $(U, V, W) \in GL_4^3$, applying an algorithm to $U^{-1}AV$ and $V^{-1}BW$ returns $U^{-1}ABW$, so

$$(L, R, P) \;\mapsto\; \big(L\,(U^{-1} \otimes V^{\mathsf{T}}),\; R\,(V^{-1} \otimes W^{\mathsf{T}}),\; (U \otimes W^{-\mathsf{T}})\,P\big) \tag{1}$$

is again an algorithm, since $\mathrm{Vect}(XAY) = (X \otimes Y^{\mathsf{T}})\,\mathrm{Vect}\,A$ for row-major vectorization. These are de Groote's isotropies [4]; PLinOpt's orbit tool uses the same group. With $A_i = \mathrm{Mat}(L_i)$, $B_i = \mathrm{Mat}(R_i)$ and Z_i the 4×4 matrix of column i of P, (1) reads

$$A_i \mapsto U^{-\mathsf{T}} A_i V^{\mathsf{T}}, \qquad B_i \mapsto V^{-\mathsf{T}} B_i W^{\mathsf{T}}, \qquad Z_i \mapsto U Z_i W^{-1}.$$

Two further symmetries come from the trace form

$$\textstyle\sum_i \mathrm{tr}(X_i A)\,\mathrm{tr}(Y_i B)\,\mathrm{tr}(Z_i G) = \mathrm{tr}(ABG), \qquad X_i = A_i^{\mathsf{T}},\; Y_i = B_i^{\mathsf{T}}:$$

- the cyclic rotation $(X, Y, Z) \mapsto (Y, Z, X)$;
- the input swap $(X, Y, Z) \mapsto (Y^{\mathsf{T}}, X^{\mathsf{T}}, Z^{\mathsf{T}})$, from $(AB)^{\mathsf{T}} = B^{\mathsf{T}} A^{\mathsf{T}}$.

Every $\gamma_{p,q}$ treats L and R alike, through the norms of their rows, and P differently, through its entries and the norm over outputs. So the input swap leaves the growth factors unchanged, but the rotation does not: the three *role assignments*, one for each factor set that computes the outputs, are three different algorithms for accuracy purposes. γ_2 is symmetric in the three sets, and the rotation leaves it unchanged.

2.3 Invariances

Three facts organize the search.

- Scaling one product, $(L_i, R_i, P_{*,i}) \mapsto (\alpha L_i, \beta R_i, P_{*,i}/(\alpha\beta))$, changes no $\gamma_{p,q}$, no γ_2 and not Q_0. So none depends on the scale of U, V or W.
- For orthogonal U, V, W the Kronecker factors in (1) are orthogonal, so the 2-norms of the rows of L and R and of the columns of P are unchanged, and so is γ_2. Then $\gamma_{\infty,2} = \max_{p,q} \sum_i w_i |(U Z_i W^{\mathsf{T}})_{pq}|$ with fixed weights $w_i = \|L_i\|_2 \|R_i\|_2$, and V drops out.
- Multiplying U, V or W on the left by a signed permutation permutes rows and columns of the A_i, B_i, Z_i with signs. That changes no $\gamma_{p,q}$ and not Q_0.

3 The 262-operation variant

DPS's 262-operation variant is PLinOpt's `4x4x4_48_accurate_262`, added on 13 October 2026. Its straight-line programs take 64 additions and 8 scalings for L, 67 and 11 for R, and 112 additions and no scalings for P, against Table 1's $80+4$, $68+8$ and $108+16$. Up to symmetry it is Table 1's

Table 1: Growth factors with their error-bound exponents $\log_k \gamma$ ($k = 2$ for $\langle 2 \times 2 \times 2 : 7 \rangle$, $k = 4$ for $\langle 4 \times 4 \times 4 : 48 \rangle$), γ_2, Q_0 and the operations of the published straight-line programs. The first five rows are the algorithms of [8, Table 2] (values for DPS24, whose coefficients involve $\sqrt{3}$, from that table). The last two are points of the same orbit from our search and our earlier work [16].

algorithm	$\gamma_{\infty,\infty}$	$\gamma_{\infty,2}$	$\gamma_{2,\infty}$	$\gamma_{2,2}$	γ_2	Q_0	ops
$\langle 2 \times 2 \times 2 : 7 \rangle$							
Winograd	18 (4.170)	8 (3.000)	31.24 (4.965)	14 (3.807)	17.85	13	15
Strassen	12 (3.585)	6.83 (2.772)	17.89 (4.161)	10.45 (3.386)	14.83	8	18
DPS24 [6]	17.48 (4.128)	5.97 (2.577)	27.71 (4.793)	10.01 (3.323)	–	23	–
$\langle 4 \times 4 \times 4 : 48 \rangle$							
DPS25 [7]	288 (4.085)	38.16 (2.627)	1032.0 (5.006)	139.9 (3.564)	210.5	152	315
Table 1 [8]	148 (3.605)	25.46 (2.335)	514.9 (4.504)	91.21 (3.256)	154.5	180	284
262-op. variant	**132** (3.522)	**22.63** (2.250)	**496.5** (4.478)	**90.51** (3.250)	154.5	**80**	262
orbit's $\gamma_{\infty,2}$ min., $\mathbb{Q}(\sqrt{2})$	176.6 (3.732)	21.41 (2.210)	583.8 (4.595)	85.63 (3.210)	157.3	146	–
201-op. point [16]	399 (4.320)	75.85 (3.123)	1072.5 (5.033)	225.9 (3.910)	369.7	101	201

scheme moved by an orthogonal isotropy with entries $0, \pm 1/\sqrt{2}$. Take Table 1's (L, R, P) in its published role and apply (1) with $(U, V, W) = (Q_U, Q_V, Q_W)/\sqrt{2}$, where

$$Q_U = \begin{pmatrix} 0 & -1 & 0 & -1 \\ 1 & 0 & -1 & 0 \\ 0 & -1 & 0 & 1 \\ -1 & 0 & -1 & 0 \end{pmatrix}, \quad Q_V = \begin{pmatrix} -1 & -1 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & -1 & -1 \\ 0 & 0 & -1 & 1 \end{pmatrix}, \quad Q_W = \begin{pmatrix} 1 & 0 & -1 & 0 \\ 0 & -1 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix}.$$

Each Kronecker factor in (1) is then a $\{0, \pm 1\}$ matrix divided by 2, so the result has coefficients in $\mathbb{Z}[\frac{1}{2}]$ with denominators at most 4, like Table 1. Mapping `4x4x4_48_accurate_262` through the input swap and signed permutations of U, V, W, and relabelling and rescaling the products, gives exactly this scheme.

Proposition 3.1. *The 262-operation variant has* $\gamma_{\infty,\infty} = 132$, $\gamma_{\infty,2} = 16\sqrt{2}$, $\gamma_{2,\infty} = 16\sqrt{963} \approx 496.52$, $\gamma_{2,2} = 64\sqrt{2}$, $\gamma_2 = 64(1 + \sqrt{2})$ *and* $Q_0 = 80$. *Table 1's scheme has* 148, $18\sqrt{2} \approx 25.456$, $8\sqrt{4142} \approx 514.87$, $8\sqrt{130} \approx 91.214$, $64(1 + \sqrt{2})$ *and* 180.

The values are exact. Every weight $\|L_i\|_2 \|R_i\|_2$ and every column norm of P is a rational or a rational multiple of $\sqrt{2}$, so the sums lie in $\mathbb{Q}(\sqrt{2})$, and we computed them, and the rational squares of $\gamma_{2,\infty}$ and $\gamma_{2,2}$, in exact arithmetic. PLinOpt's `growthfactor`, working in floating point, agrees to every printed digit. Table 1 sets them beside the other algorithms of [8, Table 2] and two points from our search. The error-bound exponents $\log_4 \gamma_{\infty,2}$ and $\log_4 \gamma_{2,2}$ are exactly 9/4 and 13/4. Both algorithms' published programs pass PLinOpt's `SLPchecker` against their matrices, and both pass our exact check against the matrix multiplication tensor.

4 How we found it

4.1 Role assignments

The three role assignments of Table 1's scheme have $\gamma_{\infty,2} = 25.456$, 32.971 and 28.142 ($\gamma_{\infty,\infty} = 148$, 212, 180), so the published one is already the best. For DPS25 they give 38.162, 39.282 and 33.540, and the published one is not the best. Every search below runs from all three.

4.2 Continuous search over the orbit

We minimize $\gamma_{\infty,2}$ over $(U,V,W) \in GL_4^3$. The max over the 16 outputs is smoothed by a log-sum-exp with sharpness β and $|x|$ by $\sqrt{x^2+\epsilon^2}$, the smoothed objective is minimized by BFGS with a backtracking line search, and (β,ϵ) runs through a continuation from $(0.5, 10^{-1})$ to $(32768, 10^{-6})$. After each stage each of U, V, W is rescaled to determinant ± 1, which changes nothing (Section 2.3). The gradient is exact: with $M_i = U^{-\mathsf{T}} A_i V^{\mathsf{T}}$, $N_i = V^{-\mathsf{T}} B_i W^{\mathsf{T}}$, $Z_i' = U Z_i W^{-1}$, $a_i = \|M_i\|_F$, $b_i = \|N_i\|_F$ and π the softmax weights of the outputs, the adjoints are

$$\frac{\partial f}{\partial Z_i'} = \pi \circ a_i b_i \, s'(Z_i'), \qquad \frac{\partial f}{\partial M_i} = \frac{b_i S_i}{a_i} M_i, \qquad \frac{\partial f}{\partial N_i} = \frac{a_i S_i}{b_i} N_i,$$

with $S_i = \sum_{pq} \pi_{pq} s(Z'_{i,pq})$. They are carried through the products and through each inverse by $\partial X = -X \, \partial M \, X$. The implementation (`gamma48`, in Rust) checks this gradient against central differences to 10^{-10} relative. It runs on all cores, and one start takes about 0.5 s on one core. Starts are the identity, random near-identity points, and random rotations times random perturbations of the identity.

Within the orthogonal sub-orbit (γ_2 fixed at its minimum 154.51; 12 parameters, two Cayley-parametrized rotations), the minimum is $16\sqrt{2} \approx 22.627$ for the published role and $12 + 8\sqrt{2} \approx 23.314$ for the other two. At the minimum U and W have entries $0, \pm 1/\sqrt{2}$. Over the whole orbit, 30,000 starts from Table 1 (10,000 per role) and 7,200 each from DPS25 and from our 201-operation point give the same local optima, a direct sign that the three lie in one orbit. The lowest, 21.4073, is found thousands of times, and nothing lower appears. Its points have two orthogonal factors and one with singular values $(1, 1, s, s)$, $s = 2(\sqrt{2}-1)$, so their coefficients lie in $\mathbb{Q}(\sqrt{2})$. They cannot be made dyadic: the scaling enters R as s and P as $1/s$, and no dyadic value other than a power of two has a dyadic inverse. The next optima are 21.944 (roles 1 and 2) and 22.627 (role 0).

4.3 The dyadic orthogonal sub-orbit, exhaustively

A scheme is exact in binary floating point only if its coefficients are dyadic, and multiplications by powers of two do not round. Let G be the set of orthogonal 4×4 matrices whose entries lie in $\mathbb{Z}[\frac{1}{2}]$ or in $\mathbb{Z}[\frac{1}{2}]/\sqrt{2}$. It has 2304 elements:

- the 384 signed permutations and the 768 matrices with entries $\pm 1/2$, which together form the Weyl group $W(F_4)$;
- the 1152 matrices with entries $0, \pm 1/\sqrt{2}$.

It is closed under products and inverses. For $(U,V,W) \in G^3$ the three Kronecker factors of (1) carry the factors $\sqrt{2}^{-(c_U+c_V)}$, $\sqrt{2}^{-(c_V+c_W)}$ and $\sqrt{2}^{-(c_U+c_W)}$, with $c = 1$ for the $1/\sqrt{2}$ matrices and 0 otherwise. Their product is a power of two, so scaling each product makes every coefficient dyadic. By Section 2.3 the growth factors and Q_0 depend only on the classes of U, V, W under left multiplication by signed permutations. There are six classes of 384, so the sub-orbit of each role has 216 distinct points, 648 in all, and we enumerate them.

Theorem 4.1. *Among the* 648 *points of the dyadic orthogonal sub-orbit of Table 1's scheme, the minimum of* $\gamma_{\infty,2}$ *is* $16\sqrt{2}$*, attained at* 42 *points. The minimum of* $\gamma_{\infty,\infty}$ *is* 132*, attained at exactly eight points, all in the published role, and these eight also attain* $\gamma_{\infty,2} = 16\sqrt{2}$*. Each of the eight is the* 262*-operation variant up to signed permutations of* (U,V,W)*, the input swap, and a relabelling and rescaling of the products.*

Table 2: Smallest $\gamma_{\infty,2}$ found over the whole orbit of each rational class, 2400 starts per role, and the Li–Wang–Hu invariant φ. The role numbering follows each scheme's own coordinates.

class	φ	role 0	role 1	role 2	minimum
$s(1/4)$, the DPS class	192	21.407	22.627	21.407	21.407
$s(1/2)$	336	23.144	23.654	23.144	23.144
$s(1)$	1092	25.167	26.833	25.167	25.167
$s(2)$	4161	27.852	32.485	27.852	27.852
$s(4)$	16448.25	32.677	40.319	32.677	32.677
$s(8)$	65600.06	40.134	51.351	40.134	40.134
Agarwal's scheme	–	39.619	42.924	39.619	39.619

The proof is the enumeration together with an explicit equivalence for each of the eight points (`equiv.py`). Table 1's own point has the fourth smallest $\gamma_{\infty,2}$ in the sub-orbit (25.456, after $16\sqrt{2}$, $12+8\sqrt{2}$ and 24). So within the dyadic orthogonal moves of Table 1, the 262-operation variant is, up to symmetry, the unique point that is best for both growth factors.

4.4 Other classes

Rational rank-48 schemes are not all in one orbit. Li, Wang and Hu put DPS's scheme on a curve $s(t)$ whose members are pairwise inequivalent for $t \geq 1/4$ [13], with the invariant $\varphi = 64+4/t^2+1024t^2$; the DPS class is $t = 1/4$, the minimum of φ. For $t = 2^k$ the coefficients are dyadic. Agarwal replaced twelve products of $s(2)$ to get a scheme off the curve [1]. We rebuilt both exactly from the papers' statements (checked on all 4096 Brent equations) and ran the continuous search from each, 2400 starts per role (Table 2). The smallest $\gamma_{\infty,2}$ grows steadily with t. No other class has any point, dyadic or not, as low as the 262-operation variant's 22.63; the closest, $s(1/2)$, bottoms out at 23.14.

5 Accuracy

5.1 Implementation

The measurements use our fast matrix multiplication library for Apple silicon [17]. One level of a $\langle 4\times 4\times 4 : 48\rangle$ scheme runs as three fused passes, each one streaming loop over the elements of the blocks:

- the left pass computes all 48 combinations $L\,\mathrm{Vect}\,A$ of the 16 blocks of A;
- the right pass does the same for B;
- the output pass combines the 48 block products into the 16 blocks of C.

The block products go to Accelerate's GEMM (`cblas_sgemm`, `cblas_dgemm`). The passes are generated from the published straight-line programs, operation by operation in the programs' order, so each program's rounding is the one measured. Deeper levels recurse on the block products. One, two and three levels at $n = 4096$ leave GEMMs of size 1024, 256 and 64, and at $n = 8192$ sizes 2048, 512 and 128. For comparison we include Winograd's $\langle 2\times 2\times 2 : 7\rangle$ (Strassen–Winograd, 15 additions) at one, two and four levels. Two of its levels reduce the size as much as one $\langle 4\times 4\times 4\rangle$ level.

5.2 Protocol

- Inputs are uniform on $[-1, 1)$ or standard normal, from a xorshift generator. Each setting uses five seeds, three for double precision at $n = 8192$.
- Single precision is measured against the double-precision product of the same inputs (Accelerate's `dgemm`). Its 29 extra bits make it exact for this purpose.
- Double precision is measured against a compensated product: every entry is an Ogita–Rump–Oishi Dot2 dot product [15], with each product split exactly by a fused multiply-add and the sum accumulated by error-free TwoSum steps, then rounded to double. Its own error is about the unit roundoff, two orders of magnitude below the errors measured.
- The error is $\max_{ij}|\hat{C}_{ij} - C_{ij}| / \max_{ij}|C_{ij}|$. We report it as a multiple of the classical GEMM's own error on the same inputs.

The programs are:

- Table 1's published 284-operation programs;
- the 262-operation variant's published programs;
- our representative of the same point, with programs generated by PLinOpt's `optimizer` (445 operations);
- DPS25's 315-operation programs and the 201-operation programs of [16];
- the orbit's $\gamma_{\infty,2}$ minimum, written as direct sums with its coefficients rounded to the working precision;
- Table 1 written as direct sums, a control for program style;
- two dyadic points of Table 1's orbit, written as direct sums, chosen from the enumeration of Section 4.3 to separate the two growth factors. Probe A has the 262-operation variant's $\gamma_{\infty,2} = 16\sqrt{2}$ but $\gamma_{\infty,\infty} = 256$ (role 0, classes $5, 2, 3$); probe B has $\gamma_{\infty,2} = 28.97$, larger than Table 1's, but $\gamma_{\infty,\infty} = 144$, smaller (role 2, classes $4, 4, 2$). Their Q_0 are 144 and 94.

5.3 Results

Table 3 gives each scheme's mean error as a multiple of the classical GEMM's. Table 4 gives the 262-operation variant's error divided by Table 1's on the same inputs, and Figure 1 plots the means.

- The 262-operation variant is the most accurate scheme at every depth, in both precisions and for both distributions. Its error is 0.73–0.83 of Table 1's averaged over the seeds of each setting, 0.75–0.77 averaged over all instances, and 0.62–0.90 for single instances.
- At one and two levels, where the library's plans run, it is more accurate than the classical GEMM itself (0.72–0.81 of its error), while Table 1 is about as accurate as the classical GEMM (0.94–1.05). The block products are dot products a quarter or a sixteenth as long, which saves more than the passes add.
- Single and double precision agree within a few percent, and so do the two sizes and the two distributions.

Table 3: Error against the exact reference as a multiple of the classical GEMM's own error, mean over all instances: single precision 20 instances ($n = 4096$ and 8192, uniform and normal, five seeds each); double precision 16 for the first five schemes ($n = 4096$, five seeds per distribution; $n = 8192$, three) and 10 ($n = 4096$ only) for the rest. Winograd's levels are binary, so its columns are two and four levels; "–": not measured.

	single precision			double precision		
levels	1	2	3	1	2	3
Table 1, published programs (284)	0.94	1.05	1.54	0.94	1.03	1.57
262-op. variant, published programs	0.73	0.79	1.17	0.72	0.81	1.16
262-op. variant, our programs (445)	0.71	0.83	1.33	0.73	0.83	1.32
Table 1, direct sums	0.96	1.07	1.65	0.93	1.08	1.70
probe A, direct sums	0.74	0.82	1.32	0.71	0.85	1.36
probe B, direct sums	0.78	0.82	1.26	0.74	0.86	1.36
orbit's $\gamma_{\infty,2}$ minimum, $\mathbb{Q}(\sqrt{2})$	0.86	1.00	1.61	0.88	1.01	1.57
DPS25 (315)	1.06	1.60	3.27	1.03	1.56	3.23
201-operation point	2.78	8.72	36.84	2.59	8.74	36.77
Winograd, 2 and 4 binary levels	–	3.11	10.25	–	3.00	9.93

Table 4: Error of the 262-operation variant (published programs) divided by Table 1's on the same inputs: mean over the seeds, with the range in brackets.

precision	setting	1 level	2 levels	3 levels
single	$n = 4096$, uniform	0.82 [0.75, 0.90]	0.74 [0.67, 0.81]	0.78 [0.76, 0.82]
single	$n = 4096$, normal	0.76 [0.64, 0.88]	0.81 [0.77, 0.84]	0.75 [0.66, 0.80]
single	$n = 8192$, uniform	0.77 [0.69, 0.85]	0.74 [0.67, 0.79]	0.76 [0.70, 0.78]
single	$n = 8192$, normal	0.77 [0.73, 0.81]	0.73 [0.62, 0.77]	0.75 [0.68, 0.85]
double	$n = 4096$, uniform	0.79 [0.73, 0.86]	0.78 [0.72, 0.83]	0.73 [0.70, 0.82]
double	$n = 4096$, normal	0.77 [0.71, 0.83]	0.78 [0.75, 0.85]	0.76 [0.67, 0.89]
double	$n = 8192$, uniform	0.73 [0.64, 0.83]	0.83 [0.80, 0.88]	0.74 [0.74, 0.74]
double	$n = 8192$, normal	0.75 [0.68, 0.81]	0.78 [0.75, 0.81]	0.76 [0.70, 0.83]

- Over these depths the advantage is nearly constant: from two levels to three both errors grow by a factor of 1.46–1.47 in single precision (1.52 for Table 1 and 1.43 for the variant in double). With large leaves the leaf GEMMs' error dominates, and the difference comes from the passes' constant (Q_0), not yet from the growth factor. Section 5.4 recurses down to scalars to see the growth.

5.4 Deep recursion

To see the growth factors at work we recursed down to scalars: $n = 1024$ with one to five levels (leaves 256, 64, 16, 4 and 1), as in DPS's experiments [8, Fig. 1], ten instances per precision. Table 7 gives the results.

- The 262-operation variant is the most accurate at every depth again, and its advantage over Table 1 grows with depth: its error is 0.80–0.83 of Table 1's at one level and 0.72 at five (single instances 0.67–0.80), in both precisions.

Table 5: The factors beside the measured single-precision error at three levels (a multiple of the classical GEMM's error, mean over all instances) and its growth from two levels to three. Sorted by the measured error.

scheme	$\gamma_{\infty,2}$	$\gamma_{\infty,\infty}$	Q_0	program	3 levels	growth
262-op. variant	22.63	132	80	published, 262	1.17	1.47
probe B	28.97	144	94	direct sums	1.26	1.53
probe A	22.63	256	144	direct sums	1.32	1.62
262-op. variant	22.63	132	80	PLinOpt, 445	1.33	1.60
Table 1	25.46	148	180	published, 284	1.54	1.46
orbit's $\gamma_{\infty,2}$ min.	21.41	176.6	146	direct sums	1.61	1.61
Table 1	25.46	148	180	direct sums	1.65	1.55
DPS25	38.16	288	152	published, 315	3.27	2.05
201-op. point	75.85	399	101	published, 201	36.84	4.22

Table 6: Time of the published programs in our library: the best of three alternating rounds, each the median of five runs.

precision variant	plan, n ratio	Table 1	262-op.	
single	`D48/G`, 4096	0.044 s	0.044 s	1.000
single	`D48/G6`, 4096	0.036 s	0.036 s	1.000
single	`D48+4/D48/G6`, 8192	0.236 s	0.237 s	1.004
single	`D48/G`, 8192	0.272 s	0.271 s	0.996
double	`D48/G`, 4096	0.139 s	0.140 s	1.007
double	`D48/G6`, 4096	0.177 s	0.173 s	0.977
double	`D48+4/D48/G6`, 8192	0.800 s	0.803 s	1.004
double	`D48/G`, 8192	1.054 s	1.061 s	1.007

- From four levels to five, the errors grow by 2.60–2.61 for the 262-operation variant and 2.75–2.79 for Table 1, about 6% slower per level. That is the direction $\gamma_{\infty,2}$ predicts, but less than its ratio $16\sqrt{2}/18\sqrt{2} = 8/9$ would give.
- Programs matter more at depth. The 262-operation variant's 445-operation programs fall behind Table 1's published programs from three levels on (15.5 against 13.3 times the classical error at five levels, single precision), while its published 262-operation programs are the best of all.
- Among the direct-sum programs, the dyadic probes stay 24–29% below Table 1 at five levels and the orbit's $\gamma_{\infty,2}$ minimum only about 8%: the rounding of its constants compounds with depth.
- DPS25 grows by about 3.6 per level, the 201-operation point by about 7.5, and Winograd by about 7 per two binary levels.

6 Speed

Table 6 times the published programs in our library. It covers single levels with one GEMM at a time (`D48/G`) and the plans the library picks at these sizes, which use six GEMM streams and




Figure 1: Mean error against the exact reference as a multiple of the classical GEMM's error, over all sizes, distributions and seeds of the main campaign ($n = 4096$ and 8192). Winograd's points are two and four binary levels.

pipelined passes (`D48/G6`, `D48+4/D48/G6`), in alternating rounds. Both take the same time to within the timer's resolution (0.977–1.007). The passes move the same data, and with 262 against 284 operations neither is bound by arithmetic. Our representative's 445-operation programs also ran at Table 1's speed (1.000–1.028 in an earlier run). So the more accurate algorithm costs nothing at run time, and it is the cheaper one in operation count.

7 Discussion

What predicts the measured error. Table 5 sets the factors beside the error measured at three levels in single precision, where the schemes differ most. The growth factors separate the distant schemes (DPS25, the 201-operation point, two Winograd levels), but no single factor orders the close ones.

- Probe A has the 262-operation variant's $\gamma_{\infty,2}$ and measures worse.
- Probe B has a larger $\gamma_{\infty,2}$ than Table 1 and measures better.
- The orbit's $\gamma_{\infty,2}$ minimum has the smallest $\gamma_{\infty,2}$ of all and measures like Table 1.

Among the dyadic schemes, Q_0 comes closest to the measured order: 80, 94, 144 and 180 for the 262-operation variant, probe B, probe A and Table 1. The 262-operation variant is best in every factor at once, and it is the most accurate.

Dyadic coefficients. The orbit's $\gamma_{\infty,2}$ minimum and probe A have nearly the same $\gamma_{\infty,2}$ (21.41 and 22.63) and Q_0 (146 and 144), and the former has the smaller $\gamma_{\infty,\infty}$ (176.6 against 256). Yet the former measures about 20% worse at three levels: 1.61 against 1.32 times the classical error in single precision, and 1.57 against 1.36 in double. Its constants, such as $2(\sqrt{2}-1)$, are rounded to the working precision, so the scheme itself is inexact by a unit roundoff, and each multiplication by

Table 7: Deep recursion at $n = 1024$: mean error against the exact reference as a multiple of the classical GEMM's error, over ten instances per precision (uniform and normal, five seeds each); five levels reach scalar leaves. Winograd: twice as many binary levels.

	single precision					double precision				
levels	1	2	3	4	5	1	2	3	4	5
Table 1 (284)	0.95	1.10	2.00	4.77	13.29	0.91	1.18	2.09	4.91	13.50
262 variant (262)	0.79	0.97	1.67	3.75	9.79	0.76	0.98	1.63	3.68	9.58
262 variant (445)	0.78	1.10	2.29	5.71	15.45	0.78	1.09	2.20	5.48	15.62
Table 1, direct sums	1.00	1.35	2.69	6.76	19.63	0.95	1.29	2.57	6.62	19.36
probe A	0.78	1.08	2.27	5.40	14.91	0.80	1.09	2.21	5.43	14.45
probe B	0.82	1.08	2.12	5.28	13.96	0.82	1.05	2.12	5.10	14.05
$\gamma_{\infty,2}$ min., $\mathbb{Q}(\sqrt{2})$	0.89	1.22	2.40	6.42	18.06	0.92	1.20	2.36	5.89	17.57
DPS25	1.08	1.92	4.76	15.48	56.64	1.13	1.90	4.80	15.36	55.20
201-op. point	2.75	9.86	48.87	304	2242	2.89	9.60	47.50	301	2293
Winograd	2.88	10.46	46.34	254	1726	3.38	11.14	47.75	248	1753

them rounds. Multiplications by $\pm 1/2^k$ are exact. The gap compounds with depth: at five levels the probes are 24–29% below Table 1's direct sums and the $\mathbb{Q}(\sqrt{2})$ point only 8%. A search for accuracy should therefore stay in dyadic coordinates.

Programs. The same scheme measures differently with different programs. The 262-operation variant's published programs beat our 445-operation ones, and Table 1's published programs beat its direct sums (1.17 against 1.33, and 1.54 against 1.65, times the classical error at three levels in single precision). Fewer operations mean shorter rounding chains, so sparse programs are worth having for accuracy as well as for speed.

Recommendation. Wherever Table 1's algorithm is used, the 262-operation variant is better on both counts: 22 fewer operations and about a quarter less error, in the same orbit and with the same γ_2.

Limitations.

- The measurements come from one machine and one implementation style: fused passes over a vendor GEMM, with large leaves. The inputs are i.i.d. uniform or normal matrices, the error is the max-norm error relative to $\max|C|$, and there are at most three levels at $n = 4096$ and 8192 and five at $n = 1024$. Structured or badly scaled inputs may rank the schemes differently.
- The continuous search gives no proof of global optimality. Only the dyadic orthogonal moves are enumerated exhaustively; non-orthogonal dyadic isotropies, such as shears and scalings by two, are not.

Outlook. The natural next objective combines the growth factors with Q_0 and the operation count over dyadic points, or uses the measured error of deep recursion directly; `gamma48` makes such searches cheap. Lower precisions (bf16, fp16), where every level's error is a larger share of the result, are where an accurate scheme matters most.




Figure 2: Deep recursion at $n = 1024$: mean error against the exact reference as a multiple of the classical GEMM's error, over ten instances per precision. Winograd's points are two binary levels per step.

8 Data and code

Everything is in the public repository https://github.com/gsidebottom/logic; paths below are relative to it.

- The search: `src/bin/gamma48.rs` (the continuous search) and `matmul/dps48/gamma/`, which holds the role assignments and the isotropy action (`orbit.py`), the earlier Python search (`search_orth.py`, `search_gl.py`), the exact constructions (`exact.py`), the exhaustive dyadic enumeration (`dyadic_orth.py`) and the equivalence test (`equiv.py`, `equiv262.py`).
- Growth factors: `matmul/fastmm/growth.py`, checked against PLinOpt's `growthfactor`.
- Other classes: `matmul/dps48/lwh_family.py` and `agarwal_replacement.py`, which rebuild the Li–Wang–Hu members and Agarwal's scheme from the papers.
- Measurements: `matmul/fastmm` (`bench_fmm.c`, `bench_impl.h`, `gen_kernels.py`); the build for the 262-operation programs is `build/bench_fmm_acc262`. `FMM_EXACT`, `FMM_NORMAL` and `FMM_SEED` select the reference, the distribution and the instance, and `FMM_REFCACHE` shares references between builds. The probes' programs are in `matmul/dps48/gamma/slp/`. Raw results are in `matmul/fastmm/results/` (`accuracy_campaign_2026-10-25.txt`, `accuracy_deep_2026-10-25.txt`, `gamma_search_2026-10-25.txt`, `gamma_classes_2026-10-25.txt`), and `matmul/fastmm/paper_acc262.py` turns them into this paper's tables and figures.
- The schemes: PLinOpt's `data/4x4x4_48_accurate_262` and `data/4x4x4_48_accurate` [11]; our representative is `matmul/dps48/gamma/g16s2_{L,R,P}.sms`.

Acknowledgments

This work was carried out in an interactive collaboration with Claude (Anthropic; the Opus 5.5 model), which implemented the search, measurement and verification tools and much of this text under the author's direction and review. Every claim is checked mechanically by the tools listed above; the results rest on those checks rather than on trust in either the author or the tools. The algorithm is Dumas, Pernet and Sedoglavic's: we rediscovered it, and found that it was already in PLinOpt only after constructing it.

References

[1] A. Agarwal, *Certified local rank and uniqueness barriers for a 48-term matrix-multiplication decomposition*, arXiv:2609.13596 (2026).

[2] D. Bini, G. Lotti, *Stability of fast algorithms for matrix multiplication*, Numer. Math. 36 (1980) 63–72.

[3] R. P. Brent, *Algorithms for matrix multiplication*, Technical Report STAN-CS-70-157, Stanford University (1970).

[4] H. F. de Groote, *On varieties of optimal algorithms for the computation of bilinear mappings I. The isotropy group of a bilinear mapping*, Theoret. Comput. Sci. 7 (1978) 1–24.

[5] J. Demmel, I. Dumitriu, O. Holtz, R. Kleinberg, *Fast matrix multiplication is stable*, Numer. Math. 106 (2007) 199–224.

[6] J.-G. Dumas, C. Pernet, A. Sedoglavic, *Strassen's algorithm is not optimally accurate*, Proc. ISSAC 2024, 254–263.

[7] J.-G. Dumas, C. Pernet, A. Sedoglavic, *A non-commutative algorithm for multiplying* 4×4 *matrices using* 48 *non-complex multiplications*, arXiv:2506.13242 (2025).

[8] J.-G. Dumas, C. Pernet, A. Sedoglavic, *A more accurate rational non-commutative algorithm for multiplying* 4×4 *matrices using* 48 *multiplications*, arXiv:2603.18699 (v1 19 March 2026, v3 29 July 2026).

[9] J.-G. Dumas, C. Pernet, A. Sedoglavic, *Towards automated generation of fast and accurate algorithms for recursive matrix multiplication*, J. Symbolic Comput. 134 (2026) 102524.

[10] J.-G. Dumas, C. Pernet, A. Sedoglavic, J. Stapleton, P. Tichavský, *Fast matrix multiplication via recursive* $\langle 4 \times 4 \times 4 : 48 \rangle$ *algorithms into practice*, arXiv:2609.12027 (v1 10 September 2026, v2 2 October 2026).

[11] J.-G. Dumas, B. Grenet, C. Pernet, A. Sedoglavic, *PLinOpt, a collection of C++ routines handling linear & bilinear programs*, https://github.com/jgdumas/plinopt; data files `4x4x4_48_accurate` (commit `fb753b5`, 23 June 2026) and `4x4x4_48_accurate_262` (commit `b441528`, 13 October 2026).

[12] N. J. Higham, *Accuracy and Stability of Numerical Algorithms*, 2nd ed., SIAM, 2002.

[13] X. Li, Y. Wang, S. Hu, *Substitution and quotient of the isotropy group action*, arXiv:2607.15069 (2026).

[14] A. Novikov et al., *AlphaEvolve: a coding agent for scientific and algorithmic discovery*, arXiv:2506.13131 (2025).

[15] T. Ogita, S. M. Rump, S. Oishi, *Accurate sum and dot product*, SIAM J. Sci. Comput. 26 (2005) 1955–1988.

[16] G. Sidebottom, *201 operations for rank-48* 4×4 *matrix multiplication over rings containing* $1/2$, preprint (2026); PLinOpt data file `4x4x4_48_201`.

[17] G. Sidebottom, *logic*, source repository, https://github.com/gsidebottom/logic (2026).