

Patient name: Srikrishna Gopalakrishnan
DOB: August 11, 1978
Sex: Male

Kit ID: ERW946
System version: 5.0.0


Sample Collected: March 3, 2026
Sequence Sample Received: March 11, 2026
Report Generated: April 1, 2026

Microbiome Summary Score

KEY: ■ Lowest 20% ■ Below median ■ Above median ■ Top 10%

74 /100 microbiome score

There may be imbalances in your microbiome, and we recommend taking action and resampling in the future.



ⓘ About the microbiome summary score


Your Microbiome Summary Score (0–100) combines key microbiome metrics for your age range into a single number. Higher scores indicate better microbiome health. We analyze factors like beneficial bacteria and opportunistic pathogens, then compare your results to samples from your age group (using public data and proprietary data from Tiny Health customers). Your percentile shows where you rank. Example: 41st percentile means 41% of those tested scored lower than you, 59% scored higher.

Note:


- Beta metrics aren't included in this score.
- Scores for pregnant women are calculated separately and should only be compared to other pregnant scores.

Sample at a glance

Overview



Category Overview



Top areas we recommend focusing on first



Metrics that need support and improvement across categories

Beneficial microbes	
Beneficial Bifidobacterium	
 Bifidobacterium	8.296%
Metabolic Health	
 Akkermansia	0%
Disruptive microbes	
Opportunistic Pathogens	
 Klebsiella	0.194%
 Klebsiella pneumoniae	0.127%
 Citrobacter	0.205%
 Enterobacter	0.181%
 Haemophilus parainfluenzae	0.025%
 Enterococcus faecalis	0.029%
Gut barrier & inflammation	
Hexa-LPS Index	
 Hexa-LPS index	76
Oxygen Exposure Index	
 Oxygen exposure index	0.09
Digestive capacity	
Fiber Digestion Capacity	
 Chitin	190 rpkm
 Pectin	80 rpkm
Complex Sugar Digestion Capacity	
 Isomaltooligosaccharides	5.3 rpkm
Protein Breakdown Capacity	
 Branched Chain Amino Acids	1359 rpkm
Vitamin Production Capacity	
 Vitamin B12	771 rpkm

Microbial enzymes & metabolites

Mitochondrial Health Support

 Urolithin-producing species 0.007%

Results Summary

	Metrics	Clinical indication examples
Beneficial microbes	⚠️ Beneficial Bifidobacterium ⚠️ Metabolic Health ✅ Anti-inflammatory Markers ✅ Common Probiotic Species	• Prebiotics / probiotics support • Review Akkermansia promoting supplements • Track and diversify fiber intake • Increase fermented foods • Move and stay physically active
Disruptive microbes	✅ Antibiotic Resistance Signature ⚠️ Opportunistic Pathogens ✅ Potential Stomach Inflammation ✅ Potential Fungal Overgrowth ✅ Parasites and Infection ✅ Methane Production	• Potential anti-microbial support (e.g., herbs) • Prebiotics / probiotics support • Close monitoring post-antibiotics to track recovery • Nutritional support and therapeutic diet (e.g., AIP, candida diet, etc) • Review symptoms/conditions • Consider additional testing (e.g., SIBO, OAT, nutritional labs etc)
Gut barrier & inflammation	⚠️ Hexa-LPS Index ✅ Mucus Degradation Index ✅ Hydrogen Sulfide Index ✅ Host DNA ⚠️ Oxygen Exposure Index	
Short-chain fatty acids	✅ Butyrate Capacity ✅ Propionate Capacity ✅ Acetate Capacity	• Track and diversify fiber intake • Prebiotics / probiotics / postbiotics support • Increase fermented foods • Supportive nutraceuticals (butyrate supplements, spore-based probiotics) • Detoxification supports (calcium d-glucarate, broccoli sprouts, DIM)
Digestive capacity	⚠️ Fiber Digestion Capacity ⚠️ Complex Sugar Digestion Capacity ⚠️ Vitamin Production Capacity ⚠️ Protein Breakdown Capacity	
Diversity & resilience	✅ Microbiome Diversity ✅ Gut Resilience Score ✅ Oral Microbes in the Gut ✅ Microbiome Age ✅ Major Bacterial Phyla ✅ Common Microbiome Members ✅ Gut Ratio	

Microbial enzymes & metabolites	✔ Microbial Histamine Capacity ✔ Complex Compound Breakdown ✔ Microbial GABA Capacity ✔ Modified Bile Acid Production Capacity ⚠ Mitochondrial Health Support
---------------------------------------	--

Detailed Gut Health Test Results

TAXONOMY LEGEND

(p) - Phylum (f) - Family (g) - Genus (sp) - Species

GRAPH INDEX LEGEND

Needs support

Needs improving

Okay

Great

Beneficial microbes

Beneficial microbes support microbiome health and overall health. We generally want to see high levels of these.

Beneficial Bifidobacterium

Bifidobacterium
Your levels are very high. While these bacteria support digestion and gut balance, very high levels may indicate gut imbalance in the overall ecosystem, reduced diversity, or overgrowth driven by diet or supplements. This should be monitored.

0.005%

0.054%

8.296%

7.3%

Improve

Species in your sample:

Bifidobacterium adolescentis

3.881%

Bifidobacterium animalis

4.009%

Bifidobacterium longum

0.339%

Metabolic Health

Akkermansia
We didn't detect *Akkermansia*. This needs support, and we recommend taking action. See your Action Plan.

0%

0.005%

0.62%

2.75%

4%

Support

Anti-inflammatory Markers

Faecalibacterium
Your levels are typical. This is normal, and no action is needed.

3.143%

1.9%

1.1%

6.5%

Okay

Species in your sample:

Faecalibacterium prausnitzii	0.171%
Faecalibacterium prausnitzii_A	0.444%
Faecalibacterium prausnitzii_C	1.016%
Faecalibacterium prausnitzii_D	0.636%
Faecalibacterium prausnitzii_G	0.511%
Faecalibacterium sp900539885	0.100%
Faecalibacterium sp900539945	0.174%



Common Probiotic Species


Bifidobacterium We detected this.	8.296%	Okay
Lactobacillaceae We detected this. This is normal, and no action is needed.	1.061%	Okay
Lacticaseibacillus paracasei We detected this species. That is great! No action is needed.	0.226%	Okay
Lactobacillus acidophilus We detected this species. That is great! No action is needed.	0.054%	Okay
Bifidobacterium infantis We detected this species. That is great! No action is needed.	0.010%	Okay
Bifidobacterium longum We detected this species. That is great! No action is needed.	0.339%	Okay
Bifidobacterium animalis We detected this species. That is great! No action is needed.	4.009%	Okay

Bifidobacterium adolescentis We detected this species. That is great! No action is needed.	3.881%	Okay
Streptococcus thermophilus We detected this species. That is great! No action is needed.	0.081%	Okay

Disruptive microbes

Disruptive microbes include pathogens or bad actors that hurt gut health. Low or absent levels are ideal.

Antibiotic Resistance Signature		
Abundance index This index is typical. This is normal, and no action is needed.		Okay
Richness index This index is typical. This is normal, and no action is needed.		Okay


Opportunistic Pathogens		
Enterobacteriaceae Your levels are typical. This is normal, and no action is needed.		Okay
Species in your sample: Citrobacter braakii 0.189% Enterobacter hormaechei_A 0.064% Escherichia flexneri 0.071% Klebsiella pneumoniae 0.127% Klebsiella_A michiganensis 0.052%		

Klebsiella Your levels are borderline high. This could be improved, and we recommend monitoring or taking optional next steps.		Improve
Species in your sample: Klebsiella pneumoniae 0.127% Klebsiella_A michiganensis 0.052%		

Klebsiella pneumoniae Your levels are high. This needs support, and we recommend taking action. See your Action Plan.	0.127% 0.005% Support
Klebsiella oxytoca We didn't detect <i>K. oxytoca</i> . This is great! No action is needed.	0% 0.005% Okay
Salmonella enterica We didn't detect <i>S. enterica</i> . This is great! No action is needed.	0% 0.005% Okay
Escherichia coli Your levels are typical. This is normal, and no action is needed.	0.019% 0.49%1.1% Okay
Escherichia flexneri Your levels are typical. This is normal, and no action is needed.	0.071% 0.14%0.3% Okay
Escherichia dysenteriae We didn't detect <i>E. dysenteriae</i> . This is great! No action is needed.	0% 0.005%0.015% Okay
Citrobacter Your levels are high. This needs support, and we recommend taking action. See your Action Plan.	0.205% 0.03%0.12% Support
Species in your sample: Citrobacter braakii 0.189%	
Enterobacter Your levels are borderline high. This could be improved, and we recommend monitoring or taking optional next steps.	0.181% 0.062%0.25% Improve
Species in your sample: Enterobacter hormaechei_A 0.064%	
Morganella We didn't detect this. This is great! No action is needed.	0% 0.005% Okay
Raoultella We didn't detect this. This is great! No action is needed.	0% 0.05%0.2% Okay

Streptococcus


You have typical levels. This is normal, and no action is needed.



Okay

Staphylococcus

We didn't detect this. This is great! No action is needed.



Okay

Pseudomonas aeruginosa

We didn't detect this. This is great! No action is needed.



Okay

Haemophilus influenzae


We didn't detect this. This is great! No action is needed.



Okay

Haemophilus parainfluenzae

We detected this at borderline high levels. This could be improved, and we recommend monitoring or taking optional next steps.



Improve

Enterococcus faecium

We didn't detect this. This is great! No action is needed.



Okay

Enterococcus faecalis


We detected high levels. This needs support, and we recommend taking action. See your Action Plan.



Support

Clostridioides difficile

We didn't detect *C. difficile*. This is great! No action is needed.



Okay

Clostridium perfringens

We didn't detect *C. perfringens*. This is great! No action is needed.



Okay

Acinetobacter baumannii

We didn't detect *A. baumannii*. This is great! No action is needed.



Okay

Campylobacter

We didn't detect this. This is great! No action is needed.



Okay

Potential Stomach Inflammation

Helicobacter pylori
We didn't detect this. This is great! No action is needed.

0%

0.005%

Okay

Potential Fungal Overgrowth

Candida
We didn't detect *Candida* in your sample. This is great! No action is needed.

0%

0.005%

Okay

Aspergillus
We did not detect this. This is normal, and no action is needed.

0%

0.005%

Okay

Cryptococcus
We did not detect this. This is normal, and no action is needed.

0%

0.005%

Okay

Saccharomyces
We did not detect this. This is normal, and no action is needed.

0%

0.5%

Okay

Rhodotorula
We did not detect this. This is normal, and no action is needed.

0%

0.005%

Okay

Saprochaete
We did not detect this. This is normal, and no action is needed.

0%

0.005%

Okay

Malassezia
We did not detect this. This is normal, and no action is needed.

0%

0.005%

Okay

Microsporum
We did not detect this. This is normal, and no action is needed.

0%

0.005%

Okay

Trichophyton
We did not detect this. This is normal, and no action is needed.

0%

0.005%

Okay

Parasites and Infection

Blastocystis We didn't detect <i>Blastocystis</i> . This is normal, and no action is needed.	0%  0.005%	Okay
Cryptosporidium We didn't detect this. This is great! No action is needed.	0%  0.005%	Okay
Entamoeba histolytica We didn't detect this. This is great! No action is needed.	0%  0.005%	Okay
Entamoeba dispar We didn't detect this. This is great! No action is needed.	0%  0.005%	Okay
Giardia We didn't detect this. This is great! No action is needed.	0%  0.005%	Okay
Yersinia enterocolitica We didn't detect this. This is great! No action is needed.	0%  0.005%	Okay
Vibrio We didn't detect this. This is great! No action is needed.	0%  0.005%	Okay
Vibrio cholerae We didn't detect this. This is great! No action is needed.	0%  0.005%	Okay
Cyclospora cayetanensis We didn't detect this. This is great! No action is needed.	0% 	Okay

Methane Production

Methanobrevibacter smithii We didn't detect <i>M. smithii</i> . This is normal, and no action is needed.	0%  0.15% 0.44%	Okay
Methane production capacity We did not detect genes for methane production. This is okay.	0.00 rpkm  1	Okay

Gut barrier & inflammation

Gut barrier & inflammation metrics reflect microbial activities that can weaken the gut lining and increase inflammation. Low levels are ideal.

Hexa-LPS Index

Hexa-LPS index

Your hexa-LPS index is borderline high. Too much of this bacterial molecule crossing your gut barrier can drive inflammation linked to a higher risk of multiple health conditions. While not critical, these levels could be improved or monitored.

0.1

76.8

76.6

83.2

Improve

Mucus Degradation Index

Mucus degradation index

Your mucus degradation index is typical. This is normal, and no action is needed.

28.9

41.3

62.1

65.9

Okay

Hydrogen Sulfide Index

Hydrogen sulfide index

Your hydrogen sulfide index is typical. This is normal, and no action is needed.

33

69.0

89.5

92

Okay

Host DNA

Host DNA

Your levels are typical. This is normal, and no action is needed.

0.169%

5%

Okay

Oxygen Exposure Index

Oxygen exposure index

Your microbiome oxygen exposure is borderline high. This could be improved, and we recommend monitoring or taking optional next steps.

0.08

0.09

0.11

Improve


Short-chain fatty acids

Short chain fatty acids regulate immune function and reduce inflammation. They come from fiber in diet. We want to see typical or high levels.

Butyrate Capacity

Butyrate

Your potential to produce butyrate is typical. This is normal, and no action is needed.




Okay

Propionate Capacity

Propionate

Your potential to produce propionate is high. This is great! No action is needed.




Great

Acetate Capacity

Acetate

Your potential to produce acetate is high. This is great! No action is needed.



Great


Digestive capacity

Digestive capacity markers show if your microbes have optimal potential to digest nutrients in your food and to produce vitamins and other compounds.

Fiber Digestion Capacity

Cellulose


Your potential to digest cellulose is typical. This is great! No action is needed.



Great

Resistant starch

Your potential to digest resistant starch is high. This is great! No action is needed.



Great

Improve

Support

Great

Great

Great

Improve


Okay

Great

Great

Ammonia


Your microbiome potential to produce ammonia is typical. This is great! No action is needed.



Okay

Branched Chain Amino Acids


Your gut shows a high potential to produce BCAAs, a pattern that has been associated with insulin resistance and metabolic concerns in adults. This could be improved or monitored.



Improve

p-Cresol

Your microbiome's potential to produce p-Cresol is typical. This is great! No action is needed.



Okay

Vitamin Production Capacity

Vitamin B2


Your microbiome potential to produce vitamin B2 is high. This is great! No action is needed.



Great

Vitamin B7


Your microbiome potential to produce vitamin B7 is typical. This is great! No action is needed.



Great

Vitamin B9


Your microbiome potential to produce vitamin B9 is typical. This is great! No action is needed.



Great

Vitamin B12


Your microbiome potential to produce vitamin B12 is low. This could be improved, and we recommend monitoring or taking optional next steps.



Improve

Vitamin K

Your microbiome potential to produce vitamin K is typical. This is great! No action is needed.



Great

Diversity & resilience

Diversity & resilience metrics tell about the overall community found in your microbiome. More diversity is generally a good thing.

Microbiome Diversity

Shannon diversity
Your diversity is typical. This is normal, and no action is needed.

6.0
5.1
4.8
6.1

Okay

Species richness
Your levels are high. This is great! No action is needed.

425 species
103
92
162

Great

Gut Resilience Score

Gut resilience score
Your score is typical. This is normal, and no action is needed.

0.54
0.46
0.43
0.56

Okay

Oral Microbes in the Gut

Oral microbes
Your levels are typical. This is normal, and no action is needed.

0.046%
0.5%

Okay

Microbiome Age

Microbiome age
Your microbiome age is typical for your age. This is normal, and no action is needed.

36.9 years

Okay

Major Bacterial Phyla

Bacteroidota
Your levels are typical. No action is needed.

28.876%

Okay

Firmicutes
Your child's levels are typical. No action is needed.

56.818%

Okay

Actinobacteriota

10.199%

Your levels are typical. No action is needed.

Okay

Proteobacteria

1.405%

Your levels are typical. No action is needed.

Okay

Common Microbiome Members

Bacteroides

1.884%

Your levels are low. This is normal, and no action is needed.

Okay

Bacteroides fragilis

0%

We didn't detect *B. fragilis*. This is normal, and no action is needed.

Okay

Prevotella

15.255%

Your levels are very high. This could be improved, and we recommend monitoring or taking optional next steps.

Okay

Ruminococcus

5.571%

Your levels are typical. This is great! No action is needed.

Great

Ruminococcus gnavus

0.006%

Your levels are low. This is normal, and no action is needed.

Okay

Blautia

7.029%

Your levels are high. This is great! No action is needed.

Great

Roseburia

0.972%

Your levels are typical. This is great! No action is needed.

Great

Phocaeicola dorei

0.015%

Your levels are low. This is normal, and no action is needed.

Okay

Gut Ratio

Firmicutes / Bacteroidota ratio

2.0

Your levels are typical. This is normal, and no action is needed.

Okay

Proteobacteria / Actinobacteriota ratio

Your levels are typical. This is normal, and no action is needed.

0.14

Okay

Microbial enzymes & metabolites

Microbial enzymes & metabolites are involved in metabolism, detoxification, and inflammation.

Microbial Histamine Capacity

Histamine index

Your histamine index is low. This is great! No action is needed.

1.5

19.5

70.5

76

Great

Complex Compound Breakdown

Beta-glucuronidase capacity

Your microbiome potential to produce beta-glucuronidase is typical. This is good! No action is needed.

1104

1

996

1530

1600

Great

Microbial GABA Capacity

GABA production

Your microbiome potential to produce GABA is typical. This is good! No action is needed.

10.4 rpkm

1

36.7

Okay

GABA breakdown

We didn't detect any genes associated with GABA breakdown. This is great! No action is needed.

0.00 rpkm

1

Great

Modified Bile Acid Production Capacity

Unconjugated bile acids

Your microbiome potential to produce unconjugated bile acids is typical. This is great! No action is needed.

68.0 rpkm

66

112

Great

Secondary bile acids

We didn't detect genes for secondary bile acid production. This is okay, no action is needed.

0.00 rpkm

1


4.6

Okay

Mitochondrial Health Support

Urolithin-producing species


Your levels are low. This needs support, and we recommend taking action. See your Action Plan.




Support

Major Microbial Members & Ratios


Common Bacterial Phyla




Gut Ratio



Gut Type



Bacteroides	6.019%
Ruminococcaceae	8.062%
Prevotella	18.102%
Lachnospiraceae	28.013%
Enterobacteriaceae	0.688%
Bifidobacterium	8.296%
Other	30.382%



These bars represent the proportion of the different gut types in our database for different age ranges.

The gut type of a sample is determined by the dominant type of microbe in that sample.

Type Other

We classified this gut type as 'Other' because it doesn't fit well in the other types. Instead of being dominated by *Bacteroides*, *Prevotella*, *Ruminococcaceae*, *Lachnospiraceae*, or *Bifidobacterium*, the 'Other' gut type has higher levels of a variety of species.

Having this gut type is not a bad thing; in fact, its high diversity may be beneficial for gut health. As microbiome research continues, we may be able to give you more insights about this gut type.

Microbiome Breakdown

- **Beneficial:** These support microbiome health and overall health.
- **Unfriendly:** Low levels of these microbes are expected. Higher levels for extended periods of time can cause issues. This should make up a minority of the microbiome, kept in check by beneficial and variable microbes. Any unfriendly microbes out of range will be flagged in the "Detailed Gut Health Test Results" section above.
- **Variable:** These can be beneficial or unfriendly depending on their levels and what other microbes are around them. Lower levels are usually ok, and higher levels can cause issues.
- **Unknown:** These are lesser known microbes. Not enough is known to classify them as beneficial, unfriendly or variable at this time.

Your breakdown



Top 20 species

Metric	Result	Description
● Prevotella copri_A	13.011%	<i>P. copri</i> is one the most common species in the gut. Its role in health is not quite clear. Some studies have associated <i>P. copri</i> with better blood sugar metabolism and protection from food allergy. However, high numbers of this bug in babies may contribute to development of respiratory allergies later in life. Other research indicates that high amounts of <i>P. copri</i> are linked to rheumatoid arthritis, insulin resistance, and diabetes.
● Agathobacter rectalis	5.465%	<i>A. rectalis</i> is a beneficial bacterium found in the gut of many people. It's a butyrate producer whose levels increase when babies start eating solid foods. According to one study, very low levels of <i>A. rectalis</i> may contribute to inflammatory bowel syndrome. But too much of it may be associated with obesity. So better to keep <i>A. rectalis</i> at that sweet spot. If you need to give it a boost, you can add more fiber to your diet.
● Agathobacter faecis	4.219%	<i>A. faecis</i> digests the fiber you eat and produces butyrate, a beneficial short-chain fatty acid (SCFA). SCFAs keep your gut lining healthy and help reduce inflammation. A good way to boost your levels of <i>A. faecis</i> is to follow a Mediterranean diet.
● Bifidobacterium animalis	4.009%	<i>B. animalis</i> is often detected in the gut of healthy adults, but scientists believe it is just a transient passenger. <i>B. animalis</i> subspecies <i>lactis</i> is frequently used as a probiotic to relieve symptoms of irritable bowel syndrome or diarrhea associated with antibiotics use. In pregnant women, low levels of this bug have been associated with gestational diabetes.

● Bifidobacterium adolescentis	3.881%	<i>B. adolescentis</i> can be passed from mom to baby through vaginal birth or breastfeeding. While it can be found in the baby's gut, it is more common in adults. Why? <i>B. adolescentis</i> prefers to eat carbohydrates typical of an adult diet. In fact, it can help you digest foods high in resistant starch, like oats and beans. Low levels of this bug have been associated with Type 1 diabetes. Another advantage is its production of GABA, a compound that helps us sleep well, have a good mood, and better cognition. Having <i>B. adolescentis</i> could also promote a longer life, as it has been detected in people over 100 years old.
● Ruminococcus_D bicirculans	3.228%	<i>R. bicirculans</i> are common members of the gut microbiome. They contribute to gut health by breaking down fiber and producing acetate, a beneficial short-chain fatty acid. One of their favorite fibers to digest is that of barley.
● Prevotellamassilia timonensis	2.827%	Scientists don't know enough about this species to tell you anything about it. It was detected in the human gut for the first time in 2016. This shows how new microbiome research really is! We will keep updating our information as research advances.
● Zag1 sp001765415	2.544%	
● Dialister hominis	2.528%	
● Phocaeicola plebeius_A	2.214%	<i>P. plebeius</i> can digest the fiber from seaweed. And because of that, this bacterium is usually found in people whose diet is high in seaweed. One study found that people with type 2 diabetes had higher amounts of <i>P. plebeius</i> than healthy people. So if blood sugar levels are a concern, it may be a good idea to not go crazy eating tons of seaweed.
● Coprococcus eutactus_A	1.680%	<i>C. eutactus</i> is a beneficial bacterium that produces butyrate. It is very good at digesting beta-glucans, a type of fiber present in barley, oats, whole grains, seaweed, and some mushrooms. Some studies indicate <i>C. eutactus</i> may be positively associated with language development in children and may contribute to decreasing eczema severity. Also, it may be protective against Type 1 diabetes.
● Phocaeicola vulgatus	1.525%	<i>P. vulgatus</i> is one of the most abundant bacteria in the human gut, detected very early in babies. People with cardiovascular disease have low levels of <i>P. vulgatus</i> which could indicate it is a beneficial bacterium. But women with polycystic ovary syndrome have high levels of it. And <i>P. vulgatus</i> has been associated with Crohn's disease and ulcerative colitis, although the effects depend on the strain. Also, when in high numbers throughout the first two years of life, it may contribute to the development of Type 1 diabetes.
● Fusicatenibacter saccharivorans	1.278%	<i>F. saccharivorans</i> is a beneficial bacterium that produces short-chain fatty acids and promotes the production of anti-inflammatory molecules. People with active ulcerative colitis or rheumatoid arthritis have low levels of <i>F. saccharivorans</i>.

● Alistipes putredinis	1.248%	<i>A. putredinis</i> is a normal member of the gut microbiome whose role seems to vary. It has been shown to protect against the yeast <i>Candida albicans</i> in the gut. In pregnant women, low numbers of this bug have been associated with gestational diabetes. However, one study found that samples of colon cancer had high levels of <i>A. putredinis</i>. Its levels are higher in vegetarians and vegans than in omnivores and its growth is boosted when eating cruciferous vegetables.
● Ruminococcus_E sp900314705	1.186%	We don't know much about this particular species of <i>Ruminococcus</i>. However, <i>Ruminococcus</i> are a diverse group of bacterial species in your gut. Helpful in small numbers, these bugs break down starches and fiber from your food that you can't digest on your own. This gives you and other bacteria in your gut the energy you need. Luckily, you can keep your <i>Ruminococcus</i> bacteria happy by eating plenty of fiber.
● Blautia_A obeum	1.182%	<i>Blautia</i> species contribute to a healthy gut by digesting dietary fiber and producing beneficial and anti-inflammatory molecules. This species may also protect against allergies. A plant-based diet or eating 85% dark chocolate have been shown to increase gut levels of <i>B. obeum</i> in some people.
● Clostridium saudiense	1.142%	
● Blautia_A wexlerae	1.069%	<i>B. wexlerae</i> is one of the most abundant <i>Blautia</i> species in humans. Along with a healthy diet high in fiber, it may contribute to healthy metabolism. High levels of <i>B. wexlerae</i> may protect against obesity, insulin resistance, eczema and inflammation.
● Blautia_A wexlerae_A	1.055%	<i>B. wexlerae</i> is one of the most abundant <i>Blautia</i> species in humans. Along with a healthy diet high in fiber, it may contribute to healthy metabolism. High levels of <i>B. wexlerae</i> may protect against obesity, insulin resistance, eczema and inflammation.
● Faecalibacterium prausnitzii_C	1.016%	Common in many healthy microbiomes, <i>F. prausnitzii</i> is a beneficial species. These bugs are known for curbing inflammation, keeping your gut healthy, and protecting you from disease. These bacteria produce a special molecule called butyrate. It plays a role in supporting immune system function and the structural integrity of your gut. Low <i>F. prausnitzii</i> counts are linked to type 2 diabetes, obesity and a handful of gut-related conditions. All the more reason to show these friendly partners some love.

All species

● Eubacterium_R sp003526845	1.000%	● Blautia_A sp900066335	0.999%
● Barnesiella intestinihominis	0.981%	● Agathobaculum butyriciproducens	0.977%
● Blautia_A massiliensis	0.973%	● Bacteroides ovatus	0.961%
● Alistipes shahii	0.933%	● Gemmiger formicilis	0.824%
● Ligilactobacillus ruminis	0.749%	● UBA11524 sp000437595	0.742%
● Anaerobutyricum soehngenii	0.733%	● Romboutsia timonensis	0.730%
● Bariatricus comes	0.725%	● Gemmiger qucibialis	0.670%
● KLE1615 sp900066985	0.667%	● Prevotella copri	0.641%
● Faecalibacterium prausnitzii_D	0.636%	● Dorea_A longicatena	0.558%
● Turicibacter sp001543345	0.557%	● Faecalibacterium prausnitzii_G	0.511%
● Sutterella wadsworthensis	0.508%	● Roseburia inulinivorans	0.499%
● Paraprevotella clara	0.446%	● Faecalibacterium prausnitzii_A	0.444%
● Prevotella sp900557255	0.423%	● Anaerobutyricum hallii	0.409%
● Blautia_A sp003480185	0.384%	● Odoribacter splanchnicus	0.380%
● Ruminococcus_C sp000437255	0.375%	● Blautia_A sp000436615	0.361%
● Bacteroides uniformis	0.353%	● Lachnospira eligens_A	0.346%
● Intestinibacter bartlettii	0.343%	● Lachnospira sp000437735	0.340%
● Bifidobacterium longum	0.339%	● Acetatifactor sp003447295	0.325%
● Ruminococcus_C sp900545285	0.325%	● CAG-177 sp003514385	0.323%
● Roseburia sp900552665	0.310%	● Faecalibacillus faecis	0.307%
● Acetatifactor sp900066565	0.302%	● Anaerostipes hadrus_A	0.295%
● Acetatifactor sp900066365	0.288%	● Dorea_A longicatena_B	0.286%
● Coprococcus sp900066115	0.269%	● Bacteroides eggerthii	0.262%
● UBA737 sp900543215	0.252%	● Prevotella sp900557035	0.251%
● Agathobacter sp900546625	0.249%	● Ruminococcus_A sp003011855	0.247%
● Coprococcus_A catus	0.241%	● Anaerostipes hadrus	0.239%
● CAG-110 sp003525905	0.238%	● UBA7597 sp900542935	0.228%
● Lacticaseibacillus paracasei	0.226%	● Alistipes communis	0.225%
● CAG-83 sp000435555	0.224%	● CAG-41 sp900066215	0.222%

● Agathobaculum sp900544475	0.221%	● Prevotella sp900546535	0.216%
● CAG-841 sp900544285	0.213%	● Eubacterium_F sp003491505	0.212%
● CAG-245 sp000435175	0.209%	● Mediterraneibacter faecis	0.197%
● Blautia_A schinkii	0.192%	● Blautia_A luti	0.189%
● Citrobacter braakii	0.189%	● Lactococcus lactis	0.186%
● Dorea formicigenerans	0.183%	● Lachnospira sp000436535	0.181%
● Butyricimonas paravirosa	0.180%	● Agathobacter sp900317585	0.174%
● Faecalibacterium sp900539945	0.174%	● CAG-81 sp900066785	0.173%
● ER4 sp000765235	0.171%	● Faecalibacterium prausnitzii	0.171%
● Prevotella hominis	0.170%	● CAG-110 sp000434635	0.168%
● Parabacteroides distasonis	0.157%	● Blautia_A faecis	0.156%
● CAG-1193 sp900550145	0.145%	● Collinsella sp003487125	0.143%
● CAG-288 sp000437395	0.142%	● Collinsella sp002232035	0.142%
● UBA9502 sp003480315	0.140%	● Klebsiella pneumoniae	0.127%
● CAG-81 sp900066535	0.125%	● GCA-900066135 sp900543575	0.124%
● Eubacterium_G ventriosum	0.119%	● UMGS1820 sp900545865	0.119%
● Eisenbergiella sp900066775	0.117%	● Bacteroides xylanisolvens	0.114%
● Bacteroides sp902388495	0.111%	● TF01-11 sp003524945	0.108%
● Eubacterium_I ramulus_A	0.107%	● Collinsella sp900541645	0.105%
● CAG-1427 sp000436075	0.103%	● Enterocloster sp900541315	0.102%
● Faecalibacterium sp900539885	0.100%	● Prevotella sp900544825	0.100%
● Blautia_A sp000285855	0.099%	● CAG-390 sp000437015	0.099%
● PeH17 sp000435055	0.097%	● CAG-83 sp003487665	0.093%
● Blautia_A sp900120195	0.090%	● ER4 sp900317525	0.088%
● UMGS1375 sp900066615	0.088%	● Blautia_A sp900066145	0.087%
● Gemmiger sp900539695	0.087%	● Roseburia hominis	0.086%
● UBA9502 sp003478505	0.084%	● Phocaeicola plebeius	0.082%
● Clostridium_Q sp003024715	0.081%	● Prevotella sp900551985	0.081%
● Streptococcus thermophilus	0.081%	● Bilophila wadsworthia	0.080%
● CAG-273 sp000437855	0.080%	● Faecalibacillus intestinalis	0.079%
● Hungatella effluvii	0.079%	● Alistipes_A indistinctus	0.076%

● Dysosmobacter sp001916835	0.076%	● Megasphaera sp000417505	0.076%
● CAG-826 sp000437235	0.074%	● Lactococcus lactis_E	0.074%
● Mesosutterella multiformis	0.073%	● CAG-45 sp000438375	0.072%
● Alistipes onderdonkii	0.071%	● Escherichia flexneri	0.071%
● Phocaeicola sp900553715	0.070%	● Dysosmobacter sp900550685	0.068%
● Ruminococcus_A faecicola	0.067%	● Evtepia gabavorous	0.066%
● QAMH01 sp003149935	0.066%	● Enterobacter hormaechei_A	0.064%
● Collinsella sp003439125	0.063%	● UBA3402 sp003478355	0.063%
● CAG-170 sp000432135	0.062%	● Adlercreutzia celatus_A	0.060%
● Anaerostipes hadrus_B	0.059%	● CAG-582 sp000435515	0.058%
● Lachnoclostridium_B sp900066555	0.057%	● Prevotella sp900767615	0.057%
● Prevotella disiens	0.056%	● UMGS874 sp900546315	0.056%
● CAG-110 sp900544945	0.055%	● Lactobacillus acidophilus	0.054%
● UBA11475 sp003538975	0.054%	● Erysipelatoclostridium ramosum	0.053%
● Intestinibacter sp900553485	0.052%	● Klebsiella_A michiganensis	0.052%
● Lachnospira sp900316325	0.051%	● Collinsella sp900541145	0.049%
● Prevotella sp900551275	0.049%	● Enterobacter ludwigii	0.048%
● Anaerobutyricum sp900554965	0.047%	● Collinsella sp003436275	0.047%
● Eubacterium_I ramulus	0.047%	● UBA7182 sp003481535	0.047%
● Collinsella sp003459245	0.046%	● Collinsella sp003465825	0.046%
● UBA737 sp002451855	0.046%	● Blautia_A sp900066355	0.044%
● CAG-485 sp900549375	0.043%	● SFFH01 sp900542445	0.043%
● Collinsella sp003469205	0.042%	● Prevotella sp900554835	0.042%
● CAG-83 sp900547745	0.041%	● Roseburia intestinalis	0.039%
● CAG-110 sp900546075	0.038%	● Paraprevotella xylaniphila	0.038%
● Agathobacter sp900550845	0.037%	● CAG-170 sp003516765	0.037%
● Ruminococcus_E sp003526955	0.037%	● UBA737 sp900549055	0.037%
● Blautia_A sp003471165	0.036%	● Bacteroides thetaiotaomicron	0.035%
● Clostridium sp000435835	0.035%	● Collinsella sp900549335	0.035%
● Olsenella_B sp900768455	0.035%	● Blautia_A sp900066205	0.034%
● CAG-628 sp000438415	0.034%	● Clostridium_A leptum	0.034%

● Dysosmobacter welbionis	0.034%	● Anaerosacchariphilus sp900066385	0.033%
● Prevotella pectinovora	0.033%	● Collinsella sp900551015	0.032%
● Phil1 sp001940855	0.032%	● Ruminococcus_C callidus	0.032%
● Adlercreutzia equolifaciens	0.031%	● Alistipes_A sp900240235	0.031%
● CAG-452 sp000434035	0.030%	● Clostridium sp900543325	0.030%
● Lachnospira sp003451515	0.030%	● Lawsonibacter asaccharolyticus	0.030%
● CAG-110 sp000435995	0.029%	● Enterococcus faecalis	0.029%
● Enterococcus_B lactis	0.029%	● Ruminiclostridium_E siraeum	0.029%
● CAG-103 sp000432375	0.028%	● Angelakisella sp900547385	0.027%
● Ruthenibacterium lactatiformans	0.027%	● SFFH01 sp900548125	0.027%
● Agathobacter sp900552085	0.026%	● Buttiauxella noackiae	0.026%
● UBA10281 sp900767815	0.026%	● UMGS1071 sp900542375	0.026%
● Dialister sp900555245	0.025%	● Streptococcus salivarius	0.025%
● UBA738 sp003522945	0.025%	● Blautia_A sp003477525	0.024%
● Collinsella aerofaciens_H	0.024%	● Collinsella sp003458415	0.024%
● Dialister sp000434475	0.024%	● Eubacterium_F sp000433735	0.024%
● Agathobacter sp900547695	0.023%	● NK3B98 sp900758315	0.023%
● Roseburia sp900542495	0.023%	● Collinsella sp900554595	0.022%
● Phocaeicola sp900557085	0.022%	● Bifidobacterium dentium	0.021%
● Butyrivibrio_A crossotus	0.021%	● Collinsella sp900542825	0.021%
● Collinsella sp900542945	0.021%	● Enterobacter quasiormaechei	0.021%
● RUG115 sp900066395	0.021%	● Agathobaculum sp003481705	0.020%
● Clostridium sp001916075	0.020%	● Collinsella sp900544465	0.020%
● Dysosmobacter sp014297375	0.020%	● Enterobacter bugandensis	0.020%
● GCA-900066135 sp900066135	0.020%	● Kosakonia cowanii	0.020%
● Ruminococcus_E bromii_B	0.020%	● Coprococcus eutactus	0.019%
● Dysosmobacter sp900542115	0.019%	● Escherichia coli_D	0.019%
● Flavonifractor plautii	0.019%	● Merdibacter sp900543035	0.019%
● TF01-11 sp003529475	0.019%	● Chitinophaga sp003412465	0.018%
● Collinsella aerofaciens_N	0.018%	● Collinsella sp900546115	0.018%
● Collinsella sp900551195	0.018%	● UBA1191 sp900549125	0.018%

● CAG-170 sp002404795	0.017%	● CAG-460 sp900544625	0.017%
● Collinsella sp900552425	0.017%	● Faecalibacterium prausnitzii_J	0.017%
● Gemmiger sp900540595	0.017%	● RUG013 sp001486445	0.017%
● UBA7597 sp003448195	0.017%	● Collinsella sp900549245	0.016%
● Collinsella sp900553935	0.016%	● Eubacterium_G sp900552275	0.016%
● Haemophilus_D parainfluenzae_K	0.016%	● Lawsonibacter sp014287875	0.016%
● UMGS1600 sp900553315	0.016%	● UMGS1670 sp900553995	0.016%
● CAG-170 sp900545925	0.015%	● CAG-180 sp000432435	0.015%
● Collinsella sp003437035	0.015%	● Collinsella sp900548565	0.015%
● Phocaeicola dorei	0.015%	● Ruminococcus_C sp000980705	0.015%
● Schaedlerella sp900066545	0.015%	● UBA1777 sp900547315	0.015%
● Allisonella histaminiformans	0.014%	● Blautia_A obeum_B	0.014%
● Collinsella sp900544115	0.014%	● Collinsella sp900547765	0.014%
● Lachnospira sp003537285	0.014%	● Limosilactobacillus mucosae	0.014%
● Prevotella sp900290275	0.014%	● Prevotella sp900313215	0.014%
● QALS01 sp003150575	0.014%	● 1XD42-69 sp014287635	0.013%
● CAG-81 sp900066055	0.013%	● Citrobacter freundii	0.013%
● Clostridium cuniculi	0.013%	● Coprococcus_A sp900548825	0.013%
● UBA9502 sp003506385	0.013%	● Anaerostipes sp900066705	0.012%
● Bifidobacterium pseudocatenulatum	0.012%	● Collinsella aerofaciens_A	0.012%
● ER4 sp003522105	0.012%	● Faecalibacterium prausnitzii_E	0.012%
● Gemmiger sp900540775	0.012%	● Phocaeicola coprocola	0.012%
● Bacillus_A luti	0.011%	● Collinsella sp900542085	0.011%
● Enterocloster sp000431375	0.011%	● Enterocloster sp900543885	0.011%
● Flavonifractor sp000508885	0.011%	● Gemmiger variabilis_B	0.011%
● Prevotella sp900548535	0.011%	● UBA11774 sp003507655	0.011%
● Anaerosaccharophilus sp014290175	0.010%	● Bifidobacterium infantis	0.010%
● Blautia_A sp003474435	0.010%	● CAG-177 sp003538135	0.010%
● CAG-95 sp900066375	0.010%	● Clostridium disporicum_A	0.010%
● Collinsella sp900542235	0.010%	● Collinsella sp900548255	0.010%
● Collinsella sp900756725	0.010%	● Collinsella sp900761995	0.010%

● Collinsella sp900767675	0.010%	● Eubacterium_R sp000436835	0.010%
● Faecalibacterium sp900758465	0.010%	● Haemophilus_D pittmaniae	0.010%
● Olsenella_B sp000752675	0.010%	● Prevotella sp900550035	0.010%
● Pseudoruminococcus massiliensis	0.010%	● UMGS1217 sp900549625	0.010%

Our high confidence detection limit is 0.01% abundance. Very low abundance organisms can be important members of the microbiome but also very difficult to accurately detect. Detections below 0.01% abundance have an increased risk of being false positives. However, in our testing, we have observed that most false positive identifications are closely related to a detected species.

● Bacteroides caccae	0.009%	● Bifidobacterium catenulatum	0.009%
● Birpovirus hiberniae	0.009%	● Butyrivibrio_A sp900543865	0.009%
● CAG-83 sp900549395	0.009%	● CAG-841 sp002405565	0.009%
● Clostridium_AQ innocuum	0.009%	● Collinsella aerofaciens_E	0.009%
● Collinsella sp900540995	0.009%	● Collinsella sp900552705	0.009%
● Collinsella sp900757205	0.009%	● Collinsella sp902388345	0.009%
● Eubacterium_R sp000434995	0.009%	● Faecalibacterium prausnitzii_F	0.009%
● Faecalibacterium sp900765705	0.009%	● Haemophilus_D parainfluenzae	0.009%
● Lachnospira sp900552795	0.009%	● QAMH01 sp900544245	0.009%
● Romboutsia ilealis	0.009%	● Roseburia sp003470905	0.009%
● Victivallis sp002998355	0.009%	● Blautia_A wexlerae_B	0.008%
● Collinsella sp003471585	0.008%	● Collinsella sp900541125	0.008%
● Collinsella sp900543515	0.008%	● Collinsella sp900549275	0.008%
● Collinsella sp900551205	0.008%	● Collinsella sp900556365	0.008%
● Collinsella sp900762325	0.008%	● Dialister sp900543455	0.008%
● Eubacterium_G sp000434315	0.008%	● Holdemanella massiliensis	0.008%
● Mediterraneibacter_A butyricigenes	0.008%	● Prevotella buccalis	0.008%
● Prevotellamassilia sp900541575	0.008%	● Ruminococcus_C sp000433635	0.008%
● TF01-11 sp000436755	0.008%	● UBA2821 sp902786515	0.008%
● UMGS1387 sp900551295	0.008%	● Acutalibacter sp900543555	0.007%
● CAG-841 sp002479075	0.007%	● Collinsella aerofaciens_F	0.007%
● Collinsella sp900540855	0.007%	● Collinsella sp900541745	0.007%
● Collinsella sp900544755	0.007%	● Collinsella sp900759335	0.007%
● Coprococcus sp000154245	0.007%	● Enterobacter cancerogenus	0.007%

● Enterobacter kobei	0.007%	● Enterocloster citroniae	0.007%
● F23-B02 sp001916715	0.007%	● Faecalibacterium prausnitzii_H	0.007%
● Faecalibacterium sp003449675	0.007%	● Haemophilus_D sp001815355	0.007%
● Kluyvera cryocrescens	0.007%	● Lawsonibacter sp900549405	0.007%
● Pauljensenia sp000278725	0.007%	● Prevotella bryantii	0.007%
● Ruminococcus sp900545125	0.007%	● Turicibacter sanguinis	0.007%
● UBA11452 sp003526375	0.007%	● UBA1390 sp002305315	0.007%
● UBA737 sp002431945	0.007%	● UMGS1603 sp900553265	0.007%
● UMGS1633 sp900553645	0.007%	● Agathobacter sp900550545	0.006%
● Alistipes sp900549305	0.006%	● Anaerosacchariphilus sp900553635	0.006%
● Bacteroides sp900765805	0.006%	● Bariatricus comes_A	0.006%
● Bifidobacterium ruminantium	0.006%	● Blautia stercoris	0.006%
● Blautia_A sp900548245	0.006%	● CAG-273 sp003507395	0.006%
● CAG-826 sp002477825	0.006%	● CAG-841 sp000437375	0.006%
● Collinsella sp003462685	0.006%	● Collinsella sp900542325	0.006%
● Collinsella sp900758375	0.006%	● Collinsella sp900768435	0.006%
● Faecalimonas phoceensis	0.006%	● Lachnospira sp000436475	0.006%
● Lactobacillus delbrueckii	0.006%	● OF09-33XD sp003481995	0.006%
● Oxalobacter sp900760095	0.006%	● Pauljensenia sp902373545	0.006%
● Ruminococcus_B gnavus	0.006%	● TF01-11 sp001414325	0.006%
● UBA7597 sp002474405	0.006%	● UMGS995 sp900547465	0.006%
● Zag111 sp002102825	0.006%		

References

- [1] A. Uzan-Yulzari et al., "Neonatal antibiotic exposure impairs child growth during the first six years of life by perturbing intestinal microbial colonization," *Nat Commun*, vol. 12, no. 1, p. 443, 2021, doi: 10.1038/s41467-020-20495-4.
- [2] K. E. Fujimura et al., "Neonatal gut microbiota associates with childhood multisensitized atopy and T cell differentiation," *Nat Med*, vol. 22, no. 10, pp. 1187–1191, 2016, doi: 10.1038/nm.4176.
- [3] M. B. Azad et al., "Infant gut microbiota and food sensitization: associations in the first year of life," *Clin Exp Allergy*, vol. 45, no. 3, pp. 632–643, 2015, doi: 10.1111/cea.12487.
- [4] H. M. Tun et al., "Roles of Birth Mode and Infant Gut Microbiota in Intergenerational Transmission of Overweight and Obesity From Mother to Offspring," *Jama Pediatr*, vol. 172, no. 4, p. 368, 2018, doi: 10.1001/jamapediatrics.2017.5535.
- [5] T. Vatanen et al., "The human gut microbiome in early-onset type 1 diabetes from the TEDDY study," *Nature*, vol. 562, no. 7728, pp. 589–594, 2018, doi: 10.1038/s41586-018-0620-2.
- [6] C. J. Stewart et al., "Temporal development of the gut microbiome in early childhood from the TEDDY study," *Nature*, vol. 562, no. 7728, pp. 583–588, 2018, doi: 10.1038/s41586-018-0617-x.
- [7] S. Tamburini, N. Shen, H. C. Wu, and J. C. Clemente, "The microbiome in early life: implications for health outcomes," *Nat Med*, vol. 22, no. 7, pp. 713–722, 2016, doi: 10.1038/nm.4142.
- [8] M. J. Ege, "The Hygiene Hypothesis in the Age of the Microbiome," *Ann Am Thorac Soc*, vol. 14, no. Supplement_5, pp. S348–S353, 2017, doi: 10.1513/annalsats.201702-139aw.
- [9] B. M. Henrick et al., "Bifidobacteria-mediated immune system imprinting early in life," *Cell*, 2021, doi: 10.1016/j.cell.2021.05.030.
- [10] A. Marcobal et al., "Bacteroides in the Infant Gut Consume Milk Oligosaccharides via Mucus-Utilization Pathways," *Cell Host Microbe*, vol. 10, no. 5, pp. 507–514, 2011, doi: 10.1016/j.chom.2011.10.007.
- [11] T. Wilmski et al., "Gut microbiome pattern reflects healthy ageing and predicts survival in humans," *Nat Metabolism*, vol. 3, no. 2, pp. 274–286, 2021, doi: 10.1038/s42255-021-00348-0.
- [12] M. A. Jackson et al., "Signatures of early frailty in the gut microbiota," *Genome Med*, vol. 8, no. 1, p. 8, 2016, doi: 10.1186/s13073-016-0262-7.
- [13] J. M. Pickard, M. Y. Zeng, R. Caruso, and G. Núñez, "Gut microbiota: Role in pathogen colonization, immune responses, and inflammatory disease," *Immunol Rev*, vol. 279, no. 1, pp. 70–89, 2017, doi: 10.1111/imr.12567.
- [14] O. Manor et al., "Health and disease markers correlate with gut microbiome composition across thousands of people," *Nat Commun*, vol. 11, no. 1, p. 5206, 2020, doi: 10.1038/s41467-020-18871-1.
- [15] M. Scudellari, "Cleaning up the hygiene hypothesis," *Proc National Acad Sci*, vol. 114, no. 7, pp. 1433–1436, 2017, doi: 10.1073/pnas.1700688114.
- [16] R. Debray, R. A. Herbert, A. L. Jaffe, A. Crits-Christoph, M. E. Power, and B. Koskella, "Priority effects in microbiome assembly," *Nat Rev Microbiol*, vol. 20, no. 2, pp. 109–121, 2022, doi: 10.1038/s41579-021-00604-w.
- [17] R. Debray, R. A. Herbert, A. L. Jaffe, A. Crits-Christoph, M. E. Power, and B. Koskella, "Priority effects in microbiome assembly," *Nat Rev Microbiol*, vol. 20, no. 2, pp. 109–121, 2022, doi: 10.1038/s41579-021-00604-w.
- [18] C. Huttenhower et al., "Structure, function and diversity of the healthy human microbiome," *Nature*, vol. 486, no. 7402, pp. 207–214, 2012, doi: 10.1038/nature11234.
- [19] P. I. Costea et al., "Enterotypes in the landscape of gut microbial community composition," *Nat Microbiol*, vol. 3, no. 1, pp. 8–16, 2018, doi: 10.1038/s41564-017-0072-8.
- [20] W. E. Anthony et al., "Acute and persistent effects of commonly used antibiotics on the gut microbiome and resistome in healthy adults," *Cell Reports*, vol. 39, no. 2, p. 110649, 2022, doi: 10.1016/j.celrep.2022.110649.
- [21] A. Palleja et al., "Recovery of gut microbiota of healthy adults following antibiotic exposure," *Nat Microbiol*, vol. 3, no. 11, pp. 1255–1265, 2018, doi: 10.1038/s41564-018-0257-9.
- [22] A. Bell et al., "Elucidation of a sialic acid metabolism pathway in mucus-foraging *Ruminococcus gnavus* unravels mechanisms of bacterial adaptation to the gut," *Nat Microbiol*, vol. 4, no. 12, pp. 2393–2404, 2019, doi: 10.1038/s41564-019-0590-7.
- [23] M. T. Henke, D. J. Kenny, C. D. Cassilly, H. Vlamakis, R. J. Xavier, and J. Clardy, "Ruminococcus gnavus, a member of the human gut microbiome associated with Crohn's disease, produces an inflammatory polysaccharide," *Proc National Acad Sci*, vol. 116, no. 26, pp. 12672–12677, 2019, doi: 10.1073/pnas.1904099116.
- [24] W. Li et al., "Ecological and network analyses identify four microbial species with potential significance for the diagnosis/treatment of ulcerative colitis (UC)," *Bmc Microbiol*, vol. 21, no. 1, p. 138, 2021, doi: 10.1186/s12866-021-02201-6.
- [25] F. Blachier, M. Beaumont, and E. Kim, "Cysteine-derived hydrogen sulfide and gut health," *Curr Opin Clin Nutr*, vol. 22, no. 1, pp. 68–75, 2019, doi: 10.1097/mco.0000000000000526.
- [26] S. C. Peck, K. Denger, A. Burrichter, S. M. Irwin, E. P. Balskus, and D. Schleheck, "A glycol radical enzyme enables hydrogen sulfide production by the human intestinal bacterium *Bifidobacterium wadsworthia*," *Proc National Acad Sci*, vol. 116, no. 8, pp. 3171–3176, 2019, doi: 10.1073/pnas.1815661116.
- [27] H. M. HAMER, D. JONKERS, K. VENEMA, S. VANHOUTVIN, F. J. TROOST, and R. -J. BRUMMER, "Review article: the role of butyrate on colonic function," *Aliment Pharm Therap*, vol. 27, no. 2, pp. 104–119, 2008, doi: 10.1111/j.1365-2036.2007.03562.x.
- [28] Y. Furusawa et al., "Commensal microbe-derived butyrate induces the differentiation of colonic regulatory T cells," *Nature*, vol. 504, no. 7480, pp. 446–450, 2013, doi: 10.1038/nature12721.
- [29] H. Sokol et al., "Faecalibacterium prausnitzii is an anti-inflammatory commensal bacterium identified by gut microbiota analysis of Crohn disease patients," *Proc National Acad Sci*, vol. 105, no. 43, pp. 16731–16736, 2008, doi: 10.1073/pnas.0804812105.
- [30] D. Ríos-Covián, P. Ruas-Madiedo, A. Margolles, M. Gueimonde, C. G. de los Reyes-Gavilán, and N. Salazar, "Intestinal Short Chain Fatty Acids and their Link with Diet and Human Health," *Front Microbiol*, vol. 7, p. 185, 2016, doi: 10.3389/fmicb.2016.00185.
- [31] Q. Zhou et al., "Association Between Gut Akkermansia and Metabolic Syndrome is Dose-Dependent and Affected by Microbial Interactions: A Cross-Sectional Study," *Diabetes Metabolic Syndrome Obes Targets Ther*, vol. 14, pp. 2177–2188, 2021, doi: 10.2147/dmso.s311388.