

Imaging Approaches to Mesenteric Ischemia

Frandics Chan, M.D., Ph.D.



Stanford University Medical Center



Lucile Packard Children's Hospital

Mesenteric Ischemia

- Acute type (90%)
 - Arterial embolism
 - Arterial thrombosis
 - Nonocclusive form
 - Low flow state
 - Vasospasm
 - Venous occlusion
- Chronic type (10%)



SUMC



LPCH

Department of Radiology

Acute Mesenteric Ischemia

- Increasing incident
- Nonspecific clinical presentation
- Majority from arterial embolism
- 60% mortality
- Early intervention critical
 - Vasodilator (Papaverine)
 - Thrombolysis, thrombectomy, angioplasty, stenting
 - Infarct bowel resection, revascularization



SUMC



LPCH

Department of Radiology

Chronic Mesenteric Ischemia

- Atherosclerosis (95%)
- Proximal vessel stenoses
- Two or more vessels involved
- Collateral arteries
- Clinical triad
 - Bowel angina
 - Food aversion
 - Weight loss
- Treatment: revascularization, stenting



SUMC



LPCH

Department of Radiology

Diagnostic Test

- Acute Mesenteric Ischemia
 - CT Angiography
 - Catheter Angiography + Vasodilator Tx
- Chronic Mesenteric Ischemia
 - CTA and MRA
 - Physiologic Challenge Test





SUMC



LPCH

Department of Radiology

Acute SMA Thrombosis





SUMC



LPCH

Department of Radiology

Acute Aortic Dissection




SUMC



LPCH

Department of Radiology

Acute SMV Thrombosis



SUMC



LPCH

Department of Radiology

CT Diagnosis in AMI

- Menke J. Diagnostic Accuracy of Multidetector CT in Acute Mesenteric Ischemia: Systematic Review and Meta-Analysis. RSNA 2010.
- 1996 – 2009
- 6 studies: 3 prospective, 3 retrospective
- Pooled sensitivity: 93%
- Pooled specificity: 96%



SUMC



LPCH

Department of Radiology

Diagnostic Test

- Acute Mesenteric Ischemia
 - CT Angiography
 - Catheter Angiography + Vasodilator Tx
- Chronic Mesenteric Ischemia
 - CTA and MRA
 - Physiologic Challenge Test



SUMC




LPCH


Department of Radiology

Normal Response

Fasting
State



Fed
State



SUMC




LPCH


Department of Radiology

Ischemic Response

Fasting
State



Fed
State



SUMC




LPCH

Department of Radiology

Normal Flow Pathway

Fasting State



SUMC




LPCH

Department of Radiology

Normal Flow Pathway

Fed State



SUMC




LPCH

Department of Radiology

Compensated SMA Stenosis

Fed State




SUMC



LPCH

Department of Radiology

Mesenteric Collateral Arteries



SUMC




LPCH

Department of Radiology

Chronic Mesenteric Ischemia

Fed State



SUMC



LPCH

Department of Radiology

Perfusion Measurement

- Velocity
 - Doppler ultrasound at
 - SMA
 - SMV
- Flow
 - MRI phase contrast at
 - SMA
 - SMV



SUMC




LPCH


Department of Radiology

Doppler US of SMA

Before
Meal



After
Meal



SUMC




LPCH

Department of Radiology

Doppler US of SMV

Before
Meal



After
Meal



SUMC




LPCH


Department of Radiology

Flow Area

Venous flow increases by increase in cross-sectional area.



Pre-meal



Post-meal



SUMC



LPCH

Department of Radiology

Perfusion Measurement

- Velocity
 - Doppler ultrasound at
 - SMA
 - SMV
- Flow
 - MRI phase contrast at
 - SMA
 - SMV




SUMC



LPCH

Department of Radiology




SUMC



LPCH

Department of Radiology

Pre and Postprandial Flow



SUMC



LPCH

Department of Radiology

Mesenteric Ischemia Protocol

- Fasting > 6 hrs
- Localize SMV
- Pre-meal PC
- 260 cal diet supplement
- **CE MRA**
- 30 min post meal
- Localize SMV
- Post-meal PC
- Flow Quantification




SUMC



LPCH

Department of Radiology

Normal Study



> 50% increase in
post-meal SMV flow




SUMC



LPCH

Department of Radiology

Chronic Mesenteric Ischemia



< 50% increase in
post-meal SMV flow



SUMC



LPCH

Department of Radiology

Limitations

- CE MRA is limited to proximal stenosis in the mesenteric arteries.
- Physiologic challenge is not sensitive to small segmental ischemia.
- Problems with inadequate fasting, poor GI motility, abnormal endocrine response.
- Physiologic challenge is not appropriate for acute mesenteric ischemia.



SUMC



LPCH

Department of Radiology

