

Chronic Mesenteric Ischemia: Diagnosis and Management

Christopher J. White*

Department of Cardiovascular Diseases, The John Ochsner Heart & Vascular Institute, Ochsner Clinic Foundation, New Orleans, LA 70121

Abstract

Chronic mesenteric ischemia (CMI) is the most common vascular disorder involving the intestines, however it is unusual in clinical practice. The redundancy of the visceral circulation with multiple interconnections between the superior mesenteric artery (SMA) and the inferior mesenteric artery (IMA) is the most likely explanation for the infrequent occurrence of CMI in clinical practice. Atherosclerosis is by the far the most common etiology of CMI. The increased utilization of diagnostic abdominal cross-sectional imaging has increased the recognition of atherosclerotic mesenteric stenoses. CMI is a clinical diagnosis, based upon symptoms and consistent anatomic findings. The classic setting for CMI is a female patient presenting with post-prandial abdominal discomfort that results in significant weight loss. Endovascular therapy with stenting has become the most common method chosen for revascularization having replaced open surgery with its associated morbidity and mortality. (Prog Cardiovasc Dis 2011;54:36-40)

© 2011 Elsevier Inc. All rights reserved.

Keywords:

Chronic mesenteric ischemia; Mesenteric stenosis; Mesenteric artery disease; Revascularization; Visceral ischemia; Mesenteric stent

Chronic mesenteric ischemia (CMI) is the most common vascular disorder involving the intestines; however, it is unusual in clinical practice. Atherosclerosis is by the far the most common etiology of CMI, whereas other etiologies associated with this uncommon syndrome include fibromuscular dysplasia, Buerger disease, and aortic dissection. With increased utilization of cross-sectional imaging (magnetic resonance angiography [MRA], and computerized tomographic angiography [CTA]) of the abdomen, finding atherosclerotic disease of the aorta with associated aorto-ostial stenosis of the visceral vessels becomes a common occurrence and remains an infrequent clinical problem.

A population-based prevalence study of mesenteric artery stenosis of 553 healthy Medicare beneficiaries was performed using abdominal duplex ultrasonography to

screen for evidence of mesenteric disease.¹ Severe (>50% diameter stenosis) stenosis of a mesenteric artery was detected in 17.5% of the total cohort, the great majority (>97%) of which were isolated to the celiac artery. There was no correlation with age, race, sex, or body mass index and the presence of mesenteric artery stenosis. Only 1.3% of the patients had involvement of more than one mesenteric artery.

Another natural history study reported on a group of 980 asymptomatic patients with mesenteric ischemia who were followed clinically.² Only 3 patients eventually developed symptoms, and they had severe stenosis of all 3 mesenteric vessels. The redundancy of the visceral circulation with multiple interconnections between the superior mesenteric artery (SMA) and the inferior mesenteric artery (IMA) is the most likely explanation for the infrequent occurrence of CMI in clinical practice.

Statement of Conflict of Interest: see page 40.

* Address reprint requests to Christopher J. White, MD, Chairman, Department of Cardiovascular Diseases, The John Ochsner Heart & Vascular Institute, Ochsner Clinic Foundation, 1514 Jefferson Highway, New Orleans, LA 70121.

E-mail address: cwhite@ochsner.org (C.J. White).

Clinical presentation

Women present for treatment of CMI much more commonly (70%) than men. The classic symptom for

Abbreviations and Acronyms	
CMI	= chronic mesenteric ischemia
CTA	= computerized tomographic angiography
IMA	= inferior mesenteric artery
MRA	= magnetic resonance angiography
SMA	= superior mesenteric artery

CMI is post-prandial abdominal discomfort that results in significant weight loss (Fig 1). The abdominal discomfort associated with eating causes patients to avoid food (food fear), and they typically lose weight. When a patient presents for consideration of CMI and has not had significant weight loss, consideration should be given

to alternate diagnoses including functional bowel complaints. Patients with ischemic gastropathy may also present with atypical symptoms such as vomiting, diarrhea, constipation, ischemic colitis, and lower gastrointestinal bleeding. Most patients will have evidence of atherosclerosis in other vascular beds manifested by a history of myocardial infarction, stroke, or intermittent claudication.

Patients with atypical symptoms may be very difficult to diagnose, but a high degree of suspicion for CMI in patients with other manifestations of atherosclerosis and unexplained weight loss is appropriate. Often, the diagnosis is delayed in patients who are being evaluated for a possible malignancy as an explanation for their weight loss. Patients with functional bowel complaints rarely experience significant weight loss, which helps to differentiate them from patients with CMI. Evidence of significant obstruction of 2 or more of these vessels is often found when classic symptoms and endoscopy suggest bowel ischemia,³ although single-vessel disease, usually of the SMA, has been described as a cause of symptomatic CMI, particularly if collateral connections have been disrupted by prior abdominal surgery.




Fig 1. Initial clinical presentation of patients with CMI.³⁴




Fig 2. Bar graph of mortality rates for surgery and endovascular treatment of CMI taken from the National Inpatient Sample database.⁷ PTA/S indicates percutaneous transluminal angioplasty or stent; CKD, chronic kidney disease; DM, diabetes mellitus.

Diagnosis

Chronic mesenteric ischemia is a clinical diagnosis based on symptoms and consistent anatomical findings. Patients in whom there is a high degree of clinical suspicion for CMI should undergo screening with duplex ultrasound imaging or noninvasive angiography with CTA and MRA to confirm their anatomy.

The ability to visualize the mesenteric vessels with duplex ultrasound may be technically challenging and requires a skilled and dedicated technologist. The reported accuracy of duplex imaging to identify significant stenoses of the celiac and superior mesenteric arteries approaches 90%.^{4,5} With the relatively common application of CTA and MRA imaging for abdominal pathology, it is now possible to make the “anatomic” diagnosis without performing an invasive procedure.⁶

Invasive digital subtraction angiography is also very useful for diagnosis and allows the measurement of translesional pressure gradients in lesions of equivocal severity. Angiography requires a large field (15-in image intensifier) frontal view to visualize collateral vessels (ie, wandering [meandering] artery of Drummond) and a lateral




Fig 3. Bar graph of mortality rates for surgery and endovascular treatment of CMI taken from the National Inpatient Sample database.⁷ LOS indicates length of stay.




Fig 4. A, Baseline angiogram of tight proximal SMA stenosis with brachial artery access. B, Final angiogram after deployment of balloon expandable stent (Herculink; Abbott, Abbott Park, IL).

aortogram to visualize the ostia of the mesenteric arteries (Figs 2 and 3). Occasionally, an enlarged collateral vessel connecting a branch of the IMA with the SMA (arc of Riolan) is seen on the anterior-posterior aortogram and is an indication of proximal mesenteric artery disease. When critical stenoses ($\geq 70\%$) in multiple arteries are found in symptomatic patients, revascularization is appropriate.

Treatment

Revascularization has traditionally been performed with open surgery for either endarterectomy or bypass grafting of the mesenteric vessels. However, this patient group has




Fig 5. A, Baseline aortogram showing 99% stenosis of the IMA. B, Selective imaging of the IMA stenosis. C, Post-stent angiogram of the IMA. D, Aortogram after IMA stent placement.

Table 1
The Ochsner Clinic CMI experience³⁴

Vessel Treated	n (%)
Celiac	45 (59%)
SMA	21 (28%)
IMA	8 (11%)
Common hepatic	1 (1.4%)
Vein graft	1 (1.4%)

a high incidence of underlying coronary artery disease, and the perioperative mortality ranges from 3.5% to 15% (Fig 2),^{7–24} with the highest incidence of complications occurring in patients older than 70 years (Fig 3).^{7,23}

Atherosclerotic aorto-ostial obstructions of the visceral vessels are very similar to those of the renal arteries, and the technical considerations for percutaneous transluminal angioplasty (PTA) with stent placement are similar to those for renal artery intervention (Figs 4 and 5). Analogous with renal interventions, stent placement offers a superior late patency compared with PTA alone. There is no reason to perform PTA alone on these predominantly ostial lesions, which suffer from significant recoil.²⁵ The endovascular approach does not require general anesthesia or the operative trauma associated with open surgery resulting in lower acute mortality and morbidity.^{3,19,25–36}

Because of the infrequency of CMI, there are no clinical trials directly comparing surgery with endovascular treatment. The 20-year (1977–1997) Cleveland Clinic experience in 85 patients with CMI treated with surgery demonstrated an 8% perioperative mortality rate. One third of the patients had a major complication of surgery.¹⁸ Advanced age, hypertension, coronary disease, and disease in other vascular beds correlated with surgical complications. At late follow-up, 23% (n = 18) had

objective evidence of restenosis, 21% (n = 16) had recurrent CMI symptoms, and 12% (n = 9) underwent target vessel revascularization. Interestingly, the use of vein conduit as graft material was associated with poorer patency than Dacron as a graft material. The 5-year survival rate was 64% (95% confidence interval, 53%–75%), and the 3-year symptom-free survival rate was 81% (95% confidence interval, 72%–90%).

Data from Mayo Clinic on 229 consecutive patients with demonstrated a propensity for higher risk patients with multiple comorbidities to be referred for angioplasty with stenting, and lower risk patients tended to have open surgical repair.³⁷ They found that despite the lower risk status of the surgical patients, there remained greater morbidity and prolonged hospitalization rates compared with the endovascular group. The authors noted a higher restenosis rate for endovascular therapy, but there was no difference in secondary patency between surgery and stented patients.³⁷

Our series of CMI patients from the Ochsner Medical Center includes 59 patients and 79 vessels treated percutaneously (Table 1).³⁴ The technical success rate was 96%, which yielded symptom relief in 88% of the patients (Fig 6). There was 1 perioperative death (1.7%) and 2 access site complications. At a mean follow-up of 38 ± 15 months, 17% of the patients had recurrence of their symptoms, but none developed acute mesenteric ischemia. All patients with recurrent symptoms underwent successful retreatment without complication. The in-stent restenosis rate at 14 ± 5 months with 90% of the vessels imaged by CTA, invasive angiography, or Duplex ultrasound was 29%. The rate of target vessel revascularization was 17%. The 5-year cumulative freedom from death, symptom-recurrence, or both was 72%, 79% and 57%, respectively.

Several recent studies have summarized noncontemporaneous comparisons of surgical therapy and catheter-based therapies for CMI patients.^{37–41} These reports emphasize that complete surgical revascularization has benefits in terms of midterm patency, but generally at a cost of significant periprocedural morbidity and mortality for the patients. Endovascular approaches are generally preferred by patients, including those with severe comorbidities including cachexia and a hostile abdomen.³⁷

Conclusion

The infrequent occurrence of CMI has made randomized control trials comparing treatment outcomes very difficult to perform. Case series have shown that percutaneous therapy with stent placement offers the lowest morbidity and roughly equivalent long-term outcomes when compared with surgery. The current treatment recommendation is that patients who are candidates for either surgery or percutaneous therapy should receive percutaneous therapy with stent placement.




Fig 6. Bar graph showing Ochsner Clinic results of endovascular therapy for CMI.³⁴

Statement of Conflict of Interest

The author declares that there are no conflicts of interest.

References

- Hansen KJ, Wilson DB, Craven TE, et al: Mesenteric artery disease in the elderly. *J Vasc Surg* 2004;40:45-52.
- Thomas JH, Blake K, Pierce GE, et al: The clinical course of asymptomatic mesenteric arterial stenosis. *J Vasc Surg* 1998;27: 840-844.
- Matsumoto AH, Tegtmeier CJ, Fitzcharles EK, et al: Percutaneous transluminal angioplasty of visceral arterial stenoses: results and long-term clinical follow-up. *J Vasc Interv Radiol* 1995;6: 165-174.
- Bowersox JC, Zwolak RM, Walsh DB, et al: Duplex ultrasonography in the diagnosis of celiac and mesenteric artery occlusive disease. *J Vasc Surg* 1991;14:780-786 [discussion 786-788].
- Zwolak RM, Fillinger MF, Walsh DB, et al: Mesenteric and celiac duplex scanning: a validation study. *J Vasc Surg* 1998;27:1078-1087 [discussion 1088].
- Chow LC, Chan FP, Li KC: A comprehensive approach to MR imaging of mesenteric ischemia. *Abdom Imaging* 2002;27:507-516.
- Schermerhorn ML, Giles KA, Hamdan AD, et al: Mesenteric revascularization: management and outcomes in the United States, 1988-2006. *J Vasc Surg* 2009;50:341-348.e341.
- Oderich GS, Malgor RD, Ricotta II JJ: Open and endovascular revascularization for chronic mesenteric ischemia: tabular review of the literature. *Ann Vasc Surg* 2009;23:700-712.
- Kieny R, Batellier J, Kretz JG: Aortic reimplantation of the superior mesenteric artery for atherosclerotic lesions of the visceral arteries: sixty cases. *Ann Vasc Surg* 1990;4:122-125.
- Cormier JM, Fichelle JM, Vennin J, et al: Atherosclerotic occlusive disease of the superior mesenteric artery: late results of reconstructive surgery. *Ann Vasc Surg* 1991;5:510-518.
- Cunningham CG, Reilly LM, Rapp JH, et al: Chronic visceral ischemia. Three decades of progress. *Ann Surg* 1991;214:276-287 [discussion 287-278].
- McAfee MK, Cherry KJ Jr, Naessens JM, et al: Influence of complete revascularization on chronic mesenteric ischemia. *Am J Surg* 1992; 164:220-224.
- Calderon M, Reul GJ, Gregoric ID, et al: Long-term results of the surgical management of symptomatic chronic intestinal ischemia. *J Cardiovasc Surg (Torino)* 1992;33:723-728.
- Christensen MG, Lorentzen JE, Schroeder TV: Revascularisation of atherosclerotic mesenteric arteries: experience in 90 consecutive patients. *Eur J Vasc Surg* 1994;8:297-302.
- Gentile AT, Moneta GL, Taylor LM Jr, et al: Isolated bypass to the superior mesenteric artery for intestinal ischemia. *Arch Surg* 1994; 129:926-931 [discussion 931-922].
- Johnston KW, Lindsay TF, Walker PM, et al: Mesenteric arterial bypass grafts: early and late results and suggested surgical approach for chronic and acute mesenteric ischemia. *Surgery* 1995;118:1-7.
- Moawad J, Gewertz BL: Chronic mesenteric ischemia. Clinical presentation and diagnosis. *Surg Clin North Am* 1997;77: 357-369.
- Mateo RB, O'Hara PJ, Hertzner NR, et al: Elective surgical treatment of symptomatic chronic mesenteric occlusive disease: early results and late outcomes. *J Vasc Surg* 1999;29:821-831 [discussion 832].
- Sivamurthy N, Rhodes JM, Lee D, et al: Endovascular versus open mesenteric revascularization: immediate benefits do not equate with short-term functional outcomes. *J Am Coll Surg* 2006;202:859-867.
- Foley RN, Parfrey PS, Kent GM, et al: Serial change in echocardiographic parameters and cardiac failure in end-stage renal disease. *J Am Soc Nephrol* 2000;11:912-916.
- Jimenez JG, Huber TS, Ozaki CK, et al: Durability of antegrade synthetic aortomesenteric bypass for chronic mesenteric ischemia. *J Vasc Surg* 2002;35:1078-1084.
- Kihara TK, Blebea J, Anderson KM, et al: Risk factors and outcomes following revascularization for chronic mesenteric ischemia. *Ann Vasc Surg* 1999;13:37-44.
- Park WM, Cherry KJ Jr, Chua HK, et al: Current results of open revascularization for chronic mesenteric ischemia: a standard for comparison. *J Vasc Surg* 2002;35:853-859.
- Leke MA, Hood DB, Rowe VL, et al: Technical consideration in the management of chronic mesenteric ischemia. *Am Surg* 2002;68: 1088-1092.
- Landis MS, Rajan DK, Simons ME, et al: Percutaneous management of chronic mesenteric ischemia: outcomes after intervention. *J Vasc Interv Radiol* 2005;16:1319-1325.
- Hallisey MJ, Deschaine J, Illescas FF, et al: Angioplasty for the treatment of visceral ischemia. *J Vasc Interv Radiol* 1995;6:785-791.
- Allen RC, Martin GH, Rees CR, et al: Mesenteric angioplasty in the treatment of chronic intestinal ischemia. *J Vasc Surg* 1996;24:415-421 [discussion 421-413].
- Maspe F, Mazzetti di Pietralata G, et al: Percutaneous transluminal angioplasty in the treatment of chronic mesenteric ischemia: results and 3 years of follow-up in 23 patients. *Abdom Imaging* 1998;23: 358-363.
- Nyman U, Ivancev K, Lindh M, et al: Endovascular treatment of chronic mesenteric ischemia: report of five cases. *Cardiovasc Interv Radiol* 1998;21:305-313.
- Sheeran SR, Murphy TP, Khwaja A, et al: Stent placement for treatment of mesenteric artery stenoses or occlusions. *J Vasc Interv Radiol* 1999;10:861-867.
- Kasirajan K, O'Hara PJ, Gray BH, et al: Chronic mesenteric ischemia: open surgery versus percutaneous angioplasty and stenting. *J Vasc Surg* 2001;33:63-71.
- AbuRahma AF, Stone PA, Bates MC, et al: Angioplasty/stenting of the superior mesenteric artery and celiac trunk: early and late outcomes. *J Endovasc Ther* 2003;10:1046-1053.
- Sharafuddin MJ, Olson CH, Sun S, et al: Endovascular treatment of celiac and mesenteric arteries stenoses: applications and results. *J Vasc Surg* 2003;38:692-698.
- Silva JA, White CJ, Collins TJ, et al: Endovascular therapy for chronic mesenteric ischemia. *J Am Coll Cardiol* 2006;47:944-950.
- Cognet F, Ben Salem D, Dransart M, et al: Chronic mesenteric ischemia: imaging and percutaneous treatment. *Radiographics* 2002; 22:863-879 [discussion 879-880].
- Matsumoto AH, Angle JF, Spinosa DJ, et al: Percutaneous transluminal angioplasty and stenting in the treatment of chronic mesenteric ischemia: results and long-term follow-up. *J Am Coll Surg* 2002;194:S22-S31.
- Oderich GS, Bower TC, Sullivan TM, et al: Open versus endovascular revascularization for chronic mesenteric ischemia: risk-stratified outcomes. *J Vasc Surg* 2009;49:1472-1479.e3.
- van Petersen AS, Kolkman JJ, Beuk RJ, et al: Open or percutaneous revascularization for chronic splanchnic syndrome. *J Vasc Surg* 2009;51:1309-1316.
- Davies RSM, Wall ML, Silverman SH, et al: Surgical versus endovascular reconstruction for chronic mesenteric ischemia: a contemporary UK series. *Vasc Endovascular Surg* 2009;43:157-164.
- Atkins MD, Kwolek CJ, LaMuraglia GM, et al: Surgical revascularization versus endovascular therapy for chronic mesenteric ischemia: a comparative experience. *J Vasc Surg* 2007;45:1162-1171.
- Dias NV, Acosta S, Resch T, et al: Mid-term outcome of endovascular revascularization for chronic mesenteric ischaemia. *Br J Surg* 2010;97:195-201.