



---

# Magnetic Resonance Imaging (MRI)

---

## Part 2: Image Contrast


**Rasmus M. Birn, Ph.D.**

University of Wisconsin – Madison  
Madison, WI, USA




# MRI Contrast

---






**T1-weighted**



**T2-weighted**






# MRI Contrast






# MRI Contrast – T1




Longitudinal  
Relaxation time

**T1**








# MRI Contrast – T1






# MRI Contrast – T1



Longitudinal  
Relaxation time

**T1**








# T1-weighted Imaging


---

- T1 of Brain Tissues – Shortest to Longest:
  - White Matter
  - Gray Matter
  - CSF






# MRI Contrast – T2










# MRI Contrast – T2






# MRI Contrast – T2



Transverse  
Relaxation time

**T2**






# T2-weighted Imaging


---

- T2 of Brain Tissues – Shortest to Longest:
  - White Matter
  - Gray Matter
  - CSF






# T1 & T2



*Signal for steady-state images is reduced since the longitudinal magnetization has not fully recovered between excitations.*




# Contrast is determined by TR and TE




TR = long  
TE = short

Proton-Density



TR = short  
(TE = short)

T1-weighted





(TR = long)  
TE = long

T2-weighted




# Flip Angle





# T1 & T2






Optimal flip angle for highest steady-state signal:

Ernst Angle

$$\cos(\theta) = e^{-TR/T1}$$



# Magnetic Resonance Imaging (MRI)



Sensitive to:

- # of protons ( $\text{H}_2\text{O}$ )
- Magnetic environment
  - Tissue structure