

Todo list

Either remove references to the boundaru ∂X or introduce it, and check if more hypothesis are needed for the characterisation	4
Prove that attracting flag is well-defined when there exists a singular value gap, and explain that in the bassin, g attracts flags uniformly to the attracting flag	7
Is this true for Cartan subalgebras of rank-1 Lie groups?	10
Add some theory on densities and show that the density on $\mathcal{F}_{1,p+1}(\mathbb{R}^d)$ is $SL(d, \mathbb{R})$ equivariant	11
Does this clarifying comment on the axes of the ellipsoid help?	14
Correct ellipsoid drawing	14
Correct the definition of the Busemann cocycle (see [Qui02] Lemme 6.1)	
Then add proof for Lemma 3.1.1	20
Why do we construct a measure over $\mathcal{F}_{1,d_\rho+1}(\mathbb{R}^d)$ and not the limit set? We had talked about it but I forgot.	23
Why is the section measurable?	24
Outline proof steps for counterexample to Labourie's lemma	30
Is it not the case $d = 4$ that is of importance in physics?	35
Type up the proof of the quadratic form reduction that I have already handwritten. . .	36
inline	36
Is $\text{Ad}_{\text{SU}(2,1)}$ Zariski-dense in $\text{SO}(\mathfrak{su}(2,1), K)$? If yes, then a generalisation of Theorem 5 to $\text{SO}(4,4)$ would account for the validity of the result in the case of Proposition 4.0.2	36

Limit sets of Anosov representations

Giorgos

July 30, 2024

Abstract

For a projective Anosov subgroup $\rho(\Gamma)$ of $\mathrm{SL}(d, \mathbb{R})$ whose limit set is a Lipschitz submanifold of $\mathbb{P}(\mathbb{R}^d)$, based on [PSW23], we show that the Hausdorff dimension of the limit set is equal to the critical exponent of the Falconer functional when the Zariski-closure of $\rho(\Gamma)$ is $\mathrm{SL}(d, \mathbb{R})$ or $\mathrm{SO}(2, d - 2)$ for $d \neq 4$.

Contents

1	Introduction	2
1.1	Limit sets and critical exponents	4
1.2	Gromov hyperbolic groups	4
1.3	Grassmanians and flag spaces	5
1.4	Lie groups and Anosov representations	6
1.5	Important functionals and definitions	9
1.6	Hausdorff measures and densities	11
2	Upper bound	12
2.1	Lemmata	12
2.2	Proof of bound	18
3	Lower bound	20
3.1	Busemann cocycle and Patterson–Sullivan measures	20
3.2	Proof strategy	22
3.3	Existence of Patterson–Sullivan measure	23
3.4	Proof of the main theorem	27
4	Dropping the Zariski-density assumption	29
5	Conclusion	37
A	Tangent space to the Grassmanian	38

Chapter 1

Introduction

Let $\mathbb{H}_{\mathbb{R}}^n$ be the real hyperbolic space and $\Gamma \leq \mathrm{SO}(1, n)$ be a discrete convex-cocompact subgroup of isometries. To the action $\Gamma \curvearrowright \mathbb{H}_{\mathbb{R}}^n$, we associate (Definition 1.1.2) its critical exponent δ_{Γ} , which is a dynamical invariant expressing the exponential growth rate of the Γ -orbits in $\mathbb{H}_{\mathbb{R}}^n$. Another object associated (Definition 1.1.1) to the action is the limit set Λ_{Γ} , which should be thought of as the set of accumulation points of an orbit at infinity. In [Sul79], Sullivan proved that the critical exponent δ_{Γ} is equal to the Hausdorff dimension of the limit set Λ_{Γ} , providing a stunning connection between the fractal geometry of the limit set and the dynamics of the group action:

Theorem 1 ([Sul79]). *Let $\Gamma \leq \mathrm{PSO}(1, n)$ be convex co-compact. Then the critical exponent of Γ is equal to the Hausdorff dimension of its limit set:*

$$\dim_{\mathcal{H}}(\Lambda_{\Gamma}) = \delta_{\Gamma}.$$

To accomplish this, Sullivan generalised to all hyperbolic spaces the construction done for the hyperbolic plane by Patterson in [Pat76] of a Γ -quasi-equivariant measure on the limit set of hyperbolic space. Since then, there have been numerous generalizations of this result to other settings, such as the action of other kinds of discrete groups on hyperbolic spaces (see for instance [Rob03; DOP00; Co093]), or to other kinds of spaces (for instance [Coo93] for the case of X being a Gromov hyperbolic space).

Here, we will in large part follow [PSW23], whose authors are interested in the generalization of Sullivan's result to certain subgroups of higher rank Lie groups. In particular, they focus on Anosov subgroups of $\mathrm{SL}(d, \mathbb{R})$, which are defined (in Definition 1.4.1 using the characterisation of [KLP17]) as the subgroups arising from representations into $\mathrm{SL}(d, \mathbb{R})$ with a singular value growth that is exponentially controlled by the word length of the group element. They are particularly relevant in this context because they are considered as a promising generalization of convex co-compact subgroups of rank one Lie groups to ones of higher rank, and their Gromov boundary is realised as a subset of the projective space through the boundary map (Theorem 3), which will be the analogue of the limit set Λ_{Γ} to the higher rank case. The only regularity assumption that they pose on the representations considered is that the limit set is a Lipschitz submanifold of the ambient projective space, which in particular is the case for three large classes of examples, namely maximal representations ([BIW10]), Anosov quasi-fuchsian AdS representations ([MB12]) and $\mathbb{H}^{p,q}$ -convex-cocompact representations ([DGK18]).

On second glance (more extensively described in Example 1.5.1), the critical exponent of Γ can be seen as the critical exponent (defined in Definition 1.5.2 as the radius of convergence

for the respective Poincaré series) of a functional over the Cartan subalgebra of $\mathrm{SL}(d, \mathbb{R})$, associating to each γ , its displacement $d(o, \gamma \cdot o)$ with respect to some fixed point $o \in X$. Given this, the following modified version of the result appearing in [PSW23] should be seen as a natural generalization of Sullivan's theorem to the higher rank case:

Theorem 4. Let $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{R})$ be a Zariski-dense, projective Anosov representation in $\mathrm{SL}(d, \mathbb{R})$ such that $\xi_\rho^1(\partial\Gamma)$ is a Lipschitz submanifold of $\mathbb{P}(\mathbb{R}^d)$. Then

$$\dim(\xi_\rho^1(\partial\Gamma)) = h_{\rho(\Gamma)}(F),$$

where $h_{\rho(\Gamma)}(F)$ is the critical exponent of the Falconer functional F .

The above statement differs from the one of the respective result in [PSW23] in replacing the strong irreducibility assumption on the representation with that the Zariski-density of the subgroup in $\mathrm{SL}(d, \mathbb{R})$, because of a gap that we noticed in the proof of the latter. Since this is a strong assumption that is not always satisfied in the three classes of examples presented above, we present the following result that covers certain cases of interest:

Theorem 5. Let $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{R})$ be a projective Anosov representation such that $\xi_\rho^1(\partial\Gamma)$ is a Lipschitz submanifold of $\mathbb{P}(\mathbb{R}^d)$. Assume moreover that $\dim \xi_\rho^1(\partial\Gamma) = d - 3, d > 4$ and $\rho(\Gamma)$ is Zariski-dense in $\mathrm{SO}(2, d - 2)$. Then

$$\dim \xi_\rho^1(\partial\Gamma) = h_{\rho(\Gamma)}(F).$$

It should be noted that this result is possibly not optimal, as we hope its proof generalises (at least) to other groups of the form $\mathrm{SO}(p, q)$.

In the remaining of this chapter, we will recall the definition of the Hausdorff dimension, along with all necessary background on Lie groups, Anosov representations that will be necessary in the subsequent chapters. We will also introduce the Falconer functional and the unstable Jacobian, which will be the functionals whose critical exponents capture the geometric information of the dimension of the limit set.

In Chapter 2, we will use a classical recipe to obtain the upper bound of the Hausdorff dimension, by finding an appropriate covering of the limit set $\xi_\rho^1(\partial\Gamma)$, for which the Hausdorff content is dominated by the Dirichlet series of the functional F . We would like to stress that the only part of the hypothesis utilized in that chapter is that the representation is projective Anosov, meaning that it still holds without requiring that Λ_Γ is a Lipschitz submanifold of $\mathbb{P}(\mathbb{R}^d)$ or that $\rho(\Gamma)$ is Zariski-dense.

On the other hand, to establish the upper bound in Chapter 3 of the critical exponent one uses the more elaborate machinery of Patterson–Sullivan measures as generalised into higher rank by [Qui02] and further developed in [PSW23]. For this part of the proof, the hypothesis that $\xi_\rho^1(\partial\Gamma)$ is a Lipschitz submanifold will be introduced (in Proposition 3.3.1) to construct a Patterson–Sullivan measure μ over a space of partial flags, while the Zariski-density of $\rho(\Gamma)$ is necessary (see Example 3.3.1) to show that it is μ -irreducible (that is, the measure of the set of transverse flags to each flag is positive, see Definition 3.3.1) and obtain estimates (in Lemma 3.3.2) that will be used to prove that the sum defining the critical exponent is finite.

Finally, in Chapter 4 we trace the gap in the proof of [PSW23] to lemma appearing in [Lab06] and provide two counterexamples that show it is substantial. Then we present the proof of Theorem 5, and conclude with some remarks on the generalisation of the result to other subgroups.

1.1 Limit sets and critical exponents

Definition 1.1.1. Let Γ be a discrete group of isometries of a metric space (X, d) . We define the limit set Λ_Γ of Γ to be the set of accumulation points of the orbit of some fixed $x \in X$.

Remark 1.1.1. One can show that the definition of the limit set is independent of the choice of x . When moreover Γ is nonelementary (i.e. it does not admit any finite invariant subset of $X \cup \partial X$), it is easy to see that $\Lambda_\Gamma \subseteq \partial X$ and one can show that it is the minimal closed Γ -invariant subset of ∂X (see for example [Qui06, Proposition 4.7]).

Definition 1.1.2. Let Γ be a discrete group of isometries of a metric space (X, d) . We define the critical exponent of Γ to be the asymptotic exponential growth of its orbits, i.e. the following limit:

$$\begin{aligned} \delta_\Gamma &= \limsup_{n \rightarrow \infty} \frac{\log \#\{\gamma \in \Gamma : d(x, \gamma x) \leq n\}}{n} = \\ &= \inf \left\{ s \in \mathbb{R} : \sum_{\gamma \in \Gamma} e^{-sd(x, \gamma x)} < \infty \right\} \end{aligned}$$

for some fixed $x \in X$.

Remark 1.1.2. It is not hard to show that the critical exponent is independent of the choice of x .

Later in the thesis, we will be using certain terms to refer to lattices, which we recall below:

Definition 1.1.3. Let G be a Lie group, and $\Gamma \leq G$ a discrete subgroup.

- (i) Γ is a lattice if it is co-finite, meaning that the quotient G/Γ admits a finite G -invariant measure.
- (ii) Γ is a uniform lattice if it is co-compact, meaning that the quotient G/Γ is compact.

1.2 Gromov hyperbolic groups

In this section we recall the basic definitions and facts of Gromov hyperbolic spaces and groups which were introduced by Gromov to generalise certain metric properties of hyperbolic spaces. For more details or proofs of the following facts, we refer to [GH13].

Let (X, d) be a metric space. The Gromov product of two points $x, y \in X$ with respect to a point $z \in X$ is defined as

$$(x, y)_z = \frac{d(x, z) + d(y, z) - d(x, y)}{2}.$$

For some $\delta > 0$, we say that X is δ -hyperbolic if for all $x, y, z, w \in X$ the following relation is satisfied

$$(x, y)_z \geq \min\{(x, w)_z, (w, y)_z\} - \delta.$$

A map f defined over $[0, \infty)$ or \mathbb{N} that takes values in X is called a geodesic ray if

$$d(f(t), f(s)) = |t - s|$$

Either remove references to the boundary ∂X or introduce it, and check if more hypothesis are needed for the characterisation

holds for all $t, s \in [0, \infty)$ or \mathbb{N} respectively. It is called a (c, C) -quasigeodesic ray if it satisfies

$$\frac{1}{C}|t - s| - c \leq d(f(t), f(s)) \leq C|t - s| + c$$

for all $t, s \in [0, \infty)$ or \mathbb{N} respectively.

Supposing that X is a geodesic metric space (meaning that for any $x, y \in X$, there exists a geodesic joining x and y), we define the Gromov boundary $\partial_\infty X$ as the set of equivalence classes of geodesics in X that start from a fixed base point $z \in X$ under the equivalence relation

$$f \sim g \text{ if and only if } \sup_t d(f(t), g(t)) < \infty.$$

We could consider the same equivalence relation over (c, C) -quasigeodesic rays, and the resulting space would be another model for the Gromov boundary, homeomorphic to the previous one. In any case, it is important to know that the space $\partial_\infty X$ is metrizable and compact.

Let Γ be a finitely generated group and S be a symmetric (meaning that $s^{-1} \in S$ for any $s \in S$) generating set. The Cayley graph Γ with respect to S is the graph whose vertices are the elements of Γ and there is an edge between $g, h \in \Gamma$ if and only if $g^{-1}h \in S$. It becomes a metric space with the word metric $d_S(g, h)$ defined as the length of the shortest path in the Cayley graph connecting g and h . The group Γ is called hyperbolic if the Cayley graph is δ -hyperbolic for some $\delta > 0$. The space $\hat{\Gamma} \stackrel{\text{def}}{=} \Gamma \cup \partial_\infty \Gamma$ is compact.

Finally, another object that will become useful in later chapters are the coarse cones on the boundary of a hyperbolic group Γ , which was taken from [PSW23]. For $c_0, c_1 > 0$, we define it as follows:

$$C_\infty^{c_0, c_1}(\gamma) = \left\{ [(a_j)_{j \in \mathbb{N}}] : (a_j)_j \text{ is a } (c_0, c_1)\text{-quasi-geodesic ray, } a_0 = \gamma^{-1} \text{ and } e \in \{a_j\}_j \right\}.$$

1.3 Grassmanians and flag spaces

In this section we will recall some terminology and notions related to Grassmanians and flag spaces, that will prove useful in Section 1.4 in presenting the dynamical properties of Anosov representations, as well as in Chapter 2, where we will construct an open cover for the limit set of Anosov subgroups.

For $k \in \llbracket 1, d \rrbracket$, the Grassmanian $\mathcal{G}_k(\mathbb{R}^d)$ is the space of k -dimensional subspaces of \mathbb{R}^d . For two subspaces $V \in \mathcal{G}_k(\mathbb{R}^d)$, $W \in \mathcal{G}_l(\mathbb{R}^d)$ of possibly different dimensions, we can define the notion of an angle as

$$\angle(V, W) \stackrel{\text{def}}{=} \min_{v \in V^\times, w \in W^\times} \angle(v, w) \in [0, \pi/2],$$

where $V^\times \stackrel{\text{def}}{=} V \setminus \{0\}$, while the angle between vectors $v, w \in \mathbb{R}^d$ is defined as

$$\angle(v, w) \stackrel{\text{def}}{=} \arccos \left(\frac{\langle v, w \rangle}{\langle v, v \rangle \langle w, w \rangle} \right) \in [0, \pi].$$

For $\Theta \subseteq \llbracket 1, d \rrbracket$ we define the space of partial flags

$$\mathcal{F}_\Theta(\mathbb{R}^d) = \{(x_\theta)_{\theta \in \Theta} : x_\theta \in \mathcal{G}_\theta(\mathbb{R}^d) \text{ for } \theta \text{ and } x_\theta \leq x_{\theta'} \text{ for } \theta < \theta' \text{ in } \Theta\},$$

where $\mathcal{G}_r(\mathbb{R}^d)$ is the r -th Grassmanian, that is the space of r -dimensional subspaces of \mathbb{R}^d . To each $\Theta \subseteq \llbracket 1, d \rrbracket$, we associate $i\Theta = \{d - \theta : \theta \in \Theta\}$. In this notation, two flags $(x_\theta)_\theta \in \mathcal{F}_\Theta(\mathbb{R}^d)$ and $(y_\theta)_\theta \in \mathcal{F}_{i\Theta}(\mathbb{R}^d)$ are transverse if $x_\theta \cap y_{d-\theta} = \{0\}$ for all $\theta \in \Theta$. For the sake of readability, we will often skip writing the braces around the indices of the flags and write $\mathcal{F}_{i,j}(\mathbb{R}^d)$ instead of $\mathcal{F}_{\{i,j\}}(\mathbb{R}^d)$.

1.4 Lie groups and Anosov representations

Let G be a real semisimple Lie group, whose Lie algebra is denoted by \mathfrak{g} , K is a maximal compact subgroup of G whose algebra is denoted by \mathfrak{k} . We let \mathfrak{p} be the orthogonal complement of \mathfrak{k} in \mathfrak{g} with respect to the Killing form. We denote with \mathfrak{a} a maximal abelian subalgebra of \mathfrak{p} . A non-trivial linear form $\alpha \in \mathfrak{a}^*$ is called a restricted root when the set

$$\mathfrak{g}_\alpha = \{u \in \mathfrak{g} : \text{ad}_g(a) = \alpha(a)u \text{ for all } a \in \mathfrak{a}\}$$

is not zero. The set of restricted roots is denoted by Δ and the closure of any component

$$\mathfrak{a}^+ \subseteq \mathfrak{a} \setminus \bigcup_{\alpha \in \Delta} \ker \alpha$$

is called a Weyl chamber.

The following theorem is a fundamental result in the theory of semi-simple Lie groups:

Theorem 2 (Cartan decomposition). *Let K be maximal compact subgroup, \mathfrak{a} a Cartan subalgebra of \mathfrak{g} and \mathfrak{a}^+ a Weyl chamber of \mathfrak{a} . Then G can be decomposed as $G = Ke^{\mathfrak{a}^+}K$, while for every $g \in G$, there exists a unique $\mu(g)$ such that $g \in Ke^{\mu(g)}K$.*

The map

$$\mu : G \rightarrow \mathfrak{a}^+$$

appearing in the theorem is called the Cartan projection.

For us, the only relevant case will be $G = \text{SL}(d, \mathbb{R})$, so we will fix the maximal compact subgroup $K = \text{SO}(d, \mathbb{R})$, the Cartan subalgebra

$$\mathfrak{a} = \{\text{diag}(\alpha_1, \dots, \alpha_d) : \alpha_1 + \dots + \alpha_d = 0\}$$

and its Weyl chamber

$$\mathfrak{a}^+ = \{\text{diag}(\alpha_1, \dots, \alpha_d) : \alpha_1 \geq \dots \geq \alpha_d\}.$$

Note that then, the Cartan decomposition of $\text{SL}(d, \mathbb{R})$ is the same as the The Cartan projection is then given by

$$\mu(g) = \text{diag}(\lambda_1(g), \dots, \lambda_d(g))$$

where $\log \sigma_i(g) = \lambda_i(g)$ is the i -th singular value of g , i.e. the square root of the eigenvalue of $g^t \cdot g$. The restricted roots are $\mathfrak{a}_{ij} = a_i - a_j \in \mathfrak{a}^*$ and the i -th fundamental weight is defined as $\omega_i(a) = a_1 + \dots + a_i$, for $i \in \llbracket 1, d-1 \rrbracket$.

Before going on with the definition and the properties of Anosov representations, we will briefly discuss the way that a linear transformation in $\text{SL}(d, \mathbb{R})$ acts on the euclidean space \mathbb{R}^d , namely by stretching and contracting. Let $g \in \text{SL}(d, \mathbb{R})$, and consider a Cartan decomposition $g = k_g e^{\mu(g)} l_g$. Since g acts linearly on \mathbb{R}^d , to understand its action on the Euclidean space it suffices to understand its action on the unit sphere \mathbb{S}^{d-1} , which g distorts into an ellipsoid (see Figure 1.1).

In other words, we claim that the image of the unit sphere through g is an ellipsoid with axes

$$u_i(g) = k_g e_i, \text{ for } i \in \llbracket 1, d \rrbracket$$

and respective lengths $\sigma_i(g) = e^{\lambda_i(g)}$. Indeed, if we write $w = \beta_1 u_1(g) + \dots + \beta_d u_d(g)$, then $w \in g \cdot \mathbb{S}^{d-1}$ if and only if

$$\begin{aligned} \|g^{-1}w\|^2 &= \|\beta_1 \sigma_1(g)^{-1} l_g^{-1} + \dots + \beta_d \sigma_d(g)^{-1}\|^2 = \\ &= \left(\frac{\beta_1}{\sigma_1(g)} \right)^2 + \dots + \left(\frac{\beta_d}{\sigma_d(g)} \right)^2 \leq 1, \end{aligned}$$




Figure 1.1: Linear transformations map spheres to ellipsoids.

which is precisely the defining inequality for an ellipsoid.

Recall that for a general element in $g \in \text{SL}(d, \mathbb{R})$, we may not make any assumption on the singular values $\sigma_i(g)$ except from the fact that they are in non-increasing order. In particular, certain singular values may coincide, which in the context of the above discussion means that the choice of the axes $u_i(g)$ is arbitrary. In fact, when for instance $\sigma_2(g) = \sigma_3(g)$, any rotation of the orthonormal basis on the $\mathbb{R}u_i(g) \oplus \mathbb{R}u_{i+1}(g)$ plane will give the equally valid orthonormal choice of axis $u_1(g), \cos(\theta)u_1(g) + \sin(\theta)u_2(g), -\sin(\theta)u_1(g) + \cos(\theta)u_2(g), u_3(g), \dots, u_d(g)$. This has to do with the fact that the choice of elements k_g, l_g in the Cartan decomposition of g is not unique (see Figure 1.2).

When however there is a singular value gap, i.e. $\sigma_p(g) > \sigma_{p+1}(g)$, then (noting that $\sigma_{d-p+1}(g^{-1}) < \sigma_{d-p}(g^{-1})$ as well because $\sigma_i(g^{-1}) = \sigma_{d-i+1}(g)^{-1}$) the choice of the axis $u_p(g)$ is unique up to a sign, and the attracting and repelling p -planes of g are well defined:

$$U_p(g) \stackrel{\text{def}}{=} k_g \cdot \mathbb{R}e_1 \oplus \dots \oplus \mathbb{R}e_p$$

$$U_{d-p}(g^{-1}) \stackrel{\text{def}}{=} l_g^{-1} \cdot \mathbb{R}e_{p+1} \oplus \dots \oplus \mathbb{R}e_d$$

are well defined and form a decomposition

$$g^{-1} \cdot U_p(g) \oplus U_{d-p}(g^{-1})$$

that is orthogonal with respect to the standard inner product on \mathbb{R}^d .

In particular, when $g \in \text{SL}(d, \mathbb{R})$ has a gap between the θ -th and $(\theta + 1)$ -th singular values for all $\theta \in \Theta \subseteq \llbracket 1, d - 1 \rrbracket$, we define for any $\alpha > 0$ the basin of attraction of g as

$$B_{\Theta, \alpha}(g) \stackrel{\text{def}}{=} \{x \in \mathcal{F}_{\Theta}(\mathbb{R}^d) : \angle(x^{\theta}, U_{d-\theta}(g^{-1})) < \alpha \text{ for all } \theta \in \Theta\}.$$

We may think of the basin of attraction as the set of flags that are away from the repelling flag $(U_{d-\theta}(g^{-1}))_{\theta \in \Theta}$ of g . To justify the name, we briefly mention informally the fact that flags

Prove that attract-ing flag is well-defined when there exists a singular value gap, and explain that in the basin, g attracts flags uniformly to the attract-ing flag




Figure 1.2: Equality between singular values results in axes ambiguity.

contained in the basin of attraction are uniformly attracted to the attracting flag $U_\Theta(g)$ under the action of g , and refer to [BPS19][Lemma A.6]. Often, for readability purposes, we will often skip writing the curly braces around the indices comprising Θ .

We are now ready to give one of the various equivalent definitions of Anosov representations in the literature. In light of the previous discussion, one could phrase it as: "a representation in $\mathrm{SL}(d, \mathbb{R})$ is p -Anosov if the length of the p -th axis of the ellipsoid $g \cdot \mathbb{S}^{d-1}$ is exponentially larger than the length of the $(p+1)$ -th axis, with the exponential rate controlled by the word length of the group element".

Definition 1.4.1. Let $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{R})$ be a linear representation and $p \in \llbracket 1, d-1 \rrbracket$. We say that ρ is p -Anosov if there exist constants $\mu, C > 0$ such that for all $\gamma \in \Gamma$:

$$\frac{\sigma_{p+1}}{\sigma_p}(\rho(\gamma)) \leq C e^{-\mu|\gamma|}.$$

The following theorem shows the dynamical behaviour of Anosov representations through their action on flag spaces. In particular, a p -Anosov representation ρ will have a well-defined attracting flag $\xi_\rho^p(x)$ in $\mathcal{F}_{p,d-p}(\mathbb{R}^d)$ for every $x \in \partial\Gamma$.

Theorem 3 ([KLP17; Lab06]). *Let $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{R})$ be a p -Anosov representation. Then there exists an equivariant continuous map $\xi : \hat{\Gamma} \rightarrow \mathcal{F}_{p,d-p}(\mathbb{R}^d)$ that restricts to*

$$\begin{aligned} \xi^p(\gamma) &= U_p(\rho(\gamma)) \stackrel{\text{def}}{=} k \cdot \mathbb{R}e_1 \oplus \cdots \oplus \mathbb{R}e_p \\ \xi^{d-p}(\gamma) &= U_{d-p}(\rho(\gamma)) \stackrel{\text{def}}{=} k \cdot \mathbb{R}e_1 \oplus \cdots \oplus \mathbb{R}e_{d-p}, \end{aligned}$$

for $\gamma \in \Gamma, \rho(\gamma) = ke^{\mathbf{a}l}$ and where e_1, \dots, e_d is the standard orthonormal basis for the Euclidean inner product. Moreover, it has the following transversality property on the boundary: for every $x, y \in \partial\Gamma$, the flags $\xi_\rho(x), \xi_\rho(y)$ are transverse unless $x = y$.

Again, we would like to stress that if the elements of the Anosov subgroup did not have the singular value gap, then the attracting flags U_p, U_{d-p} would not be well defined.

1.5 Important functionals and definitions

In this section we will introduce certain functionals on the Weyl chamber, whose critical exponents (see Definition 1.5.2) is going to generalise the critical exponent of the group to the higher rank case.

Definition 1.5.1. For $p \in \llbracket 2, \dots, d \rrbracket$, $s \in \mathbb{R}$ we define the functionals $\Psi_s^p : \mathfrak{a}^+ \rightarrow \mathbb{R}$, the Falconer functional $F_s : \mathfrak{a}^+ \rightarrow \mathbb{R}$, and the unstable Jacobian $J_p^u : \mathfrak{a}^+ \rightarrow \mathbb{R}$ as follows:

$$\begin{aligned}\Psi_s^p &= \mathbf{a}_{1,2} + \dots + \mathbf{a}_{1,p-1} + (s - (p-2))\mathbf{a}_{1,p} \\ F_s(a) &= \min \left\{ \sum_{j=2}^d s_j \alpha_{1,j}(a) : s_j \in (0, 1], \sum_{j=2}^d s_j = s \right\} \\ J_p^u &= (p+1)\omega_1 - \omega_{p+1} = \mathbf{a}_{1,2} + \dots + \mathbf{a}_{1,p+1}.\end{aligned}$$

Lemma 1.5.1. *The functionals defined above are related as follows:*

$$\begin{aligned}F_s(a) &= \max_{p \in \llbracket 2, d \rrbracket} \Psi_s^p(a) = \Psi_s^{p_0}(a) \text{ for } s \in [p_0 - 2, p_0 - 1]. \text{ for all } a \in \mathfrak{a}^+ \\ \frac{s}{p} J_p^u &\leq \Psi_{s+p}^{p+1} \text{ for all } s > 0\end{aligned}$$

Proof. Indeed, a quick calculation shows that for $s \geq 0, p \in \llbracket 2, d \rrbracket$ and $a \in \mathfrak{a}^+$ we have that:

$$\Psi_s^p(a) \leq \Psi_s^p(a) \text{ if and only if } s \geq p-1.$$

and that equality holds in the case $s = p-1$. Thus for $s \in [p-2, p-1]$ we have that

$$\begin{aligned}s \geq p-2, \dots, 1 \text{ implies that } \Psi_s^p(a) &\geq \dots \geq \Psi_s^2(a) \\ s \leq p, \dots, d-1 \text{ implies that } \Psi_s^p(a) &\leq \dots \leq \Psi_s^d(a)\end{aligned}$$

Another way to see this (refer to Figure 1.3) is to note that $\Psi_s^2(a), \dots, \Psi_s^d(a)$ is a sequence of functions that are affine in s , with slopes $\alpha_{12}(a) \leq \dots \leq \alpha_{1d}(a)$ and that they satisfy $\Psi_1^2(a) = \Psi_2^2(a), \Psi_2^3(a) = \Psi_3^4(a) \dots, \Psi_{d-2}^{d-1}(a) = \Psi_{d-2}^d(a)$.




Figure 1.3: Visual illustration that $\max_p \Psi_s^p(a) = \Psi_s^{p_0}(a)$ for $s \in [p_0 - 2, p_0 - 1]$.

For the inequality, the calculations has as follows:

$$\begin{aligned}
\Psi_s^{p+1} &= \mathbf{a}_{1,2} + \cdots + \mathbf{a}_{1,p+1} + (s-p)\mathbf{a}_{1,p+2} = \\
&= \mathbf{a}_{1,2} + \cdots + \mathbf{a}_{1,p+1} + p \left(\frac{s}{p} - 1 \right) \mathbf{a}_{1,p+2} \geq \\
&\geq \mathbf{a}_{1,2} + \cdots + \mathbf{a}_{1,p+1} + \left(\frac{s}{p} - 1 \right) (\mathbf{a}_{1,2} + \cdots + \mathbf{a}_{1,p+1}) = \\
&= \frac{s}{p} J_p^u.
\end{aligned}$$

□

Remark 1.5.1. The reason to consider the functionals Ψ_s^p, F_s will be apparent in Chapter 2, where we prove that the Hausdorff dimension of the limit set is bounded above by the critical exponent of the Falconer functional. There, we will see that the sums in the left-hand-side of the equality

$$\min_{p \in \llbracket 2, d \rrbracket} \left\{ \sum_{|\gamma|=T} \frac{\sigma_2}{\sigma_1} \cdots \frac{\sigma_{p-1}}{\sigma_1}(g) \left(\frac{\sigma_{p-1}}{\sigma_1}(g) \right)^{s-(p-2)} \right\} = \sum_{|\gamma|=T} e^{-F_s(a(g))}$$

correspond to the Hausdorff content of respective coverings of $\xi_\rho^1(\partial\Gamma)$ by projective ellipses of different lengths. For the use of the unstable Jacobian J_p^u we refer to Section 3.2.

As described in the begining of this section, some of the core objects that we will be working with are critical exponents and Hausdorff dimensions, which we recall below:

Definition 1.5.2. Let $\phi \in (\mathfrak{a}^+)^*$ be a functional over the Weyl chamber and $\Gamma \leq \mathrm{SL}(d, \mathbb{R})$ a discrete subgroup. We define the critical exponent of ϕ over Γ to be

$$h_\Gamma(\phi) = \inf \left\{ s > 0 : \sum_{\gamma \in \Gamma} e^{-\phi(\gamma)} < \infty \right\}.$$

Example 1.5.1. Let Γ be a uniform lattice of $\mathrm{SO}(1, d)$. Then any functional $\phi \in \mathfrak{a}^*$ is of the form

$$\phi(\mu(\gamma)) = d_\phi(\gamma \cdot o, o)$$

for some distance d_ϕ on \mathbb{H}^d . In particular, fixing some $\phi \in \mathfrak{a}^*$ such that $d_\phi = d$, we have that

$$h_\Gamma(\phi) = \delta_\Gamma.$$

Is this true for Cartan subalgebras of rank-1 Lie groups?

When $\phi = F$ is the Falconer functional, we obtain the following special case (of projective Anosov representations, i.e. $P = 1$) of a definition from [LL23]:

Definition 1.5.3. We define the Falconer dimension $\dim_F(\rho)$ of ρ to be the critical exponent of the Falconer functional:

$$\dim_F(\rho) = \inf \left\{ s > 0 : \sum_{\gamma \in \Gamma} e^{-F_s(\rho(\gamma))} < \infty \right\}.$$

Lastly, we recall the definition of the Hausdorff dimension:

Definition 1.5.4. Let (X, d) be a metric space and $A \subseteq X$ be a subset. We define the Hausdorff dimension of A to be

$$\dim_{\mathcal{H}}(A) = \inf \{s > 0 : \mathcal{H}_{\infty}^s(A) = 0\},$$

where \mathcal{H}_{∞}^s is the s -dimensional Hausdorff content of A , defined as

$$\mathcal{H}_{\infty}^s(A) = \liminf_{\delta \rightarrow 0} \left\{ \sum_{i=1}^{\infty} \text{diam}(U_i)^s : A \subseteq \bigcup_{i=1}^{\infty} U_i \right\}.$$

1.6 Hausdorff measures and densities

Definition 1.6.1. Let $V \in \mathcal{G}_{p+1}\mathbb{R}^d$ and $l \in \mathbb{P}(V)$. Using the canonical identification $T_l\mathbb{P}(V) \simeq \text{hom}(l, V/l)$, we define the density $|\Omega_{l,V}| : (T_l\mathbb{P}(V))^p \rightarrow \mathbb{R}^+$ by

$$|\Omega_{l,V}|(\phi_1, \dots, \phi_p) = \frac{\|v \wedge \tilde{\phi}_1(v) \wedge \dots \wedge \tilde{\phi}_p(v)\|}{\|v\|^{p+1}}$$

for any $v \in l - \{0\}$, where $\tilde{\phi}_1, \dots, \tilde{\phi}_p \in \text{hom}(l, V)$ are such that $\phi_i = \tilde{\phi}_i + \text{hom}(l, l)$ and $\|\cdot\|$ denotes the norm on $\bigwedge^{p+1} \mathbb{R}^d$ induced by the standard euclidean inner product.

Add some theory on densities and show that the density on $\mathcal{F}_{1,p+1}(\mathbb{R}^d)$ is $\text{SL}(d, \mathbb{R})$ equivariant

Chapter 2

Upper bound

In this section we prove the upper bound on the Hausdorff dimension, namely that for a projective Anosov representation in $\mathrm{SL}(d, \mathbb{R})$, the Hausdorff dimension of the limit set is bounded above by the Falconer dimension:

Lemma 2.2.1 (Upper bound for dimension). Let $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{R})$ be a projective Anosov representation. Then:

$$\dim_{\mathcal{H}}(\xi^1(\partial\Gamma)) \leq h_{\rho(\Gamma)}(F).$$

The idea of the proof is to find a covering whose Hausdorff content is dominated by the Dirichlet series of some functional Ψ_s^p , which will in turn imply that $\dim_{\mathcal{H}}(\xi^1(\partial\Gamma)) \leq h_{\rho(\Gamma)}(\Psi^p)$. Choosing then the most “effective” cover (i.e. the one which yields the smallest Hausdorff content up to a constant) we obtain that

$$\dim_{\mathcal{H}}(\xi^1(\partial\Gamma)) \leq h_{\rho(\Gamma)}(F) = h_{\rho(\Gamma)}\left(\max_p \Psi^p\right)$$

To obtain this we first cover $\xi^1(\partial\Gamma)$ by the shifted bassins of attraction $\rho(\gamma) \cdot B_{\alpha_1, \alpha}(\rho(\gamma))$ for $\gamma \in \Gamma$ satisfying $|\gamma| = T$. Then, for each $p \in \llbracket 2, d \rrbracket$, the set $\rho(\gamma) \cdot B_{\alpha_1, \alpha}(\rho(\gamma))$ is contained in an ellipsoid of axes lengths

$$\beta_2 = \frac{1}{\sin(\alpha)} \frac{\sigma_2}{\sigma_1}(\rho(\gamma)), \dots, \beta_d = \frac{1}{\sin(\alpha)} \frac{\sigma_d}{\sigma_1}(\rho(\gamma)).$$

which in turn is covered by roughly $\beta_2 \cdots \beta_{p-1} / \beta_p^{p-2}$ balls of radius $\sqrt{d-1} \beta_p$. Using this cover to bound the Hausdorff content of $\xi_p^1(\partial\Gamma)$, we obtain the bound

$$\mathcal{H}^s(\xi^1(\partial\Gamma)) \lesssim \sum_{|\gamma|=T} e^{-\Psi_s^p(\rho(\gamma))} \leq \sum_{|\gamma|=T} e^{-F_s(\rho(\gamma))}.$$

If we assume that $s > h_{\rho(\Gamma)}(F)$, then the right-hand side of the inequality converges to zero as we let $T \rightarrow \infty$, which means that $\mathcal{H}^s(\xi_p^1(\partial\Gamma)) = 0$, giving us the desired bound. Finally for each $q \in \llbracket 2, d \rrbracket$, we cover each ellipsoid by balls of some of radius β_q .

2.1 Lemmata

To simplify the proof of the bound, we present a few preliminary facts in the form of lemmata, whose proofs constitute the matter of this section. Our first goal will be to obtain an open

cover for the limit set using basins of attraction of elements in the Anosov subgroup at a fixed distance from the identity. It appears as [PSW23, Proposition 3.5].

Lemma 2.1.1. *Let $\rho : \Gamma \rightarrow SL(d, \mathbb{R})$ be projective Anosov. Then for $\alpha > 0$ small enough, there exists some $T_0 > 0$ such that for all $T \geq T_0$ the family*

$$\mathcal{U}_T = \{\rho(\gamma)B_{\alpha_1, \alpha}(\rho(\gamma)) : |\gamma| = T\}$$

is an open covering of $\xi^1(\partial\Gamma)$.

To prove this, we will need the following lemma ([PSW23, Lemma 2.4]). It essentially tells us that if we evaluate an Anosov representation at two points of a geodesic passing from $e \in \Gamma$ that are far apart, then the attracting line at one point is uniformly (with respect to geodesics) far apart from the attracting hyperplane at the other point.

Lemma 2.1.2. *Let $\rho : \Gamma \rightarrow SL(d, \mathbb{R})$ be a projective Anosov representation. For $\alpha > 0$ small enough, there exists $T > 0$ such that for any geodesic ray $(a_j)_{j \in \mathbb{N}}$ through e we have:*

$$\angle(U_1(\rho(a_i)), U_{d-1}(\rho(a_0))) > \alpha$$

when $|a_i|, |a_0| > T$.

Proof. Assume the contrary for the sake of contradiction. Then (see Figure 2.1) for each $n > 0$, there exists a geodesic ray $(a^n)_{j \in \mathbb{N}}$ through e such that

$$|a_n^n|, |a_0^n| > n \text{ and } \angle(U_1(\rho(a_n^n)), U_{d-1}(\rho(a_0^n))) < \frac{1}{n}.$$

Due to compactness of $\Gamma \cup \partial\Gamma$ we can find some subsequence k_n and $x, y \in \partial\Gamma$ such that $a_{k_n}^{k_n} \rightarrow x$, $a_0^{k_n} \rightarrow y$. Since there exists a geodesic joining $a_{k_n}^{k_n}, a_0^{k_n}$ passing from e , we also know that $x \neq y$. Also, the limit map being continuous (Theorem 3), we have that

$$\angle(\xi^1(x), \xi^{d-1}(y)) = 0,$$

which contradicts its transversality property. \square




Figure 2.1: Situation in Lemma 2.1.2

We may now proceed with the proof of Lemma 2.1.1:

Proof of Lemma 2.1.1. Let $\alpha, T > 0$ be as in the statement of Lemma 2.1.2 and $x \in \partial\Gamma$ be represented by a geodesic ray $(\gamma_j)_{j \geq 0}$ starting from e . Then $(\gamma_T^{-1}\gamma_j)_j$ is a geodesic ray starting from $(\gamma_T)^{-1}$ that passes through e , so

$$\angle(U_1(\rho(\gamma_T^{-1}\gamma_j)), U_{d-1}(\rho(\gamma_T^{-1}))) > \alpha$$

as implied by Lemma 2.1.2. Taking the limit $j \rightarrow \infty$ and using the equivariance of the limit map, we obtain

$$\angle(\rho(\gamma_T^{-1})\xi^1(x), U_{d-1}(\rho(\gamma_T^{-1}))) > \alpha$$

and thus $\xi^1(x) \in \rho(\gamma_T) \cdot B_{\alpha_1, \alpha}(\rho(\gamma_T))$. \square

The next step will be to cover each of the sets in \mathcal{U}_T by ellipsoids in the projective space, which are merely the projectivisation of the usual ellipsoids in \mathbb{R}^d . Note that while in the case of the latter, the axes are lines (for instance $\mathbb{R}e_1, \dots, \mathbb{R}e_d$), their projective analogue should be planes (for instance $\mathbb{R}u_1 \oplus \mathbb{R}u_2, \dots, \mathbb{R}u_1 \oplus \mathbb{R}u_d$).

Definition 2.1.1. Let V be a finite-dimensional \mathbb{R} -vector space. We consider a decomposition

$$V = \mathbb{R}u_1 \oplus \dots \oplus \mathbb{R}u_d$$

that is orthogonal with respect to a fixed inner-product over V . Given $\beta_2 \geq \dots \beta_d > 0$, we define an ellipsoid with axes $(u_1 \oplus u_p)_{p \in [2, d]}$ and lengths $(\beta_p)_{p \in [2, d]}$ to be the image of

$$\left\{ v = \sum_1^d v_i u_i \in V : \sum_2^d \left(\frac{v_j}{\beta_j} \right)^2 \leq 1 \right\}$$

through the projection $V \rightarrow \mathbb{P}(V)$.

Does this clarifying comment on the axes of the ellipsoid help?




Figure 2.2: Depiction in \mathbb{R}^3 of an ellipsoid of $\mathbb{P}(\mathbb{R}^2)$

When we want to distinguish between the projective ellipsoid of the above definition and the usual ellipsoid in \mathbb{R}^d , we will refer to the latter as a euclidean ellipsoid. When there is no ambiguity, we will simply refer to the projective ellipsoid as an ellipsoid.

The following is [PSW23, Proposition 3.8] and tells us that each set of the covering \mathcal{U}_T from Lemma 2.1.1 is contained in a projective ellipsoid, whose axes correspond to the axes of the euclidean ellipsoid $g \cdot \mathbb{S}^{d-1}$, and whose axes depend on the singular values of g . Recalling the discussion preceding Definition 1.4.1, it should not come as a surprise since the bassin of attraction is a set of flags that are stretched and contracted at a uniform degree under the action of g .

Correct ellipsoid drawing

Proposition 2.1.1. *Let $g \in \mathrm{SL}(d, \mathbb{R})$, $\alpha > 0$, fix a Cartan decomposition $g = k_g e^{\mu(g)} l_g$ and consider the corresponding choice of axes $u_i(g) = k_g e_i$, $u_i(g^{-1}) = l_g^{-1} e_{d-i+1}$ for $i \in \llbracket 1, d \rrbracket$, along with the respective flags $U_i(g), U_i(g^{-1})$. Denoting with*

$$B_{a_1, \alpha}(g) = \{x \in \mathbb{P}(\mathbb{R}^d) : \angle(x, U_{d-1}(g^{-1})) > \alpha\}$$

the bassin of attraction of g , the set $g \cdot B_{a_1, \alpha}(g)$ lies in the ellipsoid with axes $(u_1(g) \oplus u_p(g))_{p \in \llbracket 2, d \rrbracket}$ of lengths

$$\frac{1}{\sin \alpha} \cdot \frac{\sigma_p(g)}{\sigma_1(g)}$$

Proof. Using the definition of the basin of attraction (see Figure 2.3), for $w \in \mathbb{R}^d$, we have (up to considering $-w$ instead of w) that:

$$\angle(\mathbb{R}w, U_{d-1}(g^{-1})) = \angle(w, w') = \frac{|w_d|}{|w|},$$

when

$$\begin{aligned} w &= w_1 u_1(g^{-1}) + \cdots + w_d u_d(g^{-1}) \\ &= w_1 l_g^{-1} e_1(g) + \cdots + w_d l_g^{-1} e_d(g) \end{aligned}$$

and

$$\begin{aligned} w' &= w - w_d u_d(g^{-1}) = \\ &= w_1 u_1(g^{-1}) + \cdots + w_{d-1} u_{d-1}(g^{-1}) \\ &= w_1 l_g^{-1} e_1(g) + \cdots + w_{d-1} l_g^{-1} e_{d-1}(g). \end{aligned}$$

Then $\mathbb{R}w \in B_{a_1, \alpha}(g)$ if and only if

$$w_d^2 \geq (\sin \alpha)^2 \sum_{i=1}^d w_i^2.$$

Considering now some $v = v_1 u_1(g) + \cdots + v_d u_d(g)$ such that $\mathbb{R}v \in g \cdot B_{a_1, \alpha}(g)$ we have that

$$\begin{aligned} w &= g^{-1} v = v_1 \sigma_1(g)^{-1} l_g^{-1} e_1(g) + \cdots + v_d \sigma_d(g)^{-1} l_g^{-1} e_d(g) \\ &= v_1 \sigma_1(g)^{-1} u_d(g^{-1}) + \cdots + v_d \sigma_d(g)^{-1} u_1(g^{-1}) \end{aligned}$$

where we used that $u_p(g^{-1}) = l_g^{-1} e_{d+1-p}$. Hence

$$\sigma_1(g)^{-2} \cdot v_1^2 \geq (\sin \alpha)^2 \sum_{i=1}^d \sigma_i(g)^{-2} v_i^2.$$

Since the only condition on v is that the line through v lies in $g \cdot B_{a_1, \alpha}(g)$, we may choose v such that $v_1 = \sigma_1(g)$. Indeed, the only case that this would not be possible is when

$$v \in \mathbb{R}(u_2(g) \oplus \cdots \oplus u_d(g)) = g U_{d-1}(g^{-1}),$$

which is not possible since this would imply that $g^{-1}v \in U_{d-1}(g^{-1})$. Thus, the inequality above becomes precisely the condition for $\mathbb{R}w$ to lie in the projective ellipsoid. \square




Figure 2.3: Aid for Proposition 2.1.1

The following is [PSW23, Lemma 3.7]:

Lemma 2.1.3. *For any $p \in \llbracket 2, d \rrbracket$, an ellipsoid in $\mathbb{P}(\mathbb{R}^d)$ of axes lengths $\beta_2 \geq \dots \geq \beta_d$ is covered by*

$$2^{p-2} \frac{\beta_2 \cdots \beta_{p-1}}{\beta_p^{p-2}}$$

many (projected) balls of radius $\sqrt{d-1}\beta_p$.

Proof. We assume that E is an ellipsoid about $\mathbb{R}e_1$, so it suffices to cover its intersection $E_1 = E \cap U_1 \subseteq \mathbb{R}^{d-1}$ with the affine chart $U_1 = \{[x_1 : \dots : x_d] \in \mathbb{P}(\mathbb{R}^d) : x_1 \neq 0\}$. Clearly $E_1 \subseteq [-\beta_2, \beta_2] \times \dots \times [-\beta_d, \beta_d]$, so we proceed by covering the rectangle with side-lengths $2\beta_2, \dots, 2\beta_d$. Clearly each interval $(-\beta_j, \beta_j)$ is contained in the union of $\lceil \beta_j/\beta_p \rceil$ intervals of length $2\beta_p$, thus E_1 is contained in the union of

$$\left\lceil \frac{\beta_2}{\beta_p} \right\rceil \cdots \left\lceil \frac{\beta_{p-1}}{\beta_p} \right\rceil = \left\lceil \frac{\beta_2}{\beta_p} \right\rceil \cdots \left\lceil \frac{\beta_d}{\beta_p} \right\rceil$$

many squares of side-length $2\beta_p$. Since each such product is contained in a $(d-1)$ -ball of radius $\sqrt{d-1}\beta_p$ we may use at most

$$\left\lceil \frac{\beta_2}{\beta_p} \right\rceil \cdots \left\lceil \frac{\beta_{p-1}}{\beta_p} \right\rceil \leq \sum_{i \in \{0,1\}^{p-2}} \prod_{j=2}^{p-1} \left(\frac{\beta_j}{\beta_p} \right)^{i_j} \leq 2^{p-2} \frac{\beta_2}{\beta_p} \cdots \frac{\beta_{p-1}}{\beta_p}$$

many $(d-1)$ -balls of radius $\sqrt{d-1}\beta_p$ to cover E_1 . \square

The following can be found in [PSW23, Proposition 3.3]. It will not be used until Chapter 2, but we include it here to stress the fact that (similarly to all results of this chapter and in contrast to the next one) it holds for any projective Anosov representation, without any more assumptions (like Zariski-density of the subgroup or regularity of the limit set).

Proposition 2.1.2. *Let $\rho : \Gamma \rightarrow SL(d, \mathbb{R})$ be projective Anosov and $\alpha > 0$. Then there exist $c_0, c_1 > 0$ that depends only on α and ρ such that for all $\gamma \in \Gamma$:*

$$(\xi^1)^{-1}(B_{\alpha_1, \alpha}(\rho(\gamma))) \subseteq C_{c_0, c_1}^\infty(\gamma)$$

Proof. We begin by noting that it suffices to show this for all but finitely many $\gamma \in \Gamma$, since then we may alter the constants to satisfy the wanted inclusion for the finitely many remaining $\gamma \in \Gamma$ as well. Hence, we may assume that $|\gamma| \geq l_0$ where $l_0 > 0$ is such that

$$Ce^{-\mu l_0} < 1 \text{ and } a_1(\gamma) \geq C|\gamma| - L.$$

Suppose $x \in \partial\Gamma$ such that $\xi^1(x) \in B_{\alpha_1, \alpha}(\rho(\gamma))$, and consider a geodesic ray $a_j \rightarrow x$ starting from $a_0 = e$. To prove the result, it suffices to find constants c_0, c_1 independent of γ and for which there exists a (c_0, c_1) -quasi-geodesic from γ^{-1} to x that passes through e and stays at a bounded distance from $(a_j)_{j=0}^\infty$.

Using the exponential convergence rate of $\xi^1(a_j) \rightarrow \xi^1(x)$ and the definition of $B_{\alpha_1, \alpha}(\rho(\gamma))$ we have that:

$$\begin{aligned} d(\xi^1(a_j), \xi^1(\gamma)) &\geq d(\xi^1(x), U_1(\rho(\gamma^{-1})) - d(\xi^1(a_j), \xi^1(x))) \geq \\ &\geq d(\xi^1(x), U_{d-1}(\rho(\gamma^{-1})) - d(\xi^1(a_j), \xi^1(x))) \geq \sin \alpha - Ce^{-\mu j} \end{aligned}$$

which along with the uniform continuity of $\xi^1 : \Gamma \cup \partial\Gamma \rightarrow \mathbb{P}(\mathbb{R}^d)$ implies that there exists some $\alpha' > 0$ and $L > 0$ such that for all $j \geq L$:

$$d(a_j, \gamma^{-1}) \geq \alpha'.$$

Upon considering a large L , we may also assume that $|a_L| = L > l_0$. Note that both α' and L do not depend on each γ but only on ρ and α .

Using a coarse geometric argument, we can show that for all $j \geq L$

$$d(\gamma^{-1}, a_j) > \alpha' \Rightarrow d_{\mathcal{H}}([\gamma^{-1}, a_j], e) < \alpha''$$

for some α'' that depends only on Γ and α' , where $[a_j, \gamma^{-1}]$ denotes any geodesic segment connecting γ^{-1} and a_j , and $d_{\mathcal{H}}$ denotes the Hausdorff distance between sets. Indeed, [GH13, Lemme 2.17] states that $d([\gamma^{-1}, a_j]) \leq (\gamma_j^{-1}, a_j)_e + \delta$ where δ is the hyperbolicity constant of Γ . Thus

$$d([\gamma^{-1}, a_j]) \leq \delta + \frac{d(a_j, e) + d(\gamma^{-1}, e) + d(a_j, \gamma^{-1})}{2} \leq \delta + \frac{L + d(\gamma^{-1}, e) + \alpha'}{2}.$$

Consider the concatenation $(a'_j)_{j=L-K}^\infty$ of $[\gamma^{-1}, a_L]$ and $[a_L, x]$. To find quasi-geodesic-constants that are uniform in γ , we note that for any $c_0 \geq 1, c_1 \geq 0$:

$$c_0^{-1}|i - j| - c_1 \leq d(a'_i, a'_j) = d(a_i, a_j) \leq c_0|i - j| + c_1 \text{ when } i, j \geq L \text{ or } i, j \leq L$$

and that the upper bound follows trivially by the triangle inequality.

For the lower bound we proceed in two steps. First we bound the distance of $\gamma^{-1} = a'_{L-K}$ to a_{L+j} for $j \geq 0$:

$$\begin{aligned} d(a'_{L-K}, a'_{L+j}) &\geq \nu(|a_{L+j}| - |\gamma^{-1}|) - c'_0 - c'_1 |\log(d(U_1(\rho(a_{L+j})), U_1(\rho(\gamma^{-1}))))| \geq \\ &\geq \nu((L+j) + (K-L)) - c'_0 - c'_1 |\log(\sin a)| \geq \\ &= c_0^{-1}(j+K) - c_1 \end{aligned}$$


for $c_0 = \nu^{-1}, c_1 = c'_0 + c'_1 |\log(\sin \alpha)|$. The first inequality comes from [PSW23, Lemma 3.9]. For the second inequality we estimate $|\gamma^{-1}|$ from below using the triangle inequality. We are now ready to show that the concatenation $(a'_j)_j$ is indeed a (c_0, c_1) -geodesic:

$$\begin{aligned} d(a_{L+j}, a_{L-i}) &\geq d(a_{L+j}, a_{L_K}) - d(a_{L_K}, a_{L-i}) \geq c_0^{-1}(j+K) - c_1 - (K-i) \geq \\ &\geq c_0^{-1}(j+i) - c_1. \end{aligned}$$

Note however that (a'_j) does not necessarily lie in $C_{\infty}^{c_0, c_1}$ since it may not pass through e . For this reason we some $L - K \leq i_0 \leq L$ such that $|a_{i_0}| < \alpha''$, the existence of which is guaranteed by the fact that $d([\gamma^{-1}, a_L], e) < \alpha''$. We then consider alter (a'_j) at i_0 so that it passes through e to obtain

$$a''_j = \begin{cases} a_j & \text{for } j \neq i_0 \\ e & \text{for } j = i_0 \end{cases}$$

which is a $(c_0, c_1 + \alpha'')$ -quasigeodesic passing from e and converging to x . \square



2.2 Proof of bound

We are now ready to prove the upper bound of the Hausdorff dimension of the limit set by formalizing the proof strategy outlined in the beginning of this chapter.

Lemma 2.2.1 (Upper bound for dimension). *Let $\rho : \Gamma \rightarrow \text{SL}(d, \mathbb{R})$ be a projective Anosov representation. Then:*

$$\dim_{\mathcal{H}}(\xi^1(\partial\Gamma)) \leq h_{\rho(\Gamma)}(F).$$

Proof of Lemma 2.2.1. Let $p \in \llbracket 2, d \rrbracket$. Then using Proposition 2.1.1, Lemma 2.1.1, and Lemma 2.1.3 we have that for $T > 0$ large enough, $\xi^1(\partial\Gamma)$ is covered by the family

$$\mathcal{U}_T = \{\rho(\gamma)B_{\alpha_1, \alpha}(\rho(\gamma)) : |\gamma| = T\},$$

and that each basin $\rho(\gamma)B_{\alpha_1, \alpha}(\rho(\gamma))$ is in turn covered by

$$2^{p-2} \cdot \frac{\sigma_2(g) \cdots \sigma_{p-1}(g)}{\sigma_p(g)^{p-2}}$$

many balls of radius

$$\sqrt{d-1} \frac{1}{\sin \alpha} \frac{\sigma_p(g)}{\sigma_1(g)}.$$

By the definition of the Hausdorff measure, for $s \geq 0$:

$$\begin{aligned}
\mathcal{H}^s(\xi^1(\partial\Gamma)) &\leq \sum_{|\gamma|=T} 2^{p-2} \cdot \frac{\sigma_2(\rho(\gamma))}{\sigma_1(\rho(\gamma))} \dots \frac{\sigma_{p-1}(\rho(\gamma))}{\sigma_1(\rho(\gamma))} \left(\frac{\sigma_p(\rho(\gamma))}{\sigma_1(\rho(\gamma))} \right)^{-(p-2)} \left(\sqrt{d-1} \frac{1}{\sin \alpha} \frac{\sigma_p(\rho(\gamma))}{\sigma_1(\rho(\gamma))} \right)^s = \\
&= 2^{p-2} \cdot \left(\frac{\sqrt{d-1}}{\sin \alpha} \right)^s \sum_{|\gamma|=T} \frac{\sigma_2(\rho(\gamma))}{\sigma_1(\rho(\gamma))} \dots \frac{\sigma_{p-1}(\rho(\gamma))}{\sigma_1(\rho(\gamma))} \left(\frac{\sigma_p(\rho(\gamma))}{\sigma_1(\rho(\gamma))} \right)^{s-(p-2)} = \\
&= 2^{p-2} \cdot \left(\frac{\sqrt{d-1}}{\sin \alpha} \right)^s \sum_{|\gamma|=T} e^{-(\alpha_{12} + \dots + \alpha_{1(p-1)} + (s-(p-2))\alpha_{1p})\rho(\gamma)} \\
&= 2^{p-2} \cdot \left(\frac{\sqrt{d-1}}{\sin \alpha} \right)^s \sum_{|\gamma|=T} e^{-\Psi_s^p(\rho(\gamma))}
\end{aligned}$$

and thus

$$\mathcal{H}^s(\xi^1(\partial\Gamma)) \leq 2^{p-2} \cdot \left(\frac{\sqrt{d-1}}{\sin \alpha} \right)^s \sum_{|\gamma|=T} e^{-\max_p \Psi_s^p(\rho(\gamma))} \lesssim \sum_{|\gamma|=T} e^{-F_s(\rho(\gamma))}.$$

To see that the above implies the upper bound, consider some $s > h_{\rho(\Gamma)}(F)$. By the definition of the Falconer dimension, this implies that the Dirichlet series corresponding to the Falconer functional converges:

$$\sum_{\gamma \in \Gamma} e^{-F_s(\rho(\gamma))} < \infty$$

and in particular

$$\mathcal{H}^s(\xi^1(\partial\Gamma)) \leq \lim_{T \rightarrow \infty} \sum_{|\gamma|=T} e^{-F_s(\rho(\gamma))} = 0.$$

□

Looking again at the proof above, one could wonder whether the cover of the ellipsoid with axes β_2, \dots, β_d by balls of radius $\sqrt{d-1}\beta_p$ is optimal, in the sense that it gives the smallest Hausdorff content. To show that this is the case, consider the cover by balls of radius $r > 0$. This yields bounds of the form $\mathcal{H}^s(\xi_\rho^1(\partial\Gamma)) \lesssim \Phi_{r,s}$, where

$$\Phi_{r,s} = \begin{cases} \sum_{|\gamma|=T} r^s, & \text{for } r > \beta_2 \\ \sum_{|\gamma|=T} \frac{\beta_2 \dots \beta_{p-1}}{r^{p-2}} r^s, & \text{for } \beta_{p-1} < r < \beta_p, p \in \llbracket 3, d \rrbracket \\ \vdots \\ \sum_{|\gamma|=T} \frac{\beta_2 \dots \beta_d}{r^{d-1}} r^s, & \text{for } 0 < r < \beta_d. \end{cases}$$

The optimality of $r = \beta_p$ can be seen that for $s \in [0, 1]$, $\Phi_{r,s}$ is increasing in $[\beta_2, \infty]$, and decreasing in $[0, \beta_2]$, so it is minimized as a function of r for $r = \beta_2$, and thus gives the optimal Hausdorff content when $s \in [0, 1]$. Proceeding similarly, we see that the analogue holds for each of the other intervals as well.

Chapter 3

Lower bound

3.1 Busemann cocycle and Patterson–Sullivan measures

We denote with Π the set of simple positive roots, and for $\Theta \subseteq \Pi$ we consider the Levi-Anosov subspace of \mathfrak{a}

$$\mathfrak{a}_\Theta = \bigcap_{\alpha \notin \Theta} \ker \alpha,$$

which in particular admits $\{\omega_{\alpha_i} : \alpha_i \in \Theta\}$ as a basis.

Definition 3.1.1. Let $\Theta \subseteq \Pi$. We define the Busemann cocycle

$$b : \mathrm{PSL}(d, \mathbb{R}) \times \mathcal{F}_\Theta \rightarrow \mathfrak{a}_\Theta$$

as the unique element $b(g, kP_\Theta) \in \mathfrak{a}_\Theta$ such that

$$gk \in Ke^{b(g, kP_\Theta)}N.$$

where $N = \{n \in \mathrm{SL}(d, \mathbb{R}) : n_{ij} = 0 \text{ for } i > j, n_{ii} = 1 \text{ for all } i\}$ is the unipotent group of upper subgroup of upper triangular matrices with 1s on the diagonal, and P_Θ is the parabolic subgroup of $\mathrm{PSL}(d, \mathbb{R})$ corresponding to Θ , i.e. $\mathcal{F}_\Theta = \mathrm{PSL}(d, \mathbb{R})/P_\Theta$.

Lemma 3.1.1. For $y \in \mathcal{F}_\Theta(\mathbb{R}^d)$ we have

$$\omega_{\alpha_i}(b_\Theta(g, x)) = \log \frac{\|gv_1 \wedge \cdots gv_i\|}{\|v_1 \wedge \cdots v_i\|} \text{ for all } \alpha_i \in \Theta$$

for any basis v_1, \dots, v_i of $x^i \in \mathcal{G}_i(\mathbb{R}^d)$, where $\|\cdot\|$ denotes the norm on $\bigwedge^i \mathbb{R}^d$ induced by the euclidean inner product on \mathbb{R}^d , i.e. $\langle v_1 \wedge \cdots \wedge v_k, w_1 \wedge \cdots \wedge w_k \rangle := \det(\langle v_i, w_j \rangle)$.

Definition 3.1.2. We define

$$\Lambda^k : \mathrm{SL}(d, \mathbb{R}) \rightarrow \mathrm{SL}(\Lambda^k \mathbb{R}^d), \quad \Lambda^k : \mathcal{G}_k(\mathbb{R}^d) \rightarrow \mathbb{P}(\Lambda^k \mathbb{R}^d)$$

as

$$\Lambda^k(g)(v_1 \wedge \cdots \wedge v_k) = gv_1 \wedge \cdots \wedge gv_k, \quad \Lambda^k(\mathbb{R}v_1 \oplus \cdots \mathbb{R}v_k) = v_1 \wedge \cdots \wedge v_k$$

Lemma 3.1.2. Let $g \in \mathrm{SL}(d, \mathbb{R})$, $\alpha > 0$ and $k \in \llbracket 2, d \rrbracket$.

Correct the definition of the Busemann cocycle (see [Qui02] Lemme 6.1) Then add proof for Lemma 3.1.1

(i) $\omega_1(a(\Lambda^k g)) = \omega_k(a(g))$ and $\omega_1(b(\Lambda^k g, \Lambda^k y)) = \omega_k(b(g, y))$.

(ii) There exists some $\alpha' > 0$ independent of g such that $\Lambda^k B_{\mathbf{a}_k, \alpha}(\Lambda^k g) \subseteq B_{\mathbf{a}_1, \alpha'}(\Lambda^k g)$.

Proof. (i) Follows from the definitions of the fundamental weights and the Cartan projection.

(ii) Let $g = k_g e^{a(g)} l_g$ be the Cartan decomposition of g . Then using the definitions of the respective subspaces:

$$\begin{aligned} U_{d-k}(g^{-1} l_g^{-1}) &= \mathbb{R} e_{k+1} \oplus \cdots \oplus \mathbb{R} e_d \\ x_0 := U_{d-1}(\Lambda^k g^{-1} l_g^{-1}) &= \bigoplus_{\substack{i_1 < \cdots < i_k \\ (i_1, \dots, i_k) \neq (1, \dots, k)}} \mathbb{R} e_{i_1} \oplus \cdots \oplus \mathbb{R} e_{i_k} \end{aligned}$$

The first equality implies that

$$y \in B_{\mathbf{a}_k, \alpha}(g) \Leftrightarrow l_g y \in B_{\mathbf{a}_k, \alpha}(g l_g^{-1}) = B_{\mathbf{a}_k, \alpha}(\text{Id}),$$

so for every $y \in B_{\mathbf{a}_k, \alpha}(g)$ we have that

$$l_g y = l U_k(\text{Id}) \text{ for some } l \in L$$

where

$$L = \{l \in \text{SO}(d, \mathbb{R}) : l U_k(\text{Id}) \in B_{\mathbf{a}_k, \alpha}(\text{Id})\}.$$

Note that L is compact, being a closed subset of a compact group. Moreover, the fact that $\Lambda^k(y) \notin U_{d-1}(\Lambda^k g^{-1})$ implies that

$$0 < \angle(\Lambda^k(y), U_{d-1}(\Lambda^k g^{-1})) = \angle(\Lambda^k(l_g y), U_{d-1}(\Lambda^k g^{-1} l_g^{-1})) = \angle(\Lambda^k(l) U_k(\text{Id}), x_0)$$

The right-hand side is in the image of the compact set L under a continuous map, so it is bounded below by a positive number $\alpha' > 0$.

(iii) Follows from the definition of the Cartan projection and the Busemann cocycle. \square

Definition 3.1.3. For a discrete subgroup $H < \text{PSL}(d, \mathbb{R})$, $\phi \in (\alpha_\Theta)^*$, a (Γ, ϕ) -Patterson Sullivan measure on \mathcal{F}_Θ is a finite Radon measure μ such that for every $h \in H$

$$\frac{dh_* \mu}{d\mu}(x) = e^{-\phi(b_\Theta(h^{-1}, x))}, \text{ for all } x \in \mathcal{F}_\Theta(\mathbb{R}^d).$$

Lemma 3.1.3. Let $\alpha > 0, \Theta \subseteq \Pi$. There exists $K = K(\alpha) > 0$ such that for each $g \in \text{SL}(d, \mathbb{R})$, $\mathbf{a}_i \in \Theta$, $\mathbf{y} \in B_{\Theta, \alpha}(g)$, $\phi \in \mathfrak{a}_\Theta$

$$|\phi(a(g) - b(g, y))| \leq K.$$

Proof. We begin by noting that it suffices to consider the case where $\phi = \mathbf{a}_k$ for $\mathbf{a}_k \in \Theta$, since $\{\omega_i\}_{\mathbf{a}_i \in \Theta}$ is a basis for \mathfrak{a}_Θ^* .

We first consider the case where $k = 1$. We recall that the first component of the Cartan projection coincides with the spectral norm of g , i.e.

$$a_1(g) = \log \sup_{v \neq 0} \frac{\|gv\|}{\|v\|} = \log \|g k_2^{-1} e_1\|$$

where $g = k_1 e^{a(g)} k_2$ is the Cartan decomposition of g . Let $v = v_1 k_2^{-1} e_1 + \cdots + v_d k_2^{-1} e_d \in \mathbb{R}^d$ be such that $\|v\| = 1$ and $y = \mathbb{R}v$, we have

$$\begin{aligned} |\omega_1(a(g) - b(g, y))| &= |\log \|gk_2^{-1}e_1\| - \log \|gv\|| = \\ &= |\log |e^{a_1(g)}| - \log \|e^{a_1(g)}v_1k_1e_1 + \cdots + e^{a_d(g)}v_dk_1e_d\|| = \\ &= \left| \log \left\| v_1k_1e_1 + e^{-a_{12}(g)}v_2k_1e_2 + \cdots + e^{-a_{1d}(g)}v_dk_1e_d \right\| \right| \leq \\ &\leq |\log \|v_1\|| = |\log \sin(\angle(v, U_{d-1}(g^{-1})))| \leq |\log \sin \alpha|. \end{aligned}$$

For the case where $\Theta = \{\mathbf{a}_k\}$, we consider $\alpha' > 0$ such that

$$\Lambda^k(B_{\mathbf{a}_k, \alpha}(g)) \subseteq B_{\mathbf{a}_k, \alpha'}(\Lambda^k g)$$

Then using the case $k = 1$ we have that

$$|\omega_k(a(g) - b(g, y))| = |\omega_1(a(\Lambda^k g) - b(\Lambda^k g, \Lambda^k y))| \leq |\log \sin \alpha'|.$$

□

3.2 Proof strategy

Denoting with $d_\rho = \dim_{\mathcal{H}} \xi_\rho^1(\partial\Gamma)$ the Hausdorff dimension of the limit set, we shall outline how to obtain its lower bound by the Falconer dimension

$$d_\rho \geq h_{\rho(\Gamma)}(F).$$

First we recall that $F_s(a) = \max\{\Psi_s^p(a) : p \in \llbracket 2, d \rrbracket\}$, so the lower bound will follow once we have shown that

$$d_\rho \geq h_{\rho(\Gamma)}(\Psi^{d_\rho+1}),$$

since $h_{\rho(\Gamma)}(F) \leq h_{\rho(\Gamma)}(\Psi^{d_\rho+1})$.

But in Lemma 1.5.1, we showed that $\frac{s}{d_\rho} J_{d_\rho}^u \leq \Psi_{s+d_\rho}^{d_\rho+1}$, so the above bound will follow as soon as we have shown that

$$h_{\rho(\Gamma)}(J_{d_\rho}) \leq 1. \tag{LB}$$

which will be obtained using the method of Patterson–Sullivan–Quint. To apply the latter, apart from showing the existence of a $(J_{d_\rho}^u, \rho)$ -Patterson–Sullivan measure μ , we will also need to show that there exists a collection of open sets $\{U_\gamma\}_\gamma \in \Gamma$ such that

$$\mu(U_\gamma) \sim e^{-J_{d_\rho}^u(a(\rho(\gamma)))} \text{ and } M \stackrel{\text{def}}{=} \sup_{n \in \mathbb{N}} \max \left\{ \#A : A \subseteq \Gamma_n, \bigcap_{\gamma \in A} U_\gamma \neq \emptyset \right\} < \infty \tag{MP}$$

where $\Gamma_n = \{\gamma \in \Gamma : |\gamma| = n\}$. For the proof of the existence of a $(J_{d_\rho}^u, \rho)$ -Patterson–Sullivan measure that satisfies the property (MP) we refer to Section 3.3, noting that the Zariski-density assumption is necessary only for the equivalence appearing on the left hand side of Equation (MP). Assuming that for the time being, below we outline the Patterson–Sullivan–Quint method of obtaining Equation (LB).

Indeed, we first obtain the uniform in n bound:

$$\sum_{\gamma \in \Gamma_n} e^{-J_{d_\rho}^u(a(\rho(\gamma)))} \lesssim \sum_{\gamma \in \Gamma_n} \mu(U_\gamma) \leq \frac{1}{M} \mu(\mathcal{F}_\Theta(\mathbb{R}^d)) < \infty$$

along with the bound implied by the Anosov property of ρ :

$$J_{d_\rho}(a(\rho(\gamma))) \geq \mathbf{a}_{12}(a(\rho(\gamma))) \geq C|\gamma| - b$$

to conclude that

$$\sum_{\gamma \in \Gamma} e^{-(s+1)J_{d_\rho}^u(a(\rho(\gamma)))} = \sum_{n \geq 0} \sum_{\gamma \in \Gamma_n} e^{-sJ_{d_\rho}^u(a(\rho(\gamma)))} e^{-J_{d_\rho}^u(a(\rho(\gamma)))} \lesssim \sum_{n \geq 0} e^{-s(Cn-b)} < \infty$$

which holds for any $s > 0$, and thus Equation (LB) holds.

3.3 Existence of Patterson–Sullivan measure

Let $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{R})$ be a projective Anosov representation such that $\xi_\rho^1(\partial\Gamma)$ is a Lipschitz submanifold of dimension d_ρ . A naive approach would be to attempt to construct a Patterson–Sullivan measure over the limit set $\xi_\rho^1(\partial\Gamma)$. However, complications can arise in the Lipschitz regularity case, because the limit set does not admit a well-defined tangent space at every point. To overcome this difficulty, proceed as follows. To almost every $\xi \in \xi_\rho^1(\partial\Gamma)$, we have a well-defined tangent space $T_\xi \xi_\rho^1(\partial\Gamma)$. The latter, being a subspace of the tangent space $T_\xi(\mathbb{P}(\mathbb{R}^d))$ to the projective space $\mathbb{P}(\mathbb{R}^d)$, can be seen as the projectivisation (or a hyperplane) of a subspace of \mathbb{R}^d . In this way we can associate to almost every $\xi \in \xi_\rho^1(\partial\Gamma)$ an element of the flag space $\mathcal{F}_{1,d_\rho+1}(\mathbb{R}^d)$, thus defining a measurable section ζ_ρ . Restricting the density over the flag space defined in Section 1.6, we obtain an almost everywhere defined density

$$\xi \mapsto |\Omega_{\zeta_\rho(\xi)}|.$$

The following is [PSW23, Proposition 6.4]:

Proposition 3.3.1. *Let $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{R})$ be a projective Anosov representation, and assume that $\xi_\rho^1(\partial\Gamma)$ is a Lipschitz submanifold of dimension d_ρ . Then there exists a $(\rho(\Gamma), J_{d_\rho}^u)$ -Patterson–Sullivan measure on $\mathcal{F}_{1,d_\rho+1}(\mathbb{R}^d)$, which is supported on the image of a ρ -equivariant measurable section*

$$\zeta_\rho : \xi_\rho^1(\partial\Gamma) \rightarrow \mathcal{F}_{1,d_\rho+1}(\mathbb{R}^d)$$

of the projection $\mathrm{pr}_1 : \mathcal{F}_{1,d_\rho+1}(\mathbb{R}^d) \rightarrow \mathbb{P}(\mathbb{R}^d)$.

Proof. By Rademacher’s theorem, $\xi_\rho^1(\partial\Gamma)$ has a well-defined Lebesgue measure class, and Lebesgue-almost every $\xi_\rho^1(x) \in \xi_\rho^1(\partial\Gamma)$ admits a well-defined tangent space $T_{\xi_\rho^1(x)} \xi_\rho^1(\partial\Gamma)$. Considering such a $\xi_\rho^1(x)$ we let

$$\pi : \mathrm{hom}(\xi_\rho^1(x), \mathbb{R}^d) \rightarrow \mathrm{hom}(\xi_\rho^1(x), \mathbb{R}^d / \xi_\rho^1(x)) \simeq T_{\xi_\rho^1(x)} \mathbb{P}(\mathbb{R}^d),$$

and

$$x^{d_\rho+1} = \pi^{-1}(T_{\xi_\rho^1(x)} \xi_\rho^1(\partial\Gamma)) \xi_\rho^1(x) \in \mathcal{G}_{d_\rho+1}(\mathbb{R}^d).$$

Why do we construct a measure over $\mathcal{F}_{1,d_\rho+1}(\mathbb{R}^d)$ and not the limit set? We had talked about it but I forgot.

The calculation of the dimension follows from the rank-nullity theorem on π restricted to $\pi^{-1}(T_{\xi_\rho^1(x)}\xi_\rho^1(\partial\Gamma))$ and the fact that $\xi_\rho^1(\partial\Gamma) \leq \pi^{-1}(T_{\xi_\rho^1(x)}\xi_\rho^1(\partial\Gamma))$. Then Proposition A.0.2 implies that

$$T_{\xi_\rho^1(x)}\xi_\rho^1(\partial\Gamma) = \text{hom}(\xi_\rho^1(x), x^{d_\rho+1}/\xi_\rho^1(x)),$$

In this notation, we shall define (Lebesgue-almost everywhere) the map

$$\zeta_\rho : \xi_\rho^1(\partial\Gamma) \rightarrow \mathcal{F}_{1,d_\rho+1}(R^d), \quad \zeta_\rho(\xi_\rho^1(x)) = (\xi_\rho^1(x), x^{d_\rho+1}).$$

which is a section in the sense that the projection $\text{pr}_1 : \mathcal{F}_{1,d_\rho+1} \rightarrow \mathbb{P}(\mathbb{R}^d)$ to the first coordinate satisfies $\text{pr}_1 \circ \zeta_\rho = \text{Id}$. Moreover, it is $\rho(\Gamma)$ equivariant, since for $\gamma \in \Gamma$ we have

$$\begin{aligned} \rho(\gamma)\zeta_\rho(\xi_\rho^1(x)) &= \left(\rho(\gamma)\xi_\rho^1(x), \rho(\gamma)\pi^{-1}(T_{\xi_\rho^1(x)}\xi_\rho^1(\partial\Gamma)) \right) = \\ &= \left(\rho(\gamma)\xi_\rho^1(x), \rho(\gamma)\pi^{-1}(T_{\xi_\rho^1(x)}\xi_\rho^1(\partial\Gamma))\rho(\gamma)^{-1}\rho(\gamma)\xi_\rho^1(x) \right) = \\ &= \left(\rho(\gamma)\xi_\rho^1(x), \pi^{-1}(\rho(\gamma)T_{\xi_\rho^1(x)}\xi_\rho^1(\partial\Gamma)) \right) = \\ &= \left(\rho(\gamma)\xi_\rho^1(x), \pi^{-1}(T_{\rho(\gamma)\xi_\rho^1(x)}\xi_\rho^1(\partial\Gamma)) \right) = \\ &= \zeta_\rho(\rho(\gamma)\xi_\rho^1(x)) = \zeta_\rho(\xi_\rho^1(\gamma x)). \end{aligned}$$

We now define the non-negative density on $\xi_\rho^1(\partial\Gamma)$

$$\mu_{\xi_\rho^1(x)} = |\Omega_{\zeta_\rho(\xi_\rho^1(x))}|$$

Why is the section measurable?

which satisfies

$$\frac{d\rho(\gamma)_*\mu}{d\mu}(\xi) = \frac{d(\rho(\gamma)^{-1})^*\mu}{d\mu}(\xi) = e^{-J_{d_\rho+1}^u(b_\Theta(\rho(\gamma)^{-1}, \zeta_\rho(x)))},$$

where the term on the left-hand side involves measures, the term between the two equalities involves densities, and $\Theta = \{1, d_\rho+1\}$. The first equality holds for any density and its respective measure, while for the second one we consider $\phi_1, \dots, \phi_{d_\rho} \in T_{\xi_\rho^1(x)}\xi_\rho^1(\partial\Gamma)$ and calculate:

$$\begin{aligned} &(\rho(\gamma)^*\mu)_{\xi_\rho^1(x)}(\phi_1, \dots, \phi_{d_\rho}) \\ &= \mu_{\rho(\gamma)\xi_\rho^1(x)}(\rho(\gamma)\phi_1\rho(\gamma)^{-1}, \dots, \rho(\gamma)\phi_{d_\rho}\rho(\gamma)^{-1}) \\ &= \frac{\|\rho(\gamma)\xi_\rho^1(x) \wedge \rho(\gamma)\phi_1(\xi_\rho^1(x)) \wedge \dots \wedge \rho(\gamma)\phi_{d_\rho}(\xi_\rho^1(x))\|}{\|\rho(\gamma)\xi_\rho^1(x)\|^{d_\rho+1}} \\ &= \frac{\|\rho(\gamma)\xi_\rho^1(x) \wedge \rho(\gamma)\phi_1(\xi_\rho^1(x)) \wedge \dots \wedge \rho(\gamma)\phi_{d_\rho}(\xi_\rho^1(x))\|}{\|\xi_\rho^1(x) \wedge \phi_1(\xi_\rho^1(x)) \wedge \dots \wedge \phi_{d_\rho}(\xi_\rho^1(x))\|} \\ &\quad \cdot \frac{\|\xi_\rho^1(x) \wedge \phi_1(\xi_\rho^1(x)) \wedge \dots \wedge \phi_{d_\rho}(\xi_\rho^1(x))\|}{\|\rho(\gamma)\xi_\rho^1(x)\|^{d_\rho+1}} \cdot \frac{\|\rho(\gamma)\xi_\rho^1(x)\|^{d_\rho+1}}{\|\xi_\rho^1(x)\|^{d_\rho+1}} \\ &= e^{\omega_{d_\rho}(b_\Theta(\rho(\gamma), \zeta_\rho(\xi_\rho^1(x))))} \cdot \mu_{\xi_\rho^1(x)}(\phi_1, \dots, \phi_{d_\rho}) \\ &\quad \cdot e^{-(d_\rho+1)\omega_1(b_\Theta(\rho(\gamma), \zeta_\rho(\xi_\rho^1(x))))} \\ &= e^{-J_{d_\rho+1}^u(b_\Theta(\rho(\gamma)^{-1}, \zeta_\rho(\xi_\rho^1(x))))} \mu_{\xi_\rho^1(x)}(\phi_1, \dots, \phi_{d_\rho}). \end{aligned}$$

Finally, we let $\nu = \zeta_{\rho*}\mu$, which is the wanted Patterson–Sullivan measure on $\mathcal{F}_{1,d_{\rho}+1}(\mathbb{R}^d)$, since for $f \in C_c(\mathcal{F}_{1,d_{\rho}+1}(\mathbb{R}^d))$:

$$\begin{aligned} \int_{\mathcal{F}_{1,d_{\rho}+1}(\mathbb{R}^d)} f \, d(\rho(\gamma)_*\zeta_{\rho*}\mu) &= \int_{\xi_{\rho}^1(\partial\Gamma)} f \circ \rho(\gamma) \circ \zeta_{\rho} \, d\mu = \int_{\xi_{\rho}^1(\partial\Gamma)} f \circ \zeta_{\rho} \circ \rho(\gamma) \, d\mu = \\ &= \int_{\xi_{\rho}^1(\partial\Gamma)} f \circ \zeta_{\rho}(\xi_{\rho}^1(x)) e^{-J_{d_{\rho}}^u(b_{\Theta}(\rho(\gamma)^{-1}, \zeta_{\rho}(\xi_{\rho}^1(x)))} \, d\mu(\xi_{\rho}^1(x)) = \\ &= \int_{\mathcal{F}_{1,d_{\rho}+1}(\mathbb{R}^d)} f(y) e^{-J_{d_{\rho}}^u(b_{\Theta}(\rho(\gamma)^{-1}, y)} \, d(\zeta_{\rho*}\mu)(y) \end{aligned}$$

□

The next lemma is should regarded as an analog of Lemma 2.1.1 and Proposition 2.1.2 to arbitrary flag varieties, and relies only on the Anosov property of ρ , and the fact that ζ_{ρ} is a section of $\pi_{a_1} : \mathcal{F}_{1,d_{\rho},\Gamma+1}(\mathbb{R}^d) \rightarrow \mathbb{P}(\mathbb{R}^d)$.

Lemma 3.3.1. *Let $\rho : \Gamma \rightarrow SL(d, \mathbb{R})$ be projective Anosov and $\Theta = \{a_1, a_{d_{\rho},\Gamma}\} \subseteq \Pi$. Then for $\alpha > 0$ small enough, there exists some $C, T_0 > 0$ such that for all $T \geq T_0$ the family*

$$\mathcal{U}_T = \{\rho(\gamma)B_{\Theta,\alpha}(\rho(\gamma)) : |\gamma| = T\}$$

is an collection of open subsets of $\mathcal{F}_{\Theta}(\mathbb{R}^d)$ such that every $\zeta_{\rho}(\xi_{\rho}^1(x))$ is contained in at most C many sets in \mathcal{U}_T .

Proof. Suppose $\zeta_{\rho}(\xi_{\rho}^1(x)) \in \rho(\gamma)B_{\Theta,\alpha}(\rho(\gamma)) \cap \rho(\eta)B_{\Theta,\alpha}(\rho(\eta))$ for some $\gamma, \eta \in \Gamma_T$. Then $\xi_{\rho}^1(x) \in \rho(\gamma)B_{a_1,\alpha}(\rho(\gamma)) \cap \rho(\eta)B_{a_1,\alpha}(\rho(\eta))$. But using Proposition 2.1.2 we have that

$$\begin{aligned} x &\in (\xi_{\rho}^1)^{-1}(\rho(\gamma)B_{a_1,\alpha}(\rho(\gamma)) \cap \rho(\eta)B_{a_1,\alpha}(\rho(\eta))) = \\ &= \gamma(\xi_{\rho}^1)^{-1}(B_{a_1,\alpha}(\rho(\gamma))) \cap \eta(\xi_{\rho}^{-1})(B_{a_1,\alpha}(\rho(\eta))) \subseteq \\ &\subseteq \gamma C_{c_0,c_1}(\gamma) \cap \eta C_{c_0,c_1}(\eta). \end{aligned}$$

Thus x is represented by (c_0, c_1) -quasi-geodesic rays $(a_j)_0^{\infty}, (b_j)_0^{\infty}$, that start from e and pass from γ and η respectively. By Morse’s lemma, we know that there exists some geodesic ray starting from e and some $A > 0$ depending only on c_0, c_1 and the hyperbolicity constant of Γ such that the Hausdorff distance of the geodesic ray to each of the quasi-geodesics is bounded by A . Let ϵ_0, ϵ_1 be two points on the geodesic ray such that $d(\gamma, \epsilon_0), d(\eta, \epsilon_1) \leq A$. Then we have that

$$\begin{aligned} d(\gamma, \eta) &\leq d(\gamma, \epsilon_0) + d(\epsilon_0, \epsilon_1) + d(\epsilon_1, \eta) \leq 2A + \|\epsilon_0\| - \|\epsilon_1\| \leq \\ &\leq 2A + \|\epsilon_0\| - |\gamma| + |\gamma| - |\eta| + \|\epsilon_1\| - |\eta| \leq 4A. \end{aligned}$$

In particular, any γ' such that $\zeta_{\rho}(\xi_{\rho}^1(\gamma')) \in \xi_{\rho}^1(\rho(\gamma')B_{\Theta,\alpha}(\rho(\gamma')))$, will lie in a ball of radius $4A$ around γ . Since Γ is finitely generated, there exists some $C > 0$ such that the ball of radius $4A$ around γ contains at most C elements of Γ . □

Before giving the next definition, we recall that the annihilator of an element $y \in \mathcal{F}_F i\Theta(\mathbb{R}^d)$ is the set of partial flags that are not transverse to y , that is:

$$\text{Ann}(y) = \{x \in \mathcal{F}_{\Theta}(\mathbb{R}^d) : x^{\theta} \cap y^{d-\theta} \neq 0 \text{ for some } \theta \in \Theta\}.$$

Definition 3.3.1. Let $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{R})$ be a linear representation, $\Theta \subseteq \Pi$ and μ a $(\rho(\Gamma), \phi)$ -Patterson–Sullivan measure over $\mathcal{F}_\Theta(\mathbb{R}^d)$. We say that ρ is μ -irreducible there is no element in $\mathcal{F}_{i\Theta}(\mathbb{R}^d)$, whose annihilator is of full measure, i.e. for all $y \in \mathcal{F}_{i\Theta}(\mathbb{R}^d)$:

$$\mu(\mathrm{Ann}(y)) < \mu(\mathcal{F}_\Theta(\mathbb{R}^d)).$$

Example 3.3.1. If $\rho(\Gamma)$ is Zariski-dense in $\mathrm{SL}(d, \mathbb{R})$, then ρ is μ -irreducible for any ρ -quasi-equivariant measure μ , and in particular for any $(\rho(\Gamma), \phi)$ -Patterson–Sullivan measure.

Remark 3.3.1. The reason that we introduce the concept of μ -irreducibility is that for any μ -irreducible representation $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{R})$, there exist $\alpha, \kappa > 0$ such that $\mu(B_{\Theta, \alpha}(\rho(\gamma))) \geq \kappa$ for all $\gamma \in \Gamma$.

Indeed, if this were not the case, then there would exist a sequence $\alpha_n \searrow 0$ and $\gamma_n \in \Gamma$ such that

$$\mu(B_{\Theta, \alpha_n}(\rho(\gamma_n))) \leq \frac{1}{n}.$$

Due to the compactness of $\mathcal{F}_\Theta(\mathbb{R}^d)$, up to considering a subsequence, we may assume that the repelling flags or $\rho(\gamma_n)$ converge to some $\xi \in \mathcal{F}_\Theta(\mathbb{R}^d)$:

$$(U_{d-i}(\rho(\gamma_n)^{-1}))_{a_i \in \Theta} \rightarrow \xi$$

In that case, the complements $B_{\Theta, \alpha_n}^c(\rho(\gamma_n))$ will converge to the annihilator of ξ , in the sense:

$$\limsup_n B_{\Theta, \alpha_n}^c(\rho(\gamma_n)) \subseteq \mathrm{Ann}(\xi).$$

Indeed, let $y \in \limsup_n B_{\Theta, \alpha_n}^c(\rho(\gamma_n))$ and consider a subsequence k_n such that $y \in B_{\Theta, \alpha_n}^c(\rho(\gamma_{k_n}))$. By the very definition of $B_{\Theta, \alpha_n}^c(\rho(\gamma_n))$, there exists some p such that up to considering a subsequence of k_n ,

$$\angle(y^p, U_{d-p}(\rho(\gamma_n)^{-1})) \leq \alpha_n$$

holds. Taking the limit as $n \rightarrow \infty$, we have that $y^p \cap \xi^{d-p} \neq 0$ and hence $y \in \mathrm{Ann}(\xi)$.

Using a measure-theoretic argument we conclude that $\mathrm{Ann}(\xi)$ is of full measure, which contradicts the μ -irreducibility of ρ :

$$\mu(\mathrm{Ann}(\xi)) \geq \mu(\limsup_n B_{\Theta, \alpha_n}^c(\rho(\gamma_{k_n}))) \geq \limsup_n \mu(B_{\Theta, \alpha_n}^c(\rho(\gamma_{k_n}))) = \mu(\mathcal{F}_\Theta(\mathbb{R}^d)).$$

Lemma 3.3.2. Let $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{R})$ be a representation and μ^ϕ be a $(\rho(\Gamma), \phi)$ -Patterson–Sullivan measure. If $\rho(\Gamma)$ is μ -irreducible, then there exists some $\alpha_0 > 0$, such that for any $\alpha \in (0, \alpha_0)$, there's some $k = k(\alpha) > 0$ for which

$$\frac{1}{k} e^{-\phi(a(\rho(\gamma)))} \leq \mu^\phi(\rho(\gamma) B_{\Theta, \alpha}(\rho(\gamma))) \leq k e^{-\phi(a(\rho(\gamma)))}$$

for all $\gamma \in \Gamma$.

Proof. Let $\alpha_0, k > 0$ be as in the remark preceeding the statement of the lemma. As noted in Lemma 3.1.3, there exists some $K = K(\alpha_0, \phi) > 0$ such that for any $\alpha \in (0, \alpha_0)$ and $y \in B_{\Theta, \alpha}(\rho(\gamma))$:

$$|\phi(a(\rho(\gamma))) - b(\rho(\gamma), y)| \leq K,$$

from which we obtain the upper bound

$$\begin{aligned} \mu^\phi(\rho(\gamma) B_{\Theta, \alpha}(\rho(\gamma))) &= (\rho(\gamma)^{-1})_* \mu^\phi(B_{\Theta, \alpha}(\rho(\gamma))) = \int_{\mathcal{F}_\Theta(\mathbb{R}^d)} e^{-\phi(b(\rho(\gamma), y))} d\mu^\phi(y) \leq \\ &\leq e^{-K} \mu^\phi(\mathcal{F}_\Theta(\mathbb{R}^d)) e^{-\phi(a(\rho(\gamma)))}. \end{aligned}$$

Similarly we obtain the lower bound. □

3.4 Proof of the main theorem

In this section we shall prove the main theorem, which we restate for the reader's convenience:

Theorem 4. *Let $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{R})$ be a Zariski-dense, projective Anosov representation such that $\xi_\rho^1(\partial\Gamma)$ is a Lipschitz submanifold of $\mathbb{P}(\mathbb{R}^d)$. Then the dimension of the limit set $\xi_\rho^1(\partial\Gamma)$ equals the Falconer dimension of ρ :*

$$\dim \xi_\rho^1(\partial\Gamma) = h_{\rho(\Gamma)}(F)$$

where $F_s(a) = \max\{\Psi_s^p(a) : p \in \llbracket 2, d \rrbracket\}$ is the Falconer functional.

Proof. We have already seen in Lemma 2.2.1 that $d_{\rho, \Gamma} \leq h_{\rho(\Gamma)}(F)$. For the opposite inequality, we merely need to piece together the results of the previous sections as outlined in Section 3.2. There we have seen that $h_{\rho(\Gamma)}(F) \leq h_{\rho(\Gamma)}(\Psi^{d_{\rho, \Gamma}+2})$ since $F_s \geq \Psi_s^{d_{\rho, \Gamma}+2}$, so may as well show that $h_{\rho(\Gamma)}(\Psi^{d_{\rho, \Gamma}+2}) \leq d_{\rho, \Gamma}$, i.e. that

$$\sum_{\gamma \in \Gamma} e^{-\Psi_s^{d_{\rho, \Gamma}+2}(\rho(\gamma))} < \infty$$

for all $s \geq d_{\rho, \Gamma}$. This will follow as soon as we have shown that $h_{\rho(\Gamma)}(J_{d_{\rho, \Gamma}}) \leq 1$, since

$$\begin{aligned} \Psi_s^{d_{\rho, \Gamma}+1} \circ a &= \mathbf{a}_{12} + \cdots + \mathbf{a}_{1(d_{\rho, \Gamma}+1)} + (s - d_{\rho, \Gamma})\mathbf{a}_{1(d_{\rho, \Gamma}+2)} = \\ &= \mathbf{a}_{12} + \cdots + \mathbf{a}_{1(d_{\rho, \Gamma}+1)} + d_{\rho, \Gamma} \left(\frac{s}{d_{\rho, \Gamma}} - 1 \right) \mathbf{a}_{1(d_{\rho, \Gamma}+2)} \geq \\ &\geq \mathbf{a}_{12} + \cdots + \mathbf{a}_{1(d_{\rho, \Gamma}+1)} + \left(\frac{s}{d_{\rho, \Gamma}} - 1 \right) (\mathbf{a}_{12} + \cdots + \mathbf{a}_{1(d_{\rho, \Gamma}+1)}) = \\ &= \frac{s}{d_{\rho, \Gamma}} J_{d_{\rho, \Gamma}}^u. \end{aligned}$$

Using the Anosov property of ρ we have that

$$J_{d_{\rho}}(a(\rho(\gamma))) \geq \mathbf{a}_{12}(a(\rho(\gamma))) \geq C|\gamma| - b$$

for certain $C, b > 0$ which when we break up the sum defining the critical exponent into the sum over the sets $\Gamma_T = \{\gamma \in \Gamma : |\gamma| = T\}$ gives us:

$$\begin{aligned} \sum_{\gamma \in \Gamma} e^{-(s+1)J_{d_{\rho}, \Gamma}^u(a(\rho(\gamma)))} &= \sum_{T \geq 0} \sum_{\gamma \in \Gamma_T} e^{-sJ_{d_{\rho}, \Gamma}^u(a(\rho(\gamma)))} e^{-J_{d_{\rho}, \Gamma}^u(a(\rho(\gamma)))} = \\ &= \sum_{T \geq 0} e^{-sJ_{d_{\rho}, \Gamma}^u(a(\rho(\gamma)))} \sum_{\gamma \in \Gamma_T} e^{-J_{d_{\rho}, \Gamma}^u(a(\rho(\gamma)))} \leq \\ &\leq \sum_{T \geq 0} e^{-s(CT-b)} \sum_{\gamma \in \Gamma_T} e^{-J_{d_{\rho}, \Gamma}^u(a(\rho(\gamma)))} \end{aligned}$$

To obtain a bound on the inner sums that is uniform in T , we recall Proposition 3.3.1. There we saw that $\xi_\rho^1(\partial\Gamma)$ being a Lipschitz submanifold of $\mathbb{P}(\mathbb{R}^d)$ implies the existence of a $(\rho(\Gamma), J_{d_{\rho, \Gamma}}^u)$ -Patterson–Sullivan measure μ on $\zeta_\rho^1(\xi^1(\partial\Gamma)) \subseteq \mathcal{F}_{1, d_{\rho, \Gamma}+1}(\mathbb{R}^d)$. By Lemma 3.3.1 we have that for $\alpha > 0$ small enough, there exists some $M, T_0 > 0$ such that for all $T \geq T_0$ the family

$$\mathcal{U}_T = \{\rho(\gamma)B_{\alpha, \Theta}(\rho(\gamma)) : |\gamma| = T\}$$

is an open covering of $\zeta_\rho(\xi_\rho^1(\partial\Gamma))$ for which

$$\max \left\{ \#A : A \subseteq \Gamma_T, \bigcap_{\gamma \in A} \rho(\gamma)B_{\alpha,\Theta}(\rho(\gamma)) \neq \emptyset \right\} \leq M.$$

But μ is in particular ρ -quasi-equivariant which along with the Zariski-density of $\rho(\Gamma)$ implies that ρ is μ -irreducible, as we have seen in Example 3.3.1. Hence the bound in Lemma 3.3.2 applies and we have that

$$\begin{aligned} \sum_{\gamma \in \Gamma_T} e^{-J_{d_{\rho,\Gamma}}^u(a(\rho(\gamma)))} &\leq \sum_{\gamma \in \Gamma_T} \mu(\rho(\gamma)B_{\alpha,\Theta}(\rho(\gamma))) \leq \\ &\leq \frac{1}{M} \mu(\mathcal{F}_{1,d_{\rho,\Gamma}+1}(\mathbb{R}^d)) < \infty. \end{aligned}$$

□

Chapter 4

Dropping the Zariski-density assumption

Since many of interesting examples of Anosov subgroups are not Zariski-dense, one could certainly argue that the Zariski-density assumption in Theorem 4 is not cheap. The reason that it's needed is that otherwise one can't guarantee that it is μ -irreducible for a Patterson–Sullivan measure μ . In fact, Zariski-density is not assumed in [PSW23], where the authors suggest a way to obtain μ -irreducibility through the following lemma which as we will see below is false.

Lemma 4.0.1 (Lemma 6.8 in [PSW23]). *Let Γ be a hyperbolic group and $\eta : \Gamma \rightarrow \mathrm{PGL}(d, \mathbb{R})$ be a strongly irreducible projective Anosov representation such that $\xi_\eta(\partial\Gamma)$ is homeomorphic to \mathbb{S}^{d_ρ} , and which admits a measurable η -equivariant section $\zeta : \partial\Gamma \rightarrow \mathcal{F}_{\{\mathbf{a}_1, \mathbf{a}_{d_\rho+1}\}}(\mathbb{R}^d)$. Then η is μ -irreducible for any $(\eta(\Gamma), \phi)$ -Patterson–Sullivan measure μ on $\zeta(\partial\Gamma) \subseteq \mathcal{F}_{\{\mathbf{a}_1, \mathbf{a}_{d_\rho+1}\}}(\mathbb{R}^d)$.*

For convenience, we recall that a linear representation $\rho : \Gamma \rightarrow \mathrm{GL}(d, \mathbb{R})$ is strongly irreducible if there is no proper $\rho(\Gamma)$ -invariant subspace of \mathbb{R}^d , and it is μ -irreducible if there is no element in $\mathcal{F}_{i\Theta}(\mathbb{R}^d)$, whose annihilator is of full measure.

More specifically, in the proof of the preceeding lemma, the authors argue by contradiction that if the result is not true, then there exists subspaces $W_0 \in \mathcal{G}_{d-d_\rho-1}$ and $V \in \mathcal{G}_{d-d_\rho-1}$ such that

$$\eta(\gamma)V \cap W_0 \neq 0$$

for all $\gamma \in \Gamma$. Indeed, if this were not the case, then there exist $(W_0, P_0) \in \mathcal{F}_{\mathbf{a}_{d-d_\rho-1}, \mathbf{a}_{d-1}}$ such that $\mathrm{Ann}(W_0, P_0)$ is of full μ measure. In fact, η -equivariance of ζ implies that one can choose W_0, P_0 such that $P_0 \in \xi_\eta^{d-1}(\partial\Gamma)$, meaning that for μ -almost every $(\xi_\eta^1(x), \zeta(x)^{d_\rho+1}) \in \zeta(\partial\Gamma)$, we have that $\zeta(x)^{d_\rho+1} \cap W_0 \neq 0$. Again using the η -equivariance of ζ , we have that for all $\gamma \in \Gamma$, the set $\{\zeta(x) \in \zeta(\partial\Gamma) : \zeta(x)^{d_\rho+1} \cap \gamma W_0 \neq 0\}$ is of full measure, which implies that their intersection is non-empty, and we can take $V = \zeta(x)^p$ for any $x \in \partial\Gamma$ such that $\zeta(x)$ lies in this intersection.

Finally to conclude, the authors assert that this is absurd by using [Lab06, Proposition 10.3]:

Proposition 4.0.1 (Labourie). *Let $G \leq \mathrm{SL}(d, \mathbb{R})$ be an algebraic group. If $C \in \mathcal{G}_k(\mathbb{R}^d)$, $B \in \mathcal{G}_{d-k}(\mathbb{R}^d)$ such that*

$$gC \cap B \neq 0, \tag{4.1}$$

then the connected component of G containing the identity element is not irreducible.

However, the results of Lemma 4.0.1 and Proposition 4.0.1 are false, leaving room for further investigation. In particular, below we present a direct counterexample to Proposition 4.0.1, followed by a construction which shows that Equation (4.1) does not contradict strong irreducibility, and where the representation η is not μ -irreducible, thus disproving Lemma 4.0.1. Nevertheless, if we proceed to calculate the dimension of the respective limit set and the critical exponent of the Falconer functional, we notice that they coincide and hence do not provide a counterexample to the main result presented in [PSW23], which suggests that the result may still hold at least for certain special cases.

Example 4.0.1. Consider the space $\mathbb{R}^{d,d}$, that is \mathbb{R}^{2d} with the bilinear form $\langle x, y \rangle = x_1 y_1 + \dots + x_d y_d - x_{d+1} y_{d+1} - \dots - y_{2d}$, and let $S \subseteq \mathcal{G}_d(\mathbb{R}^{2d})$ to be the set of maximal isotropic subspaces of $\mathbb{R}^{d,d}$. Assuming that d is odd, one can show that for any subspaces V, W in the same component of S , we have $gV \cap W \neq 0$ for all $g \in \mathrm{SO}(d, d)$. However, for $d > 0$ of any parity, we have that $\mathrm{SO}_0(d, d)$ acts irreducibly on $\mathrm{SO}(d, d)$, which means that the choice of any V, W in the same connected component of S provides a counterexample to Proposition 4.0.1 for $G = \mathrm{SL}(2d, \mathbb{R})$.

Outline
proof steps
for coun-
terex-
ample to
Labourie's
lemma

After providing a counterexample to Proposition 4.0.1 as above, one could object that this does not necessarily imply that the result of Lemma 4.0.1 is false. For this, one would need to provide another counterexample for Lemma 4.0.1 as well, which is the goal of the next pages and is summarised in Proposition 4.0.2. Before giving the statement of the latter, we recall that a uniform lattice Γ of a locally compact group G is a discrete subgroup of G that is co-compact, i.e. G/Γ is compact.

Proposition 4.0.2. *Let $\Gamma \leq \mathrm{SU}(2, 1)$ be a uniform lattice and $\eta : \Gamma \rightarrow \mathrm{SL}(\mathfrak{su}(2, 1))$ be the restriction of the adjoint representation. Then*

- (i) η is strongly irreducible,
- (ii) η is projective Anosov
- (iii) η admits a measurable η -equivariant section:

$$\zeta : \partial\Gamma \rightarrow \mathcal{F}_{\{1,4\}}(\mathfrak{su}(2, 1)) \simeq \mathcal{P}$$

$$x \mapsto (\xi^1(x), T_{\xi^1(x)}\xi^1(\partial\Gamma)) \simeq (\xi^1(x), (d_{\xi^1(x)}p)^{-1}(T_{\xi^1(x)}\xi^1(\partial\Gamma))\xi^1(x)).$$

where $d_{\xi^1(x)}p : \mathrm{hom}(\xi^1(x), \mathfrak{su}(2, 1)) \rightarrow \mathrm{hom}(\xi^1(x), \mathfrak{su}(2, 1)/\xi^1(x))$ is the canonical projection.

- (iv) For all $x, y \in \partial\Gamma : \zeta(x)^4 \cap \zeta(y)^4 \neq 0$.
- (v) For any $y_0 \in \mathrm{SU}(2, 1)/P_0$ and $W_0 \in \mathcal{G}_7(\mathbb{R}^4)$ that contains $\zeta(y_0)^4$, we have that $\mathrm{Ann}(\zeta(y_0)^4, W_0) \supseteq \zeta(\mathrm{SU}(2, 1)/P_0)$ and is in particular of full μ -measure, for any $(\eta(\Gamma), \phi)$ -Patterson-Sullivan measure μ supported over $\zeta(\partial\Gamma)$.
- (vi)

$$h_{\rho(\Gamma)}(F) = \dim_{\mathcal{H}}(\xi_{\rho}^1(\partial\Gamma)) = 3.$$

For the sake of readability, we have broken up the proof of Proposition 4.0.2 into multiple propositions which are then combined.

Remark 4.0.1. When $G = \mathrm{Isom}(X)$ is the isometry group of a complete Riemannian manifold X , and Γ is a uniform lattice of G , then it acts properly discontinuously and cocompactly on X .

We begin by showing proving the Anosov property of η .

Proposition 4.0.3. *Let Γ be a uniform lattice of $\mathrm{SU}(2, 1)$, and $\eta : \Gamma \rightarrow \mathrm{SL}(\mathfrak{su}(2, 1))$ be the restriction of the adjoint representation: $\eta(\gamma) = \mathrm{Ad}_\gamma$ for all $\gamma \in \Gamma$. Then η is projective Anosov.*

Proof. Let $\gamma \in \Gamma$. Since $\gamma \in \mathrm{SU}(2, 1)$, we have that

$$\gamma = k_1 \exp(r(\gamma)x_0) k_2$$

for x_0 a fixed non-zero in the Weyl-chamber $\mathbb{R}x_0$ of $\mathfrak{su}(2, 1)$, $r(\gamma) \in \mathbb{R}$ and $k_1, k_2 \in \mathrm{U}(2)$. Then by the definition of a uniform lattice, we have that Γ acts properly discontinuously and cocompactly, which means that the inclusion $\Gamma \hookrightarrow \mathrm{SU}(2, 1)$ is projective Anosov (since $\mathrm{SU}(2, 1)$ is of rank 1). Thus there exist constants $L \geq 1, b \geq 0$ such that for all $\gamma \in \Gamma$:

$$r(\gamma) \geq \mathfrak{a}_1(x_0)^{-1}(L|\gamma| - b) = L'|\gamma| - b'.$$

Note that $\mathfrak{a}_1(x_0) > 0$ since x_0 is in the interior of the Weyl-chamber $\mathbb{R}x_0$.

Letting $k'_1 = \mathrm{Ad}_{k_1}, k'_2 = \mathrm{Ad}_{k_2}$ and $K' \leq \mathrm{SL}(\mathfrak{su}(2, 1))$ be a maximal compact subgroup containing them, we have that:

$$\eta(\gamma) = \mathrm{Ad}_\gamma = k'_1 \mathrm{Ad}_{\exp(r(\gamma)x_0)} k'_2 = k'_1 \exp(r(\gamma) \mathrm{ad}_{x_0}) k'_2.$$

Thus

$$\mathfrak{a}_1(\mu(\eta(\gamma))) = r(\gamma) \mathfrak{a}_1(\mathrm{ad}_{x_0}) \geq (L'|\gamma| - b') \mathfrak{a}_1(x_0)$$

which is Anosov because $\mathfrak{a}_1(\mathrm{ad}_{x_0}) > 0$, as can be seen by concrete calculations. \square

Before giving an expression for the projective part of the limit map of η , we make a few observations regarding Gromov boundary of Γ . In particular, we claim that since Γ is a uniform lattice of $\mathrm{SU}(2, 1)$, we have that $\partial\Gamma$ is homeomorphic to $\mathrm{SU}(2, 1)/P$, where P is a parabolic subgroup of $\mathrm{SU}(2, 1)$, and it coincides with the stabilizer of some isotropic line $l \in \partial_\infty \mathbb{H}_{\mathbb{C}}^2$.

Indeed, for a uniform lattice Γ of the isometry group G of a homogenous G -space X , the Milnor-Švarc lemma implies that for any $x_0 \in X$, the map $\Gamma \rightarrow X, \gamma \mapsto \gamma x_0$ is a quasi-isometry. In our case $G = \mathrm{SU}(2, 1)$ and $X = \mathbb{H}_{\mathbb{C}}^2$ is a hyperbolic metric space, so the quasi-isometry extends to a homeomorphism $\partial\Gamma \rightarrow \partial\mathbb{H}_{\mathbb{C}}^2$ of the Gromov-boundaries. On the other hand, the action of $\mathrm{SU}(2, 1)$ on $\partial_\infty \mathbb{H}_{\mathbb{C}}^2$ is transitive, so we have that $\partial\mathbb{H}_{\mathbb{C}}^2 \simeq \mathrm{SU}(2, 1)/P$ where P is the stabilizer of a point in $\partial\mathbb{H}_{\mathbb{C}}^2$. In fact, we have that P is a parabolic subgroup of $\mathrm{SU}(2, 1)$. The combination of the above, along with the fact that the geometric and the Gromov boundaries agree in the case of $\mathbb{H}_{\mathbb{C}}^2$, we deduce that $\partial\Gamma \simeq \mathrm{SU}(2, 1)/P$.

To calculate the projective part of the limit map, we shall show that there exists a unique $\mathrm{SU}(2, 1)$ -equivariant map from $\partial\Gamma$ to $\mathbb{P}(\mathbb{R}^d)$. The uniqueness follows from the following characterisation of limit maps:

Lemma 4.0.2. *Let $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{R})$ be a strongly irreducible projective Anosov representation, and denote with $\xi_\rho : \partial\Gamma \rightarrow \mathbb{P}(\mathbb{R}^d)$ its limit map. Then ξ_ρ^1 is the unique continuous, $\rho(\Gamma)$ -equivariant map from $\partial\Gamma$ to $\mathbb{P}(\mathbb{R}^d)$.*

Proof. Let $\eta^1 : \partial\Gamma \rightarrow \mathbb{P}(\mathbb{R}^d)$ be a continuous, $\rho(\Gamma)$ -equivariant map. Since the action of Γ on its boundary $\partial\Gamma$ has dense orbits, it suffices to show that it agrees with ξ_ρ^1 on at least one boundary point.

Suppose for the sake of contradiction that η^1 does not coincide with ξ^1 and let $z \in \partial\Gamma, y \in \partial\Gamma \setminus \{z\}$. Then for any $x \in \partial\Gamma \setminus \{y\}$ we may find some quasi-geodesic $\{\gamma_n\}_n$ such that $\gamma_n \rightarrow$

$x, \gamma_{-n} \rightarrow y$ as $n \rightarrow \infty$. Then since $z \neq y$ we know that $\gamma_n z \rightarrow z$ as $n \rightarrow \infty$ and continuity of η^1 implies that $\eta^1(\gamma_n z) \rightarrow \eta^1(z)$. But equivariance of η^1 and the fact that ξ^1 is dynamics-preserving implies that $\eta(\gamma_n z) = \rho(\gamma_n)\eta(z) \rightarrow \xi^1(x)$, unless $\eta^1(z) \in \xi^{d-1}(y)$. But if in fact $\eta^1(z) \notin \xi^{d-1}(y)$, then the limits would agree, i.e. $\eta^1(x) = \xi^1(x)$ which is a contradiction. Thus we have that $\eta^1(z) \in \xi^{d-1}(y)$ and since y was an arbitrary points of $\partial\Gamma \setminus \{y\}$, we have that

$$\eta^1(z) \in \bigcap_{y \in \partial\Gamma \setminus \{z\}} \xi^{d-1}(y) \subseteq \bigcap_{y \in \partial\Gamma \setminus \Gamma \cdot z} \xi^{d-1}(y).$$

In particular, the set appearing on the right hand side is ρ -equivariant, non-empty proper subset of \mathbb{R}^d , which contradicts the strong irreducibility assumption of ρ . \square

Given the lemma above, it suffices to find an $\mathrm{SU}(2, 1)$ -equivariant map from $\partial\Gamma$ to $\mathbb{P}(\mathbb{R}^d)$.

Proposition 4.0.4. *Let Γ be a uniform lattice of $\mathrm{SU}(2, 1)$, and $\eta : \Gamma \rightarrow \mathrm{SL}(\mathfrak{su}(2, 1))$ be the restriction of the adjoint representation: $\eta(\gamma) = \mathrm{Ad}_\gamma$ for all $\gamma \in \Gamma$. The projective part of the limit map of η is given by*

$$\xi_\eta : \partial\Gamma = \mathrm{SU}(2, 1)/P_0 \rightarrow \mathbb{P}(\mathfrak{su}(2, 1)), \quad \xi_\eta(gP_0) = \mathbb{R} \mathrm{Ad}_\gamma x_0.$$

where $P_0 = \mathrm{St}_{\mathrm{SU}(2, 1)}[1 : 0 : 0]$ and

$$x_0 = i \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \in \mathfrak{su}(2, 1).$$

Its derivative satisfies:

$$d_x \xi(T_x \mathrm{SU}(2, 1)/P_0) = \pi(\mathrm{ad}_{\xi^1(x)} \mathfrak{su}(2, 1))$$

where $\pi : \mathrm{hom}(\xi^1(x), \mathfrak{su}(2, 1)) \rightarrow \mathrm{hom}(\xi^1(x), \mathfrak{su}(2, 1)/\xi^1(x))$ is the canonical projection.

Proof. Since by Lemma 4.0.2 the limit map is the unique continuous ρ -equivariant from the boundary of Γ to the projective space, it suffices to show that there exists an η -equivariant map $\xi^1 : \mathrm{SU}(2, 1)/P_0 \rightarrow \mathbb{P}(\mathfrak{su}(2, 1))$, since it will then restrict to the limit map on $\partial\Gamma$.

We consider the parabolic subgroup $P_0 = \mathrm{St}_{\mathrm{SU}(2, 1)}[1 : 0 : 0]$ of $\mathrm{SU}(2, 1)$. Then its Lie algebra is given by:

$$\mathfrak{p}_0 = \mathrm{St}_{\mathfrak{su}(2, 1)}[1 : 0 : 0] = \left\{ \begin{pmatrix} u - is & a & it \\ 0 & 2is & -\bar{a} \\ 0 & 0 & -u - is \end{pmatrix} : a \in \mathbb{C}, u, s, t \in \mathbb{R} \right\}.$$

Since for $\mathbb{R}x \in \mathbb{P}(\mathfrak{su}(2, 1))$ we have that P_0 fixes $\mathbb{R}x$ if and only if \mathfrak{p}_0 fixes $\mathbb{R}x$. But a quick calculation shows that the only element of $\mathfrak{su}(2, 1)$ fixed by \mathfrak{p}_0 is x_0 .

For the calculation of the image of the differential at the identity coset P , we differentiate the commutative diagram:

$$\begin{array}{ccc} \mathrm{SU}(2, 1) & \xrightarrow{\mathrm{Ad} \cdot x_0} & \mathfrak{su}(2, 1) \\ \downarrow & & \downarrow \\ \mathrm{SU}(2, 1)/P_0 & \xrightarrow{\xi^1} & \mathbb{P}(\mathfrak{su}(2, 1)) \end{array} \quad \text{to get} \quad \begin{array}{ccc} \mathfrak{su}(2, 1) & \xrightarrow{\mathrm{ad} \cdot x_0} & \mathfrak{su}(2, 1) \\ \downarrow & & \downarrow \pi \\ \mathfrak{su}(2, 1)/\mathfrak{p}_0 & \xrightarrow{d_P \xi^1} & T_{\xi^1(P)} \mathbb{P}(\mathfrak{su}(2, 1)) \end{array}$$

In the general case we use the equivariance of the limit map

$$\begin{aligned}
d_{gP}\xi^1(T_{gP}\mathrm{SU}(2,1)/P_0) &= d_{gP}\xi^1 d_P g(T_P\mathrm{SU}(2,1)/P_0) = d_{\xi^1(P)} g d_P \xi^1(T_P\mathrm{SU}(2,1)/P_0) = \\
&= d_{\xi^1(P)} g \pi(\mathrm{ad}_{\xi^1(P)} \mathfrak{su}(2,1)) = \\
&= \pi(\mathrm{Ad}_g(\mathrm{ad}_{\xi^1(P)} \mathfrak{su}(2,1))) = \pi(\mathrm{ad}_{\mathrm{Ad}_g \xi^1(P)} \mathfrak{su}(2,1)) = \\
&= \pi(\mathrm{ad}_{\xi^1(gP)} \mathfrak{su}(2,1)).
\end{aligned}$$

□

Before moving on, we recall that all parabolic subgroups of $\mathrm{SU}(2,1)$ are conjugate to each other, so we have the following identification, which will be useful in the proof of Proposition 4.0.2 and the following lemma:

$$\begin{aligned}
\mathrm{SU}(2,1)/P_0 &\leftrightarrow \{ \text{Parabolic subgroups of } \mathrm{SU}(2,1) \} \leftrightarrow \{ \text{Parabolic subalgebras of } \mathfrak{su}(2,1) \} \\
gP_0 &\leftrightarrow gP_0g^{-1} \leftrightarrow \mathrm{Ad}_g(\mathfrak{p}_0)
\end{aligned}$$

Lemma 4.0.3. $\mathrm{Ad}_{\mathrm{SU}(2,1)}$ acts transitively on pairs of distinct parabolic subalgebras of $\mathfrak{su}(2,1)$. In other words, for every distinct parabolic subalgebras $\mathfrak{p}, \mathfrak{p}' \leq \mathfrak{su}(2,1)$, there exists some $g \in \mathrm{SU}(2,1)$ such that $\mathrm{Ad}_g(\mathfrak{p}) = \mathfrak{p}_0$ and $\mathrm{Ad}_g \mathfrak{p}' = \mathfrak{p}_0^t$, where $\mathfrak{p}_0, \mathfrak{p}_0^t$ are the subalgebras of the parabolic subgroups $P_0 = \mathrm{St}_{\mathrm{SU}(2,1)}[1:0:0]$ and $P_0^t = \{g^t : g \in P_0\} = \mathrm{St}_{\mathrm{SU}(2,1)}[0:0:1]$ respectively.

Proof. Recall that we have defined $P_0 = \mathrm{St}_{\mathrm{SU}(2,1)}[1:0:0]$. To see why $P_0^t = \mathrm{St}_{\mathrm{SU}(2,1)}[0:1:0]$, we note that $P_0^t = g_0 P_0 g_0^{-1} = g_0 P_0 g_0^{-1} = \mathrm{St}_{\mathrm{SU}(2,1)} g_0 [1:0:0] = \mathrm{St}_{\mathrm{SU}(2,1)} [0:0:1]$ for

$$g_0 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

Thus $\mathfrak{p}_0 = \mathrm{St}_{\mathfrak{su}(2,1)}[1:0:0]$ and $\mathfrak{p}_0^t = \mathrm{St}_{\mathfrak{su}(2,1)}[0:0:1]$. Let now $l, l' \in \partial_\infty \mathbb{H}_{\mathbb{C}}^2$ be the isotropic lines corresponding to $\mathfrak{p}, \mathfrak{p}'$ respectively, i.e. $\mathfrak{p} = \mathrm{St}_{\mathfrak{su}(2,1)} l$ and $\mathfrak{p}' = \mathrm{St}_{\mathfrak{su}(2,1)} l'$. Denoting with $P = \mathrm{St} l, P' = \mathrm{St} l'$ the respective parabolic subgroups, Witt's theorem guarantees that there exists some $g \in \mathrm{SU}(2,1)$ such that $gl = \mathbb{R}e_1, gl' = \mathbb{R}e^3$. Then we have that $gPg^{-1} = \mathrm{St}_{\mathrm{SU}(2,1)} gl = P_0$ and $gP'g^{-1} = \mathrm{St}_{\mathrm{SU}(2,1)} gl' = P_0^t$. □

We are now ready to fill in the gaps, and provide a proof of Proposition 4.0.2.

Proof of Proposition 4.0.2. (i) Follows from the fact that $\mathrm{SU}(2,1)$ is a simple Lie group.

(ii) Shown in Proposition 4.0.3.

(iii) Follows from the fact that ξ^1 is $\mathrm{SU}(2,1)$ -equivariant and the equivariant identification of $\mathcal{F}_{\{1,4\}}(\mathfrak{su}(2,1)) \simeq \mathcal{P}$.

(iv) Letting $g \in \mathrm{SU}(2,1)$ be as in Lemma 4.0.3, we have that $\mathrm{Ad}_g(\mathfrak{p}_0) = \mathfrak{p}$ and $\mathrm{Ad}_g(\mathfrak{p}_0^t) = \mathfrak{p}'$. Thus $\zeta(x)^4 \cap \zeta(y)^4 \neq \emptyset$ if and only if

$$\begin{aligned}
\emptyset \neq \mathrm{Ad}_g(\zeta(x)^4 \cap \zeta(y)^4) &= \mathrm{Ad}_g \zeta(x)^4 \cap \mathrm{Ad}_g \zeta(y)^4 = \zeta(gx)^4 \cap \zeta(gy)^4 = \zeta(\mathfrak{p}_0)^4 \cap \zeta(\mathfrak{p}_0^t)^4 = \\
&= \pi \left(\mathbb{R} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right).
\end{aligned}$$

For the last equality, we use Proposition 4.0.4 and the fact that $\mathfrak{p}_0^t = \text{Ad}_g \mathfrak{p}_0$ for

$$g = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix},$$

to conclude that

$$\zeta(\mathfrak{p}_0) = \zeta(P_0) = \pi \left(\left\{ \begin{pmatrix} u & a & it \\ 0 & 0 & -\bar{a} \\ 0 & 0 & -u \end{pmatrix} : a \in \mathbb{C}, u, t \in \mathbb{R} \right\} \right),$$

and

$$\zeta(\mathfrak{p}_0^t) = \zeta(gP_0) = \text{Ad}_g \zeta(P_0) = \pi \left(\left\{ \begin{pmatrix} u & 0 & 0 \\ a & 0 & 0 \\ it & -\bar{a} & 0 \end{pmatrix} : a \in \mathbb{C}, u, t \in \mathbb{R} \right\} \right).$$

- (v) This is evident from the preceeding point of the proposition.
- (vi) Since $\Gamma \leq \text{SU}(2, 1)$ is a uniform lattice,, we know that $\partial\Gamma \simeq \partial_\infty H_{\mathbb{C}}^2 \simeq \mathbb{S}^3$, hence $d_\Gamma = 3$. Thus, by the arguments presented in Chapter 2, it suffices to show that $h_\eta(J_3^u) = 1$. Since $\text{SU}(2, 1)$ is of rank-1, one can show that

$$J_3^u(\mu(\eta(\gamma))) = 2d(\gamma[0 : 0 : 1], [0 : 0 : 1]),$$

where

$$\cosh^2 \left(\frac{d(x, y)}{2} \right) = \frac{\langle x, y \rangle \langle y, x \rangle}{\langle x, x \rangle \langle y, y \rangle},$$

and $\langle x, y \rangle = x_1 \bar{y}_1 + x_2 \bar{y}_2 + x_3 \bar{y}_3$ for $x, y \in \mathbb{H}_{\mathbb{C}}^2$, which implies that the critical exponent of J_3^u is equal to half of the critical exponent δ_Γ of Γ , i.e.

$$h_\eta(J_3^u) = \frac{\delta_\Gamma}{2} := \lim_{R \rightarrow \infty} \frac{\sharp\{\gamma : d(\gamma[0 : 0 : 1], [0 : 0 : 1]) \leq R\}}{R}.$$

But the critical exponent of Γ is equal to the volume entropy of the metric d , which can be explicitly computed:

$$\delta_\Gamma = \lim_{R \rightarrow \infty} \frac{\log \text{Vol}(B_R([0 : 0 : 1]))}{R} = 2.$$

□

As can be seen in the Proposition 4.0.2, the above construction provides a counterexample to Lemma 4.0.1 and to Proposition 4.0.2, but a direct calculation shows that it verifies the result of the main theorem. For this reason, it is natural to look for special cases, where Equation (4.1) constitutes indeed a contradiction. In particular, we have found that this is indeed true if we add the assumption that $\eta(\Gamma)$ is Zariski-dense in $\text{SO}(2, p)$ for some p different than 2, which covers a particularly interesting source of examples of discrete subgroups of $\text{SO}(2, p)$, namely quasi-Fuchsian representations which are intimately connected with the construction of globally hyperbolic maximally compact Cauchy manifolds (see [MST23]). This provides a modified version of the main theorem:

Theorem 5. Let $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{R})$ be a projective Anosov representation such that $\xi_\rho^1(\partial\Gamma)$ is a Lipschitz submanifold of $\mathbb{P}(\mathbb{R}^d)$. Assume moreover that $\dim \xi_\rho^1(\partial\Gamma) = d - 3, d > 4$ and $\rho(\Gamma)$ is Zariski-dense in $\mathrm{SO}(2, d - 2)$. Then

$$\dim \xi_\rho^1(\partial\Gamma) = h_{\rho(\Gamma)}(F).$$

The proof of the above theorem will in part be based on reducing it to the analogous result for the case of $\mathrm{SO}(1, d - 1)$, which we will first state and prove.

Lemma 4.0.4. Let $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{R})$ be a projective Anosov representation such that $\xi_\rho^1(\partial\Gamma)$ is a Lipschitz submanifold of $\mathbb{P}(\mathbb{R}^d)$. Assume moreover that $\dim \xi_\rho^1(\partial\Gamma) = d - 2, d \geq 3$, and $\rho(\Gamma)$ is Zariski-dense in $\mathrm{SO}(1, d - 1)$. Then

$$\dim \xi_\rho^1(\partial\Gamma) = h_{\rho(\Gamma)}(F).$$

Proof. The proof is analogous to the proof of Theorem 4, with the only difference that μ -irreducibility does not follow trivially from Zariski-density. Nevertheless, we still follow the strategy of [PSW23] and argue by contradiction. Up to swapping V and W we can assume that $\dim V = 1$ and $\dim W = d - 1$. Then the arguments following the statement of Lemma 4.0.1 show that it suffices to show that for any $V \in \mathcal{G}_1(\mathbb{R}^d), W \in \mathcal{G}_{d-1}(\mathbb{R}^d)$ there exists some $g \in \mathrm{SO}(1, d - 1)$ such that

$$gV \cap W = 0.$$

Using Witt's theorem, we see that this is equivalent to showing that for any $V \in \mathcal{G}_1(\mathbb{R}^d), W \in \mathcal{G}_{d-1}(\mathbb{R}^d)$ there exists some $V' \in \mathcal{G}_1(\mathbb{R}^d)$ such that

$$\mathrm{sgn} \omega|_V = \mathrm{sgn} \omega|_{V'} \quad \text{and} \quad V' \cap W = 0,$$

where ω is the standard form on $\mathbb{R}^{1, d-1}$. In other words, we need to show that the complement of every hyperplane $W \subseteq \mathbb{R}^d$ contains vectors of all signatures. In fact, we will show that it contains two zero-vectors, infinitely many positive vectors and infinitely many negative vectors.

Let $w \in W$ be a non-zero vector. Such a vector exists because otherwise W would be an isotropic subspace of dimension $d - 1 \geq 2$ which is not possible for a form of signature $(1, d - 1)$. Let $u \in W^\perp$ be a vector of the opposite sign to w , which exists since otherwise ω would be either positive or negative on \mathbb{R}^d . Then the affine line $L = \{u_t := u + tw : t \in \mathbb{R}\}$ does not intersect W and contains vectors of all signatures. Indeed, assume without loss of generality that $\omega(u, u) > 0 > \omega(w, w)$. Then

$$\omega(u_t, u_t) = \omega(w, w)t^2 + 2\omega(u, w)t + \omega(u, u).$$

The discriminant of the above quadratic is $4\omega(u, w)^2 - 4\omega(u, u)\omega(w, w)$, which is positive since u and w have opposite signs. Let $t_1 < t_2$ be the two roots of the above quadratic. Then

$$\omega(u_t, u_t) \begin{cases} > 0 & \text{if } t \in (t_1, t_2) \neq \emptyset \\ = 0 & \text{if } t = t_1 \text{ or } t = t_2 \\ < 0 & \text{if } t \notin [t_1, t_2] \end{cases} \quad (4.2)$$

□

To reduce the proof of Theorem 5 to the proof of Lemma 4.0.4, we will use the following lemma:

Is it not the case $d = 4$ that is of importance in physics?

Lemma 4.0.5. *Let ω be a non-degenerate form on \mathbb{R}^d and v a zero-vector for ω . Then ω induces the form $\tilde{\omega}$ on $\perp v/\mathbb{R}v$ given by*

$$\tilde{\omega}(x + \mathbb{R}v, y + \mathbb{R}v) = \omega(x, y),$$

and which has signature $\text{sgn}(\tilde{\omega}) = \text{sgn}(\omega) - (1, 1)$.

We are now ready to provide a proof for Theorem 5:

Proof of Theorem 5. The same arguments as in the proof of Lemma 4.0.4 show that it suffices to show that for any $V \in \mathcal{G}_2(\mathbb{R}^d)$, $W \in \mathcal{G}_{d-2}(\mathbb{R}^d)$ there exists some $V' \in \mathcal{G}_2(\mathbb{R}^d)$ that is transverse to W and has the same signature as V .

For the case where V is non-degenerate, we can perturb V to a non-degenerate hyperplane V' that is transverse to W without changing its signature.

The case where V is degenerate is more subtle. Assuming that $\dim(V \cap V^\perp) \geq 1$, we first show that there exists some zero vector $v \in W^c$ such that $W \not\subseteq v^\perp$. Indeed, the same arguments as in the Lemma 4.0.4 show that there exist $w \in W, u \in W^c$ such that $\omega(w, w) > 0 > \omega(u, u)$ and that there exist $t_1 < t_2$ such that Equation (4.2) holds for the affine line $L = \{u_t := u + tw : t \in \mathbb{R}\} \subseteq W^c$. In particular u_{t_1}, u_{t_2} are not contained in W . Moreover, at least one of u_{t_1}, u_{t_2} is not orthogonal to W , since otherwise $w = (u_{t_2} - u_{t_1})^{-1} \in \langle u_{t_1}, u_{t_2} \rangle$ would be orthogonal to W , which is absurd because w was assumed to be non-zero.

Let $p : v^\perp \rightarrow v^\perp/\mathbb{R}v$ be the canonical projection. We claim that for $W' = p(W \cap v^\perp)$,

W' is a hyperplane of $v^\perp/\mathbb{R}v$ and ω induces a form of signature $(1, d-3)$ on $v^\perp/\mathbb{R}v$.

The signature of the induced form follows from Lemma 4.0.5. To calculate the dimension of W' , note that $\dim p(W \cap v^\perp) = \dim(W \cap v^\perp)$ since $\ker p \cap W = \mathbb{R}v \cap W = 0$, by the fact that v is not contained in W . On the other hand $\dim(W \cap v^\perp) = \dim(\ker W \ni x \mapsto (v, x)) = \dim W - 1 = d - 3$, while $\dim(v^\perp/\mathbb{R}v) = \dim v^\perp - 1 = (d-1) - 1 = d-2$.

For any $x \in v^\perp$ such that $p(x)$ is transverse to W' , we have that $V' = \langle v, x \rangle$ is a plane that is transverse to W . To show transversality, we let $r, s \in \mathbb{R}$ such that $rv + sx \in W$. Then $p(rv + sx) = sp(x) \in W'$, which implies that $s = 0$ by the choice of x . But then $rv \in W$ which means that $r = 0$ by the choice of v . In fact, the same arguments show that x and v are linearly independent and hence V is a plane.

Finally, we show that by picking the appropriate x , we can ensure that V' has the same signature as V . Indeed, $\dim V = 2$, so $\text{sgn}(\omega|_V) \in \{(1, 1, 0), (0, 2, 0), (0, 1, 1)\}$. Letting x be a positive, negative or zero vector respectively, we have that

$$\omega|_{V'} \in \left\{ \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, 0, \begin{bmatrix} 0 & 0 \\ 0 & -1 \end{bmatrix} \right\} \text{ in the basis } \{v, x\}$$

respectively. □

Try to generalize the above in other choices of $\text{SO}(p, q)$

Is $\text{Ad}_{\text{SU}(2,1)}$ Zariski-dense in $\text{SO}(\mathfrak{su}(2, 1), K)$? If yes, then a generalisation of Theorem 5 to $\text{SO}(4, 4)$ would account for the validity of the result in the case of Proposition 4.0.2

Type up the proof of the quadratic form reduction that I have already handwritten.

inline

Chapter 5

Conclusion

Anosov representations have been a subject of interest ever since their introduction in [Lab06], and are now considered as a natural generalisation of convex-cocompact representations into Lie groups of rank higher than 1. Their limit set, aside from being a natural object to study in its own right, has been shown to have applications in such diverse fields as anti-de Sitter geometry, being important in the characterisation of globally hyperbolic maximally compact Cauchy manifolds.

In this thesis, based upon the work of [PSW23], we have explored different situations where that the Falconer functional of an Anosov subgroup is equal to the Hausdorff dimension of its limit set, provided that the latter is a Lipschitz submanifold of the projective space. In Chapter 2 we have done this under the assumption that the Anosov subgroup is Zariski-dense in $SL(d, \mathbb{R})$ and in Chapter 4 we have shown that this is not necessary, provided that the Anosov subgroup is Zariski-dense in $SO(2, p)$ for some $p \neq 2$.

However, we have yet no reason to believe that this last result is optimal, and it would be interesting to investigate whether it can be generalised for instance to the case where the group is Anosov and dense in $SO(p, q)$, or even to the non-Anosov case.

Appendix A

Tangent space to the Grassmanian

Let V be a d -dimensional real vector space. We denote with $\mathcal{G}_k(V)$ the Grassmanian of k -dimensional subspaces of V . Our first objective is to find a convenient way to express its tangent space.

Proposition A.0.1. *We have the following canonical identification:*

$$\begin{aligned} \text{hom}(W, V/W) &\simeq T_W \mathcal{G}_k(V) \\ \phi &\mapsto \left. \frac{d}{dt} \right|_{t=0} \Gamma(t\phi) \end{aligned}$$

where $\Gamma(\phi) = (Id + \phi)(W)$ is the graph of ϕ .

Proof. We will consider the map

$$F : \text{Injhom}(W, V) \rightarrow \mathcal{G}_k(V), \quad \phi \mapsto \phi(W).$$

whose derivative is given by:

$$d_I F(\phi) = F \left(\left. \frac{d}{dt} \right|_{t=0} (I + t\phi) \right) = \left. \frac{d}{dt} \right|_{t=0} (I + t\phi(W)) = \left. \frac{d}{dt} \right|_{t=0} \Gamma(t\phi).$$

The result will follow as soon as we have shown that $d_I F$ is surjective and that $\ker d_I F = \text{hom}(W, W)$.

To show that it is surjective, we consider a $(d-k)$ -dimensional subspace $W' \in \mathcal{G}_{d-k}(V)$ that is complementary to W , i.e. $V = W \oplus W'$. Denoting with $U_{W'} = \{Z \in \mathcal{G}_k(V) : Z \cap W' = 0\}$, we recall the corresponding chart:

$$\begin{aligned} \Psi : \text{hom}(W, W') &\rightarrow U_{W'} \\ \phi &\mapsto \Gamma(\phi). \end{aligned}$$

Surjectivity of $d_I F$ now follows by the fact that

$$d_I F(\phi) = \left. \frac{d}{dt} \right|_{t=0} \Gamma(t\phi) = d_0 \Psi(\phi).$$

To show that $\ker d_I F = \text{hom}(W, W)$, we first note that clearly $\ker d_I F \supseteq \text{hom}(W, W)$. Equality then follows by the fact that $\dim \text{hom}(W, W) = \dim \ker d_I F$, which is a direct consequence of the surjectivity. \square

Note that another way to prove the above identification throught the fact that the Grassmanian is a homogeneous space of $\mathrm{GL}(d, \mathbb{R})$, giving us the diffeomorphism

$$\begin{aligned} \mathrm{GL}(V)/\mathrm{St}_{\mathrm{GL}(V)}W &\rightarrow \mathcal{G}_k(V) \\ [g] &\mapsto gW, \end{aligned}$$

where $\mathrm{St}_{\mathrm{GL}(V)}W = \{g \in \mathrm{GL}(V) : gW = W\}$ is the stabilizer of W . Thus an expression for the tangent space at W may be obtained by differentiating the map above at the identity coset:

$$\mathrm{hom}(W, V/W) \simeq \mathrm{hom}(V, V)/\mathrm{hom}(W, W) \simeq T_W \mathcal{G}_k(V).$$

In particular the above gives us a direct proof that the kernel of the map constructed in the previous proof is indeed $\mathrm{hom}(W, W)$.

Our second objective is to identify subspaces of $T_l \mathbb{P}(V)$ with subspaces of V , by considering the first as projectivisation of the second. More concretely, we shall consider the space

$$\mathcal{P} = \{(l, P) : l \in \mathbb{P}(V), P \in \mathcal{G}_k(T_l \mathbb{P}(V))\}$$

as a homogenous space of $\mathrm{SL}(V)$, where the action is given by

$$g \cdot (l, P) = (gl, d_l g(P) = g\pi^{-1}(P)g^{-1} + \mathrm{hom}(gl, gl)).$$

where we use the identification of $T_l \mathbb{P}(V)$ with $\mathrm{hom}(l, V/l)$ as above and denote with $\pi : \mathrm{hom}(l, V) \rightarrow \mathrm{hom}(l, V/l)$ the canonical projection. For the sake of completeness, we outline the calculation of the differential:

$$\begin{aligned} \mathrm{hom}(l, V/l) &\rightarrow T_l \mathbb{P}(V) && \rightarrow T_{gl} \mathbb{P}(V) && \rightarrow \mathrm{hom}(gl, V/gl) \\ \phi &\mapsto \left. \frac{d}{dt} \right|_{t=0} g(I + t\tilde{\phi})(l) && \mapsto \left. \frac{d}{dt} \right|_{t=0} (I + tg\tilde{\phi}g^{-1})(gl) && \mapsto g\tilde{\phi}g^{-1} + \mathrm{hom}(gl, gl) \end{aligned}$$

where $\phi \in \mathrm{hom}(l, V/l)$, $\tilde{\phi} \in \mathrm{hom}(l, V)$ such that $\tilde{\phi} + \mathrm{hom}(l, l) = \phi$.

Another subtle point is that conjugation by g helps us pass from $\mathcal{G}_k(T_x \mathbb{P}(V))$ to $\mathcal{G}_k(T_{gx} \mathbb{P}(V))$:

Lemma A.0.1. *Let $g \in \mathrm{SL}(V)$. Then for every $x \in \mathbb{P}(V)$, $P \in \mathcal{G}_k(T_x \mathbb{P}(V))$, we have:*

$$\pi^{-1}(gPg^{-1}) = g\pi^{-1}(P)g^{-1},$$

where π denotes the projection $\mathrm{hom}(gx, \mathbb{R}^d) \rightarrow T_{gx} \mathbb{P}(\mathbb{R}^d)$ on the left hand side of the equality, and the projection $\mathrm{hom}(x, \mathbb{R}^d) \rightarrow T_x \mathbb{P}(\mathbb{R}^d)$ on the right hand side.

Proof. We denote with $C_g : \mathrm{hom}(x, \mathbb{V}) \rightarrow \mathrm{hom}(gx, \mathbb{V})$ the isomorphism by conjugation $C_g(T) = gTg^{-1}$. Since the following diagram commutes

$$\begin{array}{ccc} \mathrm{hom}(x, V) & \xrightarrow{C_g} & \mathrm{hom}(gx, V) \\ \downarrow \pi & & \downarrow \pi \\ \mathrm{hom}(x, V/x) & \xrightarrow{\tilde{C}_g} & \mathrm{hom}(gx, V/gx) \end{array}$$

C_g induces an isomorphism \tilde{C}_g between $T_x(\mathbb{P}(V))$ and $T_{gx}(\mathbb{P}(V))$. We now have

$$\pi^{-1}(gPg^{-1}) = \pi^{-1}(\tilde{C}_g(P)) = (\tilde{C}_g^{-1}\pi)^{-1}(P) = (\pi C_g^{-1})^{-1}(P) = C_g(\pi^{-1}(P)),$$

which is a rephrasing of the desired equality. \square

We are now ready to express the needed identification:

Proposition A.0.2. *We have the following $\mathrm{SL}(V)$ equivariant identification:*

$$\begin{aligned}\mathcal{P} &\rightarrow \mathcal{F}_{1,k+1}(V) \\ (l, P) &\mapsto (l, \pi^{-1}(P)l) \\ (l, \mathrm{hom}(l, Q/l)) &\hookleftarrow (l, Q).\end{aligned}$$

where $\pi : \mathrm{hom}(l, V) \rightarrow \mathrm{hom}(l, V/l)$ is the canonical projection.

Proof. We begin by showing that the left-to-right direction of the map is well-defined. For this, we first need to check that for $(l, P) \in \mathcal{P}$, we have that $\dim \pi^{-1}(P)l = k + 1$. Indeed, we have that $\dim \pi^{-1}(P) = k + 1$ as implied by the rank-nullity theorem for $\pi : \pi^{-1}(P) \rightarrow P$. The result then follows by the fact that $\pi^{-1}(P)l = T_1(l) \oplus \cdots \oplus T_{k+1}(l)$ for any base T_1, \dots, T_{k+1} of $\pi^{-1}(P)$. The second thing to check is that $l \leq \pi^{-1}(P)l$, which holds since $\ker \pi = \mathrm{hom}(l, l) \leq \pi^{-1}(P)$.

To see that the two directions above are inverse to each other, we begin by examining the right-to-left-to-right composition:

$$(l, Q) \mapsto (l, \pi(\mathrm{hom}(l, Q))) \mapsto (l, \pi^{-1}\pi(\mathrm{hom}(l, Q))l) = (l, \mathrm{hom}(l, Q)l) = (l, Q),$$

where we did the calculation $\pi^{-1}\pi(\mathrm{hom}(l, Q)l) = \mathrm{hom}(l, Q) + \ker \pi = \mathrm{hom}(l, Q)$ in the second to last inequality. For the left-to-right-to-left composition we have

$$(l, P) \mapsto (l, \pi^{-1}(P)l) \mapsto (l, \mathrm{hom}(l, \pi^{-1}(P)l/l)),$$

so it suffices to show that $\mathrm{hom}(l, \pi^{-1}(P)l/l) = P$. Indeed, for $\pi^{-1}(P) = \mathbb{R}T_1 \oplus \cdots \oplus \mathbb{R}T_k$, we have that

$$\begin{aligned}\mathrm{hom}(l, \pi^{-1}(P)l/l) &= \mathrm{hom}(l, \pi^{-1}(P)l) / \mathrm{hom}(l, l) = (\oplus_i \mathrm{hom}(l, T_i(l)) / \mathrm{hom}(l, l)) = \\ &= (\oplus_i \mathbb{R}T_i) / \mathrm{hom}(l, l) = \pi^{-1}(P) / \mathrm{hom}(l, l) = P.\end{aligned}$$

For the equivariance, the calculations has as follows:

$$\begin{array}{ccc} (l, P) & \xrightarrow{\quad\quad\quad} & (l, \pi^{-1}(P)l) \\ \downarrow g & & \downarrow g \\ (gl, g\pi^{-1}(P)g^{-1} + \mathrm{hom}(gl, gl)) & \xrightarrow{\quad\quad\quad} & (gl, (g\pi^{-1}(P)g^{-1})(gl)) = (gl, g\pi^{-1}(P)l) \end{array} \quad \square$$

Index

- Anosov
 - boundary maps, 8
- Busemann cocycle, 20
- Cartan
 - decomposition, 6
 - projection, 6
- critical exponent
 - of functional, 10
 - subgroup, 4
- ellipsoid, 14
- Falconer
 - dimension, 10
 - functional, 9
- Hausdorff
 - dimension, 11
- irreducible representation
 - strongly, 29
 - with respect to measure, 26
- limit set, 4
- Patterson–Sullivan measure, 21
- singular value, 6
- uniform lattice, 30
- unstable Jacobian, 9

Bibliography

- [LL23] François Ledrappier and Pablo Lessa. “Dimension gap and variational principle for Anosov representations”. In: (Dec. 2023). arXiv:2310.13465 [math] version: 3 (cit. on p. 10).
- [MST23] Daniel Monclair, Jean-Marc Schlenker, and Nicolas Tholozan. “Gromov-Thurston manifolds and anti-de Sitter geometry”. In: *arXiv preprint arXiv:2310.12003* (2023) (cit. on p. 34).
- [PSW23] Beatrice Pozzetti, Andrés Sambarino, and Anna Wienhard. “Anosov representations with Lipschitz limit set”. In: *Geometry & Topology* 27.8 (Nov. 2023). arXiv:1910.06627 [math], pp. 3303–3360. ISSN: 1364-0380, 1465-3060 (cit. on pp. 1–3, 5, 13, 14, 16, 17, 23, 29, 30, 35, 37).
- [BPS19] Jairo Bochi, Rafael Potrie, and Andrés Sambarino. “Anosov representations and dominated splittings”. In: *Journal of the European Mathematical Society* 21.11 (2019), pp. 3343–3414 (cit. on p. 8).
- [DGK18] Jeffrey Danciger, François Guéritaud, and Fanny Kassel. “Convex cocompactness in pseudo-Riemannian hyperbolic spaces”. In: *Geometriae Dedicata* 192 (2018), pp. 87–126 (cit. on p. 2).
- [KLP17] M Kapovich, B Leeb, and J Porti. “Anosov subgroups: dynamical and geometric characterizations”. In: *European Journal of Mathematics* 3.4 (2017), pp. 808–898 (cit. on pp. 2, 8).
- [GH13] Etienne Ghys and Pierre de la Harpe. *Sur les groupes hyperboliques d’après Mikhael Gromov*. Vol. 83. Springer Science & Business Media, 2013 (cit. on pp. 4, 17).
- [MB12] Quentin Mérigot and Thierry Barbot. “Anosov AdS representations are quasi-Fuchsian”. In: *Groups, Geometry, and Dynamics* 6.3 (2012), pp. 441–483 (cit. on p. 2).
- [BIW10] Marc Burger, Alessandra Iozzi, and Anna Wienhard. “Surface group representations with maximal Toledo invariant”. In: *Annals of mathematics* (2010), pp. 517–566 (cit. on p. 2).
- [Lab06] François Labourie. “Anosov flows, surface groups and curves in projective space”. en. In: *Inventiones mathematicae* 165.1 (July 2006), pp. 51–114. ISSN: 1432-1297 (cit. on pp. 3, 8, 29, 37).
- [Qui06] Jean-François Quint. “An overview of Patterson-Sullivan theory”. In: *Workshop The barycenter method, FIM, Zurich*. 2006 (cit. on p. 4).
- [Rob03] Thomas Roblin. “Ergodicité et équidistribution en courbure négative”. In: (2003) (cit. on p. 2).
- [Qui02] J-F Quint. “Mesures de Patterson—Sullivan en rang supérieur”. In: *Geometric & Functional Analysis GAFA* 12.4 (2002), pp. 776–809 (cit. on pp. 3, 20).

- [DOP00] Françoise Dal’Bo, Jean-Pierre Otal, and Marc Peigné. “Séries de Poincaré des groupes géométriquement finis”. In: *Israel Journal of Mathematics* 118 (2000), pp. 109–124 (cit. on p. 2).
- [Coo93] Michel Coornaert. “Mesures de Patterson-Sullivan sur le bord d’un espace hyperbolique au sens de Gromov”. In: *Pacific Journal of Mathematics* 159.2 (1993), pp. 241–270 (cit. on p. 2).
- [Sul79] Dennis Sullivan. “The density at infinity of a discrete group of hyperbolic motions”. In: *Publications Mathématiques de l’IHÉS* 50 (1979), pp. 171–202 (cit. on p. 2).
- [Pat76] SJ Patterson. “The limit set of a Fuchsian group”. In: *Acta Mathematica* 136 (1976), pp. 241–273 (cit. on p. 2).