



A. Q1: the OXPHOS-to-configuration coupling, within each tissue

Partial Spearman, 10,000 PATIENT-level bootstrap resamples. All three tumour comparators, always. Normals carry no quadrant and none is used here.




B. tumour minus normal, with the declared positive control

The control is PRE-SPECIFIED BUT NOT BLIND. Fe-S mitochondrial is NOT a control - it is mitochondrial, so coupling to OXPHOS is near-guaranteed - and carries no interval, being a companion.



C. Q2: tumour MYC minus that patient's OWN normal, 113 pairs. DESCRIPTIVE - no verdict

Point is the median paired difference, bar the IQR. The label is the % of pairs in which the tumour exceeds its own normal, with a Wilson interval. Immune to the collection-site skew by construction, and unaffected by the Q1 stop condition.



E36. EXPLORATORY, nothing pre-registered. TCGA-BRCA only; SCAN-B carries no normals, so this is UNREPLICATED. Joint snapshot built by E37 over 1208 samples (1095 tumour, 113 normal); no value here is comparable with one from data/from_validation/. The normals are FIELD-ADJACENT tissue from breasts that grew a tumour, and all 113 come from patients whose tumour is in the same analysis - the design is PAIRED. They come from 6 of 40 collection sites, which is why every across-person comparison is shown on three nested tumour sets and why the within-patient contrast in panel C is the one immune to it. Cross-sectional, bulk, transcript-level: N3, nothing here is 'primed', and the configuration is not lethality. No product term; no outcome variable. VERDICT: UNINTERPRETABLE.