


Is the OXPPOS-to-configuration coupling derived by genomic burden?

Reading on the rule fixed at 74507b1 before any estimate: SATURATED - the declared expected outcome TCGA only; SCAN-B has no CNV, so no replication cohort exists. EXPLORATORY; not pre-registered.



Standardised means from lm within each burden stratum: configuration ~ quadrant + PAM50 + purity + leukocyte fraction (m1), and the same without PAM50 within combined Luminal (m2L). The configuration is E33's signed mean-z on log2(linear + 1). STRATIFIED, NOTADJUSTED: there is no product term and no continuous interaction anywhere, and no line joins the strata because they are not exchangeable groups - burden tracks proliferation (rho +0.375 to +0.568) and subtype, so the lowest tertile is three quarters LumA. THE ENDPOINT IS THE COUPLING, NOT KILLING: there is no death readout in TCGA. Burden is a PROXY for drive, not a measure of it, and the configuration is not lethality. TCGA only; SCAN-B carries no CNV, so no replication cohort exists. Genome doublings is shown on its three native levels, not tertiles, and its level 2 is omitted as not readable (minimum quadrant cell 4 against a declared floor of 20). Only strata clearing that floor are drawn. NOTHING HERE LICENSES A TREATMENT-SELECTION CLAIM.