
BCL-2 family, OXPHOS level
ρ, OXPHOS-proliferation
Proliferation – OXPHOS (95% CI)
BBC3 . TCGA
0.24
-0.73 (-0.84, -0.62)
BBC3 . SCAN-B
0.40
-0.39 (-0.46, -0.32)
BCL2L1 . TCGA
0.24
-0.55 (-0.65, -0.44)
BCL2L1 . SCAN-B
0.40
-0.54 (-0.60, -0.47)
MCL1 . TCGA
0.24
0.10 (-0.02, 0.22)
MCL1 . SCAN-B
0.40
0.38 (0.31, 0.44)
lower expression
higher expression
-0.6
-0.4
-0.2
0.0
0.2
0.4
0.6
Association with expression (SD per SD of each score)
Chemotherapy response
ρ, OXPHOS-proliferation
Proliferation / OXPHOS ratio of ORs (95% CI)
GSE194040
0.38
1.72 (1.28, 2.31)
GSE164458
0.49
2.07 (1.42, 3.04)
GSE25066
0.36
2.40 (1.44, 4.00)
Pooled
1.93 (1.56, 2.39)
lower pCR
higher pCR
0.33
0.50
0.75
1.00
1.50
2.00
3.00
Odds ratio for pCR per within-cohort SD
Breast-cancer-specific survival
ρ, OXPHOS-proliferation
Proliferation / OXPHOS ratio of HRs (95% CI)
METABRIC, ER+ (IHC)
0.38
1.37 (1.13, 1.67)
longer survival
shorter survival
0.67
0.80
1.00
1.25
1.50
Hazard ratio for breast-cancer death per within-cohort SD
OXPHOS, without the proliferation term
OXPHOS, with the proliferation term
Proliferation