



Symbolic Methods in Bifurcation Analysis

Thomas Sturm

University of Lorraine, CNRS, Inria, and LORIA, Nancy, France
MPI Informatics and Saarland University, Saarbrücken, Germany

3rd BLOSS Days, Montpellier, France, March 13, 2017

Overview

2008–2012 Hopf Bifurcations in a Gene Regulatory Network

Real Quantifier Elimination by Virtual Substitution

2012–2016 Hopf Bifurcations with Larger Models, Including MBO and MAPK

Convex Coordinates

Subtropical Real Root Finding


2016– Parametric Saddle-node Bifurcations with Another Model of MAPK

Cylindrical Algebraic Decomposition






Graph-theoretical Preprocessing

2008–2012

Hopf Bifurcations in a Gene Regulatory Network



The Reaction System



Dynamics of the Reaction System

$$\dot{G} = \vartheta \cdot (\gamma_0 - G) - \alpha GP_n$$

$$\dot{M} = \varrho_f G + \varrho_b \cdot (\gamma_0 - G) - \delta_M M$$

$$\dot{P} = \beta M - \delta_P P + 2A_1 + A_2 + \dots + A_{n-1}$$

$$\dot{P}_i = -A_{i-1} + A_i \quad (2 \leq i \leq n-1)$$

$$\dot{P}_n = -A_{n-1} + \vartheta \cdot (\gamma_0 - G) - \alpha GP_n$$

where

$$A_i = \frac{1}{\varepsilon} (\kappa_{i+1}^- P_{i+1} - \kappa_{i+1}^+ P_i P)$$

Simplified Dynamics

Approximating

$$\dot{P} = \beta M - \delta_P P + n(\vartheta(\gamma_0 - G) - \alpha G P_n), \quad P_n = \bar{\alpha} P^n \quad \text{with} \quad \bar{\alpha} = \frac{\kappa_1^+ \cdots \kappa_{n-1}^+}{\kappa_1^- \cdots \kappa_{n-1}^-}$$

yields

$$\begin{aligned}\dot{G} &= \vartheta \cdot (\gamma_0 - G) - \alpha \bar{\alpha} G P^n \\ \dot{M} &= \varrho_f G + \varrho_b \cdot (\gamma_0 - G) - \delta_M M \\ \dot{P} &= n\vartheta(\gamma_0 - G) - n\alpha \bar{\alpha} G P^n + \beta M - \delta_P P.\end{aligned}$$

Many more simplifications yield

$$\begin{aligned}\dot{G} &= \vartheta \cdot (\gamma_0 - G - G P^n) \\ \dot{M} &= \lambda G + \gamma_0 \mu - M \\ \dot{P} &= n\alpha \cdot (\gamma_0 - G - G P^n) + \delta \cdot (M - P).\end{aligned}$$

Translation into First-Order Logic over the Reals

yields φ_n for fixed $n \in \mathbb{N}$.

$$\begin{aligned}\varphi_9 \doteq & \exists v_1 \exists v_2 \exists v_3 (v_1 > 0 \wedge v_2 > 0 \wedge v_3 > 0 \wedge \vartheta > 0 \wedge \gamma_0 > 0 \wedge \mu > 0 \wedge \delta > 0 \wedge \alpha > 0 \wedge \\ & \vartheta(\gamma_0 - v_1 - v_1 v_3^9) = 0 \wedge \lambda v_1 + \gamma_0 \mu - v_2 = 0 \wedge \\ & 9\alpha(\gamma_0 - v_1 - v_1 v_3^9) + \delta(v_2 - v_3) = 0 \wedge \\ & \Delta_2 = 0 \wedge \Delta_1 > 0),\end{aligned}$$

where

$$\begin{aligned}\Delta_2 \doteq & 162\vartheta v_3^{17}\alpha v_1 + 162\vartheta\alpha v_1 v_3^8 + 162\alpha v_1 v_3^8\delta + \vartheta + 2\vartheta v_3^9\delta + \vartheta^2 v_3^{18}\delta + \vartheta v_3^9\vartheta\delta \\ & + 81\alpha v_1 v_3^8\vartheta\delta + 81\alpha v_1 v_3^{17}\vartheta\delta + \delta^2 + \vartheta\delta^2 + \vartheta^2\delta + \vartheta^2 + 2\vartheta^2 v_3^9 + \vartheta^2 v_3^{18} \\ & + 6561\alpha^2 v_1^2 v_3^{16} + 2\vartheta^2 v_3^9\delta + \delta + 81\alpha v_1 v_3^8 + \vartheta v_3^9\delta^2 - 9\lambda\vartheta v_1 v_3^8\delta, \\ \Delta_1 \doteq & \vartheta\delta + \vartheta v_3^9\delta + 9\lambda\vartheta v_1 v_3^8\delta.\end{aligned}$$

Hopf bifurcation for some $n \in \mathbb{N} \iff \varphi_n$ holds

Real Quantifier Elimination by Virtual Substitution

Principal Strategy

prenex formula, $\forall x_1 \varphi \longleftrightarrow \neg \exists x_1 \neg \varphi$,

$$\exists x_n \dots \exists x_2 \exists x_1 \varphi \longleftrightarrow \exists x_n \dots \exists x_2 \text{ simplify } \left(\bigvee_{(\gamma, t) \in E} \gamma \wedge \varphi[x_1 // t] \right).$$

$E = \{ \dots, (\gamma, t), \dots \}$ is a finite **elimination set**

Real Quantifier Elimination by Virtual Substitution

Principal Strategy

prenex formula, $\forall x_1 \varphi \longleftrightarrow \neg \exists x_1 \neg \varphi$,

$$\exists x_n \dots \exists x_2 \exists x_1 \varphi \longleftrightarrow \exists x_n \dots \exists x_2 \text{ simplify } \left(\bigvee_{(\gamma, t) \in E} \gamma \wedge \varphi[x_1 // t] \right).$$

$E = \{\dots, (\gamma, t), \dots\}$ is a finite **elimination set**, e.g.,

$$ax^2 - 3x + 7 \leq 0 \rightsquigarrow \left(a \neq 0 \wedge (-3)^2 - 4 \cdot a \cdot 7 \geq 0, \frac{3 + \sqrt{(-3)^2 - 4 \cdot a \cdot 7}}{2 \cdot a} \right) \in E.$$

Real Quantifier Elimination by Virtual Substitution

Principal Strategy

prenex formula, $\forall x_1 \varphi \longleftrightarrow \neg \exists x_1 \neg \varphi$,

$$\exists x_n \dots \exists x_2 \exists x_1 \varphi \longleftrightarrow \exists x_n \dots \exists x_2 \text{ simplify } \left(\bigvee_{(\gamma, t) \in E} \gamma \wedge \varphi[x_1 // t] \right).$$

$E = \{ \dots, (\gamma, t), \dots \}$ is a finite **elimination set**, e.g.,

$$ax^2 - 3x + 7 \leq 0 \rightsquigarrow \left(a \neq 0 \wedge (-3)^2 - 4 \cdot a \cdot 7 \geq 0, \frac{3 + \sqrt{(-3)^2 - 4 \cdot a \cdot 7}}{2 \cdot a} \right) \in E.$$

More Generally

substitute $\pm\infty$, **nonstandard** $t \pm \varepsilon$ with $<$, **abstract roots** for higher degrees

Real Quantifier Elimination by Virtual Substitution

Principal Strategy

prenex formula, $\forall x_1 \varphi \longleftrightarrow \neg \exists x_1 \neg \varphi$,

$$\exists x_n \dots \exists x_2 \exists x_1 \varphi \longleftrightarrow \exists x_n \dots \exists x_2 \text{ simplify } \left(\bigvee_{(y,t) \in E} y \wedge \varphi[x_1 // t] \right).$$

$E = \{\dots, (y, t), \dots\}$ is a finite **elimination set**, e.g.,

$$ax^2 - 3x + 7 \leq 0 \rightsquigarrow \left(a \neq 0 \wedge (-3)^2 - 4 \cdot a \cdot 7 \geq 0, \frac{3 + \sqrt{(-3)^2 - 4 \cdot a \cdot 7}}{2 \cdot a} \right) \in E.$$

More Generally

substitute $\pm\infty$, **nonstandard** $t \pm \varepsilon$ with $<$, **abstract roots** for higher degrees

$[x // t]$: atomic formulas \rightarrow quantifier-free formulas

Some Concrete Virtual Substitutions

Conventions: $f \in \mathbb{Z}[\mathbf{y}][x]$, f_i , g_i , $g_i^* \in \mathbb{Z}[\mathbf{y}]$

Quotients

$$(f_1 x + f_0 \leq 0) [x // \frac{g_1}{g_2}] \equiv f_1 \frac{g_1}{g_2} + f_0 \leq 0 \equiv f_1 g_1 g_2 + f_0 g_2^2 \leq 0$$

Some Concrete Virtual Substitutions

Conventions: $f \in \mathbb{Z}[\mathbf{y}][x]$, f_i , g_i , $g_i^* \in \mathbb{Z}[\mathbf{y}]$

Quotients

$$(f_1 x + f_0 \leq 0) [x // \frac{g_1}{g_2}] \equiv f_1 \frac{g_1}{g_2} + f_0 \leq 0 \equiv f_1 g_1 g_2 + f_0 g_2^2 \leq 0$$

Infinity

$$(f_2 x^2 + f_1 x + f_0 < 0) [x // \infty] \equiv f_2 < 0 \vee (f_2 = 0 \wedge f_1 < 0) \vee (f_2 = 0 \wedge f_1 = 0 \wedge f_0 < 0)$$

Some Concrete Virtual Substitutions

Conventions: $f \in \mathbb{Z}[\mathbf{y}][x]$, $f_i, g_i, g_i^* \in \mathbb{Z}[\mathbf{y}]$

Quotients

$$(f_1 x + f_0 \leq 0)[x // \frac{g_1}{g_2}] \equiv f_1 \frac{g_1}{g_2} + f_0 \leq 0 \equiv f_1 g_1 g_2 + f_0 g_2^2 \leq 0$$

Infinity

$$(f_2 x^2 + f_1 x + f_0 < 0)[x // \infty] \equiv f_2 < 0 \vee (f_2 = 0 \wedge f_1 < 0) \vee (f_2 = 0 \wedge f_1 = 0 \wedge f_0 < 0)$$

Positive infinitesimals

$$(3x^2 + 6x - 3 > 0)[x // t - \varepsilon] \equiv 3t^2 + 6t - 3 > 0 \vee (3t^2 + 6t - 3 = 0 \wedge 6t + 6 \leq 0)$$

Some Concrete Virtual Substitutions

Conventions: $f \in \mathbb{Z}[\mathbf{y}][x]$, f_i , g_i , $g_i^* \in \mathbb{Z}[\mathbf{y}]$

Quotients

$$(f_1 x + f_0 \leq 0) [x // \frac{g_1}{g_2}] \equiv f_1 \frac{g_1}{g_2} + f_0 \leq 0 \equiv f_1 g_1 g_2 + f_0 g_2^2 \leq 0$$

Infinity

$$(f_2 x^2 + f_1 x + f_0 < 0) [x // \infty] \equiv f_2 < 0 \vee (f_2 = 0 \wedge f_1 < 0) \vee (f_2 = 0 \wedge f_1 = 0 \wedge f_0 < 0)$$

Positive infinitesimals

$$(3x^2 + 6x - 3 > 0) [x // t - \varepsilon] \equiv 3t^2 + 6t - 3 > 0 \vee (3t^2 + 6t - 3 = 0 \wedge 6t + 6 \leq 0)$$

Formal solutions of quadratic equations

$$(f = 0) \left[x // \frac{g_1 + g_2 \sqrt{g_3}}{g_4} \right] \equiv \frac{g_1^* + g_2^* \sqrt{g_3}}{g_4^*} = 0 \equiv g_1^{*2} - g_2^{*2} g_3 = 0 \wedge g_1^* g_2^* \leq 0$$

Some Concrete Virtual Substitutions

Conventions: $f \in \mathbb{Z}[\mathbf{y}][x]$, $f_i, g_i, g_i^* \in \mathbb{Z}[\mathbf{y}]$

Quotients

$$(f_1 x + f_0 \leq 0)[x // \frac{g_1}{g_2}] \equiv f_1 \frac{g_1}{g_2} + f_0 \leq 0 \equiv f_1 g_1 g_2 + f_0 g_2^2 \leq 0$$

Infinity

$$(f_2 x^2 + f_1 x + f_0 < 0)[x // \infty] \equiv f_2 < 0 \vee (f_2 = 0 \wedge f_1 < 0) \vee (f_2 = 0 \wedge f_1 = 0 \wedge f_0 < 0)$$

Positive infinitesimals

$$(3x^2 + 6x - 3 > 0)[x // t - \varepsilon] \equiv 3t^2 + 6t - 3 > 0 \vee (3t^2 + 6t - 3 = 0 \wedge 6t + 6 \leq 0)$$

Formal solutions of quadratic equations

$$(f = 0) \left[x // \frac{g_1 + g_2 \sqrt{g_3}}{g_4} \right] \equiv \frac{g_1^* + g_2^* \sqrt{g_3}}{g_4^*} = 0 \equiv g_1^{*2} - g_2^{*2} g_3 = 0 \wedge g_1^* g_2^* \leq 0$$

Košta 2016 (PhD thesis)

- ▶ method generalizes to higher degree bounds
- ▶ generic implementation with a degree bound of 3 is newly available

Some Complexity Results

Upper bound on asymptotic worst-case complexity

doubly exponential in the input word length (and this is optimal for the problem)

In the linear case

doubly exponential	in	# quantifier alternations
singly exponential	in	# quantifiers
polynomial	in	# parameters (= unquantified variables)
polynomial	in	# atomic formulas

particularly good for

low degrees and many parameters

For comparison: Cylindrical Algebraic Decomposition (CAD)

[Collins 1973, Arnon, Hong, Brown, ...] doubly exponential in #all variables

Extended Quantifier Elimination

$$\text{Generalize } \exists x \varphi \longleftrightarrow \bigvee_{(\gamma, t) \in E} \gamma \wedge \varphi[t//x] \text{ to } \exists x \varphi \rightsquigarrow \left[\begin{array}{cc} \vdots & \vdots \\ \gamma \wedge \varphi[t//x] & x = t \\ \vdots & \vdots \end{array} \right]$$

A simple example

$$\varphi \equiv \exists x (ax^2 + bx + c = 0) \rightsquigarrow \left[\begin{array}{cc} a \neq 0 \wedge b^2 - 4ac \geq 0 & x = \frac{-b - \sqrt{b^2 - 4ac}}{2a} \\ a = 0 \wedge b \neq 0 & x = -\frac{c}{b} \\ a = 0 \wedge b = 0 \wedge c = 0 & x = \infty_1 \end{array} \right]$$

Semantics (for fixed parameters)

Whenever some left hand side condition holds, then $\exists x \varphi$ holds and the corresponding right hand side term is **one** sample solution.

[M. Kosta, T.S., A. Dolzmann, J. Symb. Comput. 2016]

For fixed choices of parameters, standard values can be efficiently computed for all ∞_i and ε_i in a post-processing step.

Quantifier Elimination Results for Our Problem

- **Positive quantifier elimination** exploits positivity of all variables.
- Successful not on φ_n but on $\exists\varphi_n$ ($n = 2, \dots, 10$):

n	$\exists\varphi_n$	$\exists\varphi_n[\lambda \leftarrow -\lambda]$	$\exists\varphi_n[\lambda \leftarrow 0]$	time (s)
2	false	false	false	< 0.01
3	false	false	false	19.28
4	false	false	false	21.58
5	false	false	false	19.09
6	false	false	false	23.72
7	false	false	false	23.89
8	false	false	false	22.35
9	true	false	false	0.17
10	true	false	false	0.17

- **Extended positive QE** delivers also sample solutions, e.g., for $n = 9$:

$$\alpha = 1$$

$$\lambda = 17617230.5528$$

$$v_1 = 0.000000170287832189$$

$$\delta = 1$$

$$\mu = 0$$

$$v_2 = 3$$

$$\gamma_0 = 0.0100554964908$$

$$\vartheta = 0.0000211443608455$$

$$v_3 = 1.24573093962$$

2012–2016

Hopf Bifurcations with Larger Models

Including MBO and MAPK

Switching to Convex Coordinates

In the previous example

- variables: concentrations of species
- parameters: reaction rates

Switching to Convex Coordinates

In the previous example

- ▶ variables: concentrations of species
- ▶ parameters: reaction rates

Stoichiometric network analysis (Clarke, 1980)

- ▶ We analyze system dynamics in flux space instead of concentration space.
- ▶ We represent the space of steady states with a combination of subnetworks.
- ▶ The subnetworks form a convex cone in flux space.
- ▶ Decomposing of the cone allows to search for lower dimensional facets (first).

Switching to Convex Coordinates

In the previous example

- ▶ variables: concentrations of species
- ▶ parameters: reaction rates

Stoichiometric network analysis (Clarke, 1980)



- ▶ We analyze system dynamics in flux space instead of concentration space.
- ▶ We represent the space of steady states with a combination of subnetworks.
- ▶ The subnetworks form a convex cone in flux space.
- ▶ Decomposing of the cone allows to search for lower dimensional facets (first).

This gives us


- ▶ certain Hurwitz conditions $\Delta_{n-1} = 0$, $\Delta_{n-2} > 0$, $\Delta_{n-3} > 0$ for Hopf bifurcation,
- ▶ furthermore $\Delta_{n-4} > 0, \dots, \Delta_1 > 0$ for empty unstable manifold,
- ▶ and positivity conditions on the variables and parameters.
- ▶ The steady state approximation (vector field = 0) is not explicit anymore.
- ▶ One reaction yields many such problems of various dimensions (in flux space).
- ▶ Δ_{n-1} is generally much larger than the $\Delta_{n-2}, \dots, \Delta_1$.

An Incomplete Subtropical Decision Procedure


► $f = -2x_1^5 + x_1^2 x_2 - 3x_1^2 - x_2^3 + 2x_2^2$




An Incomplete Subtropical Decision Procedure




- ▶ $f = -2x_1^5 + x_1^2x_2 - 3x_1^2 - x_2^3 + 2x_2^2$
- ▶ $f(1,1) = -3 < 0$




An Incomplete Subtropical Decision Procedure



- ▶ $f = -2x_1^5 + x_1^2x_2 - 3x_1^2 - x_2^3 + 2x_2^2$
- ▶ $f(1, 1) = -3 < 0$
- ▶ find $p \in]0, \infty[^2$ with $g(p) > 0$
and use intermediate value theorem




An Incomplete Subtropical Decision Procedure




- ▶ $f = -2x_1^5 + x_1^2x_2 - 3x_1^2 - x_2^3 + 2x_2^2$
- ▶ $f(1, 1) = -3 < 0$
- ▶ find $p \in]0, \infty[^2$ with $g(p) > 0$
and use intermediate value theorem

Computation of a Positive Point

- ▶ $\text{supp}^+(f) = \{(2, 1), (0, 2)\},$
 $\text{supp}^-(f) = \{(2, 0), (5, 0), (0, 3)\}$




An Incomplete Subtropical Decision Procedure




- ▶ $f = -2x_1^5 + x_1^2x_2 - 3x_1^2 - x_2^3 + 2x_2^2$
- ▶ $f(1, 1) = -3 < 0$
- ▶ find $p \in]0, \infty[^2$ with $g(p) > 0$
and use intermediate value theorem

Computation of a Positive Point

- ▶ $\text{supp}^+(f) = \{(2, 1), (0, 2)\}$,
 $\text{supp}^-(f) = \{(2, 0), (5, 0), (0, 3)\}$
- ▶ linear programming:
 $(0, 2) \in \text{newton}(f) \cap \text{supp}^+(f)$
 $(-3, -2)$ is normal of a separating
hyperplane oriented towards $(0, 2)$




An Incomplete Subtropical Decision Procedure



- ▶ $f = -2x_1^5 + x_1^2x_2 - 3x_1^2 - x_2^3 + 2x_2^2$
- ▶ $f(1, 1) = -3 < 0$
- ▶ find $p \in]0, \infty[^2$ with $g(p) > 0$
and use intermediate value theorem

Computation of a Positive Point

- ▶ $\text{supp}^+(f) = \{(2, 1), (0, 2)\}$,
 $\text{supp}^-(f) = \{(2, 0), (5, 0), (0, 3)\}$
- ▶ linear programming:
 $(0, 2) \in \text{newton}(f) \cap \text{supp}^+(f)$
 $(-3, -2)$ is normal of a separating
hyperplane oriented towards $(0, 2)$
- ▶ positive point at (t^{-3}, t^{-2})
for sufficiently large t



An Incomplete Subtropical Decision Procedure

$$f = -2x_1^5 + x_1^2x_2 - 3x_1^2 - x_2^3 + 2x_2^2, \quad f(1, 1) < 0 < f(2^{-3}, 2^{-2}) = f\left(\frac{1}{8}, \frac{1}{4}\right)$$

An Incomplete Subtropical Decision Procedure

$$f = -2x_1^5 + x_1^2x_2 - 3x_1^2 - x_2^3 + 2x_2^2, \quad f(1, 1) < 0 < f(2^{-3}, 2^{-2}) = f\left(\frac{1}{8}, \frac{1}{4}\right)$$

Computation of a zero

$$-2x_1^5 + x_1^2x_2 - 3x_1^2 - x_2^3 + 2x_2^2 = 0$$

$$x_1 = \frac{1}{8} + y \cdot \left(1 - \frac{1}{8}\right)$$

$$x_2 = \frac{1}{4} + y \cdot \left(1 - \frac{1}{4}\right), \quad y \in]0, 1[$$

An Incomplete Subtropical Decision Procedure

$$f = -2x_1^5 + x_1^2x_2 - 3x_1^2 - x_2^3 + 2x_2^2, \quad f(1, 1) < 0 < f(2^{-3}, 2^{-2}) = f\left(\frac{1}{8}, \frac{1}{4}\right)$$

Computation of a zero

$$-2x_1^5 + x_1^2x_2 - 3x_1^2 - x_2^3 + 2x_2^2 = 0$$

$$x_1 = \frac{1}{8} + y \cdot \left(1 - \frac{1}{8}\right)$$

$$x_2 = \frac{1}{4} + y \cdot \left(1 - \frac{1}{4}\right), \quad y \in]0, 1[$$

$$(1) \quad \bar{f} = f\left(\frac{1}{8} + y \cdot \left(1 - \frac{1}{8}\right), \frac{1}{4} + y \cdot \left(1 - \frac{1}{4}\right)\right)$$

$$= (-16807y^5 - 12005y^4 - 934y^3 - 20778y^2 + 285y + 1087)/D, \quad D \in \mathbb{N}$$

An Incomplete Subtropical Decision Procedure

$$f = -2x_1^5 + x_1^2x_2 - 3x_1^2 - x_2^3 + 2x_2^2, \quad f(1, 1) < 0 < f(2^{-3}, 2^{-2}) = f\left(\frac{1}{8}, \frac{1}{4}\right)$$

Computation of a zero

$$-2x_1^5 + x_1^2x_2 - 3x_1^2 - x_2^3 + 2x_2^2 = 0$$

$$x_1 = \frac{1}{8} + y \cdot \left(1 - \frac{1}{8}\right)$$

$$x_2 = \frac{1}{4} + y \cdot \left(1 - \frac{1}{4}\right), \quad y \in]0, 1[$$

$$\begin{aligned} (1) \quad \bar{f} &= f\left(\frac{1}{8} + y \cdot \left(1 - \frac{1}{8}\right), \frac{1}{4} + y \cdot \left(1 - \frac{1}{4}\right)\right) \\ &= (-16807y^5 - 12005y^4 - 934y^3 - 20778y^2 + 285y + 1087)/D, \quad D \in \mathbb{N} \end{aligned}$$

$$(2) \quad \text{real root isolation yields } y \in]0.2, 0.3[$$

An Incomplete Subtropical Decision Procedure

$$f = -2x_1^5 + x_1^2x_2 - 3x_1^2 - x_2^3 + 2x_2^2, \quad f(1, 1) < 0 < f(2^{-3}, 2^{-2}) = f\left(\frac{1}{8}, \frac{1}{4}\right)$$

Computation of a zero

$$-2x_1^5 + x_1^2x_2 - 3x_1^2 - x_2^3 + 2x_2^2 = 0$$

$$x_1 = \frac{1}{8} + y \cdot \left(1 - \frac{1}{8}\right)$$

$$x_2 = \frac{1}{4} + y \cdot \left(1 - \frac{1}{4}\right), \quad y \in]0, 1[$$

$$\begin{aligned} (1) \quad \bar{f} &= f\left(\frac{1}{8} + y \cdot \left(1 - \frac{1}{8}\right), \frac{1}{4} + y \cdot \left(1 - \frac{1}{4}\right)\right) \\ &= (-16807y^5 - 12005y^4 - 934y^3 - 20778y^2 + 285y + 1087)/D, \quad D \in \mathbb{N} \end{aligned}$$

(2) real root isolation yields $y \in]0.2, 0.3[$

(3) back-substitute real algebraic number $\langle \bar{f},]0.2, 0.3[$:

$$x_1 = \langle 686x^5 - 78x^3 + 584x^2 - 150x - 13,]0.32, 0.33[\rangle$$

$$x_2 = \langle 16807x^5 - 12005x^4 + 2026x^3 + 9122x^2 - 4609x + 323,]0.42, 0.43[\rangle$$

An Incomplete Subtropical Decision Procedure

Some Details on the LP Part


$$\text{supp}^+(f) = \{(2, 1), (0, 2)\} \rightsquigarrow B^+ = \begin{bmatrix} 2 & 1 & -1 \\ 0 & 2 & -1 \end{bmatrix}$$

$$\text{supp}^-(f) = \{(2, 0), (5, 0), (0, 3)\} \rightsquigarrow B^- = \begin{bmatrix} 2 & 0 & -1 \\ 5 & 0 & -1 \\ 0 & 3 & -1 \end{bmatrix}.$$

$$1. \quad \begin{bmatrix} -2 & -1 & 1 \\ 0 & 2 & -1 \\ 2 & 0 & -1 \\ 5 & 0 & -1 \\ 0 & 3 & -1 \end{bmatrix} \cdot (\mathbf{n}, c)^T \leq \begin{bmatrix} -1 \\ -1 \\ -1 \\ -1 \\ -1 \end{bmatrix} \quad \text{infeasible}$$

$$2. \quad \begin{bmatrix} 0 & -2 & 1 \\ 2 & 0 & -1 \\ 5 & 0 & -1 \\ 0 & 3 & -1 \end{bmatrix} \cdot (\mathbf{n}, c)^T \leq \begin{bmatrix} -1 \\ -1 \\ -1 \\ -1 \end{bmatrix} \quad \text{feasible with } \mathbf{n} = (-3, -2) \text{ and } c = 5$$

Methylene Blue Oscillator (MBO)



Reduction to 6 dimensions with essential species O_2 , O_2^- , HS , MB^+ , MB , MBH

Application to the Methylene Blue Oscillator (MBO)

Characteristics of a typical input polynomial (among 496) for MBO:

- ▶ 7 variables
- ▶ degree in each variable between 4 and 9
- ▶ around 6000 summands (monomials)
- ▶ essentially irreducible

Application to the Methylene Blue Oscillator (MBO)

Characteristics of a typical input polynomial (among 496) for MBO:

- ▶ 7 variables
- ▶ degree in each variable between 4 and 9
- ▶ around 6000 summands (monomials)
- ▶ essentially irreducible

Result Summary

- ▶ polynomial has zero: 30%
- ▶ polynomial has no zero: 67%
- ▶ incomplete method failed 3%
- ▶ computation time: 90 s on 60 cores (2.4 GHz Intel Xeon E5-4640)

Application to the Methylene Blue Oscillator (MBO)

Characteristics of a typical input polynomial (among 496) for MBO:

- ▶ 7 variables
- ▶ degree in each variable between 4 and 9
- ▶ around 6000 summands (monomials)
- ▶ essentially irreducible

Result Summary

- ▶ polynomial has zero: 30%
- ▶ polynomial has no zero: 67%
- ▶ incomplete method failed 3%
- ▶ computation time: 90 s on 60 cores (2.4 GHz Intel Xeon E5-4640)

We did not observe any unsatisfied Hurwitz inequalities.

Application to a Single MAPK Layer

Model with 12 reactions and 9 species taken from Conradi et al. (2008)

Typical MBO Polynomial

- ▶ 7 variables
- ▶ degrees between 4 and 9
- ▶ ~ 6000 monomials

Large MAPK Polynomial

- ▶ 10 variables
- ▶ degrees between 5 and 12
- ▶ ~ 863000 monomials (30 MB)

Application to a Single MAPK Layer

Model with 12 reactions and 9 species taken from Conradi et al. (2008)

Typical MBO Polynomial

- ▶ 7 variables
- ▶ degrees between 4 and 9
- ▶ ~ 6000 monomials

Large MAPK Polynomial

- ▶ 10 variables
- ▶ degrees between 5 and 12
- ▶ ~ 863000 monomials (30 MB)

MAPK Timings

- ▶ 10 s for finding positive and negative point
- ▶ 5 s for exact solving (with real algebraic numbers)

Application to a Single MAPK Layer

Model with 12 reactions and 9 species taken from Conradi et al. (2008)

Typical MBO Polynomial

- ▶ 7 variables
- ▶ degrees between 4 and 9
- ▶ ~ 6000 monomials

Large MAPK Polynomial

- ▶ 10 variables
- ▶ degrees between 5 and 12
- ▶ ~ 863000 monomials (30 MB)

MAPK Timings

- ▶ 10 s for finding positive and negative point
- ▶ 5 s for exact solving (with real algebraic numbers)

Some Trivia

- ▶ 25 s for reading and parsing the polynomial into Reduce/Redlog
- ▶ 2 s for determining the degrees of the 10 variables
- ▶ 3000+ pages when printing in a \LaTeX article

Application to a Single MAPK Layer

Model with 12 reactions and 9 species taken from Conradi et al. (2008)

Typical MBO Polynomial

- ▶ 7 variables
- ▶ degrees between 4 and 9
- ▶ ~ 6000 monomials

Large MAPK Polynomial

- ▶ 10 variables
- ▶ degrees between 5 and 12
- ▶ ~ 863000 monomials (30 MB)

MAPK Timings

- ▶ 10 s for finding positive and negative point
- ▶ 5 s for exact solving (with real algebraic numbers)

Some Trivia

- ▶ 25 s for reading and parsing the polynomial into Reduce/Redlog
- ▶ 2 s for determining the degrees of the 10 variables
- ▶ 3000+ pages when printing in a \LaTeX article

But the Hurwitz determinant inequalities were never satisfied!

2016–

Parametric Saddle-node Bifurcations with Another Model of MAPK

BIOMD0000000026 (MAPK)

$$\dot{x}_1 = k_2 x_6 + k_{15} x_{11} - k_1 x_1 x_4 - k_{16} x_1 x_5$$

$$\dot{x}_2 = k_3 x_6 + k_5 x_7 + k_{10} x_9 + k_{13} x_{10} - x_2 x_5 (k_{11} + k_{12}) - k_4 x_2 x_4$$

$$\dot{x}_3 = k_6 x_7 + k_8 x_8 - k_7 x_3 x_5$$

$$\dot{x}_4 = x_6 (k_2 + k_3) + x_7 (k_5 + k_6) - k_1 x_1 x_4 - k_4 x_2 x_4$$

$$\dot{x}_5 = k_8 x_8 + k_{10} x_9 + k_{13} x_{10} + k_{15} x_{11} - x_2 x_5 (k_{11} + k_{12}) - k_7 x_3 x_5 - k_{16} x_1 x_5$$

$$\dot{x}_6 = k_1 x_1 x_4 - x_6 (k_2 + k_3)$$

$$\dot{x}_7 = k_4 x_2 x_4 - x_7 (k_5 + k_6)$$

$$\dot{x}_8 = k_7 x_3 x_5 - x_8 (k_8 + k_9)$$

$$\dot{x}_9 = k_9 x_8 - k_{10} x_9 + k_{11} x_2 x_5$$

$$\dot{x}_{10} = k_{12} x_2 x_5 - x_{10} (k_{13} + k_{14})$$

$$\dot{x}_{11} = k_{14} x_{10} - k_{15} x_{11} + k_{16} x_1 x_5$$

Values for the rate constants

$k_1 = 0.02$	$k_2 = 1$	$k_3 = 0.01$	$k_4 = 0.032$	$k_5 = 1$	$k_6 = 15$
$k_7 = 0.045$	$k_8 = 1$	$k_9 = 0.092$	$k_{10} = 1$	$k_{11} = 0.01$	$k_{12} = 0.01$
$k_{13} = 1$	$k_{14} = 0.5$	$k_{15} = 0.086$	$k_{16} = 0.0011$		

BIOMD0000000026 (MAPK)

Steady-state-approximation and plugging in

$$-200x_1x_4 - 11x_1x_5 + 860x_{11} + 10000x_6 = 0$$

$$500x_{10} - 16x_2x_4 - 10x_2x_5 + 5x_6 + 500x_7 + 500x_9 = 0$$

$$-9x_3x_5 + 3000x_7 + 200x_8 = 0$$

$$-10x_1x_4 - 16x_2x_4 + 505x_6 + 8000x_7 = 0$$

$$-11x_1x_5 + 10000x_{10} + 860x_{11} - 200x_2x_5 - 450x_3x_5 + 10000x_8 + 10000x_9 = 0$$

$$2x_1x_4 - 101x_6 = 0$$

$$4x_2x_4 - 2000x_7 = 0$$

$$45x_3x_5 - 1092x_8 = 0$$

$$5x_2x_5 + 46x_8 - 500x_9 = 0$$

$$-150x_{10} + x_2x_5 = 0$$

$$11x_1x_5 + 5000x_{10} - 860x_{11} = 0$$

BIOMD0000000026 (MAPK)

Conservation laws

$$x_5 + x_8 + x_9 + x_{10} + x_{11} = k_{17}$$

$$x_4 + x_6 + x_7 = k_{18}$$

$$x_1 + x_2 + x_3 + x_6 + x_7 + x_8 + x_9 + x_{10} + x_{11} = k_{19}$$

Some realistic values for those new constants

$$k_{17} = 100$$

$$k_{18} = 50$$

$$k_{19} \in \{200, 500\}$$

BIOMD0000000026 (MAPK)

Conservation laws

$$x_5 + x_8 + x_9 + x_{10} + x_{11} = k_{17}$$

$$x_4 + x_6 + x_7 = k_{18}$$

$$x_1 + x_2 + x_3 + x_6 + x_7 + x_8 + x_9 + x_{10} + x_{11} = k_{19}$$

Some realistic values for those new constants

$$k_{17} = 100$$

$$k_{18} = 50$$

$$k_{19} \in \{200, 500\}$$

Saddle-node Bifurcation

The number of solutions for x_1, \dots, x_{11} changes from unique to non-unique.

Cylindrical Algebraic Decomposition

$\varphi(f_1, f_2)$ is a Boolean combination of constraints with left hand sides f_1, f_2 and right hand sides 0.


$$f_1(x, y) = 2y^2 - 2x^3 - 3x^2$$

$$f_1(A) = -1 < 0$$

$$f_1(B) = 2 > 0$$

$$f_1(C) = -5 < 0$$

$$f_1(D) = 0$$



Cylindrical Algebraic Decomposition

$\varphi(f_1, f_2)$ is a Boolean combination of constraints with left hand sides f_1, f_2 and right hand sides 0.

$$f_1(x, y) = 2y^2 - 2x^3 - 3x^2$$


$$f_1(A) = -1 < 0$$

$$f_1(B) = 2 > 0$$

$$f_1(C) = -5 < 0$$

$$f_1(D) = 0$$

$$f_2(x, y) = y^2 + x^2 - 1$$



Cylindrical Algebraic Decomposition

$\varphi(f_1, f_2)$ is a Boolean combination of constraints with left hand sides f_1, f_2 and right hand sides 0.

$$f_1(x, y) = 2y^2 - 2x^3 - 3x^2$$


$$f_1(A) = -1 < 0$$

$$f_1(B) = 2 > 0$$

$$f_1(C) = -5 < 0$$

$$f_1(D) = 0$$

$$f_2(x, y) = y^2 + x^2 - 1$$



Cylindrical Algebraic Decomposition

$\varphi(f_1, f_2)$ is a Boolean combination of constraints with left hand sides f_1, f_2 and right hand sides 0.

$$f_1(x, y) = 2y^2 - 2x^3 - 3x^2$$

$$f_1(A) = -1 < 0$$

$$f_1(B) = 2 > 0$$

$$f_1(C) = -5 < 0$$


$$f_1(D) = 0$$

$$f_2(x, y) = y^2 + x^2 - 1$$

$$g(x) = -2x^3 - 3x^2$$

...

projection polynomials



Application of CAD to Our MAPK Model

First Without Parameters

Recall $k_{17} = 100$, $k_{18} = 50$, $k_{19} \in \{200, 500\}$; **plug in** k_{17}, k_{18} .

For $k_{19} = 200$:

$$x^{(200)} = (90.6, 2.6, 10.4, 17.8, 35.9, 32.0, 0.0, 15.5, 2.3, 0.6, 45.4)$$

For $k_{19} = 500$:

$$x_1^{(500)} = (17.6, 6.9, 367.5, 36.6, 5.5, 12.8, 0.5, 83.4, 8.0, 0.2, 2.7)$$

$$x_2^{(500)} = (122.0, 14.6, 234.9, 14.5, 7.1, 35.0, 0.4, 69.4, 7.4, 0.7, 15.2)$$

$$x_3^{(500)} = (323.7, 9.4, 37.1, 6.7, 13.6, 43.1, 0.1, 20.8, 3.2, 0.8, 61.4)$$

- Float approximations for convenience — we have exact real algebraic numbers

Application of CAD to Our MAPK Model

Now with Parametric k_{19}

- ▶ Eliminate x_1, x_3, \dots, x_{11} by virtual substitution.
- ▶ Then CAD with variables k_{19} and x_2
- ▶ For all $k_{19} > 0$ there is at least one positive solution for x_2
- ▶ The system changes from one to three solutions around $k_{19} = 409.3$.

Application of CAD to Our MAPK Model

Now with Parametric k_{19}

- ▶ Eliminate x_1, x_3, \dots, x_{11} by virtual substitution.
- ▶ Then CAD with variables k_{19} and x_2
- ▶ For all $k_{19} > 0$ there is at least one positive solution for x_2
- ▶ The system changes from one to three solutions around $k_{19} = 409.3$.

The exact break point

is the only real zero in the interval $(409, 410)$ of $\sum_{i=0}^{10} c_i k_{19}^i$ with

$$c_{10} = 351590934502740290936895033267017158736060313940693076650155371250411$$

$$c_9 = -213699072852157674283997527746395583273033983170426080574800781989093156$$

$$c_8 = 25374851641220554774259605635053469432582109883965015804077119110958034090$$

$$c_7 = 12972493018300022707027639267804259251235991618029852880330004508564391594000$$

$$c_6 = -8468945963692802414226427249726123493448372439778349029355636316929687020660000$$

$$c_5 = 2231098270337406450670301663172664333421440833875848621423683265663846533079600000$$

$$c_4 = -376265008904112258290319173193792052014899485528994925965885895511831873444245100000$$

$$c_3 = 39262101548790869407057994985320156500968958361396178908180026842806643766783104000000$$

$$c_2 = -2492623990743029234974354081270296106309603462451517057779877596842448287799337600000000$$

$$c_1 = 70978850735887473459176997186175978425873267246760023212940616924643171868478080000000000$$

$$c_0 = -1062871192838985876948077114923898204990434138901495394834749613184670362810368000000000000$$

The Combined System Once More

$$-200x_1x_4 - 11x_1x_5 + 860x_{11} + 10000x_6 = 0$$

$$500x_{10} - 16x_2x_4 - 10x_2x_5 + 5x_6 + 500x_7 + 500x_9 = 0$$

$$-9x_3x_5 + 3000x_7 + 200x_8 = 0$$

$$-10x_1x_4 - 16x_2x_4 + 505x_6 + 8000x_7 = 0$$

$$-11x_1x_5 + 10000x_{10} + 860x_{11} - 200x_2x_5 - 450x_3x_5 + 10000x_8 + 10000x_9 = 0$$

$$2x_1x_4 - 101x_6 = 0$$

$$4x_2x_4 - 2000x_7 = 0$$

$$45x_3x_5 - 1092x_8 = 0$$

$$5x_2x_5 + 46x_8 - 500x_9 = 0$$

$$-150x_{10} + x_2x_5 = 0$$


$$11x_1x_5 + 5000x_{10} - 860x_{11} = 0$$

$$-k_{17} + x_{10} + x_{11} + x_5 + x_8 + x_9 = 0$$

$$-k_{18} + x_4 + x_6 + x_7 = 0$$

$$-k_{19} + x_1 + x_{10} + x_{11} + x_2 + x_3 + x_6 + x_7 + x_8 + x_9 = 0$$

A Minimum Vertex Cover



This Yields

$$\begin{aligned} &1062444k_{18}x_4^2x_5 + 23478000k_{18}x_4^2 + 1153450k_{18}x_4x_5^2 + 2967000k_{18}x_4x_5 \\ &\quad + 638825k_{18}x_5^3 + 49944500k_{18}x_5^2 - 5934k_{19}x_4^2x_5 - 989000k_{19}x_4x_5^2 \\ &\quad - 1062444x_4^3x_5 - 23478000x_4^3 - 1153450x_4^2x_5^2 - 2967000x_4^2x_5 \\ &\quad - 638825x_4x_5^3 - 49944500x_4x_5^2 = 0 \end{aligned}$$

$$\begin{aligned} &1062444k_{17}x_4^2x_5 + 23478000k_{17}x_4^2 + 1153450k_{17}x_4x_5^2 + 2967000k_{17}x_4x_5 \\ &\quad + 638825k_{17}x_5^3 + 49944500k_{17}x_5^2 - 1056510k_{19}x_4^2x_5 - 164450k_{19}x_4x_5^2 \\ &\quad - 638825k_{19}x_5^3 - 1062444x_4^2x_5^2 - 23478000x_4^2x_5 - 1153450x_4x_5^3 \\ &\quad - 2967000x_4x_5^2 - 638825x_5^4 - 49944500x_5^3 = 0 \end{aligned}$$

- ▶ We managed to compute a CAD with 2 parameters.
- ▶ Also a real triangularization
- ▶ This is where we are standing right now.

The story continues ...