



**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



**APPROVED  
LABEL**



**foxtrot**®


**SELECTIVE HERBICIDE**

ACTIVE CONSTITUENTS: 69 g/L FENOXAPROP-P-ETHYL  
34.5 g/L CLOQUINTOCET-MEXYL

SOLVENT: 390 g/L LIQUID HYDROCARBONS

**GROUP A HERBICIDE**

For the selective control of wild oats and annual phalaris in wheat, barley, triticale and cereal rye crops



0104wcd



**Ospray Pty Ltd** ABN 23 110 199 169

50 Middleton Road

Cottonvale QLD 4375 AUSTRALIA

Phone: 1800 624 597

[www.ospray.com.au](http://www.ospray.com.au)

Part of the **CHEMINOVA** Group

**Contents 10L**





## DIRECTIONS FOR USE

**Restraints:** DO NOT apply if rainfall is expected within 4 hours.

DO NOT apply to weeds or crop under stress due to, for example, very dry, very wet, nutrient deficient, frosty or diseased conditions.

CROP	WEED	RATE (mL/ha)	CRITICAL COMMENTS
Wheat, Barley, Triticale, Cereal rye	Wild oats ( <i>Avena</i> spp.)	475	Apply 3 to 4 weeks after sowing when cereal has 2 to 5 true leaves (Z12 to Z15). Do not use this rate when wild oats are past the four leaf stage (Z14) or when there are more than 200 plants per square metre.
		560	Apply generally 3 to 6 weeks after sowing when cereal is usually 2 leaf to early tillering (Z12 to Z22). Aim to apply to actively growing young weeds up to 5 total leaves (1 tiller) (Z14, Z21).
		635	Apply up to approximately 10 weeks after sowing to actively growing weeds up to mid tillering (Z22 to Z24)
	Annual phalaris ( <i>Phalaris minor</i> )	635	Apply generally 3 to 4 weeks after sowing when cereal usually has 2 to 5 true leaves (Z12 to Z15). Apply only to weed densities below 200 plants per square metre. Aim to apply to actively growing young weeds up to 4 total leaves (Z14).
		800	Apply generally 4 to 6 weeks after sowing when cereal has usually reached early tillering (Z21 to Z22). Aim to apply to actively growing young weeds up to 5 total leaves (1 tiller) (Z14, Z21).

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

## WITHHOLDING PERIODS

### Harvest

**DO NOT HARVEST FOR 10 WEEKS AFTER APPLICATION**

## GENERAL INSTRUCTIONS:

Results are best under good growing conditions. Application to weeds or crop under stress (eg. due to continuous severe frosts, nutrient deficient, diseased, dry or waterlogged conditions) should be avoided. Generally the best yield advantage will come from controlling weeds as early as possible.

### Resistant Weeds Warning

**GROUP A HERBICIDE**

FOXTROT Selective Herbicide is a member of the aryloxyphenoxypropionate group of herbicides. FOXTROT is a herbicide which inhibits production of acetyl CoA carboxylase. For weed resistance management FOXTROT is a Group A herbicide. Some naturally occurring weed biotypes resistant to FOXTROT, and other Group A herbicides, may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by FOXTROT or other Group A herbicides. Since occurrence of resistant weeds is difficult to detect prior to use, Ospray Pty Ltd accepts no liability for any losses that may result from the failure of FOXTROT to control resistant weeds.

DO NOT make more than one application of a Group A Herbicide to a crop per season. For further information on herbicide resistance management consult Ospray representative, CropLife,

trained agricultural consultant or local department agronomist.

### Export of Treated Produce

Growers should note that suitable MRLs or import tolerances may not be established in all markets for produce treated with FOXTROT Selective Herbicide. If you are growing produce for export, please check with Ospray Pty Ltd for the latest information on MRLs and import tolerances BEFORE using FOXTROT Selective Herbicide.

### Application

Aim for complete coverage of weeds with FOXTROT. Follow directions under "Equipment".

### Mixing

Add the required quantity of FOXTROT directly to the spray tank when about two-thirds full. Top up with water to the required volume, with agitators in motion, after adding broadleaf herbicides if required.

### Equipment

**Ground sprayers** - Standard boom sprays only are recommended and must be fitted with by-pass or mechanical agitation. It is recommended to use approximately 50 to 100 L of water/ha and a droplet size of 200 to 300 microns. Do not apply with boomless jets or misters.

**Aircraft** - FOXTROT may be applied by aircraft. Best results are obtained using 25 to 30 L of water per hectare with a steady cross wind. Do not exaggerate the swath width. Aim for a droplet size of 200 to 300 microns.

### Grazing

**DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 3 WEEKS AFTER APPLICATION**

## Compatibility

Mixtures of FOXTROT with broadleaf herbicides may lead to a reduction in grass weed control. This reduction can be minimised by closely following all use recommendations and restrictions.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to the aquatic environment. DO NOT contaminate streams, rivers or waterways with this product or the used container.

### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply in weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT plant susceptible crops (eg. corn, sorghum, oats or barley) for 3 weeks after application. **Undersown Clovers and Medics** - FOXTROT does not affect undersown clovers or medics or other broadleaf crops.

### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If

no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale vapour. When preparing spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and face shield or goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat and elbow-length PVC gloves. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126. If swallowed, do NOT induce vomiting. Give a glass of water.

### MSDS

Additional information is listed in the Material Safety Data Sheet.

### EXCLUSION OF LIABILITY

This product must be used strictly as directed, and in accordance with all instructions appearing on the label. Ospray Pty Ltd accepts no liability or responsibility for loss or damage arising from incorrect storage, handling or use.

® Trademark of Cheminova A/S, DENMARK

APVMA Approval No.

61410/10L/0607

BATCH No.

DOM:



0104wcd



**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



**APPROVED  
LABEL**

**foxtrot®**

**SELECTIVE HERBICIDE**

**ACTIVE CONSTITUENTS:** 69 g/L FENOXAPROP-P-ETHYL  
34.5 g/L CLOQUINTOCET-MEXYL

**SOLVENT:** 390 g/L LIQUID HYDROCARBONS

**GROUP A HERBICIDE**

For the selective control of wild oats and annual phalaris in wheat, barley, triticale and cereal rye crops



0104wcd



**Ospray Pty Ltd** ABN 23 110 199 169

50 Middleton Road

Cottonvale QLD 4375 AUSTRALIA

Phone: 1800 624 597

[www.ospray.com.au](http://www.ospray.com.au)

Part of the  **CHEMINOVA** Group

**Contents 20L**





## DIRECTIONS FOR USE

**Restrictions:** DO NOT apply if rainfall is expected within 4 hours.

DO NOT apply to weeds or crop under stress due to, for example, very dry, very wet, nutrient deficient, frosty or diseased conditions.

CROP	WEED	RATE (mL/ha)	CRITICAL COMMENTS
Wheat, Barley, Triticale, Cereal rye	Wild oats ( <i>Avena</i> spp.)	475	Apply 3 to 4 weeks after sowing when cereal has 2 to 5 true leaves (Z12 to Z15). Do not use this rate when wild oats are past the four leaf stage (Z14) or when there are more than 200 plants per square metre.
		560	Apply generally 3 to 6 weeks after sowing when cereal is usually 2 leaf to early tillering (Z12 to Z22). Aim to apply to actively growing young weeds up to 5 total leaves (1 tiller) (Z14, Z21).
		635	Apply up to approximately 10 weeks after sowing to actively growing weeds up to mid tillering (Z22 to Z24)
	Annual phalaris ( <i>Phalaris minor</i> )	635	Apply generally 3 to 4 weeks after sowing when cereal usually has 2 to 5 true leaves (Z12 to Z15). Apply only to weed densities below 200 plants per square metre. Aim to apply to actively growing young weeds up to 4 total leaves (Z14).
		800	Apply generally 4 to 6 weeks after sowing when cereal has usually reached early tillering (Z21 to Z22). Aim to apply to actively growing young weeds up to 5 total leaves (1 tiller) (Z14, Z21).

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

## WITHHOLDING PERIODS

### Harvest

**DO NOT HARVEST FOR 10 WEEKS AFTER APPLICATION**

### Grazing

**DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 3 WEEKS AFTER APPLICATION**

## GENERAL INSTRUCTIONS:

Results are best under good growing conditions. Application to weeds or crop under stress (eg. due to continuous severe frosts, nutrient deficient, diseased, dry or waterlogged conditions) should be avoided. Generally the best yield advantage will come from controlling weeds as early as possible.

### Resistant Weeds Warning

**GROUP A HERBICIDE**

FOXTROT Selective Herbicide is a member of the aryloxyphenoxypropionate group of herbicides. FOXTROT is a herbicide which inhibits production of acetyl CoA carboxylase. For weed resistance management FOXTROT is a Group A herbicide. Some naturally occurring weed biotypes resistant to FOXTROT, and other Group A herbicides, may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by FOXTROT or other Group A herbicides. Since occurrence of resistant weeds is difficult to detect prior to use, Ospray Pty Ltd accepts no liability for any losses that may result from the failure of FOXTROT to control resistant weeds. DO NOT make more than one application of a Group A Herbicide to a crop per season. For further information on herbicide resistance management consult Ospray representative, CropLife,

trained agricultural consultant or local department agronomist.

### Export of Treated Produce

Growers should note that suitable MRLs or import tolerances may not be established in all markets for produce treated with FOXTROT Selective Herbicide. If you are growing produce for export, please check with Ospray Pty Ltd for the latest information on MRLs and import tolerances BEFORE using FOXTROT Selective Herbicide.

### Application

Aim for complete coverage of weeds with FOXTROT. Follow directions under "Equipment".

### Mixing

Add the required quantity of FOXTROT directly to the spray tank when about two-thirds full. Top up with water to the required volume, with agitators in motion, after adding broadleaf herbicides if required.

### Equipment

**Ground sprayers** - Standard boom sprays only are recommended and must be fitted with by-pass or mechanical agitation. It is recommended to use approximately 50 to 100 L of water/ha and a droplet size of 200 to 300 microns. Do not apply with boomless jets or misters.

**Aircraft** - FOXTROT may be applied by aircraft. Best results are obtained using 25 to 30 L of water per hectare with a steady cross wind. Do not exaggerate the swath width. Aim for a droplet size of 200 to 300 microns.

## Compatibility

Mixtures of FOXTROT with broadleaf herbicides may lead to a reduction in grass weed control. This reduction can be minimised by closely following all use recommendations and restrictions.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to the aquatic environment. DO NOT contaminate streams, rivers or waterways with this product or the used container.

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply in weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT plant susceptible crops (eg. corn, sorghum, oats or barley) for 3 weeks after application. **Undersown Clovers and Medics** - FOXTROT does not affect undersown clovers or medics or other broadleaf crops.

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If

no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale vapour. When preparing spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and face shield or goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat and elbow-length PVC gloves. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, do NOT induce vomiting. Give a glass of water.

## MSDS

Additional information is listed in the Material Safety Data Sheet.

## EXCLUSION OF LIABILITY

This product must be used strictly as directed, and in accordance with all instructions appearing on the label. Ospray Pty Ltd accepts no liability or responsibility for loss or damage arising from incorrect storage, handling or use.

® Trademark of Cheminova A/S, DENMARK

APVMA Approval No.

61410/20L/0607

BATCH No.

DOM:



0104wcd