

Product Name: NOMINEE HERBICIDE
APVMA Approval No: 63233/111688



Label Name:	NOMINEE HERBICIDE
Signal Headings:	
Constituent Statements:	100 g/L BISPYRIBAC-SODIUM
Mode of Action:	GROUP B HERBICIDE
Statement of Claims:	For the control and the suppression of seed heads of Poa annua in Creeping Bentgrass, Tall Fescue, Couch and Perennial Ryegrass
Net Contents:	1L 500mL 5L
Restraints:	<p>DO NOT apply in autumn and winter DO NOT apply to plants that are stressed by moisture or temperature extremes. DO NOT apply to plants stressed by lack of nutrition DO NOT apply to plants stressed by human or mechanical traffic DO NOT apply if rain is expected within twelve hours of application. DO NOT irrigate for 24 hours after application DO NOT mow for 24 hours after application DO NOT apply more than twice per year. DO NOT sow Creeping Bentgrass, Perennial Ryegrass, Tall Fescue or Couch for at least 42 days after application of NOMINEE. DO NOT apply NOMINEE for at least 42 days after seeding of Creeping Bentgrass, Perennial Ryegrass, Tall Fescue or Couch</p> <p>SPRAY DRIFT RESTRAINTS DO NOT apply with spray droplets smaller than a MEDIUM spray droplet size category according to the nozzle manufacturer specifications that refer to the ASABE S572 or the BCPC Guideline. DO NOT apply when wind speed is less than 3 or more than 20 kilometres per hour as measured at the application site.</p>

DO NOT apply during surface temperature inversion conditions at the application site. Users of this product MUST make an accurate written record of the details of each spray application within 24 hours following application and KEEP this record for a minimum of 2 years. The spray application details that must be recorded are: 1 date with start and finish times of application; 2 location address and paddock/s sprayed; 3 full name of this product; 4 amount of product used per hectare and number of hectares applied to; 5 crop/situation and weed/pest; 6 wind speed and direction during application; 7 air temperature and relative humidity during application; 8 nozzle brand, type, spray angle, nozzle capacity and spray system pressure measured during application; 9 name and address of person applying this product. (Additional record details may be required by the state or territory where this product is used.

MANDATORY NO-SPRAY ZONES

DO NOT apply if there are aquatic and wetland areas including aquacultural ponds or surface streams and rivers downwind from the application area and within the mandatory no-spray zones shown in the table below:

FOR GROUND APPLICATION (medium droplet size

Windspeed Range at Time of Application - 3 to 20 kilometres per hour

Downwind Mandatory No-Spray Zone - 5 metres

DO NOT apply if there are sensitive crops, gardens, landscaping vegetation, protected native vegetation or protected animal habitat within the mandatory no-spray zones shown in the table below:

FOR GROUND APPLICATION (medium droplet size

Windspeed Range at Time of Application - 3 to 20 kilometres per hour

Downwind Mandatory No-Spray Zone - 15 metres

Directions for Use:

Other Limitations:

Withholding Periods:

DO NOT GRAZE TREATED AREAS OR FEED TREATED GRASS CLIPPINGS TO LIVESTOCK.

Trade Advice:

not applicable

General Instructions:

Resistance Warning:

GROUP B HERBICIDE

NOMINEE Herbicide is a member of the B group of herbicides. NOMINEE Herbicide has the inhibition of aceto-lactase synthase (ALS) mode of action. For weed resistance management, NOMINEE Herbicide is a Group B Herbicide. Some naturally-occurring weed biotypes resistant to NOMINEE Herbicide and other Group B Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by NOMINEE Herbicide or other Group B Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Sumitomo Chemical Australia Pty Ltd accepts no liability for any losses that may result from the failure of NOMINEE Herbicide to control resistant weeds.

Precautions:	
Protections:	<p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT</p> <p>Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with the chemical or used containers.</p>
Storage and Disposal:	<p>Keep out of reach of children. Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Protect from frost.</p> <p>Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.</p> <p>If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations.</p> <p>DO NOT burn empty containers or product.</p>
Safety Directions:	
First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia Tel. 131126; New Zealand 0800 764 766.
First Aid Warnings:	

NOMINEE HERBICIDE – 63233

Label Variation – DC14-14897452

DIRECTIONS FOR USE

SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
			<p>Note: NOMINEE is slow acting. The <i>Poa</i> will take several weeks before dying. This allows the Bentgrass, Ryegrass, Fescue or Couch to grow into the space vacated by the dying <i>Poa</i>. NOMINEE also provides useful seedhead suppression of <i>Poa</i> plants that survive the applications.</p> <p>Bentgrass: For best results, two consecutive applications should be applied with intervals dependent on rate.</p> <p>250mL/ha - Apply two applications at 7-14 day intervals 500mL/ha - Apply two applications at 7-21 day intervals 750mL-1L/ha - Apply two applications at 14-21 day intervals</p> <p>CAUTION: Two applications of NOMINEE can result in transient yellowing of the Bentgrass. Symptoms appear about two weeks after application, but the Bentgrass will fully recover approximately 4 to 6 weeks after the second application.</p> <p>NOMINEE can inhibit germination and growth of young Bentgrass seedlings.</p>
<u>Golf course greens</u> planted with Bentgrass (<i>Agrostis stolonifera</i>)	<i>Poa annua</i> (winter grass)	250mL-1.0L/ha (2.5- 10mL/100m ²)	For best results, two consecutive applications should be applied 7-21 days apart.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

NOMINEE HERBICIDE – 63233

Label Variation – DC14-14897452

GENERAL INSTRUCTIONS

General information

NOMINEE Herbicide is a selective post-emergence herbicide that, when used in accordance with the label, will selectively control winter grass (*Poa annua*) that is growing within established Creeping Bentgrass, Tall Fescue, Perennial Ryegrass and Couch. NOMINEE will also suppress seed head production by winter grass. NOMINEE displays excellent activity against emerged weeds, but has almost no pre-emergent activity and will therefore not control weeds that germinate after application.

Uptake and performance of NOMINEE is influenced by environmental conditions, cultural practices and spray coverage. For best results, apply NOMINEE only when both, turf and weeds are actively growing. Grass must not be under any stress (including moisture, temperature, nutrition and stress from trafficking or mowing). This can be an important consideration in early spring, if cold conditions persist. Irrigation is not required to activate NOMINEE. Turfgrass and sport fields should not be mowed or irrigated for 24 hours after application in order to allow time for the active ingredient to be absorbed and translocate within foliage.

NOMINEE has not been evaluated under all microclimates or against all biotypes of winter grass. Therefore, performance may be less effective in some locations, and against some biotypes of this weed species.

Tolerance to NOMINEE

NOMINEE is a very active herbicide, and users should use exercise good judgment and caution until familiarity is obtained with the product.

When applied in accordance with the label, NOMINEE has not caused commercially unacceptable injury to creeping bentgrass, tall fescue, perennial ryegrass or couch. However, treated turf may exhibit temporary chlorosis (yellowing) and mild growth regulation. These symptoms take longer to appear under cool and cloudy conditions, but are more persistent because the turfgrass is growing less vigorously.

In situations with high densities of wintergrass, the removal of wintergrass using NOMINEE may leave bare patches which will require time for the preferred grass species to grow into and fill these patches, or these patches will need to be resown/replanted.

NOMINEE may injure grass that is not well established. Observe the waiting periods recommended by this label.

Due to natural variability, growth stages and environmental conditions, potential users are encouraged to evaluate this product under typical growing conditions in a small area, and evaluate treated turf for 28 days for phytotoxicity, before widespread application.