

Product Name: 4FARMERS 24DB HERBICIDE  
APVMA Approval No: 95587/146456



Label Name:	4FARMERS 24DB HERBICIDE
-------------	-------------------------

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	--

Constituent Statements:	500 g/L 2,4-DB present as the dimethylamine salt
-------------------------	--

Mode of Action:	GROUP 4 HERBICIDE
-----------------	-------------------

Statement of Claims:	For the control of broadleaf weeds in seedling and established Lucerne, medic and clover pastures, peanuts and cereal crops undersown with Lucerne, medic or clover as specified in the Directions for Use table.
----------------------	---

Net Contents:	5 L - 1000 L
---------------	--------------

Restraints:	<p>DO NOT apply if rain is expected within 4 hours</p> <p>DO NOT apply to crop or weeds under stress (not actively growing) due to periods of extreme cold or heat, moisture stress (water logged or drought affected), poor nutrition or previous herbicide treatments as increased levels of crop effect and/or reduced levels of weed control may result.</p> <p>DO NOT use with hard water (&gt;500 ppm) or water containing suspended matter.</p> <p>SPRAY DRIFT RESTRAINTS:</p> <p>Specific definitions for terms used in this section of the label can be found at <a href="http://apvma.gov.au/spraydrift">apvma.gov.au/spraydrift</a></p> <p>DO NOT allow bystanders to come into contact with the spray cloud.</p> <p>DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.</p>
-------------	--

	<p>DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application (See further instructions in General Instructions for Ground and Aerial applications).</p> <p>DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.</p>
--	---

Directions for Use:	This section contains file attachment.
---------------------	--

Other Limitations:	
--------------------	--

Withholding Periods:	DO NOT GRAZE OR CUT FOR STOCK FEED FOR 7 DAYS AFTER APPLICATION
----------------------	---

Trade Advice:	
---------------	--

General Instructions:	This section contains file attachment.
-----------------------	--

Resistance Warning:	<p>4FARMERS 24DB Herbicide is a member of the Phenoxys group of herbicides. 4FARMERS 24DB Herbicide has the disruptors of plant cell growth mode of action. For weed resistance management 4FARMERS 24DB Herbicide is a Group 4 herbicide. Some naturally-occurring weed biotypes resistant to 4FARMERS 24DB Herbicide and other disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by 4FARMERS 24DB Herbicide or other disruptors of plant cell growth herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, 4 FARMERS AUSTRALIA PTY LTD accepts no liability for any losses that may result from the failure of 4FARMERS 24DB Herbicide to control resistant weeds.</p>
---------------------	---

Precautions:	THIS IS A PHENOXY HERBICIDE THAT CAN CAUSE SEVERE DAMAGE TO NATIVE VEGETATION AND SUSCEPTIBLE CROPS SUCH AS COTTON, GRAPES, TOMATOES, OILSEED CROPS AND ORNAMENTALS.
--------------	--

Protections:	<p>PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS</p> <p>Drift Warning: DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray drift onto nearby susceptible plants, adjacent crops, crop lands or pastures. Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.</p> <p>PROTECTION OF LIVESTOCK</p> <p>SPRAY-GRAZING WARNING:</p> <p>Some pasture weeds can be controlled by Spray-Grazing a combination of herbicide treatment followed by heavy grazing 7-10 days later. Sprayed weeds become more palatable to stock and a higher intake of some weed species after spraying can cause</p>
--------------	--

	<p>stock deaths and other problems. Increased quantities of poisonous plants may be eaten by stock when the Spray-Grazing technique is being used. Take care if pastures are dominated By Caltrop, Capeweed, Cape Tulip, Paterson's curse (Salvation Jane) or Variegated Thistle. With Paterson's Curse preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. DO NOT graze horses or pigs in Paterson's curse (Salvation Jane) infested pasture.</p> <p>PROTECTION OF WILDLIFE, FISH CRUSTACEA AND ENVIRONMENT DO NOT contaminate streams, rivers or watercourses with this product or used containers.</p>
--	---

Storage and Disposal:	<p>Keep out of reach of children.</p> <p>Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.</p> <p>Empty contents fully into application equipment. Close all valves and return to [point of supply/designated collection point/other specific collection details] for refill or storage. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.</p> <p>If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p>
-----------------------	---

Safety Directions:	<p>Poisonous if swallowed. May irritate the eyes and skin. This product is alkaline. Avoid contact with the eyes and skin. DO NOT inhale spray mist. When preparing spray, wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and face shield. When using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.</p>
--------------------	---

First Aid Instructions:	<p>If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 13 11 26.</p>
-------------------------	---

First Aid Warnings:	
---------------------	--

## DIRECTIONS FOR USE

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Lucerne - seedling and established	<p>Amaranths <i>Amaranthus</i> spp</p> <p><b>Burrs</b></p> <ul style="list-style-type: none"> <li>- Bathurst <i>Xanthium spinosum</i></li> <li>- Californian <i>Xanthium orientale</i></li> <li>- Noogoora <i>Xanthium strumarium</i></li> </ul> <p>Capeweed <i>Arctotheca calendula</i></p> <p>Charlock <i>Sinapis arvensis</i></p> <p>Dock (seedlings) <i>Rumex</i> spp</p> <p>Evening primrose <i>Oenothera stricta</i></p> <p>Fat hen <i>Chenopodium album</i></p> <p>Fumitory spp <i>Fumaria</i> spp</p> <p>Marshmallow <i>Malva parviflora</i></p> <p>Mexican poppy <i>Argemone</i> spp</p> <p>Mustards <i>Sisymbrium</i> spp</p> <p>Paterson's curse (Salvation Jane) <i>Echium plantagineum</i></p> <p>Plantains (Flat weeds) <i>Plantago</i> spp</p> <p>Prickly lettuce <i>Lactuca serriola</i></p> <p>Shepherd's purse <i>Capsella bursapastoris</i></p> <p>Sowthistle <i>Sonchus oleraceus</i></p> <p>Spiny emex (Doublegee) <i>Emex australis</i></p> <p>Stinging nettle <i>Urtica urens</i></p> <p><b>Thistles</b></p> <ul style="list-style-type: none"> <li>- Saffron <i>Carthamus lanatus</i></li> <li>- Slender <i>Carduus pycnocephalus</i> - Soldier</li> <li>- <i>Picnemon acarna</i></li> <li>- Spear <i>Cirsium vulgare</i></li> <li>- Variegated <i>Silybum marianum</i></li> </ul> <p>Turnip weed <i>Rapistrum rugosum</i> Wild (Mediterranean)</p> <p>turnip <i>Brassica tournefortii</i></p> <p>Wireweed <i>Polygonum aviculare</i></p>	All States	1 - 3.2 L	Spray seedling weeds at the 2-6 leaf stage when seedling lucerne has between 1-8 trifoliate leaves. Use lower end of rate range only for 2-4 leaf weeds under good growing conditions or for spray grazing. Small weeds which have been growing slowly under stress and are therefore physiologically older may not be well controlled. To avoid leaf distortion cut or graze established lucerne before spraying. Unless "Spray-Grazing" DO NOT cut, mow or graze for 4 weeks after application to maximise weed control.
<b>Clover and Medic Pastures,</b> - seedling and established	Refer to Weeds Controlled in Lucerne	All States	1 - 3.2 L	Spray weeds at the 2-6 leaf stage when seedling medic and clover has between 1-8 trifoliate leaves. Use lower end of rate range only for 2-4 leaf weeds under good growing conditions or for spray grazing.

**Legumes  
showing  
acceptable  
tolerance  
Clover**

- Arrow leaf cvs  
Cefalu, Zulu II
- Balansa cvs  
Balta,  
Frontier,  
Paradana
- Berseem cv  
Elite II
- Persian cvs  
Nitro, Lazer,  
Leeton, Prolific,  
Shaftal
- Rose cv Hykon
- Red
- Sub cvs  
Coolamon,  
Dalkeith, Gosse,  
Goulburn,  
June,ee,  
Nungarin,  
Riverina,  
Seaton Park,  
Trikkala, Urana
- White cv  
NuSiral etc

**Medic**

- Barrel cvs  
Caliph,  
Parabinga
- Burr  
(polymorpha)  
cv Circle valley
- Gama cvs  
Paraponto,  
Sapo
- Murex cv  
Zodiac
- Snail cvs  
Robinson, Sava

**Serradella**

- French cv  
Erica

**Medic and Clover safety**

**4FARMERS 24DB**

HERBICIDE may reduce  
initial vigour of most  
pasture legumes.

Depending on inherent  
tolerance this may not  
result in reduced spring  
growth or seed yield.

Tolerance can vary  
between seasons, legume  
type and cultivars.

Because of the continuing  
introduction of new pasture  
legumes and cultivars  
caution is advised before  
applying 4FARMERS  
24DB HERBICIDE to any  
new legume. Caution is  
especially advised if a  
treated crop is intended for  
seed production.

Application beyond the 8  
trifoliolate leaf stage can  
increase the chance of  
reduced vigour and  
production. The use of  
higher rates can also  
increase vigour reduction.  
DO NOT exceed 3L/ha  
when spraying Robinson or  
Sava snail Medic. Do not  
exceed 2.5 L/ha when  
spraying Sapo or Paraponto  
gama medic. DO NOT use  
on Berseem clover or on  
Red clover when it is being  
harvested for seed. Growth  
of Red clover may be  
retarded.

**4FARMERS 24DB**

HERBICIDE is unsafe to  
use on Biserrula, Gland  
clover, Popany vetch and  
Orion, Santiago & Serena  
medics due to the risk of  
excessive growth and seed.

**4FARMERS 24DB**

HERBICIDE is unsafe to  
use on chicory and plantain.

Sown Pasture containing Clover or Medics	Refer to Weeds Controlled in Lucerne	All States	1 - 3.2 L	Apply when clovers or medics are at the 1-8 trifoliate leaf stage. Grass seedlings should have 3 or more leaves. Medic and Clover safety refer to the Critical Comments above.
Sub. clover pasture	Doublegee <i>Emex australis</i>	WA only	1.2 L	Spray when Doublegee has less than 14 leaves: best results are obtained when weed has 6-8 leaves. Clover should have 2-8 trifoliate leaves. DO NOT apply to pastures when one or more false breaks in the season occur, as uneven germination will result. Doublegee which emerge after spraying will not be controlled. Medic and Clover safety refer to the Critical Comments above
Peanuts - Runner types - Virginia types - Spanish types	<i>Ipomoea</i> spp including - Bellvine <i>Ipomoea plebeia</i> - Morning glory <i>Ipomoea purpurea</i> - Red convolvulus <i>Ipomoea hederifolia</i> Spiny emex <i>Emex australis</i> Thornapples <i>Datura</i> spp	Qld only	1.3 - 1.7 L	Apply between 1 and 12 weeks after seedlings have emerged. Use the lower end of the rate range for control of <i>Ipomoea</i> spp weeds only.
Wheat, Barley, Oats, Triticale & Cereal  Rye undersown with lucerne, medic or clover	Refer to Weeds Controlled in Lucerne	All states	2.1 - 3.2 L	Spray weeds at the 2-6 leaf <b>stage</b> when seedling lucerne, medic and clover has between 1-8 trifoliate leaves. Use lower rate for 2-4 leaf weeds under good growing conditions. Apply to cereals after the "double ridge" stage which is normally through tillering after the main shoot has five leaves until booting (Z1533). The time to "double ridge" stage will vary depending on growing conditions and variety. Medic and Clover safety

				refer to the Critical Comments above.
Spot spraying (All Situations)	Refer to Weeds Controlled in Lucerne		200 - 400 mL/100 L water	Use lower rate for easy to kill weeds eg Fat hen or Variegated thistle. For harder to kill weeds eg Capeweed, Marshmallow, use higher rate. Where difficult to wet, hairy or very thick weed populations are present, add Chemwet 1000 to the spray mixture at a rate of 65 mL/100 L water.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS  
AUTHORISED UNDER APPROPRIATE LEGISLATION.**

## GENERAL INSTRUCTIONS:

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions.

## SPRAY APPLICATIONS AND DRIFT RISK ASSESSMENT

DO NOT apply when wind speed is less than 3 or more than 20 kilometres per hour (ground application) as measured at the application site.

DO NOT apply when wind speed is less than 3 or more than 15 kilometres per hour (aerial application) as measured at the application site.

USE ONLY COARSE or larger spray quality according to the definition for standard nozzles.

For aerial application it is recommended where possible for this product to be applied by an aerial applicator business that holds current accreditation for the Aerial Improvement Management System – “AIMS”, issued by the Aerial Agricultural Association of Australia Ltd.

## Checklist:

- Have you cleaned/decontaminated your boom sprayer?
- Have you contacted your neighbour prior to spraying?
- Is your sprayer set-up correctly for the particular application?
- Check – boom calibration
- At nozzle – nozzle choice
- Low drift/what spray quality
- Coarse or larger spray quality?
- Boom height – speed of intended application
- Water volume
- You must check, determine and record the weather conditions immediately prior to, and immediately after the spray application is made.
- Record – Temperatures
- Relative Humidity
- Delta T
- Wind Speed
- Is there a temperature inversion
- Night spraying – Extra care is required to ensure that inversion conditions are not present.

Use smoke generator to determine wind direction and presence of inversion conditions

For Further information contact 4 FARMERS AUSTRALIA PTY LTD and consult [spraywisedecisions.com.au](http://spraywisedecisions.com.au) – this is an online weather forecasting program and is recommended for use when planning your pesticide application.

When spraying in or near a cotton area, check online at [cottonmap.com.au](http://cottonmap.com.au) for the proximity of cotton fields.

**MIXING:** Mix with the required amount of water in the spray tank or mixing vessel. Apply using a low-volume boom spray fitted with fan jets, or by aircraft. Use mains or rain water for mixing. If not



available, water with a hardness of less than 500 ppm (expressed as calcium carbonate equivalent) can be used but is not preferred. DO NOT tank mix Ammonium Sulphate with 4FARMERS 24DB HERBICIDE in hard water for this may cause nozzle blockages.

APPLICATION: Apply using boom or spot spray equipment. Application must be uniform. Nozzle type, output, speed and pressure should be adjusted to deliver a Coarse spray quality in 110 - 220 litres per hectare by ground application and a minimum of 40 litres of water per hectare by aerial application.