

Product Name: Farmalinx DSMA 100 Herbicide
APVMA Approval No: 88844/124657



Label Name:	ProForce Dodger 100 SL Herbicide
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	100 g/L DISODIUM METHYLARSONATE = DSMA
Mode of Action:	GROUP Z HERBICIDE
Statement of Claims:	A selective post-emergent contact herbicide for the control of certain broadleaf weeds and grasses in sports turf and lawn as specified in the Directions for Use.
Net Contents:	500 mL - 200 L
Restraints:	DO NOT spray on windy days because of possible drift and damage to desirable plants. DO NOT apply with misters.
Directions for Use:	This section contains file attachment. File Name: ProForce Dodger 100 SL Herbicide DIRECTIONS FOR USE.pdf File Size: 91816 bytes
Other Limitations:	NOT TO BE USED IN THE HOME GARDEN.
Withholding Periods:	
Trade Advice:	

General Instructions:	
Resistance Warning:	<p>RESISTANT WEEDS WARNING GROUP Z HERBICIDE</p> <p>ProForce DODGER 100 SL Herbicide is a member of the Organoarsenic group of herbicides. DODGER 100 SL is a herbicide with diverse sites of action. For weed resistance management DODGER 100 SL is a Group Z herbicide. Some naturally occurring weed biotypes resistant to DODGER 100 SL and other Group Z herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by DODGER 100 SL or other Group Z herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Indigo Specialty Products Pty Ltd accepts no liability for any losses that may result from the failure of DODGER 100 SL to control resistant weeds.</p>
Precautions:	
Protections:	<p>PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants, adjacent cropping lands or pastures.</p> <p>PROTECTION OF LIVESTOCK DO NOT feed treated grass clippings from treated areas to poultry or other animals.</p> <p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Dangerous to fish. DO NOT contaminate streams, rivers or watercourses with the chemical or used containers.</p>
Storage and Disposal:	<p>Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers.</p> <p>Triple rinse container before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, State or Territory government regulations. DO NOT burn empty container or product.</p>
Safety Directions:	<p>Poisonous if absorbed by skin contact or swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat and rubber gloves. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.</p>

First Aid Instructions:	If poisoning occurs contact a doctor or Poisons Information Centre Phone Australia 13 11 26, New Zealand 0800 764 766.
-------------------------	---

First Aid Warnings:	
---------------------	--

DIRECTIONS FOR USE

CROP/SITUATION	WEEDS CONTROLLED	STATE	RATE
Turf and lawn grasses of; Common couch, Indian couch	Crab Grass (NSW), Kyllinga Weed, Mullumbimby Couch, Nut Grass,	Qld, NSW, Vic, SA only	Use 145 mL in 10 L of water over 10 m ² .
		WA only	Use 63 mL in 10 L of water over 10 m ² .
Bent and Fescue	Paspalum, Sedges, Various species,	Qld, NSW, Vic, SA only	Use 88 mL in 10 L of water over 10 m ² .
		WA only	Use 63 mL in 10 L of water over 10 m ² .
Buffalo	Summer Grasses		

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**