



Company Name: SPALDING HOLDINGS PTY. LTD.
Product Name: Spalding Caperjet Selective Herbicide
APVMA Approval No: 83255/107848

Label Name:	Spalding Caperjet Selective Herbicide
-------------	---------------------------------------

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	--------------------------------------------------------------------------------------------

Constituent Statements:	340 g/L MCPA (present as the dimethylamine salt) 80 g/L DICAMBA (present as the dimethylamine salt)
-------------------------	--------------------------------------------------------------------------------------------------------

Mode of Action:	GROUP I HERBICIDE
-----------------	-------------------

Statement of Claims:	For the Control of Certain Broadleaf Weeds in Winter Cereals, Pastures, Turf and Non-Crop Areas as per the Directions for Use
----------------------	-------------------------------------------------------------------------------------------------------------------------------

Net Contents:	1, 5, 10, 20 litres
---------------	---------------------

Restraints:	DO NOT spray when rain seems likely to occur with 4 hours. DO NOT spray when weeds are wet with dew or rain or under stress from drought, low soil fertility, extreme cold or water logging. DO NOT spray outside recommended crop growth as crop damage may result. DO NOT apply to crops or pastures with clover, lucerne or medics. DO NOT apply to legume pastures. DO NOT apply to plants suffering from stress.
-------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Directions for Use:	This section contains file attachment.
---------------------	----------------------------------------

Other Limitations:	
--------------------	--

Withholding Periods:	DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION HARVEST WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED
----------------------	----------------------------------------------------------------------------------------------------------------------------------

Trade Advice:	
---------------	--

General Instructions:	<p>MIXING Add the required amount of Spalding Caperjet directly to the tank and mix well.</p> <p>APPLICATION For Boom Application: Apply by a properly calibrated boom spray using not less than 50 litres of water per hectare unless indicated otherwise in Critical Comments column. For Aerial Application: Apply at least 25 litres total spray volume per hectare. The aircraft should fly as low as practical under the prevailing conditions to minimise drift. For High Volume Application: Apply minimum 1500 litres per hectare. For Knapsack Application: A 15 litre knapsack treats 150 square metres. NOTE: Spray when weeds are young and actively growing. A temporary wilting may be evident in crops after application. If seasonal conditions are unusually late or dry seek further advice as damage may result from spraying under these conditions. Growers should seek advice before spraying recently released cereal varieties. There is nil tolerance of pea seed contaminates in cereal grain and less than 100% control of Volunteer Peas could result in dockage.</p> <p>COMPATABILITY Spalding Caperjet is compatible with flowable terbutryn.</p>
-----------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Resistance Warning:	<p>GROUP I HERBICIDE Spalding Caperjet Selective Herbicide is a member of the Benzoic acid and Phenoxy groups of herbicides. Spalding Caperjet Selective Herbicide has the disruptors of plant cell growth mode of action. For weed resistance management Spalding Caperjet is a Group I Herbicide. Some naturally occurring weed biotypes resistant to Spalding Caperjet Selective Herbicide and other Disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Spalding Caperjet Selective Herbicide or other Group I Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Spalding Holdings Pty Ltd accepts no liability for any losses that may result from the failure of Spalding Caperjet Selective Herbicide to control resistant weeds.</p>
---------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Precautions:	
--------------	--

Protections:	<p>PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS Spray equipment must be calibrated accurately prior to working in crops. Minimise spray drift by using low pressure and nozzles which DO NOT give a fine droplet size. DO NOT use in high wind. DO NOT apply this product on or near desirable trees or plants or in locations where the chemical may be washed or moved into contact with their roots. Keep container sealed (airtight) when not in use. All spray equipment must be thoroughly washed out after use and must not be used for spraying horticultural crops. DO NOT apply under weather conditions, or from spraying equipment, that may cause drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, vines, lupins, fruit trees and ornamentals.</p>
--------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

	<p>PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT</p> <p>DO NOT use container for any other purpose.</p> <p>DO NOT contaminate streams, rivers or waterways with the chemical or used containers.</p>
--	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Storage and Disposal:	<p>Store in the closed, original container in a dry, cool, well-ventilated area away from children, animals, food, feedstuffs, seed and fertilizers. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point.</p> <p>If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in accordance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.</p>
-----------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Safety Directions:	<p>Harmful if absorbed by skin contact, inhaled or swallowed. Will irritate the eyes and skin. DO NOT inhale vapour. Repeated exposure may cause allergic disorders. When opening container and preparing spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves, face shield or goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat and elbow length PVC gloves. If product in eyes, wash out immediately with water. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.</p>
--------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

First Aid Instructions:	<p>If poisoning occurs, contact a doctor, or Poisons Information Centre (Telephone Australia 13 11 26).</p>
-------------------------	-------------------------------------------------------------------------------------------------------------

First Aid Warnings:	
---------------------	--

DIRECTIONS FOR USE

Crop & Situation	Weeds	State	Rate/ha	Critical Comments
Barley, Commercial Rye, Oats, Triticale, Wheat From early tillering stage (Zadoks 21)	Capeweed, Charlock, Common Ice-Plant, seedling Docks, Doublegee (Three cornered Jack, Spiny Emex) Hexham Scent (Melilotus), Mustards, Solider Thistle, Sorrel Seedlings, Tares, Tree Hogweed, Turnip Weed (<i>Rapistrum rugosum</i>), Vetches, Ward's Weed, Wild Radish, Wild Turnip (<i>Brassica tournefortii</i>), Wireweed (Hogweed)	All States	1L	Spray when most weeds are in the 2 to 4 leaf stage. In some instances suppression only of Wireweed (Hogweed), Capeweed and Wild Radish will occur.
	Seedling Cotton Thistle, Fat-Hen	Tas		
Barley, Commercial Rye, Oats, Triticale, Wheat From early tillering stage (Zadoks 21) until the fully tillered stage and before jointing occurs (Zadoks 30)	Cotton Thistle to bud formation	Tas	1.7L	Spray when most weeds are still in the rosette or seedling stage. N.B. Wild radish may need to be resprayed (with 2,4-D or MCPA) if future germinations occur after spraying. In some instances, suppression only of wireweed (hogweed), hoary cress and climbing buckwheat will be obtained.
	Mintweed, New Zealand Spinach	Qld, NSW		
	Capeweed, Charlock, Chickweed, Climbing Buckwheat (Black Bindweed), Clover, Common Ice-Plant, Seedling Docks, Doublegee (Three cornered Jack, Spiny Emex), Fat-Hen, Hexham Scent (Melilotus), Seedling Hoary Cress, Mouse-eared Chickweed, Mustards, Narrow-leaved Plantain, Saffron Thistle, Spurry, Swinecress, Tares, Tree Hogweed. Turnip Weed (<i>Rapistrum rugosum</i>), Variegated Thistle, Vetch, Wild Radish, Wild Turnip (<i>Brassica tournefortii</i>), Wireweed (Hogweed)	Qld, NSW, Tas, WA, Vic		
		SA	1.4L	
Barley, Commercial Rye, Triticale From early tillering stage (Zadoks 21) until the fully tillered stage before jointing occurs (Zadoks 30)	Volunteer Peas	SA	1L plus 60mL/ha Spalding Dicamba 500 Selective Herbicide	Apply when most peas are 5-7cm high.
Oats, Wheat From early tillering stage (Zadoks 21) until the fully tillered stage before jointing occurs (Zadoks 30)			1L/ha plus 120mL/ha Spalding Dicamba 500 Selective Herbicide	

Situation	Weeds Controlled	State	Rate			Critical Comments
			Per ha	Per 100L	Per 15L Knap-sack	
Grass Pastures	Bathurst Burr, Caltrop (Yellow Vine), Chicory, Crofton Weed, Cudweed, Dandelion, Deadnettle, Fleabanes, Fumitory, Groundsel, Heliotrope, Inkweed, Mintweed, Nettles, Noogoora Burr, Pennycress, Potato Weed, Ragwort, Ripplewort, Shepherd's Purse, Smartweed, Soft Rush, Stinkwort, Thistles including Cockspur, Scotch (Spear), Slender, Star, Variegated, Wireweed (Hogweed)	All States	2.8-4L	190mL – 270mL	60mL	Use the higher rate on larger weeds.
	Cotton Thistle	Tas				
		Buttercups, Hoary Cress seedlings, Horehound, Khaki Weed, Purple-top, Veined Verbena	All States	4-7L	470mL	100mL
Turf	Bindy-Eye, Capeweed, Catsear, Chickweed, Clovers, Cotula, Creeping Oxalis, Cudweeds, Dandelion, Dock, Fleabanes, Fumitory, Jo-Jo (Onehunga), Lamb's Tongue, Medics, Mustards, Narrow-leaved Plantain, Pearlwort, Pennywort, Peppergrass, Radish, Scarlet Pimpernel, Slender Thistle, Soldier Thistle, Spear Thistle, Toad Rush, Variegated Thistle, Wild Turnip, Wireweed (Hogweed)	All States	Golf Tee, Golf Green: 100mL in 15L water per 150m ² Bowling Green: 900mL in 200L water per 1400m ² Large Turf Area: (Field, Park, Golf Fairway) 6.5L in 250-400L water per 10,000m ² (1 hectare)			Apply to actively growing weeds and moist soil. DO NOT use on Lippia or Strawberry Clover or Buffalo grass. DO NOT mow turf for 2 days before or after application or fertilize within two weeks of spraying.
	Fat-Hen, Purple – Top, Red Flowered Mallow, Stagger Weed, Swinecress	Qld, NSW, Tas, SA, WA,				
Non-Crop Situations	Artichoke, Bindweed, Fennel, Hemlock, Thistle seedlings	All States	5.2L	350mL	80mL	A repeat spray may be necessary.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**