

Product Name: AMGROW ORGANIX WEED BLITZ CONCENTRATE HERBICIDE
APVMA Approval No: 68676/117712



Label Name:	AMGROW ORGANIX WEED BLITZ CONCENTRATE HERBICIDE
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	680 g/L PINE OIL
Statement of Claims:	For non-selective weed control of seedlings and young annual weeds and grasses in Commercial, Industrial and Public Service Areas, around Agricultural buildings and other non-crop farm situations. For suppression of established weeds and perennial species on paths, pavers and concrete.
Net Contents:	5 L - 942 L
Restraints:	DO NOT allow spray to contact crop plants. DO NOT apply if rain is likely within 2 hours of treatment. DO NOT spray onto rubber.
Directions for Use:	
General Instructions:	
Precautions:	PRECAUTIONS Re-Entry Period: Do not allow entry into treated areas until the spray has dried unless wearing cotton overalls (or equivalent clothing) and chemical resistant gloves.

Protections:	<p>PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby areas, including susceptible plants/crops, cropping lands or pastures, streams/rivers, wetlands or watercourses. The use of this product may temporarily affect earthworm populations in the treated areas. Field experience shows that this effect is transient and the earthworm population recovers.</p> <p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT apply Amgrow Organix Weed Blitz Concentrate Herbicide if water bodies, water courses, or wetlands are within 30m down-wind of the application area.</p>
--------------	---

Storage and Disposal:	<p>STORAGE AND DISPOSAL Store in the closed, original container in a cool, dry place out of the reach of children. Do not store in direct sunlight.</p> <p>For non-refillable containers: Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p> <p>For refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply or designated collection point for refill or storage.</p>
-----------------------	---

Safety Directions:	<p>SAFETY DIRECTIONS Will irritate the nose, throat and skin. Will damage the eyes. Avoid contact with eyes and skin. Do not inhale vapour or spray mist. When opening the container, preparing spray and using the prepared spray, if applying by low pressure hand wand, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat, elbow-length PVC gloves, goggles and a halfpiece respirator. When using the prepared spray if applying by boomspray equipment and high pressure hand wand, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat, elbow-length PVC gloves and goggles. If product on skin, immediately wash with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves, goggles and contaminated clothing and respirator and if rubber wash with detergent and warm water.</p>
--------------------	---

First Aid Instructions:	<p>If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia 13 11 26. If swallowed, do NOT induce vomiting. Give a glass of water.</p>
-------------------------	--

First Aid Warnings:	
---------------------	--

DIRECTIONS FOR USE

Crop	Weeds Controlled		Rate	Critical Comments
Orchards & Vineyards	Prince of Wales feather	<i>Amaranthus powellii</i>	200 mL/L	Ensure spray contacts all target area and that all weed foliage is covered.
	Scarlet pimpernel	<i>Anagallis arvensis</i>		
Commercial, Industrial and Public Service Areas, Agricultural buildings and other farm situations. For suppression of established weeds and perennial species on paths, patios, pavers and concrete.	Volunteer oats	<i>Avena sativa</i>	200 mL/L	DO NOT apply more than 2 applications to the treated area, per 12 months. Control: Apply at early vegetative stage of weeds. Best results are obtained at 2-4 true leaf stage when weed canopy is less than 15 cm in height. Suppression: Established weeds may require more than one application. Repeat spray after 7-10 days if required. Perennial weeds and grasses may only be suppressed and not fully controlled and will require repeated sprays to deter re-growth. Apply preferably during Spring and late Autumn when weeds are young, actively growing and weather is calm. Spray over all weed foliage to ensure full foliage penetration and coverage. Partial foliage penetration or coverage will result in partial control. Best results achieved in warm, sunny conditions.
	Cobblers pegs	<i>Bidens pilosa</i>		
	Wild Turnip	<i>Brassica rapa ssp. silvestris</i>		
	Shepherd's purse	<i>Capsella bursa-pastoris</i>		
	Chickweed	<i>Cerastium fontanum ssp. trivale</i>		
	Fat hen	<i>Chenopodium album</i>		
	Couch grass	<i>Cynodon dactylon</i>		
	Summer grass	<i>Digitaria ciliaris</i>		
	Crab grass	<i>Digitaria sanguinalis</i>		
	Willow herb	<i>Epilobium billardierianum</i>		
	Storkbill	<i>Erodium moschatum</i>		
	Pinkweed	<i>Fumaria muralis</i>		
	Toad grass	<i>Juncus bufonius</i>		
	Winter grass	<i>Poa annua</i>		
	Wireweed or hogweed	<i>Polygonum aviculare</i>		
	Common pigweed	<i>Portulacca oleracea</i>		
	Wild radish	<i>Raphanus raphanistrum</i>		
	Sorrel	<i>Rumex acetosella</i>		
	Hairy nightshade	<i>Solanum chenopodioides</i>		
	Blackberry nightshade	<i>Solanum nigrum</i>		
	Prickly sow thistle	<i>Sonchus asper</i>		
	Milk thistle	<i>Sonchus oleraceus</i>		
	Stagger weed	<i>Stachys arvensis</i>		
	Stinking roger	<i>Tagetes minuta</i>		
	White clover	<i>Trifolium repens</i>		
	Stinging nettle	<i>Urtica urens</i>		
	Ivy leaf speedwell	<i>Veronica hederifolia</i>		
	Scrambling speedwell	<i>Veronica persica</i>		

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Amgrow Organix Weed Blitz Concentrate Herbicide is a natural contact weed control that acts on the foliage, causing it to quickly dehydrate and collapse. Results often can be seen within hours of application. Amgrow Organix Weed Blitz Concentrate Herbicide will control many weed species and is non-selective.

The natural composition of Amgrow Organix Weed Blitz Concentrate Herbicide may require repeat applications to control some difficult species. Cultivating the dehydrated weeds will act as a compost adding valuable nutrients to your soil.

MIXING

Thoroughly clean out spray container with clean water. Use correct measuring device and do not mix more than is needed. To prevent foaming and ensure adequate mixing, add part of water to spray tank, then Amgrow Organix Weed Blitz Concentrate Herbicide, mix and add remaining water. Agitate whilst mixing and spraying.

APPLICATION

Ensure that all weed foliage is totally covered with spray as the product is a contact spray only. Partial coverage will give partial control. Use a foaming nozzle or spraying tips TG2 or TG3 for best results. Otherwise, use a flat fan nozzle and coarse droplets. Optimum application volumes are related to weed maturity and density. A water volume of 700 L/ha is recommended. Do not spray on to rubber. Avoid area until dry. Amgrow Weed Blitz Concentrate Herbicide is not known to cause staining on hard surfaces, but it is advisable to test a small area prior to further application.

Amgrow Organix Weed Blitz Concentrate Herbicide can be applied by ground boom, low pressure hand-wand and a high pressure hand wand. Note that hand wand = Handheld equipment not carried on the back of the user; 'low pressure' hand wand = manually pressurised spraying equipment, 'high pressure' hand wand = spraying equipment pressurised by a motor.

For Ground boom spray:

Apply as a band spray 400 mm wide on each side of the crop row. Ensure that all the weed foliage is totally covered with spray as manually pressurised is a contact spray only. Partial cover will give partial control. Use a foaming nozzle or spray tips TG2 or TG3 for best results. Otherwise use a flat fan nozzle and COARSE droplets. Optimum application volumes are related to weed maturity and density. A water volume of 700 L/ha is recommended. Do not spray onto rubber. Avoid area until dry.

For Hand Wand Spray:

Mix thoroughly and spray as close as possible onto foliage, ensuring full coverage of the weed. FOR BEST RESULTS: Handheld sprayers should be adjusted to give a COARSE spray. Hand wand sprayers should be fitted with a solid cone nozzle (1.5 – 2.0 L/minute, 40 – 60 psi). Spraying Systems Co – TG2 is recommended or equivalent. When finished, remove unused product from sprayer and rinse thoroughly with water. Dispose of excess spray on weeds.

1 litre of prepared spray covers approximately 13 square meters. Avoid prolonged contact with aluminium or rubber. Pavers treated with protective coating should be avoided. Test for colour fastness on sample area before spraying. Application area may become slippery when wet. Avoid area until dry. Contain any spillage with absorbent material and dispose in garbage.
