



Product Name: KELPIE PROMIX 550 GLY HERBICIDE
APVMA approval No.: 88261 / 125459

Label Name:	KELPIE PROMIX 550 GLY HERBICIDE
Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	550 g/L GLYPHOSATE present as the potassium salt and monoethanolamine salts
Mode of Action:	<div>GROUP M HERBICIDE</div>
Statement of Claims:	Non-selective herbicide for the control of many annual and perennial weeds.
Net Contents:	1 L - 1000 L
Restrains:	DO NOT disturb weeds by cultivation, sowing or grazing for six hours of daylight following treatment of Annual Weeds and seven days for Perennial Weeds to ensure herbicide absorption, unless specified otherwise in CRITICAL COMMENTS.
Directions for Use:	This section contains file attachment.
Other Limitations:	
Withholding Periods:	NOT REQUIRED WHEN USED AS DIRECTED
Trade Advice:	

General Instructions:	This section contains file attachment.
Resistance Warning:	<p>KELPIE® PROMIX 550 GLY HERBICIDE is a member of the Glycines group of herbicides. KELPIE® PROMIX 550 GLY HERBICIDE has the inhibition of EPSP synthase mode of action. For weed resistance management, KELPIE® PROMIX 550 GLY HERBICIDE is a Group M herbicide.</p> <p>Some naturally occurring weed biotypes resistant to KELPIE® PROMIX 550 GLY HERBICIDE and other Group M herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by KELPIE® PROMIX 550 GLY HERBICIDE or other Group M herbicides.</p> <p>Since the occurrence of resistant weeds is difficult to detect prior to use, Sinochem International Australia accepts no liability for any losses that may result from the failure of KELPIE® PROMIX 550 GLY HERBICIDE to control resistant weeds.</p>
Precautions:	
Protections:	<p>PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS Avoid contact with foliage, green stems or fruit of crops, desirable plants and trees, as severe injury or destruction may result.</p> <p>DO NOT apply under weather conditions, or from spray equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.</p> <p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT contaminate dams, rivers or streams with the product or used container. DO NOT apply to weeds growing in or over water. DO NOT spray across open bodies of water.</p>
Storage and Disposal:	<p>Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site.</p> <p>If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p> <p>For REFILLABLE containers: Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.</p>
Safety Directions:	Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container, preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat, elbow length PVC or nitrile

	gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each days use, wash gloves, face shield or goggles and contaminated clothing.
--	---

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.
-------------------------	--

First Aid Warnings:	
---------------------	--

CONSERVATION TILLAGE

SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
SOUTHERN AUSTRALIA FULL SOIL DISTURBANCE Prior to sowing a crop or pasture with full soil disturbance by cultivation or sowing with a tyned implement.	Barley Grass, Brome Grass, Volunteer Cereals, Wild Oats	330-650 mL/ha pre-tillering 650 mL-825 mL/ha post-tillering	<p>Rate Selection Use higher rates for advanced weed growth or when treating under cold/overcast conditions. Cultivation or planting may proceed from 1 hour of daylight after application to seedling annual weeds if a satisfactory seedbed can be created for crop germination and seedling establishment.</p> <p>Bentgrass Use a rate of 1.7 L/ha. Apply in late Spring following initiation of seed-head emergence. Follow up with full disturbance with a tyned implement 10-21 days after spraying.</p> <p>Silvergrass When treating dense infestations of Silvergrass, use higher rate, add Wetter TX and use water volumes of 70 L/ha or more to improve coverage.</p> <p>Perennial Weeds KELPIE PROMIX 550 GLY Herbicide will provide seasonal control and reduction in plant numbers. Control of Skeleton Weed requires addition of full soil disturbance at planting.</p> <p>Annual Ryegrass Glyphosate resistant biotypes have been detected in Australia. If glyphosate resistant weeds are known to be present, apply an additional method of control.</p> <p>In Tasmania, for perennial weeds, use 980 mL-2 L/ha.</p>
	Annual Phalaris, Annual Ryegrass, Silvergrass, Winter Grass	650 mL-825 mL/ha pre-tillering 825 mL-980 mL/ha post-tillering	
	Calomba Daisy, Capeweed, Doublegee/Spiny Emex, Fumitory, Volunteer Lupins, Volunteer Peas	330-650 mL/ha less than 8 cm dia/height 650-980 mL/ha greater than 8 cm dia/height	
	Amsinckia, Dock (seedling), Paterson's Curse, Saffron Thistle, Scotch Thistle, Spear Thistle, Variegated Thistle, Wild Turnip	650-825 mL/ha less than 12 cm dia/height 825 mL-980 mL/ha greater than 12 cm dia/height	
	Bent Grass, Perennial Phalaris, Skeleton weed, Sorrel, Sub. Clover	980 mL- 2L/ha	

SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
SOUTHERN AUSTRALIA MINIMAL SOIL DISTURBANCE To commence a fallow OR Prior to planting a crop or pasture with an implement that gives minimal soil disturbance or prior to surface seeding of pastures.	Barley Grass, Canary Grass, Wild Oats, Volunteer Cereals	650 mL—980 mL/ha	Rate Selection Use the lower rate on young weeds; increase to the higher rate where grasses reach full tillering or where broadleaf weeds commence stem elongation or budding. Use higher rates in Spring and under cold conditions. In Tasmania use 1—2 L/ha with the higher rate for control of perennial weeds. Pasture or Crop Establishment DO NOT sow into excessive trash. Excessive plant residues may be removed by grazing after treatment. Planting may proceed from 1 hour of daylight after application to seedling annual weeds if a satisfactory seedbed can be created for crop germination and seedling establishment. Aerial (or Surface) Seeding Delay seeding until trash level is reduced to allow for satisfactory placement of broadcast seed on the soil surface. Bathurst Burr For mature weeds use the higher rate. Bentgrass Use a rate of 1.7 L/ha. Apply in late Spring following initiation of seed-head emergence. Follow up with full disturbance with a tined implement 10—21 days after spraying. Couch, Kikuyu, Paspalum Use the higher rate on dense infestations. Apply sequential treatments during Summer and Autumn. Repeat application will be required for full control. For improved control, use in conjunction with cultivation. Kikuyu, Paspalum Use the low rate for suppression, the high rate for control. Dock, Flatweed Use the maximum rate for full control. Hoary cress Use at a rate of 1 L/ha. Treat from late rosette to early flowering. Silvergrass When treating dense infestations of Silvergrass, add Octyl Phenol Ethoxylate non-ionic surfactant and use water volumes of 70 L/ha or more and small droplets to improve coverage. Soursob Use at a rate of 1 L/ha. Treat at tuber exhaustion.
	Annual Ryegrass, Brome Grass, Capeweed, Hoary Cress, Paterson's Curse, Saffron Thistle, Scotch Thistle, Silvergrass, Soursob, Spear Thistle, Variegated Thistle, Wild Mustard, Wild Radish, Wild Turnip, Winter Grass	980 mL – 1.25 L/ha	
	Bathurst Burr, Bentgrass, Couch, Dock, Erodium, Flatweed, Kikuyu, Plantain, Paspalum, Perennial Phalaris, Sorrel, Sub. Clover, Yorkshire Fog	1.2 – 2.0 L/ha	
	Poa Tussock	2.0—2.65 L/ha	
Seed-Head Suppression	Bent Grass	235 – 415 mL/ha	Apply treatments late October to late November, before seedheads have emerged. Add Wetter TX. Use the higher rate where growth is excessive. Graze hard after spraying.

SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
SOUTHERN AUSTRALIA NSW, ACT, VIC, TAS only For control/ suppression prior to establishing crops or improved pasture species	Serrated Tussock	2.65 - 4.0 L/ha	Apply to actively growing and stress free plants. Best results May to October. Application Boom spray volume of 70 L/ha or more is recommended to improve plant coverage. Also see AERIAL EQUIPMENT . Surfactants Addition of 200 mL of Octyl Phenol Ethoxylate non-ionic surfactant to 100 L of spraying solution may improve control of Serrated Tussock. Site Preparation <i>Burning</i> of Serrated Tussock 10-12 months before spraying or <i>slashing / heavy grazing</i> (cell grazing) 2 weeks before spraying is essential for good results. (Note: Serrated Tussock is almost indigestible and prolonged exposure can lead to starvation and death of stock.) Rates Use lower rate on Serrated Tussock regrowth after burning (no residual dead foliage). Use higher rate on Serrated Tussock that has been slashed or grazed (may contain some residual dead foliage).
	For prevention of seed head emergence and seed formation	490 - 825 mL/ha	Apply to actively growing and stress free plants. Best results obtained during mid September – mid October. Apply prior to any seed head emergence. Also see AERIAL EQUIPMENT . Surfactants Addition of 200 mL of Octyl Phenol Ethoxylate non-ionic surfactant to 100 L of spraying solution may improve results. Rates The lower rates will be less damaging to desirable pasture species. If seed head emergence is imminent, then higher rates will give better results.
NORTHERN AUSTRALIA In fallow or prior to planting a crop. COTTON: Shielded Sprayers	Paradoxa Grass, Volunteer Cereals, Wild Oats	330 - 650 mL/ha	Rate Selection Use the lower rates on young weeds and increase to the higher rate where weeds are dense or well developed. Dense infestations of some weeds e.g. Barnyard Grass, Liverseed (<i>Urochloa</i>) Grass may need follow up treatments for complete control. Tank mixtures Read and follow all label directions, restraints, plant-back and withholding periods, regional use restrictions and safety directions for the tank mix products. Tank mixes with atrazine may give unacceptable knockdown control of certain weeds. DO NOT apply the tank-mix for control of Barnyard Grass, Liverseed Grass or Milk Thistle. Ammonium sulphate may enhance knockdown weed control where tank mixtures of atrazine are used. Shielded Sprayers Apply KELPIE® PROMIX 550 GLY HERBICIDE to weeds growing between crop rows using a shielded sprayer. DO NOT apply in Cotton less than 20 cm high. Do not allow spray or spray drift to contact any part of the cotton plant as severe injury may result. Pasture or crop establishment DO NOT sow into excessive trash. Excessive plant residues may be removed by grazing after treatment. Cultivation or planting may proceed from 1 hour of sunlight after application to seedling annual weeds if a satisfactory seedbed can be created for crop germination and seedling establishment.
	African Turnip Weed, Black Pigweed, Boggabri Weed, Caltrop (Yellow Vine), Deadnettle, Mintweed, Milk (Sow) Thistle, Stinkgrass (Lovegrass), Sweet Summer Grass, Variegated Thistle, Volunteer Sorghum	490 - 650 mL/ha up to 5 true leaves or 3 cm in diameter/height 650 mL - 1.3 L/ha greater than 5 true leaves or 3 cm in diameter/height	
	Annual Ground Cherry, Barnyard Grass, Bathurst Burr, Bladder Ketmia, Button Grass, Camel (Afgan) Melon, Caustic Weed, Columbus Grass, Liverseed Grass, Mexican Poppy, Native Millet, New Zealand Spinach, Noogoora Burr, Pigweed (up to 25 cm diameter), Spear Thistle, Stinking Goosefoot, Thornapple (Datura), Turnip Weed, Wild/Prickly Lettuce, Wireweed	650 mL - 1.3 L/ha	
	Prickly Paddy Melon	630 mL - 1.3 L/ha plus 80 mL/ha triclopyr	

SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
NORTHERN AUSTRALIA In fallow or prior to planting a crop. COTTON: Shielded Sprayers (continued)	Climbing Buckwheat (less than 12 leaves), Couch, Johnson Grass	1.3 – 2.0 L/ha	Use the higher rate on plants at the flowering/seedhead stage. For Johnson Grass apply to plants with a minimum of 30 cm new growth. For long term control of Couch and Johnson Grass, repeat applications will be required.
	Nutgrass (<i>Cyperus rotundus</i>)	2 L/ha followed by 2 L/ha	Make first application to actively growing plants when the majority of plants have reached at least the 6-8 leaf stage but preferably later. Allow for maximum re-emergence before retreating.
Sugar cane: Inter-row spraying	Annual and Perennial Grasses and Broadleaf Weeds	1.15L- 4.9L/ha	Apply to weeds growing between crop rows using a ground based hooded and shielded sprayer. Apply at early growth stage of crop before formation of the cane. Apply no more than 3 applications, to a maximum of 11.4 L/ha per crop. DO NOT allow spray or spray drift to contact any part of the crop as severe injury may result.
SUGAR CANE Ratoon spray out QLD, NSW only	Sugar cane ratoon regrowth	4 – 5.9 L/ha	Apply under good growing conditions only to actively growing ratoons 60-120 cm tall. DO NOT apply if plants are under stress from low moisture or water logging. Use the lower rate for suppression or where cultivation is to follow. Use higher rate for control.

PRE- AND POST-HARVEST USES

SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Sorghum Control	Grain Sorghum (pre-harvest)	980mL- 2L L	DO NOT apply if crop is under stress from low moisture, frost, cold or waterlogging. Apply when grain moisture is less than 25%. Use the higher rate where the crop has produced significant number of late tillers or where following crops will be established without further treatment. DO NOT apply to crops intended for seed production. Treatment may increase potential for crop lodging.
	Grain Sorghum (post-harvest)	850 mL – 1.4 L	Slashed/grazed stubble. Apply when fresh regrowth is at least 20 cm high. Use the higher rate on standing stubble or where regrowth from slashed Sorghum has advanced beyond 50 cm in height.
Cotton Pre-Harvest	Bathurst Burr, Noogoora Burr, Winter Annual Weeds	825mL- 1.65L	Treatments may be applied alone or in a tank mix with Dropp*. Apply when 60% of bolls are open. When tank mixed with conditioner/defoliant treatments, a slightly higher proportion of cotton leaf may be retained particularly where higher rates are used and conditions are unfavourable for defoliation.
PRE-HARVEST APPLICATION to reduce viable seed set of weeds in: Field Peas (<i>Pisum sativum</i>), Faba Beans (<i>Vicia faba</i>)	Annual Ryegrass (<i>Lolium rigidum</i>)	350 – 700mL	Use lower rate if Ryegrass is flowering and higher rate if Ryegrass is at milky dough stage. Application should be made at or after crop maturity. Application before this time may significantly reduce yields (in practice losses in excess of 25% can occur). Apply when the average seed moisture content is below 30%. For Faba Beans, this is indicated by the pods going black, and for field peas by the pods going yellow. DO NOT harvest within 7 days after application. DO NOT use on crops intended for seed or sprouting. Glyphosate resistant biotypes have been detected in Australia. If glyphosate resistant weeds are known to be present, apply an additional method of control.
PRE-HARVEST APPLICATION as Harvest Aid and Weed Control: Wheat (<i>Triticum aestivum</i>)	Annual Weeds	900mL – 3.5L	Apply to mature crop from late dough stage (28% moisture) onwards. The higher rate will be required when crops are heavy and leaf shading effects may occur. DO NOT harvest within 7 days after application. DO NOT use on crops intended for seed or sprouting. Where wheat is grown in rotation with any herbicide tolerant crop, management should be consistent with implementation of any management plan for herbicide tolerant crops.
PRE-HARVEST APPLICATION To desiccate a crop as a harvest aid and weed control Adzuki Beans, Chickpeas, Cowpeas, Faba Beans, Field Peas, Lentils, Mungbeans, Soybean (Application to crops intended for seed production or for sprouting may reduce germination percentage to commercially unacceptable levels)	Annual Weeds	660 mL – 1.8L	Apply with boom or by air. Use higher rates where crops or weeds are dense and where faster desiccation is required. Application should be made at or after crop maturity. Chickpeas and Lentils: Apply when physiologically mature and less than 15% green pods. Soybean: Apply only after seed pod have lost all green colour and 80-90% of leaves have dropped. Mungbeans/Adzuki Beans and Cowpea: Apply to mature crops when pods are brown/black. Field Peas: Apply when seeds turn yellow and average seed moisture content is below 30%. Faba Beans: Apply when pods turn black and average seed moisture content is below 30%. DO NOT harvest within 7 days of application. Speed of crop desiccation is dependent on crop stage, growing conditions and weather conditions during and after application.

SITUATION	CRITICAL COMMENTS READ APPLICATION CHECKLIST BEFORE USING. See Annual and Perennial Weeds section below for most appropriate rate.
GENERAL WEED CONTROL FOR GENERAL WEED CONTROL IN DOMESTIC AREAS (HOME GARDENS), COMMERCIAL, INDUSTRIAL AND PUBLIC SERVICE AREAS, AGRICULTURAL BUILDINGS AND OTHER FARM SITUATIONS. FOR SPECIFIC WEEDS REFER TO THE WEEDS CONTROLLED TABLE .	For the control of many grasses and broadleaf weeds. RATE: 7 mL/L of water. Apply when weeds are actively growing. Apply to ensure complete and uniform wetting of foliage. Visible symptoms may take from 3 to 7 days to develop.
AGRICULTURAL AREAS	KELPIE® PROMIX 550 GLY HERBICIDE may be used for control of Annual and Perennial Weeds as directed, in agricultural land prior to sowing of any edible or non-edible crop, but not prior to transplanting tomato seedlings.
DRY DRAINS AND CHANNELS ONLY	DO NOT apply to weeds growing in or over water. DO NOT spray across open bodies of water, and DO NOT allow spray to enter the water. DO NOT allow water to return to dry channels and drains within 4 days of application.
FORESTS	KELPIE® PROMIX 550 GLY HERBICIDE may be used prior to establishment of nurseries, for site preparation prior to planting and amongst established trees using a directed or shielded spray, or using selective wiper equipment. DO NOT allow wiper surface to contact any part of the tree. DO NOT allow spray or spray drift to contact foliage or green bark of desirable trees, since severe injury may result.
NON-AGRICULTURAL AREAS AROUND BUILDINGS, COMMERCIAL AND INDUSTRIAL AREAS, DOMESTIC AND PUBLIC SERVICE AREAS, RIGHT-OF-WAYS.	KELPIE® PROMIX 550 GLY HERBICIDE does not provide residual weed control. For residual control of annual weeds, KELPIE® PROMIX 550 GLY HERBICIDE may be tank mixed with certain residual herbicides. See TANK MIXTURES/COMPATIBILITY .
TREE AND VINE CROPS AVOCADO, BANANA, BLUEBERRIES, CITRUS FRUITS, CUSTARD APPLES, DUBOISIA, FIGS-DESSERT, GUAVA, HOPS, KIWIFRUIT, LITCHI, MANGO, MONSTERA-FRUIT, NUTS (INCLUDING ALMOND, PECAN, MACADAMIA, PISTACHIO AND WALNUT), OLIVES, PAWPAW, PERSIMMONS, POME FRUIT, RASPBERRIES, STONE FRUIT, TEA, VINEYARDS.	Apply as a directed or shielded spray or using wiper equipment. DO NOT apply as a spray near trees or vines less than 3 years old unless they are effectively shielded from spray and spray drift. DO NOT allow wiper surface to contact any part of the tree, vine or palm. Citrus Fruit, Nuts, Olives, Pome Fruit & Vineyards DO NOT allow spray or spray drift to contact green bark or stems, canes, laterals, suckers, fresh wounds foliage or fruit. Hops Apply in Winter, prior to crop emerging from dormancy. Tea Apply a maximum of 2.65 L/ha by shielded boom or directed off-centre nozzle or 340 mL/100L by directed hand-gun or knapsack to avoid application to the crop. All other crops DO NOT allow spray or spray drift to contact any part of the plant including the trunk. CAUTION Where split bark on Kiwifruit and green stems on Pawpaw occur, extreme care is required.

WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
ANNUAL WEEDS Amaranth, Bathurst Burr, Barley Grass, Brome Grass, Barnyard Grass, Caltrop, Canary Grass, Capeweed, Chickweed, Cobblers Peg, Deadnettle, Doublegee, Fumitory, Ground Cherry, Hedge Mustard, Lesser Swinecress, Liverseed Grass, Mintweed, Noogoora Burr, Paradoxa Grass, Paterson's Curse, Pigweed, Potato Weed, Ryegrass, Saffron Thistle, Silvergrass, Sow Thistle, Spear Thistle, Spiny Burrgrass, Spurge, Sub Clover, Thornapple, Wild Mustard, Wild Oats, Wild Turnip, Winter Grass, Variegated Thistle, Volunteer Cereals	Boom: 1.3 - 2 L/ha Handgun: 325 - 470 mL per 100 L Knapsack: 50–70 mL per 15 L	Apply to weeds whenever they are not subject to stress due to drought or frost. Use higher rate on weeds over 15 cm in height or diameter or where dense weed cover limits spray coverage. Use higher spot spraying rate when applying less than 5 L spray per 100 m ² . KELPIE® PROMIX 550 GLY HERBICIDE does not provide residual weed control. Repeat treatments may be necessary to control later germinating weeds. For residual control of annual weeds, KELPIE® PROMIX 550 GLY HERBICIDE may be tank-mixed with certain residual herbicides. See TANK MIXTURES in the GENERAL INSTRUCTIONS for directions. DO NOT use an atrazine tank-mix for control of Barnyard Grass or Liverseed Grass.
PERENNIAL WEEDS Artichoke thistle, African Lovegrass, Bent Grass, Carpet Grass, Cocksfoot, Flatweed, Johnson Grass, Kangaroo Grass, Kikuyu, Nutgrass (<i>Cyperus rotundus</i>), Paspalum, Phalaris, Plantains, Poa Tussock, Prairie Grass, Qld Blue Grass, Red-leg Grass, Rhodes Grass, Rope Twitch, Sorrel, Soursob, Yorkshire Fog	Boom: 2 - 4 L/ha Handgun: 460 - 650 mL per 100 L Knapsack: 70 - 100 mL per 15 L	Control of established perennials is best obtained when plants are at the seedhead stage. In general best control of Winter growing perennials is obtained with application during Winter-Spring. Best control of Summer growing perennials is obtained with application late Summer and Autumn. For Nutgrass in cultivated situations apply sequential low rate treatments when Nutgrass has a minimum of 6-8 leaves. Use the higher rate in uncultivated situations. For Rhodes Grass, Rope Twitch, Prairie Grass, Qld Blue Grass, Johnson Grass, Kangaroo Grass, Kikuyu, Redleg Grass, Paspalum and Sorrel, use the higher rates only.
Blady Grass, Bracken, Couch, Guinea Grass, Paragrass*, Silverleaf nightshade, Water Couch ^a ^a Use on Dry Drains and Channels ONLY (see Use Situations Critical Comments above)	Boom: 5.7 L/ha Handgun: 850 mL or 1.33 L per 100 L Knapsack: 130 or 200 mL per 15 L	For Bracken add Pulse* at 200 mL/100 L spray mix. Best control of Couch in WA and SA is obtained with Spring treatment. Most effective control of Couch in Eastern states is obtained with Summer and Autumn treatments. In cultivated situations use sequential treatments of 1.9-4.3 L/ha for control. Only use higher rates for handgun and knapsack for Silverleaf Nightshade.
WOODY WEEDS Bamboo, Bitou Bush, Boneseed, Boxthorn, Crofton Weed, Gorse, Groundsel Bush, Lantana, Mistflower	Handgun: 330 – 650 mL per 100 L Knapsack: 50 – 100 mL per 15 L	Apply to actively growing plants. DO NOT apply to drought stressed plants. Further treatment may be necessary to restrict seedling re-establishment. Bamboo: Apply when foliage/regrowth is 1 – 2 m tall. Use higher rate only. Bitou Bush / Boneseed: Apply higher rate on bushes > 1.5 m. Best results are achieved when treated at peak flower during Winter. Boxthorn: Minimum rate is 470 mL for handgun and 70 mL for knapsack. Groundsel Bush: Apply higher rate on bushes > 2m. DO NOT apply in Winter. Minimum rate is 460 mL for handgun and 70 mL for knapsack. Gorse: Always add Pulse* at 200 mL/100 L of spray mix, use higher rate only. Lantana: Use higher rate only. Addition of Pulse* (200 mL/100 L) may improve control. Boxthorn, Gorse, Lantana: Removal of bushes (after complete brownout), pasture improvement or further treatment are recommended to control seedlings and/or regrowth.

WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Blackberry, Chinese Scrub, Eucalyptus spp. (seedlings < 2m), Hawthorn, Pampas Grass, Sifton Bush, Sweet Briar, Willow (<2 m)	Handgun: 650 – 850 mL per 100 L Knapsack: 100 – 140 mL per 15 L	<p>Apply to actively growing plants. Burning (after complete brownout), pasture improvement or further treatment are recommended to control seedlings and/or regrowth.</p> <p>Blackberry: Apply from flowering to leaf fall, use higher rates on old dense infestations >2 m high, in Tasmania, DO NOT treat bushes bearing mature fruit.</p> <p>Chinese Scrub: Use higher rates on bushes > 1m.</p> <p>Eucalyptus spp.: Add Pulse* at 200 mL/100 L of spray mix.</p> <p>Hawthorn: Apply from flowering to leaf fall, use higher rates on bushes > 2m .</p> <p>Pampas Grass: Allow regrowth to reach 1 m, best results apply after flowering.</p> <p>Sifton Bush: Use higher rates on bushes > 1m.</p> <p>Sweet Briar: Apply from late flowering to leaf fall, use 1 – 1.33 L/100 L, and 150 – 200 mL/15 L.</p> <p>Use higher rates on bushes > 1.5 m.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

Product Information

KELPIE® PROMIX 550 GLY HERBICIDE is a non-volatile, non-selective, water soluble liquid herbicide for the control of annual and perennial grasses and broadleaf weeds in a wide range of agricultural and non-agricultural use situations. KELPIE® PROMIX 550 GLY HERBICIDE may be used for weed control on agricultural land prior to planting any edible or non-edible crop but not prior to transplanting tomatoes. When applying this product prior to transplanting crops into plastic mulch, care must be taken to remove residues of this product from the plastic prior to transplanting. Residues can be removed by 2 cm of natural rainfall or by applying water via a sprinkler irrigation system.

KELPIE® PROMIX 550 GLY HERBICIDE is absorbed by plant foliage and green stems. It is inactivated on clay and organic matter in soil and does not provide residual weed control. KELPIE® PROMIX 550 GLY HERBICIDE moves throughout the plant from the point of contact to and into the root system. Initial visible effects on annual weeds take 3-7 days but may not be noticeable for 2 to 3 weeks under cool cloudy conditions or on some perennial weeds.

Crop Establishment

KELPIE® PROMIX 550 GLY HERBICIDE is recommended for control of emerged weeds prior to crop establishment. Cultivation and/or planting operations which provide conditions suitable for crop emergence and establishment are required following herbicide application. Where heavy weed growth is present or soil conditions are unsuitable, planting should be delayed to allow for decay of weeds and/or development of more favourable soil conditions for the formation of a suitable seedbed. Incorporation of green or decaying vegetation may retard crop emergence under cold, wet conditions. Vegetation may be reduced by grazing and weed decay may be assisted by cultivation to leave trash on the surface.

Mixing

KELPIE® PROMIX 550 GLY HERBICIDE mixes readily with water. Reduced results may occur if water is used containing suspended clay or organic matter e.g. from dams, streams or irrigation channels, or high levels of calcium, magnesium or bicarbonate ions.

Do not mix, store or apply this product in galvanised steel or unlined steel containers or spray tanks, since a highly flammable gas mixture may be formed. Use stainless steel, aluminium, brass, copper, fibreglass, plastic or plastic lined containers or spray tanks. Spray tanks, pumps, lines and nozzles should be thoroughly cleaned with clean water following application. Ensure that the spray tank is free of any residue of other spray solutions prior to mixing. Use spray solutions promptly as a gradual loss of activity may occur over a period of days following spray preparation.

Mixing Instructions

1. Fill the spray tank 1/3 to ½ full with clean water and start agitation.
2. If adding ammonium sulphate use a 2% v/v and mix thoroughly.
3. If tank-mixing, add recommended herbicide/insecticide/additive to the spray tank and mix thoroughly.
4. Add KELPIE® PROMIX 550 GLY HERBICIDE and the remaining water. Mix thoroughly.
5. Add octyl phenol ethoxylate non-ionic surfactant, if required, near the end of the filling process.
6. Always maintain adequate agitation during application and use the tank mix promptly.

Clean all equipment after use by washing thoroughly with water.

Adjuvants – 1040 g/L Octyl phenol ethoxylate non-ionic surfactant

Octyl phenol ethoxylate non-ionic surfactant is recommended for the control of silvergrass and annual ryegrass in later Winter and Spring. Octyl phenol ethoxylate non-ionic surfactant is not a general purpose surfactant and should only be used where recommended.

Rate: 200 mL/100 L spray solution.

Adjuvants – Ammonium sulphate

Ammonium sulphate may be used as an adjuvant to alleviate the adverse effects of high levels of calcium, magnesium and bicarbonate ions in water.

Rate: 2 L/100 L spray solution.

Application

Boom Equipment: For boom application, a spray volume of 80 L/ha or less is recommended for optimum performance. Nozzles and pressure settings should be selected to deliver a MEDIUM or MEDIUM-COARSE size droplet at the target (BCPC-British Crop Protection Council Standard). The use of nozzles and/or pressure settings that produce VERY FINE or FINE droplets should be avoided as these are prone to loss or drift. In multiple product tank mixes, a minimum water volume of 50 L/ha is recommended and local advice should be sought. Correct mixing order is important as is good in-tank agitation when application is occurring.

For shielded applications a spray volume of 80 L/sprayed ha is recommended using nozzle types and pressure settings to deliver a COARSE (BCPC) size droplet at the target. Crop damage may result if spray drift occurs through incorrect nozzle and/or pressure selection, inadequate shielding and/or wind strength, high evaporation rates or excessive ground speed.

Wiper Equipment: Wiper equipment (e.g. Ropewick, canvas, felt or carpet applicators) may be used to apply KELPIE® PROMIX 550 GLY HERBICIDE. Avoid contact with desirable vegetation. Operate wiper equipment a minimum of 10 cm above the crop or pasture. Weeds should be at least 15 cm above the crop or pasture at time of application. Speed of travel should be no greater than 8 km/h. Best results are achieved at lower speeds and where two applications are made in opposite directions (double pass). Where weeds are of variable height, or occur in dense infestations or clumps, some plants may not be contacted by the herbicide solution. In these cases repeat treatment may be necessary. **RATE:** Mix 700 mL KELPIE® PROMIX 550 GLY HERBICIDE with 2.3 litres clean water. Adjust flow rate to suit equipment.

Aerial Equipment: KELPIE® PROMIX 550 GLY HERBICIDE may be applied by aircraft for control of weeds in forests, cropland or pasture prior to establishment of crops, new pastures or new forest plantings and for pre-harvest applications, up to a maximum rate of 2.651 L/ha where specified by this label. DO NOT apply treatments by aircraft in situations where drift onto sensitive crops and pastures is likely to occur.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM droplet size of 250-350 microns. In multiple product tank mixes a minimum water volume of 50 L/ha is recommended and local advice should be sought. Correct mixing order is important. Swath width should be set to take into account aircraft type, wind conditions and target height. Swath width will need to be reduced to avoid stripping under light wind conditions and/or application to tall, dense targets, e.g. pre-harvest application, treatments in heavy crop stubble. Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove herbicide residues.

Application on hilly terrain

Increase water volume to 30-80 L/ha and increase droplet diameter of output to at least 300 microns to optimise deposition of spray output onto weeds.

Air temperature and relative humidity

DO NOT apply KELPIE® PROMIX 550 GLY HERBICIDE by aircraft at temperatures above 30°C. Increase spray output to at least 30 L/ha when temperatures rise above 25°C. Avoid application when relative humidity falls below 35%.

AVOID DRIFT

DO NOT apply treatments with spraying equipment or under weather conditions which are likely to cause spray drift onto nearby susceptible crops, pastures or other sensitive plants. DO NOT apply treatments under very light winds (less than 4 km/h) or inversion conditions or where wind speeds exceed 12 km/h.

Application checklist

- Do not treat weeds under poor growing conditions due to moisture stress, waterlogging, severe frosting, insect damage etc. Reduced performance may also occur where weeds are covered with dust or silt.
- Do not add surfactants, adjuvants or other pesticides except as specifically directed on this label.
- Rain within 1 hour of application which causes run-off may require re-treatment. Rainfastness is reduced if weeds are not actively growing, under stress or conditions of low light intensity/darkness. The addition of Octyl Phenol Ethoxylate non-ionic surfactant may improve rainfastness on winter annual weeds.
- A withholding period for grazing is not required. However, it is recommended that grazing of treated plants be delayed to ensure herbicide uptake. Certain plants such as Soursob, Variegated Thistle, Sorghum and Johnson Grass may be naturally toxic to stock when eaten in large quantities under certain conditions. Where plants are known to be toxic, grazing should be delayed until complete desiccation of treated plants has occurred.
- Apply treatments to weeds which have at least one true leaf (broadleaf weeds) or two leaves (grasses) to provide an adequate surface area for herbicide uptake.
- If heavy grazing has occurred, allow regrowth to 6-8 cm before spraying and use the higher rates recommended.