

On the global linear Zarankiewicz problem

(joint with Aris Papadopoulos)

Pantelis Eleftheriou

University of Leeds

Logic Seminar, Manchester, December 10, 2025

Theme

- ▶ recognise definable algebraic structure from combinatorial data

Theme

- ▶ recognise definable algebraic structure from combinatorial data
- ▶ linear vs non-linear structure $\mathcal{M} = \langle M, +, \dots \rangle$

Theme

- ▶ recognise definable algebraic structure from combinatorial data
- ▶ linear vs non-linear structure $\mathcal{M} = \langle M, +, \dots \rangle$
- ▶ $f : X \subseteq M^m \rightarrow M^k$ is *linear* (affine) if for every $x, y, x + t, y + t \in X$,

$$f(x + t) - f(x) = f(y + t) - f(y).$$

Theme

- ▶ recognise definable algebraic structure from combinatorial data
- ▶ linear vs non-linear structure $\mathcal{M} = \langle M, +, \dots \rangle$
- ▶ $f : X \subseteq M^m \rightarrow M^k$ is *linear* (affine) if for every $x, y, x + t, y + t \in X$,

$$f(x + t) - f(x) = f(y + t) - f(y).$$

- ▶ In $\langle \mathbb{R}, <, + \rangle$, $\langle \mathbb{Z}, <, + \rangle$ every definable function is piecewise linear.

Theme

- ▶ recognise definable algebraic structure from combinatorial data
- ▶ linear vs non-linear structure $\mathcal{M} = \langle M, +, \dots \rangle$
- ▶ $f : X \subseteq M^m \rightarrow M^k$ is *linear* (affine) if for every $x, y, x + t, y + t \in X$,

$$f(x + t) - f(x) = f(y + t) - f(y).$$

- ▶ In $\langle \mathbb{R}, <, + \rangle$, $\langle \mathbb{Z}, <, + \rangle$ every definable function is piecewise linear.
- ▶ In $\langle \mathbb{R}, <, +, \cdot \upharpoonright_{(0,1)^2} \rangle$, $\langle \mathbb{R}, <, +, \mathbb{Z} \rangle$, $\langle \mathbb{R}, <, +, 2^{\mathbb{Z}} \rangle$ not, but still some linearity may be present

Global Zarankiewicz Problem

Global Zarankiewicz Problem

- ▶ M set, $d_1, \dots, d_r \in \mathbb{N}^{>0}$

Global Zarankiewicz Problem

- ▶ M set, $d_1, \dots, d_r \in \mathbb{N}^{>0}$
- ▶ $E \subseteq M^{d_1} \times \dots \times M^{d_r} = M^d$ relation

Global Zarankiewicz Problem

- ▶ M set, $d_1, \dots, d_r \in \mathbb{N}^{>0}$
- ▶ $E \subseteq M^{d_1} \times \dots \times M^{d_r} = M^d$ relation
- ▶ B grid = $B_1 \times \dots \times B_r$, where $B_i \subseteq M^{d_i}$

Global Zarankiewicz Problem

- ▶ M set, $d_1, \dots, d_r \in \mathbb{N}^{>0}$
- ▶ $E \subseteq M^{d_1} \times \dots \times M^{d_r} = M^d$ relation
- ▶ B grid = $B_1 \times \dots \times B_r$, where $B_i \subseteq M^{d_i}$
- ▶ If B finite, $n = n_B = \max_i |B_i|$

Global Zarankiewicz Problem

- ▶ M set, $d_1, \dots, d_r \in \mathbb{N}^{>0}$
- ▶ $E \subseteq M^{d_1} \times \dots \times M^{d_r} = M^d$ relation
- ▶ B grid = $B_1 \times \dots \times B_r$, where $B_i \subseteq M^{d_i}$
- ▶ If B finite, $n = n_B = \max_i |B_i|$, so $|E \cap B| \leq n^r$

Global Zarankiewicz Problem

- ▶ M set, $d_1, \dots, d_r \in \mathbb{N}^{>0}$
- ▶ $E \subseteq M^{d_1} \times \dots \times M^{d_r} = M^d$ relation
- ▶ B grid = $B_1 \times \dots \times B_r$, where $B_i \subseteq M^{d_i}$
- ▶ If B finite, $n = n_B = \max_i |B_i|$, so $|E \cap B| \leq n^r$
- ▶ Let $k \in \mathbb{N}$. Then E is k -free if it contains no k -grid B (that is, $B = B_1 \times \dots \times B_r$ with each $|B_i| = k$).

Global Zarankiewicz Problem

- ▶ M set, $d_1, \dots, d_r \in \mathbb{N}^{>0}$
- ▶ $E \subseteq M^{d_1} \times \dots \times M^{d_r} = M^d$ relation
- ▶ B grid = $B_1 \times \dots \times B_r$, where $B_i \subseteq M^{d_i}$
- ▶ If B finite, $n = n_B = \max_i |B_i|$, so $|E \cap B| \leq n^r$
- ▶ Let $k \in \mathbb{N}$. Then E is k -free if it contains no k -grid B (that is, $B = B_1 \times \dots \times B_r$ with each $|B_i| = k$).

Global Zarankiewicz (1951): E is k -free $\Rightarrow |E \cap B| = O(n^?)$

$$E \text{ is } k\text{-free} \Rightarrow |E \cap B| = O(n^?)$$

E is k -free $\Rightarrow |E \cap B| = O(n^?)$

- Kővári–Sós–Turán (1954): ($r = 2$), $O(n^{2-\frac{1}{k}})$

E is k -free $\Rightarrow |E \cap B| = O(n^r)$

- Kővári–Sós–Turán (1954): ($r = 2$), $O(n^{2-\frac{1}{k}})$, sharp $k = 2, 3$.

E is k -free $\Rightarrow |E \cap B| = O(n^?)$

- Kővári–Sós–Turán (1954): ($r = 2$), $O(n^{2-\frac{1}{k}})$, sharp $k = 2, 3$.
- Erdős (1964): $O(n^{r-\frac{1}{k^{r-1}}})$

E is k -free $\Rightarrow |E \cap B| = O(n^?)$

- Kővári–Sós–Turán (1954): ($r = 2$), $O(n^{2-\frac{1}{k}})$, sharp $k = 2, 3$.
- Erdős (1964): $O(n^{r-\frac{1}{k^{r-1}}})$
- Bohman-Keevash (2010): ($r = 2$) lower bounds (complicated)

E is k -free $\Rightarrow |E \cap B| = O(n^?)$

- Kővári–Sós–Turán (1954): ($r = 2$), $O(n^{2-\frac{1}{k}})$, sharp $k = 2, 3$.
- Erdős (1964): $O(n^{r-\frac{1}{k^{r-1}}})$
- Bohman-Keevash (2010): ($r = 2$) lower bounds (complicated)
- Szemerédi-Trotter (1983): $E \subseteq \mathbb{R}^2 \times \mathbb{R}^2$ point-line incidence, $O(n^{4/3})$, sharp

E is k -free $\Rightarrow |E \cap B| = O(n^?)$

- Kővári–Sós–Turán (1954): ($r = 2$), $O(n^{2-\frac{1}{k}})$, sharp $k = 2, 3$.
- Erdős (1964): $O(n^{r-\frac{1}{k^{r-1}}})$
- Bohman-Keevash (2010): ($r = 2$) lower bounds (complicated)
- Szemerédi-Trotter (1983): $E \subseteq \mathbb{R}^2 \times \mathbb{R}^2$ point-line incidence, $O(n^{4/3})$, sharp
- Fox-Pach-Sheffer-Suk-Zahl (2017): ($r = 2$) E semialgebraic, $O(n^{4/3})$ ($d_1 = d_2 = 2$)

E is k -free $\Rightarrow |E \cap B| = O(n^?)$

- Kővári–Sós–Turán (1954): ($r = 2$), $O(n^{2-\frac{1}{k}})$, sharp $k = 2, 3$.
- Erdős (1964): $O(n^{r-\frac{1}{k^{r-1}}})$
- Bohman-Keevash (2010): ($r = 2$) lower bounds (complicated)
- Szemerédi-Trotter (1983): $E \subseteq \mathbb{R}^2 \times \mathbb{R}^2$ point-line incidence, $O(n^{4/3})$, sharp
- Fox-Pach-Sheffer-Suk-Zahl (2017): ($r = 2$) E semialgebraic, $O(n^{4/3})$ ($d_1 = d_2 = 2$)
- Do (2018): ($r > 2$) E semialgebraic, complicated

E is k -free $\Rightarrow |E \cap B| = O(n^?)$

- Kővári–Sós–Turán (1954): ($r = 2$), $O(n^{2-\frac{1}{k}})$, sharp $k = 2, 3$.
- Erdős (1964): $O(n^{r-\frac{1}{k^{r-1}}})$
- Bohman-Keevash (2010): ($r = 2$) lower bounds (complicated)
- Szemerédi-Trotter (1983): $E \subseteq \mathbb{R}^2 \times \mathbb{R}^2$ point-line incidence, $O(n^{4/3})$, sharp
- Fox-Pach-Sheffer-Suk-Zahl (2017): ($r = 2$) E semialgebraic, $O(n^{4/3})$ ($d_1 = d_2 = 2$)
- Do (2018): ($r > 2$) E semialgebraic, complicated
- Basu-Raz (2017) / Chernikov-Galvin-Starchenko (2020): ($r = 2$) E o-minimal, $O(n^{4/3})$ ($d_1 = d_2 = 2$)

E is k -free $\Rightarrow |E \cap B| = O(n^?)$

- Kővári–Sós–Turán (1954): ($r = 2$), $O(n^{2-\frac{1}{k}})$, sharp $k = 2, 3$.
- Erdős (1964): $O(n^{r-\frac{1}{k^{r-1}}})$
- Bohman-Keevash (2010): ($r = 2$) lower bounds (complicated)
- Szemerédi-Trotter (1983): $E \subseteq \mathbb{R}^2 \times \mathbb{R}^2$ point-line incidence, $O(n^{4/3})$, sharp
- Fox-Pach-Sheffer-Suk-Zahl (2017): ($r = 2$) E semialgebraic, $O(n^{4/3})$ ($d_1 = d_2 = 2$)
- Do (2018): ($r > 2$) E semialgebraic, complicated
- Basu-Raz (2017) / Chernikov-Galvin-Starchenko (2020): ($r = 2$) E o-minimal, $O(n^{4/3})$ ($d_1 = d_2 = 2$)
- Tong (2025): ($r > 2$) E o-minimal, complicated

E is k -free $\Rightarrow |E \cap B| = O(n^?)$

- Kővári–Sós–Turán (1954): ($r = 2$), $O(n^{2-\frac{1}{k}})$, sharp $k = 2, 3$.
- Erdős (1964): $O(n^{r-\frac{1}{k^{r-1}}})$
- Bohman-Keevash (2010): ($r = 2$) lower bounds (complicated)
- Szemerédi-Trotter (1983): $E \subseteq \mathbb{R}^2 \times \mathbb{R}^2$ point-line incidence, $O(n^{4/3})$, sharp
- Fox-Pach-Sheffer-Suk-Zahl (2017): ($r = 2$) E semialgebraic, $O(n^{4/3})$ ($d_1 = d_2 = 2$)
- Do (2018): ($r > 2$) E semialgebraic, complicated
- Basu-Raz (2017) / Chernikov-Galvin-Starchenko (2020): ($r = 2$) E o-minimal, $O(n^{4/3})$ ($d_1 = d_2 = 2$)
- Tong (2025): ($r > 2$) E o-minimal, complicated
- Basit-Chernikov-Starchenko-Tao-Tran (2021): E semilinear, $O(n^{r-1})$

Global Linear Zarankiewicz Problem

- ▶ $M, d_1, \dots, d_r \in \mathbb{N}^{>0}$, $E \subseteq \prod_{i \in [r]} M^{d_i} = M^d$, B grid

Global Linear Zarankiewicz Problem

- ▶ $M, d_1, \dots, d_r \in \mathbb{N}^{>0}$, $E \subseteq \prod_{i \in [r]} M^{d_i} = M^d$, B grid
- ▶ \mathcal{C} class of grids

Global Linear Zarankiewicz Problem

- ▶ $M, d_1, \dots, d_r \in \mathbb{N}^{>0}$, $E \subseteq \prod_{i \in [r]} M^{d_i} = M^d$, B grid
- ▶ \mathcal{C} class of grids
- ▶ Let $k \in \mathbb{N}$. Say E is \mathcal{C} - k -free if it contains no k -grid $B \in \mathcal{C}$.

Global Linear Zarankiewicz Problem

- ▶ $M, d_1, \dots, d_r \in \mathbb{N}^{>0}$, $E \subseteq \prod_{i \in [r]} M^{d_i} = M^d$, B grid
- ▶ \mathcal{C} class of grids
- ▶ Let $k \in \mathbb{N}$. Say E is \mathcal{C} - k -free if it contains no k -grid $B \in \mathcal{C}$.
- ▶ E linear Z -bounds for \mathcal{C} if $\exists \alpha \in \mathbb{R}^{>0}$, $\forall_{fin} B \in \mathcal{C}$, $|E \cap B| \leq \alpha n^{r-1}$.

Global Linear Zarankiewicz Problem

- ▶ $M, d_1, \dots, d_r \in \mathbb{N}^{>0}$, $E \subseteq \prod_{i \in [r]} M^{d_i} = M^d$, B grid
 - ▶ \mathcal{C} class of grids
 - ▶ Let $k \in \mathbb{N}$. Say E is \mathcal{C} - k -free if it contains no k -grid $B \in \mathcal{C}$.
 - ▶ E linear Z -bounds for \mathcal{C} if $\exists \alpha \in \mathbb{R}^{>0}$, $\forall_{fin} B \in \mathcal{C}$, $|E \cap B| \leq \alpha n^{r-1}$.
- $Zar(E, \mathcal{C}) : (\exists k \in \mathbb{N}, E \text{ } \mathcal{C}\text{-}k\text{-free}) \Rightarrow E \text{ linear } Z\text{-bounds for } \mathcal{C}$

Global Linear Zarankiewicz Problem

- ▶ $M, d_1, \dots, d_r \in \mathbb{N}^{>0}$, $E \subseteq \prod_{i \in [r]} M^{d_i} = M^d$, B grid
 - ▶ \mathcal{C} class of grids
 - ▶ Let $k \in \mathbb{N}$. Say E is \mathcal{C} - k -free if it contains no k -grid $B \in \mathcal{C}$.
 - ▶ E linear Z -bounds for \mathcal{C} if $\exists \alpha \in \mathbb{R}^{>0}$, $\forall_{fin} B \in \mathcal{C}$, $|E \cap B| \leq \alpha n^{r-1}$.
- $Zar(E, \mathcal{C}) : (\exists k \in \mathbb{N}, E \text{ } \mathcal{C}\text{-}k\text{-free}) \Rightarrow E \text{ linear } Z\text{-bounds for } \mathcal{C}$
- Write $Zar_\alpha(E, \mathcal{C})$ if bounds are witnessed by α .

Global Linear Zarankiewicz Problem

- ▶ $M, d_1, \dots, d_r \in \mathbb{N}^{>0}$, $E \subseteq \prod_{i \in [r]} M^{d_i} = M^d$, B grid
- ▶ \mathcal{C} class of grids
- ▶ Let $k \in \mathbb{N}$. Say E is \mathcal{C} - k -free if it contains no k -grid $B \in \mathcal{C}$.
- ▶ E linear Z -bounds for \mathcal{C} if $\exists \alpha \in \mathbb{R}^{>0}$, $\forall_{fin} B \in \mathcal{C}$, $|E \cap B| \leq \alpha n^{r-1}$.
 $Zar(E, \mathcal{C}) : (\exists k \in \mathbb{N}, E \text{ } \mathcal{C}\text{-}k\text{-free}) \Rightarrow E \text{ linear } Z\text{-bounds for } \mathcal{C}$
Write $Zar_\alpha(E, \mathcal{C})$ if bounds are witnessed by α .
- ▶ $\mathcal{E} = \{E_b\}_{b \in I}$ a family of relations E_b

Global Linear Zarankiewicz Problem

- ▶ $M, d_1, \dots, d_r \in \mathbb{N}^{>0}$, $E \subseteq \prod_{i \in [r]} M^{d_i} = M^d$, B grid
- ▶ \mathcal{C} class of grids
- ▶ Let $k \in \mathbb{N}$. Say E is \mathcal{C} - k -free if it contains no k -grid $B \in \mathcal{C}$.
- ▶ E linear Z -bounds for \mathcal{C} if $\exists \alpha \in \mathbb{R}^{>0}$, $\forall_{fin} B \in \mathcal{C}$, $|E \cap B| \leq \alpha n^{r-1}$.
 $Zar(E, \mathcal{C}) : (\exists k \in \mathbb{N}, E \text{ } \mathcal{C}\text{-}k\text{-free}) \Rightarrow E \text{ linear } Z\text{-bounds for } \mathcal{C}$
Write $Zar_\alpha(E, \mathcal{C})$ if bounds are witnessed by α .
- ▶ $\mathcal{E} = \{E_b\}_{b \in I}$ a family of relations E_b
 $Zar(\mathcal{E}, \mathcal{C}) : \exists \alpha \in \mathbb{R}^{>0}$, $\forall b \in I$, $Zar_\alpha(E_b, \mathcal{C})$

Global Linear Zarankiewicz Problem

- ▶ $M, d_1, \dots, d_r \in \mathbb{N}^{>0}$, $E \subseteq \prod_{i \in [r]} M^{d_i} = M^d$, B grid
- ▶ \mathcal{C} class of grids
- ▶ Let $k \in \mathbb{N}$. Say E is \mathcal{C} - k -free if it contains no k -grid $B \in \mathcal{C}$.
- ▶ E linear Z -bounds for \mathcal{C} if $\exists \alpha \in \mathbb{R}^{>0}$, $\forall_{fin} B \in \mathcal{C}$, $|E \cap B| \leq \alpha n^{r-1}$.

$Zar(E, \mathcal{C}) : (\exists k \in \mathbb{N}, E \text{ } \mathcal{C}\text{-}k\text{-free}) \Rightarrow E \text{ linear } Z\text{-bounds for } \mathcal{C}$

Write $Zar_\alpha(E, \mathcal{C})$ if bounds are witnessed by α .

- ▶ $\mathcal{E} = \{E_b\}_{b \in I}$ a family of relations E_b
- $Zar(\mathcal{E}, \mathcal{C}) : \exists \alpha \in \mathbb{R}^{>0}, \forall b \in I, Zar_\alpha(E_b, \mathcal{C})$

Omit \mathcal{C} if it is the class of all grids.

Linear and Semibounded Zarankiewicz

Let $\mathcal{M} = \langle M, <, +, \dots \rangle$ be o-minimal.

Linear and Semibounded Zarankiewicz

Let $\mathcal{M} = \langle M, <, +, \dots \rangle$ be o-minimal.

Theorem 0 (Basit-Chernikov-Starchenko-Tao-Tran, 2021)

The following are equivalent:

Linear and Semibounded Zarankiewicz

Let $\mathcal{M} = \langle M, <, +, \dots \rangle$ be o-minimal.

Theorem 0 (Basit-Chernikov-Starchenko-Tao-Tran, 2021)

The following are equivalent:

- a. \mathcal{M} linear

Linear and Semibounded Zarankiewicz

Let $\mathcal{M} = \langle M, <, +, \dots \rangle$ be o-minimal.

Theorem 0 (Basit-Chernikov-Starchenko-Tao-Tran, 2021)

The following are equivalent:

- a. \mathcal{M} linear (\Leftrightarrow definable functions are piecewise linear
 \Leftrightarrow no infinite definable field)

Linear and Semibounded Zarankiewicz

Let $\mathcal{M} = \langle M, <, +, \dots \rangle$ be o-minimal.

Theorem 0 (Basit-Chernikov-Starchenko-Tao-Tran, 2021)

The following are equivalent:

- a. \mathcal{M} linear (\Leftrightarrow definable functions are piecewise linear
 \Leftrightarrow no infinite definable field)
- b. for every definable \mathcal{E} , $\text{Zar}(\mathcal{E})$ holds.

Linear and Semibounded Zarankiewicz

Let $\mathcal{M} = \langle M, <, +, \dots \rangle$ be o-minimal.

Theorem 0 (Basit-Chernikov-Starchenko-Tao-Tran, 2021)

The following are equivalent:

- a. \mathcal{M} linear (\Leftrightarrow definable functions are piecewise linear
 \Leftrightarrow no infinite definable field)
- b. for every definable \mathcal{E} , $\text{Zar}(\mathcal{E})$ holds.

Examples: $\langle \mathbb{R}, <, + \rangle$, $\mathbb{R}_{\text{vec}} = \langle \mathbb{R}, <, +, \{x \rightarrow rx\}_{r \in \mathbb{R}} \rangle$.

Linear and Semibounded Zarankiewicz

Let $\mathcal{M} = \langle M, <, +, \dots \rangle$ be o-minimal.

Theorem 0 (Basit-Chernikov-Starchenko-Tao-Tran, 2021)

The following are equivalent:

- a. \mathcal{M} linear (\Leftrightarrow definable functions are piecewise linear
 \Leftrightarrow no infinite definable field)
- b. for every definable \mathcal{E} , $\text{Zar}(\mathcal{E})$ holds.

Examples: $\langle \mathbb{R}, <, + \rangle$, $\mathbb{R}_{\text{vec}} = \langle \mathbb{R}, <, +, \{x \rightarrow rx\}_{r \in \mathbb{R}} \rangle$.

Theorem 1

The following are equivalent:

Linear and Semibounded Zarankiewicz

Let $\mathcal{M} = \langle M, <, +, \dots \rangle$ be o-minimal.

Theorem 0 (Basit-Chernikov-Starchenko-Tao-Tran, 2021)

The following are equivalent:

- a. \mathcal{M} linear (\Leftrightarrow definable functions are piecewise linear
 \Leftrightarrow no infinite definable field)
- b. for every definable \mathcal{E} , $\text{Zar}(\mathcal{E})$ holds.

Examples: $\langle \mathbb{R}, <, + \rangle$, $\mathbb{R}_{\text{vec}} = \langle \mathbb{R}, <, +, \{x \rightarrow rx\}_{r \in \mathbb{R}} \rangle$.

Theorem 1

The following are equivalent:

- a. \mathcal{M} semibounded

Linear and Semibounded Zarankiewicz

Let $\mathcal{M} = \langle M, <, +, \dots \rangle$ be o-minimal.

Theorem 0 (Basit-Chernikov-Starchenko-Tao-Tran, 2021)

The following are equivalent:

- a. \mathcal{M} linear (\Leftrightarrow definable functions are piecewise linear
 \Leftrightarrow no infinite definable field)
- b. for every definable \mathcal{E} , $\text{Zar}(\mathcal{E})$ holds.

Examples: $\langle \mathbb{R}, <, + \rangle$, $\mathbb{R}_{\text{vec}} = \langle \mathbb{R}, <, +, \{x \rightarrow rx\}_{r \in \mathbb{R}} \rangle$.

Theorem 1

The following are equivalent:

- a. \mathcal{M} semibounded (\Leftrightarrow definable functions are eventually linear
 \Leftrightarrow no definable field on M)

Linear and Semibounded Zarankiewicz

Let $\mathcal{M} = \langle M, <, +, \dots \rangle$ be o-minimal.

Theorem 0 (Basit-Chernikov-Starchenko-Tao-Tran, 2021)

The following are equivalent:

- a. \mathcal{M} linear (\Leftrightarrow definable functions are piecewise linear
 \Leftrightarrow no infinite definable field)
- b. for every definable \mathcal{E} , $\text{Zar}(\mathcal{E})$ holds.

Examples: $\langle \mathbb{R}, <, + \rangle$, $\mathbb{R}_{\text{vec}} = \langle \mathbb{R}, <, +, \{x \rightarrow rx\}_{r \in \mathbb{R}} \rangle$.

Theorem 1

The following are equivalent:

- a. \mathcal{M} semibounded (\Leftrightarrow definable functions are eventually linear
 \Leftrightarrow no definable field on M)
- b. for every definable \mathcal{E} , there is $m \in M^{>0}$, such that for \mathcal{C}_m the class of all m -distant grids, $\text{Zar}(\mathcal{E}, \mathcal{C}_m)$ holds.

Linear and Semibounded Zarankiewicz

Let $\mathcal{M} = \langle M, <, +, \dots \rangle$ be o-minimal.

Theorem 0 (Basit-Chernikov-Starchenko-Tao-Tran, 2021)

The following are equivalent:

- a. \mathcal{M} linear (\Leftrightarrow definable functions are piecewise linear
 \Leftrightarrow no infinite definable field)
- b. for every definable \mathcal{E} , $\text{Zar}(\mathcal{E})$ holds.

Examples: $\langle \mathbb{R}, <, + \rangle$, $\mathbb{R}_{\text{vec}} = \langle \mathbb{R}, <, +, \{x \rightarrow rx\}_{r \in \mathbb{R}} \rangle$.

Theorem 1

The following are equivalent:

- a. \mathcal{M} semibounded (\Leftrightarrow definable functions are eventually linear
 \Leftrightarrow no definable field on M)
- b. for every definable \mathcal{E} , there is $m \in M^{>0}$, such that for \mathcal{C}_m the class of all m -distant grids, $\text{Zar}(\mathcal{E}, \mathcal{C}_m)$ holds.

B is m -distant if for all i and $x, y \in B_i$, $|x - y| > m$.

Linear and Semibounded Zarankiewicz

Let $\mathcal{M} = \langle M, <, +, \dots \rangle$ be o-minimal.

Theorem 0 (Basit-Chernikov-Starchenko-Tao-Tran, 2021)

The following are equivalent:

- a. \mathcal{M} linear (\Leftrightarrow definable functions are piecewise linear
 \Leftrightarrow no infinite definable field)
- b. for every definable \mathcal{E} , $\text{Zar}(\mathcal{E})$ holds.

Examples: $\langle \mathbb{R}, <, + \rangle$, $\mathbb{R}_{\text{vec}} = \langle \mathbb{R}, <, +, \{x \rightarrow rx\}_{r \in \mathbb{R}} \rangle$.

Theorem 1

The following are equivalent:

- a. \mathcal{M} semibounded (\Leftrightarrow definable functions are eventually linear
 \Leftrightarrow no definable field on M)
- b. for every definable \mathcal{E} , there is $m \in M^{>0}$, such that for \mathcal{C}_m the class of all m -distant grids, $\text{Zar}(\mathcal{E}, \mathcal{C}_m)$ holds.

B is m -distant if for all i and $x, y \in B_i$, $|x - y| > m$.

Examples: $\langle \mathbb{R}, <, +, \cdot|_{(0,1)^2} \rangle$, $\langle \mathbb{R}_{\text{vec}}, \cdot|_{(0,1)^2} \rangle$.

Theorems

0. Linear Zarankiewicz
1. Semibounded Zarankiewicz

Theorems

0. Linear Zarankiewicz
1. Semibounded Zarankiewicz
2. Let $\mathcal{M} \models Th(\langle \mathbb{Z}, <, + \rangle)$. Then for every definable E , $Zar(E)$.

Theorems

0. Linear Zarankiewicz
1. Semibounded Zarankiewicz
2. Let $\mathcal{M} \models Th(\langle \mathbb{Z}, <, + \rangle)$. Then for every definable E , $Zar(E)$.
 - For $\mathcal{E} = \{E_t\}_{t \in \mathbb{Z}}$ with $E_t = [0, t]$, $Zar(\mathcal{E})$ does not hold.

Theorems

0. Linear Zarankiewicz
1. Semibounded Zarankiewicz
2. Let $\mathcal{M} \models Th(\langle \mathbb{Z}, <, + \rangle)$. Then for every definable E , $Zar(E)$.
 - For $\mathcal{E} = \{E_t\}_{t \in \mathbb{Z}}$ with $E_t = [0, t]$, $Zar(\mathcal{E})$ does not hold.
 - Proof by reduction to subgroups of powers of groups

Theorems

0. Linear Zarankiewicz
1. Semibounded Zarankiewicz
2. Let $\mathcal{M} \models Th(\langle \mathbb{Z}, <, + \rangle)$. Then for every definable E , $Zar(E)$.
 - For $\mathcal{E} = \{E_t\}_{t \in \mathbb{Z}}$ with $E_t = [0, t]$, $Zar(\mathcal{E})$ does not hold.
 - Proof by reduction to subgroups of powers of groups
3. Let $\mathcal{M} = \langle \mathbb{R}, <, +, \mathbb{Z} \rangle$. Then for every definable E , $Zar(E)$.

Theorems

0. Linear Zarankiewicz
1. Semibounded Zarankiewicz
2. Let $\mathcal{M} \models Th(\langle \mathbb{Z}, <, + \rangle)$. Then for every definable E , $Zar(E)$.
 - For $\mathcal{E} = \{E_t\}_{t \in \mathbb{Z}}$ with $E_t = [0, t]$, $Zar(\mathcal{E})$ does not hold.
 - Proof by reduction to subgroups of powers of groups
3. Let $\mathcal{M} = \langle \mathbb{R}, <, +, \mathbb{Z} \rangle$. Then for every definable E , $Zar(E)$.
4. Abstract Zarankiewicz for saturated \mathcal{M} , operator cl , grid class \mathcal{C} .

Theorems

0. Linear Zarankiewicz
1. Semibounded Zarankiewicz
2. Let $\mathcal{M} \models Th(\langle \mathbb{Z}, <, + \rangle)$. Then for every definable E , $Zar(E)$.
 - For $\mathcal{E} = \{E_t\}_{t \in \mathbb{Z}}$ with $E_t = [0, t]$, $Zar(\mathcal{E})$ does not hold.
 - Proof by reduction to subgroups of powers of groups
3. Let $\mathcal{M} = \langle \mathbb{R}, <, +, \mathbb{Z} \rangle$. Then for every definable E , $Zar(E)$.
4. Abstract Zarankiewicz for saturated \mathcal{M} , operator cl , grid class \mathcal{C} .

Applications:

- a. Let $\mathcal{M} \models Th(\langle \mathbb{Z}, <, + \rangle)$ saturated, \mathcal{E} definable family, $\mathcal{C}_{\mathbb{Z}}$ class of all \mathbb{Z} -distant grids. Then $Zar(\mathcal{E}, \mathcal{C}_{\mathbb{Z}})$.

Theorems

0. Linear Zarankiewicz
1. Semibounded Zarankiewicz
2. Let $\mathcal{M} \models Th(\langle \mathbb{Z}, <, + \rangle)$. Then for every definable E , $Zar(E)$.
 - For $\mathcal{E} = \{E_t\}_{t \in \mathbb{Z}}$ with $E_t = [0, t]$, $Zar(\mathcal{E})$ does not hold.
 - Proof by reduction to subgroups of powers of groups
3. Let $\mathcal{M} = \langle \mathbb{R}, <, +, \mathbb{Z} \rangle$. Then for every definable E , $Zar(E)$.
4. Abstract Zarankiewicz for saturated \mathcal{M} , operator cl , grid class \mathcal{C} .

Applications:

- a. Let $\mathcal{M} \models Th(\langle \mathbb{Z}, <, + \rangle)$ saturated, \mathcal{E} definable family, $\mathcal{C}_{\mathbb{Z}}$ class of all \mathbb{Z} -distant grids. Then $Zar(\mathcal{E}, \mathcal{C}_{\mathbb{Z}})$.
 - B is \mathbb{Z} -distant if for every i and $x, y \in B_i$, $x - y \notin \mathbb{Z}^{d_i}$.

Theorems

0. Linear Zarankiewicz
1. Semibounded Zarankiewicz
2. Let $\mathcal{M} \models Th(\langle \mathbb{Z}, <, + \rangle)$. Then for every definable E , $Zar(E)$.
 - For $\mathcal{E} = \{E_t\}_{t \in \mathbb{Z}}$ with $E_t = [0, t]$, $Zar(\mathcal{E})$ does not hold.
 - Proof by reduction to subgroups of powers of groups
3. Let $\mathcal{M} = \langle \mathbb{R}, <, +, \mathbb{Z} \rangle$. Then for every definable E , $Zar(E)$.
4. Abstract Zarankiewicz for saturated \mathcal{M} , operator cl , grid class \mathcal{C} .

Applications:

- a. Let $\mathcal{M} \models Th(\langle \mathbb{Z}, <, + \rangle)$ saturated, \mathcal{E} definable family, $\mathcal{C}_{\mathbb{Z}}$ class of all \mathbb{Z} -distant grids. Then $Zar(\mathcal{E}, \mathcal{C}_{\mathbb{Z}})$.
 - B is \mathbb{Z} -distant if for every i and $x, y \in B_i$, $x - y \notin \mathbb{Z}^{d_i}$.
- b. Zarankiewicz for stable 1-based theories with $nfcv$.

Theorems

0. Linear Zarankiewicz
1. Semibounded Zarankiewicz
2. Let $\mathcal{M} \models Th(\langle \mathbb{Z}, <, + \rangle)$. Then for every definable E , $Zar(E)$.
 - For $\mathcal{E} = \{E_t\}_{t \in \mathbb{Z}}$ with $E_t = [0, t]$, $Zar(\mathcal{E})$ does not hold.
 - Proof by reduction to subgroups of powers of groups
3. Let $\mathcal{M} = \langle \mathbb{R}, <, +, \mathbb{Z} \rangle$. Then for every definable E , $Zar(E)$.
4. Abstract Zarankiewicz for saturated \mathcal{M} , operator cl , grid class \mathcal{C} .

Applications:

- a. Let $\mathcal{M} \models Th(\langle \mathbb{Z}, <, + \rangle)$ saturated, \mathcal{E} definable family, $\mathcal{C}_{\mathbb{Z}}$ class of all \mathbb{Z} -distant grids. Then $Zar(\mathcal{E}, \mathcal{C}_{\mathbb{Z}})$.
 - B is \mathbb{Z} -distant if for every i and $x, y \in B_i$, $x - y \notin \mathbb{Z}^{d_i}$.
- b. Zarankiewicz for stable 1-based theories with *nfc*p.
- c. Zarankiewicz for locally modular regular types p in a stable theory (eg. the generic type of the solution set of the Heat differential equation), and \mathcal{C} the class of all cl_p -independent grids.

Zarankiewicz for subgroups of powers of groups

Zarankiewicz for subgroups of powers of groups

- For $j \in [r]$, let $p_j : \prod_{i \in [r]} M^{d_i} \rightarrow \prod_{i \neq j} M^{d_i}$ omit j -th coordinates

Zarankiewicz for subgroups of powers of groups

- ▶ For $j \in [r]$, let $p_j : \prod_{i \in [r]} M^{d_i} \rightarrow \prod_{i \neq j} M^{d_i}$ omit j -th coordinates
- ▶ If some $p_j|_E$ is N -to-1, then E linear Z -bounds, witnessed by N .

Zarankiewicz for subgroups of powers of groups

- ▶ For $j \in [r]$, let $p_j : \prod_{i \in [r]} M^{d_i} \rightarrow \prod_{i \neq j} M^{d_i}$ omit j -th coordinates
- ▶ If some $p_j|_E$ is N -to-1, then E linear Z -bounds, witnessed by N .
Indeed: $|E \cap B| \leq N|p_j(E) \cap p_j(B)| \leq Nn^{r-1}$

Zarankiewicz for subgroups of powers of groups

- ▶ For $j \in [r]$, let $p_j : \prod_{i \in [r]} M^{d_i} \rightarrow \prod_{i \neq j} M^{d_i}$ omit j -th coordinates
- ▶ If some $p_j|_E$ is N -to-1, then E linear Z -bounds, witnessed by N .
Indeed: $|E \cap B| \leq N|p_j(E) \cap p_j(B)| \leq Nn^{r-1}$
- ▶ Now let $\langle M, + \rangle$ be a group.

Zarankiewicz for subgroups of powers of groups

- ▶ For $j \in [r]$, let $p_j : \prod_{i \in [r]} M^{d_i} \rightarrow \prod_{i \neq j} M^{d_i}$ omit j -th coordinates
- ▶ If some $p_j|_E$ is N -to-1, then E linear Z -bounds, witnessed by N .
Indeed: $|E \cap B| \leq N|p_j(E) \cap p_j(B)| \leq Nn^{r-1}$
- ▶ Now let $\langle M, + \rangle$ be a group.
- ▶ Let $V \leq \langle M^d, + \rangle = \prod_{i \in [r]} \langle M^{d_i}, + \rangle$

Zarankiewicz for subgroups of powers of groups

- ▶ For $j \in [r]$, let $p_j : \prod_{i \in [r]} M^{d_i} \rightarrow \prod_{i \neq j} M^{d_i}$ omit j -th coordinates
- ▶ If some $p_j|_E$ is N -to-1, then E linear Z -bounds, witnessed by N .
Indeed: $|E \cap B| \leq N|p_j(E) \cap p_j(B)| \leq Nn^{r-1}$
- ▶ Now let $\langle M, + \rangle$ be a group.
- ▶ Let $V \leq \langle M^d, + \rangle = \prod_{i \in [r]} \langle M^{d_i}, + \rangle$
 - $p_j|_V$ is N -to-1 (some $N \in \mathbb{N}$) $\Leftrightarrow \ker p_j|_V$ finite

Zarankiewicz for subgroups of powers of groups

- ▶ For $j \in [r]$, let $p_j : \prod_{i \in [r]} M^{d_i} \rightarrow \prod_{i \neq j} M^{d_i}$ omit j -th coordinates
- ▶ If some $p_j|_E$ is N -to-1, then E linear Z -bounds, witnessed by N .
Indeed: $|E \cap B| \leq N|p_j(E) \cap p_j(B)| \leq Nn^{r-1}$
- ▶ Now let $\langle M, + \rangle$ be a group.
- ▶ Let $V \leq \langle M^d, + \rangle = \prod_{i \in [r]} \langle M^{d_i}, + \rangle$
 - $p_j|_V$ is N -to-1 (some $N \in \mathbb{N}$) $\Leftrightarrow \ker p_j|_V$ finite
 - Let $U_j = \pi_j(\ker p_j|_V)$.

Zarankiewicz for subgroups of powers of groups

- ▶ For $j \in [r]$, let $p_j : \prod_{i \in [r]} M^{d_i} \rightarrow \prod_{i \neq j} M^{d_i}$ omit j -th coordinates
- ▶ If some $p_j|_E$ is N -to-1, then E linear Z -bounds, witnessed by N .
Indeed: $|E \cap B| \leq N|p_j(E) \cap p_j(B)| \leq Nn^{r-1}$
- ▶ Now let $\langle M, + \rangle$ be a group.
- ▶ Let $V \leq \langle M^d, + \rangle = \prod_{i \in [r]} \langle M^{d_i}, + \rangle$
 - $p_j|_V$ is N -to-1 (some $N \in \mathbb{N}$) $\Leftrightarrow \ker p_j|_V$ finite
 - Let $U_j = \pi_j(\ker p_j|_V)$. Then $U_1 \times \cdots \times U_r \subseteq V$

Zarankiewicz for subgroups of powers of groups

- ▶ For $j \in [r]$, let $p_j : \prod_{i \in [r]} M^{d_i} \rightarrow \prod_{i \neq j} M^{d_i}$ omit j -th coordinates
- ▶ If some $p_j|_E$ is N -to-1, then E linear Z -bounds, witnessed by N .
Indeed: $|E \cap B| \leq N|p_j(E) \cap p_j(B)| \leq Nn^{r-1}$
- ▶ Now let $\langle M, + \rangle$ be a group.
- ▶ Let $V \leq \langle M^d, + \rangle = \prod_{i \in [r]} \langle M^{d_i}, + \rangle$
 - $p_j|_V$ is N -to-1 (some $N \in \mathbb{N}$) $\Leftrightarrow \ker p_j|_V$ finite
 - Let $U_j = \pi_j(\ker p_j|_V)$. Then $U_1 \times \cdots \times U_r \subseteq V$
 - Therefore $Zar(V)$ holds.

Zarankiewicz for subgroups of powers of groups

- ▶ For $j \in [r]$, let $p_j : \prod_{i \in [r]} M^{d_i} \rightarrow \prod_{i \neq j} M^{d_i}$ omit j -th coordinates
- ▶ If some $p_{j|_E}$ is N -to-1, then E linear Z -bounds, witnessed by N .
Indeed: $|E \cap B| \leq N|p_j(E) \cap p_j(B)| \leq Nn^{r-1}$
- ▶ Now let $\langle M, + \rangle$ be a group.
- ▶ Let $V \leq \langle M^d, + \rangle = \prod_{i \in [r]} \langle M^{d_i}, + \rangle$
 - $p_{j|_V}$ is N -to-1 (some $N \in \mathbb{N}$) $\Leftrightarrow \ker p_{j|_V}$ finite
 - Let $U_j = \pi_j(\ker p_{j|_V})$. Then $U_1 \times \cdots \times U_r \subseteq V$
 - Therefore $Zar(V)$ holds.
- ▶ For $C \subseteq M^d$, $Sh(C) :=$ subgroup of $\langle M^d, + \rangle$ generated by C .

Zarankiewicz for subgroups of powers of groups

- ▶ For $j \in [r]$, let $p_j : \prod_{i \in [r]} M^{d_i} \rightarrow \prod_{i \neq j} M^{d_i}$ omit j -th coordinates
- ▶ If some $p_j|_E$ is N -to-1, then E linear Z -bounds, witnessed by N .
Indeed: $|E \cap B| \leq N|p_j(E) \cap p_j(B)| \leq Nn^{r-1}$
- ▶ Now let $\langle M, + \rangle$ be a group.
- ▶ Let $V \leq \langle M^d, + \rangle = \prod_{i \in [r]} \langle M^{d_i}, + \rangle$
 - $p_j|_V$ is N -to-1 (some $N \in \mathbb{N}$) $\Leftrightarrow \ker p_j|_V$ finite
 - Let $U_j = \pi_j(\ker p_j|_V)$. Then $U_1 \times \cdots \times U_r \subseteq V$
 - Therefore $Zar(V)$ holds.
- ▶ For $C \subseteq M^d$, $Sh(C) :=$ subgroup of $\langle M^d, + \rangle$ generated by C .

General strategy for $Zar(E)$: Show that E is a finite union of sets C , each satisfying, after translating to 0:

$$(*) \quad C \text{ is } k\text{-free} \Rightarrow Sh(C) \text{ } k\text{-free}$$

(*) for linear and Presburger cells C containing 0

(*) for linear and Presburger cells C containing 0

- ▶ C k -free ($k \in \mathbb{N}$) $\Rightarrow Sh(C)$ 2-free

(*) for linear and Presburger cells C containing 0

► C k -free ($k \in \mathbb{N}$) $\Rightarrow Sh(C)$ 2-free

Linear:

(*) for linear and Presburger cells C containing 0

- ▶ C k -free ($k \in \mathbb{N}$) $\Rightarrow Sh(C)$ 2-free

Linear:

- C is open in $Sh(C)$

(*) for linear and Presburger cells C containing 0

► C k -free ($k \in \mathbb{N}$) $\Rightarrow Sh(C)$ 2-free

Linear:

- C is open in $Sh(C)$
- $Sh(C)$ not 2-free $\Rightarrow \exists U_1 \times \cdots \times U_r \subseteq Sh(C)$

(*) for linear and Presburger cells C containing 0

► C k -free ($k \in \mathbb{N}$) $\Rightarrow Sh(C)$ 2-free

Linear:

- C is open in $Sh(C)$
- $Sh(C)$ not 2-free $\Rightarrow \exists U_1 \times \cdots \times U_r \subseteq Sh(C)$,
with each $U_i \subseteq M^{d_i}$ definable infinite and close to 0

(*) for linear and Presburger cells C containing 0

► C k -free ($k \in \mathbb{N}$) $\Rightarrow Sh(C)$ 2-free

Linear:

- C is open in $Sh(C)$
- $Sh(C)$ not 2-free $\Rightarrow \exists U_1 \times \cdots \times U_r \subseteq Sh(C)$,
with each $U_i \subseteq M^{d_i}$ definable infinite and close to 0
- $C \cap (U_1 \times \cdots \times U_r)$ contains a k -grid, for any $k \in \mathbb{N}$.

(*) for linear and Presburger cells C containing 0

► C k -free ($k \in \mathbb{N}$) $\Rightarrow Sh(C)$ 2-free

Linear:

- C is open in $Sh(C)$
- $Sh(C)$ not 2-free $\Rightarrow \exists U_1 \times \cdots \times U_r \subseteq Sh(C)$,
with each $U_i \subseteq M^{d_i}$ definable infinite and close to 0
- $C \cap (U_1 \times \cdots \times U_r)$ contains a k -grid, for any $k \in \mathbb{N}$.

Presburger:

(*) for linear and Presburger cells C containing 0

► C k -free ($k \in \mathbb{N}$) $\Rightarrow Sh(C)$ 2-free

Linear:

- C is open in $Sh(C)$
- $Sh(C)$ not 2-free $\Rightarrow \exists U_1 \times \cdots \times U_r \subseteq Sh(C)$,
with each $U_i \subseteq M^{d_i}$ definable infinite and close to 0
- $C \cap (U_1 \times \cdots \times U_r)$ contains a k -grid, for any $k \in \mathbb{N}$.

Presburger:

- Let $N \in \mathbb{N}$. Say $x \in C$ is N -internal if $B_N(x) \cap Sh(C) \subseteq C$.

(*) for linear and Presburger cells C containing 0

- ▶ C k -free ($k \in \mathbb{N}$) $\Rightarrow Sh(C)$ 2-free

Linear:

- C is open in $Sh(C)$
- $Sh(C)$ not 2-free $\Rightarrow \exists U_1 \times \cdots \times U_r \subseteq Sh(C)$,
with each $U_i \subseteq M^{d_i}$ definable infinite and close to 0
- $C \cap (U_1 \times \cdots \times U_r)$ contains a k -grid, for any $k \in \mathbb{N}$.

Presburger:

- Let $N \in \mathbb{N}$. Say $x \in C$ is N -internal if $B_N(x) \cap Sh(C) \subseteq C$.
- For every $N \in \mathbb{N}$, C contains an N -internal point a .

(*) for linear and Presburger cells C containing 0

- ▶ C k -free ($k \in \mathbb{N}$) $\Rightarrow Sh(C)$ 2-free

Linear:

- C is open in $Sh(C)$
- $Sh(C)$ not 2-free $\Rightarrow \exists U_1 \times \cdots \times U_r \subseteq Sh(C)$,
with each $U_i \subseteq M^{d_i}$ definable infinite and close to 0
- $C \cap (U_1 \times \cdots \times U_r)$ contains a k -grid, for any $k \in \mathbb{N}$.

Presburger:

- Let $N \in \mathbb{N}$. Say $x \in C$ is N -internal if $B_N(x) \cap Sh(C) \subseteq C$.
- For every $N \in \mathbb{N}$, C contains an N -internal point a .
- $Sh(C)$ not 2-free $\Rightarrow \exists U_1 \times \cdots \times U_r \subseteq Sh(C)$

(*) for linear and Presburger cells C containing 0

- ▶ C k -free ($k \in \mathbb{N}$) $\Rightarrow Sh(C)$ 2-free

Linear:

- C is open in $Sh(C)$
- $Sh(C)$ not 2-free $\Rightarrow \exists U_1 \times \cdots \times U_r \subseteq Sh(C)$,
with each $U_i \subseteq M^{d_i}$ definable infinite and close to 0
- $C \cap (U_1 \times \cdots \times U_r)$ contains a k -grid, for any $k \in \mathbb{N}$.

Presburger:

- Let $N \in \mathbb{N}$. Say $x \in C$ is N -internal if $B_N(x) \cap Sh(C) \subseteq C$.
- For every $N \in \mathbb{N}$, C contains an N -internal point a .
- $Sh(C)$ not 2-free $\Rightarrow \exists U_1 \times \cdots \times U_r \subseteq Sh(C)$,
with each $U_i \cap \mathbb{Z}^{d_i}$ infinite

(*) for linear and Presburger cells C containing 0

► C k -free ($k \in \mathbb{N}$) $\Rightarrow Sh(C)$ 2-free

Linear:

- C is open in $Sh(C)$
- $Sh(C)$ not 2-free $\Rightarrow \exists U_1 \times \cdots \times U_r \subseteq Sh(C)$,
with each $U_i \subseteq M^{d_i}$ definable infinite and close to 0
- $C \cap (U_1 \times \cdots \times U_r)$ contains a k -grid, for any $k \in \mathbb{N}$.

Presburger:

- Let $N \in \mathbb{N}$. Say $x \in C$ is N -internal if $B_N(x) \cap Sh(C) \subseteq C$.
- For every $N \in \mathbb{N}$, C contains an N -internal point a .
- $Sh(C)$ not 2-free $\Rightarrow \exists U_1 \times \cdots \times U_r \subseteq Sh(C)$,
with each $U_i \cap \mathbb{Z}^{d_i}$ infinite
- $Sh(C) \cap \mathbb{Z}^d$ contains a k -grid $B \subseteq B_N(0)$ (for any $k \in \mathbb{N}$).

(*) for linear and Presburger cells C containing 0

- ▶ C k -free ($k \in \mathbb{N}$) $\Rightarrow Sh(C)$ 2-free

Linear:

- C is open in $Sh(C)$
- $Sh(C)$ not 2-free $\Rightarrow \exists U_1 \times \cdots \times U_r \subseteq Sh(C)$,
with each $U_i \subseteq M^{d_i}$ definable infinite and close to 0
- $C \cap (U_1 \times \cdots \times U_r)$ contains a k -grid, for any $k \in \mathbb{N}$.

Presburger:

- Let $N \in \mathbb{N}$. Say $x \in C$ is N -internal if $B_N(x) \cap Sh(C) \subseteq C$.
- For every $N \in \mathbb{N}$, C contains an N -internal point a .
- $Sh(C)$ not 2-free $\Rightarrow \exists U_1 \times \cdots \times U_r \subseteq Sh(C)$,
with each $U_i \cap \mathbb{Z}^{d_i}$ infinite
- $Sh(C) \cap \mathbb{Z}^d$ contains a k -grid $B \subseteq B_N(0)$ (for any $k \in \mathbb{N}$).
Let $a \in C$ N -internal.

(*) for linear and Presburger cells C containing 0

► C k -free ($k \in \mathbb{N}$) $\Rightarrow Sh(C)$ 2-free

Linear:

- C is open in $Sh(C)$
- $Sh(C)$ not 2-free $\Rightarrow \exists U_1 \times \cdots \times U_r \subseteq Sh(C)$,
with each $U_i \subseteq M^{d_i}$ definable infinite and close to 0
- $C \cap (U_1 \times \cdots \times U_r)$ contains a k -grid, for any $k \in \mathbb{N}$.

Presburger:

- Let $N \in \mathbb{N}$. Say $x \in C$ is N -internal if $B_N(x) \cap Sh(C) \subseteq C$.
- For every $N \in \mathbb{N}$, C contains an N -internal point a .
- $Sh(C)$ not 2-free $\Rightarrow \exists U_1 \times \cdots \times U_r \subseteq Sh(C)$,
with each $U_i \cap \mathbb{Z}^{d_i}$ infinite
- $Sh(C) \cap \mathbb{Z}^d$ contains a k -grid $B \subseteq B_N(0)$ (for any $k \in \mathbb{N}$).
Let $a \in C$ N -internal. Then

$$a + B \subseteq B_N(a) \cap Sh(C) \subseteq C.$$

Abstract Zarankiewicz in saturated setting

Abstract Zarankiewicz in saturated setting

Theorem (Inspired by BCSTT, 2021)

Let \mathcal{M} saturated, $\text{cl} : \mathcal{P}(\mathcal{M}) \rightarrow \mathcal{P}(\mathcal{M})$, and \mathcal{C} a class of grids that contains all cl -independent grids (each B_i is cl -independent).

Abstract Zarankiewicz in saturated setting

Theorem (Inspired by BCSTT, 2021)

Let \mathcal{M} saturated, $\text{cl} : \mathcal{P}(\mathcal{M}) \rightarrow \mathcal{P}(\mathcal{M})$, and \mathcal{C} a class of grids that contains all cl -independent grids (each B_i is cl -independent). Assume:

(DEF) Let $(a, b) \in M^{k+m}$ and $A \subseteq M$. If $a \in \text{cl}(Ab)$, then there is an A -definable set $X \subseteq M^{k+m}$ with $(a, b) \in X$, and for every $(a', b') \in X$, $a' \in \text{cl}(Ab')$.

Abstract Zarankiewicz in saturated setting

Theorem (Inspired by BCSTT, 2021)

Let \mathcal{M} saturated, $\text{cl} : \mathcal{P}(\mathcal{M}) \rightarrow \mathcal{P}(\mathcal{M})$, and \mathcal{C} a class of grids that contains all cl -independent grids (each B_i is cl -independent). Assume:

(DEF) Let $(a, b) \in M^{k+m}$ and $A \subseteq M$. If $a \in \text{cl}(Ab)$, then there is an A -definable set $X \subseteq M^{k+m}$ with $(a, b) \in X$, and for every $(a', b') \in X$, $a' \in \text{cl}(Ab')$.

(TIGHT) Let E definable, \mathcal{C} - k -free (some $k \in \mathbb{N}$). Then for every $a = (a_1, \dots, a_r) \in E$, the set $\{a_1, \dots, a_r\}$ is acl -dependent.

Abstract Zarankiewicz in saturated setting

Theorem (Inspired by BCSTT, 2021)

Let \mathcal{M} saturated, $\text{cl} : \mathcal{P}(\mathcal{M}) \rightarrow \mathcal{P}(\mathcal{M})$, and \mathcal{C} a class of grids that contains all cl -independent grids (each B_i is cl -independent). Assume:

(DEF) Let $(a, b) \in M^{k+m}$ and $A \subseteq M$. If $a \in \text{cl}(Ab)$, then there is an A -definable set $X \subseteq M^{k+m}$ with $(a, b) \in X$, and for every $(a', b') \in X$, $a' \in \text{cl}(Ab')$.

(TIGHT) Let E definable, \mathcal{C} - k -free (some $k \in \mathbb{N}$). Then for every $a = (a_1, \dots, a_r) \in E$, the set $\{a_1, \dots, a_r\}$ is acl -dependent.

(UB) Let $\mathcal{X} = \{X_b\}_{b \in I}$ definable family of sets in M^{d_i} . Then $\exists N \in \mathbb{N}$ such that $\forall b \in I$ with X_b finite, and $Y \in \mathcal{C}$, we have $|X_b \cap \pi_i(Y)| \leq N$.

Abstract Zarankiewicz in saturated setting

Theorem (Inspired by BCSTT, 2021)

Let \mathcal{M} saturated, $\text{cl} : \mathcal{P}(\mathcal{M}) \rightarrow \mathcal{P}(\mathcal{M})$, and \mathcal{C} a class of grids that contains all cl -independent grids (each B_i is cl -independent). Assume:

(DEF) Let $(a, b) \in M^{k+m}$ and $A \subseteq M$. If $a \in \text{cl}(Ab)$, then there is an A -definable set $X \subseteq M^{k+m}$ with $(a, b) \in X$, and for every $(a', b') \in X$, $a' \in \text{cl}(Ab')$.

(TIGHT) Let E definable, \mathcal{C} - k -free (some $k \in \mathbb{N}$). Then for every $a = (a_1, \dots, a_r) \in E$, the set $\{a_1, \dots, a_r\}$ is acl -dependent.

(UB) Let $\mathcal{X} = \{X_b\}_{b \in I}$ definable family of sets in M^{d_i} . Then $\exists N \in \mathbb{N}$ such that $\forall b \in I$ with X_b finite, and $Y \in \mathcal{C}$, we have $|X_b \cap \pi_i(Y)| \leq N$.

Then for every definable family \mathcal{E} , $\text{Zar}(\mathcal{E}, \mathcal{C})$.

a. $\mathcal{M} \models \text{Th}(\langle \mathbb{Z}, <, + \rangle)$. Let $\text{cl} = \text{acl}$ and \mathcal{C} all \mathbb{Z} -distant grids.

Abstract Zarankiewicz in saturated setting

Theorem (Inspired by BCSTT, 2021)

Let \mathcal{M} saturated, $\text{cl} : \mathcal{P}(\mathcal{M}) \rightarrow \mathcal{P}(\mathcal{M})$, and \mathcal{C} a class of grids that contains all cl -independent grids (each B_i is cl -independent). Assume:

(DEF) Let $(a, b) \in M^{k+m}$ and $A \subseteq M$. If $a \in \text{cl}(Ab)$, then there is an A -definable set $X \subseteq M^{k+m}$ with $(a, b) \in X$, and for every $(a', b') \in X$, $a' \in \text{cl}(Ab')$.

(TIGHT) Let E definable, \mathcal{C} - k -free (some $k \in \mathbb{N}$). Then for every $a = (a_1, \dots, a_r) \in E$, the set $\{a_1, \dots, a_r\}$ is acl -dependent.

(UB) Let $\mathcal{X} = \{X_b\}_{b \in I}$ definable family of sets in M^{d_i} . Then $\exists N \in \mathbb{N}$ such that $\forall b \in I$ with X_b finite, and $Y \in \mathcal{C}$, we have $|X_b \cap \pi_i(Y)| \leq N$.

Then for every definable family \mathcal{E} , $\text{Zar}(\mathcal{E}, \mathcal{C})$.

- $\mathcal{M} \models \text{Th}(\langle \mathbb{Z}, <, + \rangle)$. Let $\text{cl} = \text{acl}$ and \mathcal{C} all \mathbb{Z} -distant grids.
- $\text{Th}(\mathcal{M})$ stable, 1-based, nfcp. Let $\text{cl} = \text{acl}$, \mathcal{C} all grids.

Abstract Zarankiewicz in saturated setting

Theorem (Inspired by BCSTT, 2021)

Let \mathcal{M} saturated, $\text{cl} : \mathcal{P}(\mathcal{M}) \rightarrow \mathcal{P}(\mathcal{M})$, and \mathcal{C} a class of grids that contains all cl -independent grids (each B_i is cl -independent). Assume:

(DEF) Let $(a, b) \in M^{k+m}$ and $A \subseteq M$. If $a \in \text{cl}(Ab)$, then there is an A -definable set $X \subseteq M^{k+m}$ with $(a, b) \in X$, and for every $(a', b') \in X$, $a' \in \text{cl}(Ab')$.

(TIGHT) Let E definable, \mathcal{C} - k -free (some $k \in \mathbb{N}$). Then for every $a = (a_1, \dots, a_r) \in E$, the set $\{a_1, \dots, a_r\}$ is acl -dependent.

(UB) Let $\mathcal{X} = \{X_b\}_{b \in I}$ definable family of sets in M^{d_i} . Then $\exists N \in \mathbb{N}$ such that $\forall b \in I$ with X_b finite, and $Y \in \mathcal{C}$, we have $|X_b \cap \pi_i(Y)| \leq N$.

Then for every definable family \mathcal{E} , $\text{Zar}(\mathcal{E}, \mathcal{C})$.

- $\mathcal{M} \models \text{Th}(\langle \mathbb{Z}, <, + \rangle)$. Let $\text{cl} = \text{acl}$ and \mathcal{C} all \mathbb{Z} -distant grids.
- $\text{Th}(\mathcal{M})$ stable, 1-based, nfc_p . Let $\text{cl} = \text{acl}$, \mathcal{C} all grids.
- $\mathcal{M} = p(\mathcal{U})$, p locally modular regular type. Let $\text{cl}_p(B) = \{b \in M : b \not\perp B\}$, \mathcal{C} all cl_p -independent grids

$\langle \mathbb{R}, <, +, \mathbb{Z} \rangle$ and open questions

$\langle \mathbb{R}, <, +, \mathbb{Z} \rangle$ and open questions

- $\langle \mathbb{R}, <, +, \mathbb{Z} \rangle$: Special case $E = S + J + D$, where S Presburger cell, $J + D$ linear cell, J 'purely unbounded', D bounded.

$\langle \mathbb{R}, <, +, \mathbb{Z} \rangle$ and open questions

- $\langle \mathbb{R}, <, +, \mathbb{Z} \rangle$: Special case $E = S + J + D$, where S Presburger cell, $J + D$ linear cell, J 'purely unbounded', D bounded.

Adjust (*): $E = S + J + D$ k -free $\Rightarrow Sh(S) + Sh(J) + D$ k -free
 $\Rightarrow \exists j \exists N \in \mathbb{N}, p_j|_E$ N -to-1

$\langle \mathbb{R}, <, +, \mathbb{Z} \rangle$ and open questions

- ▶ $\langle \mathbb{R}, <, +, \mathbb{Z} \rangle$: Special case $E = S + J + D$, where S Presburger cell, $J + D$ linear cell, J 'purely unbounded', D bounded.
Adjust (*): $E = S + J + D$ k -free $\Rightarrow Sh(S) + Sh(J) + D$ k -free
 $\Rightarrow \exists j \exists N \in \mathbb{N}, p_j|_E$ N -to-1
- ▶ Zarankiewicz for OAGs, $\langle \mathbb{R}, <, +, 2^{\mathbb{Z}} \rangle$, $\langle \mathbb{Z}, <, +, 2^{\mathbb{N}} \rangle$, 'linear type', no definable field.

$\langle \mathbb{R}, <, +, \mathbb{Z} \rangle$ and open questions

- ▶ $\langle \mathbb{R}, <, +, \mathbb{Z} \rangle$: Special case $E = S + J + D$, where S Presburger cell, $J + D$ linear cell, J 'purely unbounded', D bounded.
Adjust (*): $E = S + J + D$ k -free $\Rightarrow Sh(S) + Sh(J) + D$ k -free
 $\Rightarrow \exists j \exists N \in \mathbb{N}, p_j|_E$ N -to-1
- ▶ Zarankiewicz for OAGs, $\langle \mathbb{R}, <, +, 2^{\mathbb{Z}} \rangle$, $\langle \mathbb{Z}, <, +, 2^{\mathbb{N}} \rangle$, 'linear type', no definable field.
- ▶ Let $\mathcal{M} = \langle \mathbb{R}, <, +, \dots \rangle$ o-minimal. For what \mathcal{C} , TFAE:
 - a. the exponential \exp is not definable in \mathcal{M} .
 - b. for every binary definable $E \subseteq M^{d_1} \times M^{d_2}$, $Zar(E, \mathcal{C})$ holds.

$\langle \mathbb{R}, <, +, \mathbb{Z} \rangle$ and open questions

- ▶ $\langle \mathbb{R}, <, +, \mathbb{Z} \rangle$: Special case $E = S + J + D$, where S Presburger cell, $J + D$ linear cell, J 'purely unbounded', D bounded.
Adjust (*): $E = S + J + D$ k -free $\Rightarrow Sh(S) + Sh(J) + D$ k -free
 $\Rightarrow \exists j \exists N \in \mathbb{N}, p_j|_E$ N -to-1
- ▶ Zarankiewicz for OAGs, $\langle \mathbb{R}, <, +, 2^{\mathbb{Z}} \rangle$, $\langle \mathbb{Z}, <, +, 2^{\mathbb{N}} \rangle$, 'linear type', no definable field.
- ▶ Let $\mathcal{M} = \langle \mathbb{R}, <, +, \dots \rangle$ o-minimal. For what \mathcal{C} , TFAE:
 - a. the exponential \exp is not definable in \mathcal{M} .
 - b. for every binary definable $E \subseteq M^{d_1} \times M^{d_2}$, $Zar(E, \mathcal{C})$ holds.
- ▶ Elekes-Szabó: recognise group structure from combinatorial data.

$\langle \mathbb{R}, <, +, \mathbb{Z} \rangle$ and open questions

- ▶ $\langle \mathbb{R}, <, +, \mathbb{Z} \rangle$: Special case $E = S + J + D$, where S Presburger cell, $J + D$ linear cell, J 'purely unbounded', D bounded.
Adjust (*): $E = S + J + D$ k -free $\Rightarrow Sh(S) + Sh(J) + D$ k -free
 $\Rightarrow \exists j \exists N \in \mathbb{N}, p_j|_E$ N -to-1
- ▶ Zarankiewicz for OAGs, $\langle \mathbb{R}, <, +, 2^{\mathbb{Z}} \rangle$, $\langle \mathbb{Z}, <, +, 2^{\mathbb{N}} \rangle$, 'linear type', no definable field.
- ▶ Let $\mathcal{M} = \langle \mathbb{R}, <, +, \dots \rangle$ o-minimal. For what \mathcal{C} , TFAE:
 - a. the exponential \exp is not definable in \mathcal{M} .
 - b. for every binary definable $E \subseteq M^{d_1} \times M^{d_2}$, $Zar(E, \mathcal{C})$ holds.
- ▶ Elekes-Szabó: recognise group structure from combinatorial data.
Let \mathcal{M} reduct of an o-minimal $\langle M, <, +, \dots \rangle$. Recover $+$.

$\langle \mathbb{R}, <, +, \mathbb{Z} \rangle$ and open questions

- ▶ $\langle \mathbb{R}, <, +, \mathbb{Z} \rangle$: Special case $E = S + J + D$, where S Presburger cell, $J + D$ linear cell, J 'purely unbounded', D bounded.
Adjust (*): $E = S + J + D$ k -free $\Rightarrow Sh(S) + Sh(J) + D$ k -free
 $\Rightarrow \exists j \exists N \in \mathbb{N}, p_j \upharpoonright_E$ N -to-1
- ▶ Zarankiewicz for OAGs, $\langle \mathbb{R}, <, +, 2^{\mathbb{Z}} \rangle$, $\langle \mathbb{Z}, <, +, 2^{\mathbb{N}} \rangle$, 'linear type', no definable field.
- ▶ Let $\mathcal{M} = \langle \mathbb{R}, <, +, \dots \rangle$ o-minimal. For what \mathcal{C} , TFAE:
 - a. the exponential \exp is not definable in \mathcal{M} .
 - b. for every binary definable $E \subseteq M^{d_1} \times M^{d_2}$, $Zar(E, \mathcal{C})$ holds.
- ▶ Elekes-Szabó: recognise group structure from combinatorial data.
Let \mathcal{M} reduct of an o-minimal $\langle M, <, +, \dots \rangle$. Recover $+$.

Thank you!