## **Overview of Models**

These models provide estimates of outdoor concentrations for six pollutants (four gases: O3, CO, SO2, NO2; Two aerosols: PM10, PM2.5) throughout the contiguous U.S. Model estimates, with the exception of ozone are annual-average values, for years with available monitoring data. Ozone model estimates are the average during May through September of the daily maximum 8-hour moving average, for years with available monitoring data.

Pollutant	Units	Available Years
$O_3$	ppb	1979-2015
$SO_2$	ppb	1979-2015
$NO_2$	ppb	1979-2015
$PM_{10}$	$\mu g/m^3$	1988-2015
CO	ppm	1990-2015
$PM_{2.5}$	$\mu g/m^3$	1999-2015

Data are available at national, state, county, census tract\*, and census block group\* levels, based on population-weighted averages of census block level predictions using 2010 census geographies.

These models are derived from publicly available concentration measurements from U.S. EPA regulatory monitors, and use information about land use (for example, locations of major and minor roads; elevation; and whether an area is urban or rural) and satellite-derived estimates of air pollution to predict concentrations at locations without measurements. Researchers generally do not use estimates of the types given here to draw conclusions about air quality at one location or a small number of specific locations; as such, we do not recommend that type of use.

## **File Format**

Parameter	Description	
Fips	Unique code for geographic unit with the following structure: $SCCCTTTTTG^1$ $^1S = 1-2$ digit state code; $C = 3$ digit county code; $T = 6$ digit tract code; $G = 1$ digit block group code This code can be used to join these data to Census data and shapefiles	
pollutant	Pollutant for a given observation (co, no2, o3, pm10, pm25, so2)	
year	Prediction year for a given observation	
pred_wght	Population-weighted concentration based on block level centroid predictions, units vary by pollutant (see above)	
state_abbr	State abbreviation for a given observation	
lat/lon	Population-weighted latitude and longitude based on block level centroid	

## Acknowledgment

Data are free and publicly available for non-commercial purposes, data may not be used for commercial purposes without prior permission. Please cite as follows:

"This article includes concentration estimates developed by the Center for Air, Climate and Energy Solutions using v1 empirical models as described in Kim et al., 2018."

Suggested citation: Kim S.-Y.; Bechle, M.; Hankey, S.; Sheppard, L.; Szpiro, A. A.; Marshall, J. D. 2018. "Concentrations of criteria pollutants in the contiguous U.S., 1979 – 2015: Role of prediction model parsimony in integrated empirical geographic regression." PLoS ONE 15(2), e0228535. DOI: 10.1371/journal.pone.0228535

<sup>\*</sup>Our models only provide estimates for tracts and block groups with a population greater than zero.