

CS665: Advanced Data Mining

Lecture#15: SVD-1

U Kang
KAIST

Outline

- ➔ ☐ **Motivation**
- ☐ Definition - properties
- ☐ Interpretation
- ☐ Complexity
- ☐ Conclusion

SVD - Motivation

- problem #1: text - LSI: find ‘concepts’
- problem #2: compression / dim. reduction

SVD - Motivation

- problem #1: text - LSI: find ‘concepts’

term document	data	information	retrieval	brain	lung
CS-TR1	1	1	1	0	0
CS-TR2	2	2	2	0	0
CS-TR3	1	1	1	0	0
CS-TR4	5	5	5	0	0
MED-TR1	0	0	0	2	2
MED-TR2	0	0	0	3	3
MED-TR3	0	0	0	1	1

SVD - Motivation

- Customer-product, for recommendation system :

	bread	lettuce	tomatos	beef	chicken
↑					
vegetarians	1	1	1	0	0
↓	2	2	2	0	0
↑	1	1	1	0	0
↓	5	5	5	0	0
↑	0	0	0	2	2
meat eaters	0	0	0	3	3
↓	0	0	0	1	1

SVD - Motivation


- problem #2: compress / reduce dimensionality

Problem - specs


- $\sim 10^6$ rows; $\sim 10^3$ columns; no updates;
- random access to any cell(s) ; small error: OK

customer	day	We	Th	Fr	Sa	Su
		7/10/06	7/11/06	7/12/06	7/13/06	7/14/06
ABC Inc.		1	1	1	0	0
DEF Ltd.		2	2	2	0	0
GHI Inc.		1	1	1	0	0
KLM Co.		5	5	5	0	0
Smith		0	0	0	2	2
Johnson		0	0	0	3	3
Thompson		0	0	0	1	1


SVD - Motivation



SVD - Motivation



Outline

- ☒ Motivation
-  ☐ **Definition - properties**
- ☐ Interpretation
- ☐ Complexity
- ☐ Conclusion

SVD - Definition

(reminder: matrix multiplication)

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix} \times \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} \\ \\ \end{bmatrix}$$

$3 \times 2 \qquad 2 \times 1$

SVD - Definition

(reminder: matrix multiplication)

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix} \times \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} \\ \\ \end{bmatrix}$$


$\begin{matrix} \xleftarrow{\hspace{10em}} & & \xrightarrow{\hspace{10em}} \\ 3 \times 2 & 2 \times 1 & 3 \times 1 \\ \xleftrightarrow{\hspace{2em}} & \xleftrightarrow{\hspace{2em}} & \end{matrix}$

SVD - Definition

(reminder: matrix multiplication)

$$\begin{array}{c}
 \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix} \times \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} -1 \end{bmatrix}
 \end{array}$$

$3 \times 2 \quad \quad 2 \times 1 \quad \quad 3 \times 1$



SVD - Definition

(reminder: matrix multiplication)

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix} \times \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

$\begin{matrix} \xleftarrow{\hspace{1.5cm}} & & \xrightarrow{\hspace{1.5cm}} \\ 3 \times 2 & 2 \times 1 & 3 \times 1 \\ \xleftrightarrow{\hspace{1cm}} & \xleftrightarrow{\hspace{1cm}} & \end{matrix}$

SVD - Definition

(reminder: matrix multiplication)

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix} \times \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} -1 \\ -1 \\ -1 \end{bmatrix}$$


SVD - Definition

$$\mathbf{A}_{[n \times m]} = \mathbf{U}_{[n \times r]} \mathbf{\Lambda}_{[r \times r]} (\mathbf{V}_{[m \times r]})^T$$

- \mathbf{A} : $n \times m$ matrix (eg., n documents, m terms)
- \mathbf{U} : $n \times r$ matrix (n documents, r concepts)
- $\mathbf{\Lambda}$: $r \times r$ diagonal matrix (strength of each ‘concept’) (r : rank of the matrix)
- \mathbf{V} : $m \times r$ matrix (m terms, r concepts)

SVD - Definition

$$\mathbf{A}_{[n \times m]} = \mathbf{U}_{[n \times r]} \mathbf{\Lambda}_{[r \times r]} (\mathbf{V}_{[m \times r]})^T$$



SVD - Properties

THEOREM [Press+92]: always possible to decompose matrix \mathbf{A} into $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{V}^T$, where

- $\mathbf{U}, \mathbf{\Lambda}, \mathbf{V}$: unique (*)
- \mathbf{U}, \mathbf{V} : column orthonormal (ie., columns are unit vectors, orthogonal to each other)
 - $\mathbf{U}^T \mathbf{U} = \mathbf{I}; \mathbf{V}^T \mathbf{V} = \mathbf{I}$ (\mathbf{I} : identity matrix)
- $\mathbf{\Lambda}$: singular are positive, and sorted in decreasing order

SVD - Example

■ $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{V}^T$ - example:

$$\begin{array}{c}
 \begin{array}{c} \uparrow \\ \text{CS} \\ \downarrow \end{array} \\
 \begin{array}{c} \uparrow \\ \text{MD} \\ \downarrow \end{array}
 \end{array}
 \begin{array}{c}
 \text{data} \quad \text{inf.} \quad \text{retrieval} \\
 \downarrow \quad \text{brain} \quad \text{lung}
 \end{array}
 \begin{bmatrix}
 1 & 1 & 1 & 0 & 0 \\
 2 & 2 & 2 & 0 & 0 \\
 1 & 1 & 1 & 0 & 0 \\
 5 & 5 & 5 & 0 & 0 \\
 0 & 0 & 0 & 2 & 2 \\
 0 & 0 & 0 & 3 & 3 \\
 0 & 0 & 0 & 1 & 1
 \end{bmatrix}
 =
 \begin{bmatrix}
 0.18 & 0 \\
 0.36 & 0 \\
 0.18 & 0 \\
 0.90 & 0 \\
 0 & 0.53 \\
 0 & 0.80 \\
 0 & 0.27
 \end{bmatrix}
 \times
 \begin{bmatrix}
 9.64 & 0 \\
 0 & 5.29
 \end{bmatrix}
 \times
 \begin{bmatrix}
 0.58 & 0.58 & 0.58 & 0 & 0 \\
 0 & 0 & 0 & 0.71 & 0.71
 \end{bmatrix}$$

SVD - Example

■ $A = U \Lambda V^T$ - example:

$$\begin{array}{c}
 \begin{array}{c} \uparrow \\ \text{CS} \\ \downarrow \end{array} \\
 \begin{array}{c} \uparrow \\ \text{MD} \\ \downarrow \end{array}
 \end{array}
 \begin{bmatrix}
 1 & 1 & 1 & 0 & 0 \\
 2 & 2 & 2 & 0 & 0 \\
 1 & 1 & 1 & 0 & 0 \\
 5 & 5 & 5 & 0 & 0 \\
 0 & 0 & 0 & 2 & 2 \\
 0 & 0 & 0 & 3 & 3 \\
 0 & 0 & 0 & 1 & 1
 \end{bmatrix}
 =
 \begin{bmatrix}
 0.18 & 0 \\
 0.36 & 0 \\
 0.18 & 0 \\
 0.90 & 0 \\
 0 & 0.53 \\
 0 & 0.80 \\
 0 & 0.27
 \end{bmatrix}
 \times
 \begin{bmatrix}
 9.64 & 0 \\
 0 & 5.29
 \end{bmatrix}
 \times
 \begin{bmatrix}
 0.58 & 0.58 & 0.58 & 0 & 0 \\
 0 & 0 & 0 & 0.71 & 0.71
 \end{bmatrix}$$

retrieval
 data inf. ↓ brain lung
 CS-concept
 MD-concept

SVD - Example

■ $A = U \Lambda V^T$ - example:

doc-to-concept
similarity matrix

$$\begin{array}{c}
 \begin{array}{c} \uparrow \\ \text{CS} \\ \downarrow \\ \uparrow \\ \text{MD} \\ \downarrow \end{array}
 \end{array}
 \begin{array}{c}
 \text{data} \quad \text{inf.} \downarrow \quad \text{retrieval} \quad \text{brain} \quad \text{lung} \quad \text{CS-concept} \quad \text{MD-concept} \\
 \begin{bmatrix}
 1 & 1 & 1 & 0 & 0 \\
 2 & 2 & 2 & 0 & 0 \\
 1 & 1 & 1 & 0 & 0 \\
 5 & 5 & 5 & 0 & 0 \\
 0 & 0 & 0 & 2 & 2 \\
 0 & 0 & 0 & 3 & 3 \\
 0 & 0 & 0 & 1 & 1
 \end{bmatrix}
 =
 \begin{bmatrix}
 0.18 & 0 \\
 0.36 & 0 \\
 0.18 & 0 \\
 0.90 & 0 \\
 0 & 0.53 \\
 0 & 0.80 \\
 0 & 0.27
 \end{bmatrix}
 \times
 \begin{bmatrix}
 9.64 & 0 \\
 0 & 5.29
 \end{bmatrix}
 \times
 \begin{bmatrix}
 0.58 & 0.58 & 0.58 & 0 & 0 \\
 0 & 0 & 0 & 0.71 & 0.71
 \end{bmatrix}
 \end{array}$$

SVD - Example

■ $A = U \Lambda V^T$ - example:

retrieval
inf. ↓ brain lung

‘strength’ of CS-concept

↑

CS

↓

↑

MD

↓

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}$$

=

$$\begin{bmatrix} 0.18 & 0 \\ 0.36 & 0 \\ 0.18 & 0 \\ 0.90 & 0 \\ 0 & 0.53 \\ 0 & 0.80 \\ 0 & 0.27 \end{bmatrix}$$

X

$$\begin{bmatrix} 9.64 & 0 \\ 0 & 5.29 \end{bmatrix}$$

X

$$\begin{bmatrix} 0.58 & 0.58 & 0.58 & 0 & 0 \\ 0 & 0 & 0 & 0.71 & 0.71 \end{bmatrix}$$

SVD - Example

■ $A = U \Lambda V^T$ - example:

term-to-concept
similarity matrix

↑

CS

↓

↑

MD

↓

	data	inf.	brain	lung
1	1	1	0	0
2	2	2	0	0
1	1	1	0	0
5	5	5	0	0
0	0	0	2	2
0	0	0	3	3
0	0	0	1	1

=

0.18	0
0.36	0
0.18	0
0.90	0
0	0.53
0	0.80
0	0.27

X

9.64	0
0	5.29

X

0.58	0.58	0.58	0	0
0	0	0	0.71	0.71

CS-concept

SVD - Example

■ $A = U \Lambda V^T$ - example:

term-to-concept
similarity matrix

retrieval
inf. ↓
data brain lung

CS
↑
↓
MD
↑
↓

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 0.18 & 0 \\ 0.36 & 0 \\ 0.18 & 0 \\ 0.90 & 0 \\ 0 & 0.53 \\ 0 & 0.80 \\ 0 & 0.27 \end{bmatrix} \times \begin{bmatrix} 9.64 & 0 \\ 0 & 5.29 \end{bmatrix} \times \begin{bmatrix} 0.58 & 0.58 & 0.58 & 0 & 0 \\ 0 & 0 & 0 & 0.71 & 0.71 \end{bmatrix}$$

CS-concept

Outline

☒ Motivation

☒ Definition - properties

 ☐ **Interpretation**

#1: documents/terms/concepts

#2: dim. reduction

#3: picking non-zero, rectangular clusters

☐ Complexity

☐ Conclusion

SVD - Interpretation #1

‘documents’, ‘terms’ and ‘concepts’:

- U : document-to-concept similarity matrix
- V : term-to-concept sim. matrix
- Λ : its diagonal elements: ‘strength’ of each concept

SVD – Interpretation #1

‘documents’, ‘terms’ and ‘concepts’:

Q: if \mathbf{A} is the document-to-term matrix, what is $\mathbf{A}^T \mathbf{A}$?

A:

Q: $\mathbf{A} \mathbf{A}^T$?

A:

SVD – Interpretation #1

‘documents’, ‘terms’ and ‘concepts’:

Q: if \mathbf{A} is the document-to-term matrix, what is $\mathbf{A}^T \mathbf{A}$?

A: term-to-term ($[m \times m]$) similarity matrix

Q: $\mathbf{A} \mathbf{A}^T$?

A: document-to-document ($[n \times n]$) similarity matrix

SVD properties

- \mathbf{V} are the eigenvectors of the *covariance matrix* $\mathbf{A}^T \mathbf{A}$
- \mathbf{U} are the eigenvectors of the *Gram (inner-product) matrix* $\mathbf{A} \mathbf{A}^T$


Further reading:

1. Ian T. Jolliffe, *Principal Component Analysis* (2nd ed), Springer, 2002.
2. Gilbert Strang, *Linear Algebra and Its Applications* (4th ed), Brooks Cole, 2005.

SVD - Interpretation #2

- best axis to project on: ('best' = min sum of squares of projection errors)


SVD - Motivation



SVD - interpretation #2

SVD: gives
best axis to project

- minimum RMS error



SVD - Interpretation #2

customer	day	We	Th	Fr	Sa	Su
		7/10/96	7/11/96	7/12/96	7/13/96	7/14/96
ABC Inc.		1	1	1	0	0
DEF Ltd.		2	2	2	0	0
GHI Inc.		1	1	1	0	0
KLM Co.		5	5	5	0	0
Smith		0	0	0	2	2
Johnson		0	0	0	3	3
Thompson		0	0	0	1	1

SVD - Interpretation #2

- $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{V}^T$ - example:

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 0.18 & 0 \\ 0.36 & 0 \\ 0.18 & 0 \\ 0.90 & 0 \\ 0 & 0.53 \\ 0 & 0.80 \\ 0 & 0.27 \end{bmatrix} \times \begin{bmatrix} 9.64 & 0 \\ 0 & 5.29 \end{bmatrix} \times \begin{bmatrix} \boxed{0.58 \ 0.58 \ 0.58 \ 0 \ 0} \\ 0 \ 0 \ 0 \ 0.71 \ 0.71 \end{bmatrix} \quad \text{v1}$$

SVD - Interpretation #2

■ $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{V}^T$ - example:

variance ('spread') on the v_1 axis

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 0.18 & 0 \\ 0.36 & 0 \\ 0.18 & 0 \\ 0.90 & 0 \\ 0 & 0.53 \\ 0 & 0.80 \\ 0 & 0.27 \end{bmatrix} \times \begin{bmatrix} 9.64 & 0 \\ 0 & 5.29 \end{bmatrix} \times \begin{bmatrix} 0.58 & 0.58 & 0.58 & 0 & 0 \\ 0 & 0 & 0 & 0.71 & 0.71 \end{bmatrix}$$

SVD - Interpretation #2

■ $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{V}^T$ - example:

□ $\mathbf{U} \mathbf{\Lambda}$ gives the coordinates of the points in the projection axis

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 0.18 & 0 \\ 0.36 & 0 \\ 0.18 & 0 \\ 0.90 & 0 \\ 0 & 0.53 \\ 0 & 0.80 \\ 0 & 0.27 \end{bmatrix} \times \begin{bmatrix} 9.64 & 0 \\ 0 & 5.29 \end{bmatrix} \times \begin{bmatrix} 0.58 & 0.58 & 0.58 & 0 & 0 \\ 0 & 0 & 0 & 0.71 & 0.71 \end{bmatrix}$$

SVD - Interpretation #2

- More details
- Q: how exactly is dim. reduction done?

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 0.18 & 0 \\ 0.36 & 0 \\ 0.18 & 0 \\ 0.90 & 0 \\ 0 & 0.53 \\ 0 & 0.80 \\ 0 & 0.27 \end{bmatrix} \times \begin{bmatrix} 9.64 & 0 \\ 0 & 5.29 \end{bmatrix} \times \begin{bmatrix} 0.58 & 0.58 & 0.58 & 0 & 0 \\ 0 & 0 & 0 & 0.71 & 0.71 \end{bmatrix}$$

SVD - Interpretation #2

- More details
- Q: how exactly is dim. reduction done?
- A: set the smallest singular values to zero:

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 0.18 & 0 \\ 0.36 & 0 \\ 0.18 & 0 \\ 0.90 & 0 \\ 0 & 0.53 \\ 0 & 0.80 \\ 0 & 0.27 \end{bmatrix} \times \begin{bmatrix} 9.64 & 0 \\ 0 & 5.29 \end{bmatrix} \times \begin{bmatrix} 0.58 & 0.58 & 0.58 & 0 & 0 \\ 0 & 0 & 0 & 0.71 & 0.71 \end{bmatrix}$$

SVD - Interpretation #2

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} \sim \begin{bmatrix} 0.18 & 0 \\ 0.36 & 0 \\ 0.18 & 0 \\ 0.90 & 0 \\ 0 & 0.53 \\ 0 & 0.80 \\ 0 & 0.27 \end{bmatrix} \times \begin{bmatrix} 9.64 & 0 \\ 0 & 0 \end{bmatrix} \times \begin{bmatrix} 0.58 & 0.58 & 0.58 & 0 & 0 \\ 0 & 0 & 0 & 0.71 & 0.71 \end{bmatrix}$$

SVD - Interpretation #2

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} \sim \begin{bmatrix} 0.18 & 0 \\ 0.36 & 0 \\ 0.18 & 0 \\ 0.90 & 0 \\ 0 & 0.53 \\ 0 & 0.80 \\ 0 & 0.27 \end{bmatrix} \times \begin{bmatrix} 9.64 & 0 \\ 0 & 0 \end{bmatrix} \times \begin{bmatrix} 0.58 & 0.58 & 0.58 & 0 & 0 \\ 0 & 0 & 0 & 0.71 & 0.71 \end{bmatrix}$$

SVD - Interpretation #2

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} \sim \begin{bmatrix} 0.18 \\ 0.36 \\ 0.18 \\ 0.90 \\ 0 \\ 0 \\ 0 \end{bmatrix} \times \begin{bmatrix} 9.64 \end{bmatrix} \times \begin{bmatrix} 0.58 & 0.58 & 0.58 & 0 & 0 \end{bmatrix}$$

SVD - Interpretation #2

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

SVD - Interpretation #2

Exactly equivalent:

‘spectral decomposition’ of the matrix:

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 0.18 & 0 \\ 0.36 & 0 \\ 0.18 & 0 \\ 0.90 & 0 \\ 0 & 0.53 \\ 0 & 0.80 \\ 0 & 0.27 \end{bmatrix} \times \begin{bmatrix} 9.64 & 0 \\ 0 & 5.29 \end{bmatrix} \times \begin{bmatrix} 0.58 & 0.58 & 0.58 & 0 & 0 \\ 0 & 0 & 0 & 0.71 & 0.71 \end{bmatrix}$$

SVD - Interpretation #2

Exactly equivalent:

‘spectral decomposition’ of the matrix:

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} | & | \\ u_1 & u_2 \\ | & | \end{bmatrix} \times \begin{bmatrix} \lambda_1 & \emptyset \\ \emptyset & \lambda_2 \end{bmatrix} \times \begin{bmatrix} \text{---} v_1 \text{---} \\ \text{---} v_2 \text{---} \end{bmatrix}$$

SVD - Interpretation #2

Exactly equivalent:

‘spectral decomposition’ of the matrix:

$$\begin{array}{c} \updownarrow \\ n \end{array} \begin{array}{c} \leftarrow m \rightarrow \\ \left[\begin{array}{ccccc} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{array} \right] \end{array} = \lambda_1 \underbrace{u_1}_{\text{column vector}} \underbrace{v_1^T}_{\text{row vector}} + \lambda_2 \underbrace{u_2}_{\text{column vector}} \underbrace{v_2^T}_{\text{row vector}} + \dots$$

SVD - Interpretation #2

Exactly equivalent:

‘spectral decomposition’ of the matrix:

$$\begin{array}{c} \updownarrow \\ n \end{array} \begin{array}{c} \leftarrow m \rightarrow \\ \left[\begin{array}{ccccc} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{array} \right] \end{array} = \begin{array}{c} \leftarrow r \text{ terms} \rightarrow \\ \lambda_1 \begin{array}{c} \nearrow \\ u_1 \\ \nwarrow \\ n \times 1 \end{array} \begin{array}{c} \nearrow \\ v_1^T \\ \nwarrow \\ 1 \times m \end{array} + \lambda_2 \begin{array}{c} u_2 \end{array} \begin{array}{c} v_2^T \end{array} + \dots \end{array}$$

SVD - Interpretation #2

approximation / dim. reduction:
by keeping the first few terms (Q: how many?)

$$\begin{array}{c} \updownarrow \\ n \end{array} \begin{array}{c} \leftarrow m \rightarrow \\ \left[\begin{array}{ccccc} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{array} \right] \end{array} = \lambda_1 \mathbf{u}_1 \mathbf{v}_1^T + \lambda_2 \mathbf{u}_2 \mathbf{v}_2^T + \dots$$

assume: $\lambda_1 \geq \lambda_2 \geq \dots$

SVD - Interpretation #2

A (heuristic - [Fukunaga]): keep 80-90% of 'energy'
y' (= sum of squares of λ_i 's)

$$\begin{array}{c} \updownarrow n \end{array} \begin{array}{c} \leftarrow m \rightarrow \\ \left[\begin{array}{ccccc} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{array} \right] \end{array} = \lambda_1 \mathbf{u}_1 \mathbf{v}_1^T + \lambda_2 \mathbf{u}_2 \mathbf{v}_2^T + \dots$$

assume: $\lambda_1 \geq \lambda_2 \geq \dots$

Pictorially: matrix form of SVD


$$\mathbf{A} \approx \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T = \sum_i \sigma_i \mathbf{u}_i \circ \mathbf{v}_i$$

The diagram illustrates the matrix form of SVD. Matrix \mathbf{A} (blue, $m \times n$) is approximated by the product of matrix \mathbf{U} (green, $m \times k$), matrix $\mathbf{\Sigma}$ (white, $k \times k$), and matrix \mathbf{V}^T (green, $k \times n$). Matrix \mathbf{U} is shown as a tall rectangle with k columns. Matrix $\mathbf{\Sigma}$ is shown as a small square with k rows and k columns. Matrix \mathbf{V}^T is shown as a wide rectangle with k rows and n columns. The approximation symbol \approx is placed between \mathbf{A} and the product $\mathbf{U}\mathbf{\Sigma}\mathbf{V}^T$.

- Best rank- k approximation in L2

Pictorially: matrix form of SVD

$$\mathbf{A} \approx \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T = \sum_i \sigma_i \mathbf{u}_i \circ \mathbf{v}_i$$



- Best rank- k approximation in L2

SVD - Interpretation #3

- finds non-zero clusters in a data matrix

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 0.18 & 0 \\ 0.36 & 0 \\ 0.18 & 0 \\ 0.90 & 0 \\ 0 & 0.53 \\ 0 & 0.80 \\ 0 & 0.27 \end{bmatrix} \times \begin{bmatrix} 9.64 & 0 \\ 0 & 5.29 \end{bmatrix} \times \begin{bmatrix} 0.58 & 0.58 & 0.58 & 0 & 0 \\ 0 & 0 & 0 & 0.71 & 0.71 \end{bmatrix}$$

SVD - Interpretation #3


- finds non-zero clusters in a data matrix

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ \hline 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 0.18 & 0 \\ 0.36 & 0 \\ 0.18 & 0 \\ 0.90 & 0 \\ 0 & 0.53 \\ 0 & 0.80 \\ 0 & 0.27 \end{bmatrix} \times \begin{bmatrix} 9.64 & 0 \\ 0 & 5.29 \end{bmatrix} \times \begin{bmatrix} 0.58 & 0.58 & 0.58 & 0 & 0 \\ 0 & 0 & 0 & 0.71 & 0.71 \end{bmatrix}$$

SVD - Interpretation #3

- finds non-zero clusters in a data matrix =
- ‘communities’ (bi-partite cores, here)

$$\left[\begin{array}{ccc|cc} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 5 & 5 & 5 & 0 & 0 \\ \hline 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{array} \right]$$



SVD - Interpretation #3

- Exercise: find the SVD, ‘by inspection’!
- Q: rank = ??

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} \quad \quad \quad \\ \quad \quad \quad \\ \quad \quad \quad \\ \quad \quad \quad \\ \quad \quad \quad \end{bmatrix} \times \begin{bmatrix} \quad \quad \quad \\ \quad \quad \quad \\ \quad \quad \quad \end{bmatrix} \times \begin{bmatrix} \quad \quad \quad \\ \quad \quad \quad \\ \quad \quad \quad \end{bmatrix}$$

SVD - Interpretation #3

- A: rank = 2 (2 linearly independent rows/cols)

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} | & | \\ ?? & ?? \\ | & | \end{bmatrix} \times \begin{bmatrix} ?? & 0 \\ 0 & ?? \end{bmatrix} \times \begin{bmatrix} \text{---} & ?? & \text{---} \\ \text{---} & ?? & \text{---} \end{bmatrix}$$

SVD - Interpretation #3

- A: rank = 2 (2 linearly independent rows/cols)

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 1 & 0 \\ 0 & 1 \\ 0 & 1 \end{bmatrix}$$

orthogonal??

$$X \begin{bmatrix} ?? & 0 \\ 0 & ?? \end{bmatrix} X$$

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}$$

SVD - Interpretation #3

- column vectors: are orthogonal - but not unit vectors:

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1/\sqrt{3} & 0 \\ 1/\sqrt{3} & 0 \\ 1/\sqrt{3} & 0 \\ 0 & 1/\sqrt{2} \\ 0 & 1/\sqrt{2} \end{bmatrix} \times \begin{bmatrix} ?? & 0 \\ 0 & ?? \end{bmatrix} \times \begin{bmatrix} 1/\sqrt{3} & 1/\sqrt{3} & 1/\sqrt{3} & 0 & 0 \\ 0 & 0 & 0 & 1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix}$$

SVD - Interpretation #3

- and the singular values are:

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1/\sqrt{3} & 0 \\ 1/\sqrt{3} & 0 \\ 1/\sqrt{3} & 0 \\ 0 & 1/\sqrt{2} \\ 0 & 1/\sqrt{2} \end{bmatrix} \times \begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix} \times \begin{bmatrix} 1/\sqrt{3} & 1/\sqrt{3} & 1/\sqrt{3} & 0 & 0 \\ 0 & 0 & 0 & 1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix}$$

SVD - Interpretation #3


- Q: How to check we are correct?

$$\begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1/\sqrt{3} & 0 \\ 1/\sqrt{3} & 0 \\ 1/\sqrt{3} & 0 \\ 0 & 1/\sqrt{2} \\ 0 & 1/\sqrt{2} \end{bmatrix} \times \begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix} \times \begin{bmatrix} 1/\sqrt{3} & 1/\sqrt{3} & 1/\sqrt{3} & 0 & 0 \\ 0 & 0 & 0 & 1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix}$$

SVD - Interpretation #3

- A: SVD properties:
 - matrix product should give back matrix \mathbf{A}
 - matrix \mathbf{U} should be column-orthonormal, i.e., columns should be unit vectors, orthogonal to each other
 - same for matrix \mathbf{V}
 - matrix $\mathbf{\Lambda}$ should be diagonal, with positive values

Outline

- ☒ Motivation
- ☒ Definition - properties
- ☒ Interpretation
-  ☐ **Complexity**
- ☐ Conclusion

SVD - Complexity

- $O(n * m * m)$ or $O(n * n * m)$ (whichever is less)
- less work, if we just want singular values
- or if we want first k singular vectors
- or if the matrix is sparse [Berry]
- Implemented: in any linear algebra package (LAPACK, matlab, Splus/R, mathematica ...)
 - Hadoop Implementation: HEigen [Kang+ PAKDD 2011]

Outline

- ☒ Motivation
- ☒ Definition - properties
- ☒ Interpretation
- ☒ Complexity

 ☐ **Conclusion**

SVD - conclusions so far

- SVD: $A = U \Lambda V^T$: unique (*)
- U : document-to-concept similarities
- V : term-to-concept similarities
- Λ : strength of each concept
- dim. reduction: keep the first few strongest singular values (80-90% of ‘energy’)
- SVD: picks up linear correlations
- SVD: picks up non-zero clusters

References

- Berry, Michael: <http://www.cs.utk.edu/~lsi/>
- Fukunaga, K. (1990). Introduction to Statistical Pattern Recognition, Academic Press.
- Press, W. H., S. A. Teukolsky, et al. (1992). Numerical Recipes in C, Cambridge University Press.