



A Performance and Energy Study of the Hyperbolic PDE Solver Engine ExaHyPE

Master's Thesis in Computational Science and Engineering

Fabian Gura

Department of Informatics
Technische Universität München

August 22, 2016

Eexaminers: Univ.-Prof. Dr. Michael Bader
Dr. Tobias Weinzierl
Supervisor: Dr. Vasco Varduhn

I hereby declare that this thesis is entirely the result of my own work except where otherwise indicated. I have only used the resources given in the list of references.

Place: _____ Date: _____ Signature: _____

Abstract

... ..

Contents

Contents	iii
1 Introduction	1
2 Theory	3
2.1 A D -dimensional ADER-DG scheme with MUSCL-Hancock a-posteriori subcell limiting for non-linear hyperbolic conser- vation laws	3
2.1.1 Notation	3
2.1.2 Hyperbolic Conservation Laws	4
2.1.3 Space and Time Discretization	4
2.1.4 Element-local weak formulation	5
2.1.5 Restriction to Finite-Dimensional Function Spaces . . .	6
2.1.6 Space-time predictor	8
2.1.7 Reference Elements and Mappings	9
2.1.8 Orthogonal Bases for the Finite-Dimensional Function Spaces	10
2.1.9 Basis Functions in Global Coordinates	14
2.1.10 A Fully-discrete Iterative Method for the Space-time Predictor	15
2.1.11 A Fully-discrete Update Scheme for the Time-discrete Solution	23
2.1.12 A Posteriori Subcell Limiting	28
2.2 Profiling and Energy-aware Computing	32
3 A profiling infrastructure for ExaHyPE	33
4 Preliminary profiling results, case studies	35
5 Conclusion and Outlook	37

6 Acknowledgment	39
-------------------------	-----------

Chapter 1

Introduction

- Challenges of exascale
- The ExaHyPE project (numerics, resilience, profiling) as an answer
- On the importance of profiling and performance measuring

Chapter 2

Theory

2.1 A D -dimensional ADER-DG scheme with MUSCL-Hancock a-posteriori subcell limiting for non-linear hyperbolic conservation laws

Arbitrary High Order Derivatives Discontinuous Galerkin (ADER-DG)

The sections on unlimited ADER-DG are to varying degree based on work by Etienne

Additional contribution: Precise index notation, which directly leads to the original FORTRAN implementation used to create the results presented in ??

2.1.1 Notation

We use vector notation whenever possible. Advantage: Complete derivation, direct conversion to code.

Details: Lai, Rubin, Kemper

Syntactic sugar to make notation more concise while maintaining preciseness. Motivated by numerical computing environments such as MATLAB[®] or Octave ([?]).

Sequence: Notation $1 : N$ denotes the sequence of integers from 1 to N , i.e. $1 : N := \{1, 2, \dots, N\}$.

Sequenced vector access: $[\alpha]_{j:k} := [[\alpha]_j, [\alpha]_{j+1}, \dots, [\alpha]_k]$ a vector of length $j - k + 1$ for $j \leq k$, otherwise the vector has length 0.

Vector access:

Vector concatenation: $\alpha = [\alpha]_{1:N} = [[\alpha]_{1:k}, [\alpha]_{k+1:N}]$ for $k \in 1 : N$.

Clarify ddx.i and [x].i

2.1.2 Hyperbolic Conservation Laws

A D -dimensional conservation law in a system with V quantities is described mathematically by a partial differential equation (PDE) of the form

$$\frac{\partial}{\partial t} [u]_v + \frac{\partial}{\partial x_d} [F(u)]_{vd} = [S(u)]_v \quad \text{on } \Omega \times [0, T] \quad (2.1)$$

together with initial conditions

$$[u(x, 0)]_v = [u_0(x)]_v \quad \forall x \in \Omega, \quad (2.2)$$

and boundary conditions

$$[u(x, t)]_v = [u_B(x, t)]_v \quad \forall x \in \partial\Omega, t \in [0, T], \quad (2.3)$$

for all $v \in \mathcal{V}$, where we define the index set $\mathcal{V} = \{1, 2, \dots, V\}$. $[0, T]$ is the time interval of interest and $\Omega \subset \mathbb{R}^D$ denotes the spatial domain. The function $F : \mathbb{R}^V \rightarrow \mathbb{R}^{V \times D}$, $u \mapsto F(u) = [f_1(u), f_2(u), \dots, f_D(u)]$ is called the flux function. For the problem to be hyperbolic we require that all Jacobian matrices $A_d(u)$, $d \in \{1, 2, \dots, D\} := \mathcal{D}$, defined as

$$[A_d]_{ij} = \frac{\partial [f_d]_i}{\partial u_j}, \quad (2.4)$$

have D real eigenvalues in each admissible state $u \in \mathbb{R}^V$.

2.1.3 Space and Time Discretization

Let \mathcal{K}_h be a quadrilateral partition of Ω , i.e.

$$K \cap J = \emptyset \quad \forall K, J \in \mathcal{K}_h, K \neq J, \quad (2.5)$$

$$\bigcup_{K \in \mathcal{K}_h} K = \Omega. \quad (2.6)$$

For the index set $\mathcal{I} := \{0, 1, \dots, I-1\}$ let $\{t_i\}_{i \in \mathcal{I}}$ be an I -fold partition of the time interval $[0, T]$ such that

$$0 = t_0 < t_1 < \dots < t_I = T. \quad (2.7)$$

For $i \in \mathcal{I}$ we furthermore define

$$\Delta t_i = t_{i+1} - t_i, \quad (2.8)$$

so that the subinterval $[t_i, t_{i+1}]$ can be written as $[t_i, t_i + \Delta t_i]$.

Without loss of generality we can solve the original PDE (2.1) on $\Omega \times [0, T]$ simply by solving the PDE locally for each element $K \in \mathcal{K}_h$ in the time interval $[t_0, t_0 + \Delta t_0]$ and then proceeding to the next time interval until we have reached the final time T . This gives rise to an element-local formulation on a subinterval in time which we will focus in the following.

2.1.4 Element-local weak formulation

Let $L^2(\Omega)^V$ be the space of vector-valued, square-integrable functions on Ω , i.e.

$$L^2(\Omega)^V = \left\{ \mathbf{w} : \Omega \rightarrow \mathbb{R}^V \mid \int_{\Omega} \|\mathbf{w}\|^2 dx < \infty \right\}. \quad (2.9)$$

Let $\mathbf{w} \in L^2(\Omega)^V$ be a spatial test function. Multiplication of the original PDE (2.1) and integration over a space-time cell $K \times [t_i, t_i + \Delta t_i]$ yields a element-local weak formulation of the problem,

$$\begin{aligned} \int_{t_i}^{t_i+\Delta t_i} \int_K \frac{\partial}{\partial t} [\mathbf{u}]_v [\mathbf{w}]_v dx dt + \int_{t_i}^{t_i+\Delta t_i} \int_K \frac{\partial}{\partial x_d} [F(\mathbf{u})]_{vd} [\mathbf{w}]_v dx dt = \\ \int_{t_i}^{t_i+\Delta t_i} \int_K [S(\mathbf{u})]_v [\mathbf{w}]_v dx dt, \end{aligned} \quad (2.10)$$

which we require to hold for all $v \in \mathcal{V}$, $\mathbf{w} \in L^2(\Omega)^V$, $K \in \mathcal{K}_h$ and $i \in \mathcal{I}$.

Integration by parts of the spatial integral in the second term yields

$$\begin{aligned} \int_K \frac{\partial}{\partial x_d} [F(\mathbf{u})]_{vd} [\mathbf{w}]_v dx = \\ \int_K \frac{\partial}{\partial x_d} ([F(\mathbf{u})]_{vd} [\mathbf{w}]_v) dx - \int_K [F(\mathbf{u})]_{vd} \frac{\partial}{\partial x_d} [\mathbf{w}]_v dx. \end{aligned} \quad (2.11)$$

Application of the divergence theorem to the first term on the right-hand side of (2.11) yields

$$\int_K \frac{\partial}{\partial x_d} ([F(\mathbf{u})]_{vd} [\mathbf{w}]_v) dx = \int_{\partial K} [F(\mathbf{u})]_{vd} [\mathbf{w}]_v [\mathbf{n}]_d ds(x), \quad (2.12)$$

where $\mathbf{n} \in \mathbb{R}^D$ is the unit-length, outward-pointing normal vector at a point x on the surface of K , which we denote by ∂K .

Inserting eqs. (2.11) and (2.12) into eq. (2.10) yields the following more favorable element-local weak formulation of the original equation (2.1):

$$\begin{aligned} \int_{t_i}^{t_i+\Delta t_i} \int_K \frac{\partial}{\partial t} [\mathbf{u}]_v [\mathbf{w}]_v dx dt - \int_{t_i}^{t_i+\Delta t_i} \int_K [F(\mathbf{u})]_{vd} \frac{\partial}{\partial x_d} [\mathbf{w}]_v dx dt + \\ \int_{t_i}^{t_i+\Delta t_i} \int_{\partial K} [F(\mathbf{u})]_{vd} [\mathbf{w}]_v [\mathbf{n}]_d ds(x) dt = \int_{t_i}^{t_i+\Delta t_i} \int_K [S(\mathbf{u})]_v [\mathbf{w}]_v dx dt. \end{aligned} \quad (2.13)$$

Again we require the weak formulation to hold for all $v \in \mathcal{V}$, $\mathbf{w} \in L^2(\Omega)^V$, $K \in \mathcal{K}_h$ and $i \in \mathcal{I}$.

2.1.5 Restriction to Finite-Dimensional Function Spaces

To discretize eq. (2.13) we need to impose the restriction that both test and ansatz functions come from a finite-dimensional function space. First, let $\mathcal{Q}_N(K)^V$ and $\mathcal{Q}_N(K \times [t_i, t_i + \Delta t_i])^V$ be the space of vector-valued, multivariate polynomials of degree less or equal than N in each variable on K and $K \times [t_i, t_i + \Delta t_i]$, respectively. We can then define the following finite-dimensional function spaces:

- For spatial functions we define

$$\mathbb{W}_h = \left\{ w_h \in L^2(\Omega)^V \mid w_h|_K := w_h^K \in \mathcal{Q}_N(K)^V \forall K \in \mathcal{K}_h \right\}. \quad (2.14)$$

- For space-time functions on the time subinterval $[t_i, t_i + \Delta t_i]$, $i \in \mathcal{I}$ we define

$$\begin{aligned} \tilde{\mathbb{W}}_h^i = \left\{ \tilde{w}_h^i \in L^2(\Omega \times [t_i, t_i + \Delta t_i]) \mid \right. \\ \left. \tilde{w}_h^i|_K := \tilde{w}_h^{K,i} \in \mathcal{Q}_N(K \times [t_i, t_i + \Delta t_i]) \forall K \in \mathcal{K}_h \right\}. \end{aligned} \quad (2.15)$$

Replacing w by $w_h \in \mathbb{W}_h$ and u by $\tilde{u}_h^i \in \tilde{\mathbb{W}}_h^i$ in eq. (2.13), i.e. restricting ourselves to test and ansatz functions from finite-dimensional function spaces, yields an approximation of the weak formulation,

$$\begin{aligned} & \int_{t_i}^{t_i+\Delta t_i} \int_K \frac{\partial}{\partial t} [\tilde{u}_h^{K,i}]_v [w_h^K]_v dx dt - \int_{t_i}^{t_i+\Delta t_i} \int_{\partial K} [F(\tilde{u}_h^{K,i})]_{vd} \frac{\partial}{\partial x_d} [w_h^K]_v dx dt + \\ & \int_{t_i}^{t_i+\Delta t_i} \int_{\partial K} [\mathcal{G}(\tilde{u}_h^{K,i}, \tilde{u}_h^{K'+i}, n)]_v [w_h^K]_v ds(x) dt = \\ & \int_{t_i}^{t_i+\Delta t_i} \int_K [S(\tilde{u}_h^{K,i})]_v [w_h^K]_v dx dt, \end{aligned} \quad (2.16)$$

which now has to hold for all $w_h \in \mathbb{W}_h$, $K \in \mathcal{K}_h$ and $i \in \mathcal{I}$. Since for a cell $K \in \mathcal{K}_h$ and one of its Voronoi neighbors $K' \in V(K)$ in general it holds that

$$\tilde{u}_h^{K,i}(x^*) \neq \tilde{u}_h^{K',i}(x^*) \quad (2.17)$$

for $x^* \in K \cap K'$, i.e. \tilde{u}_h^i is double-valued at the interface between K and K' , in order to compute the surface integral we need to introduce the numerical flux function $\mathcal{G}(\tilde{u}_h^{K,i}, \tilde{u}_h^{K',i}, n)$. The numerical flux at a position $x^* \in K \cap K'$ on the interface is obtained by (approximately) solving a Riemann problem in normal direction.

2.1. A D -dimensional ADER-DG scheme with MUSCL-Hancock a-posteriori subcell limiting for non-linear hyperbolic conservation laws

Excursus: The Riemann Problem

Let \mathbf{x}^* be a point on interface ∂K between a cell $K \in \mathcal{K}_h$ and its Voronoi neighbor $K' \in V(K)$ and let \mathbf{n} be the outward pointing unit normal vector at this point. Then to obtain the numerical flux we need to solve the initial boundary value problem (“Riemann problem”)

$$\frac{\partial}{\partial t} [\mathbf{g}]_v + \sum_{d=1}^D \frac{\partial}{\partial x_d} [F(\mathbf{g})]_{vd} [\mathbf{n}]_d = 0 \quad (2.18)$$

along the line $\mathbf{x} = \mathbf{x}^* + \alpha \mathbf{n}$ for $\alpha \in \mathbb{R}$ with discontinuous initial conditions

$$\mathbf{g}(\mathbf{x}^* + \alpha \mathbf{n}, 0) = \begin{cases} \tilde{\mathbf{u}}_h^{K,i} \big|_{\mathbf{x}^*} & \text{if } \alpha < 0 \\ \tilde{\mathbf{u}}_h^{K',i} \big|_{\mathbf{x}^*} & \text{if } \alpha > 0. \end{cases} \quad (2.19)$$

We then evaluate the similarity solution $\tilde{\mathbf{g}}(\alpha/t)$ of the problem and define

$$\left[\mathcal{G} \left(\tilde{\mathbf{u}}_h^{K,i}, \tilde{\mathbf{u}}_h^{K',i}, \mathbf{n} \right) \right]_v := \left[\tilde{\mathbf{g}} \big|_0 \right]_v. \quad (2.20)$$

TODO: In practice use approximate Riemann solvers. See bla for an overview on state of the art approximate Riemann solvers.

Continuing with eq. (2.16), integration by parts in time of the first term and noting that \mathbf{w}_h is constant in time yields the following one-step update scheme for the cell-local time-discrete solution $\tilde{\mathbf{u}}_h^{K,i}$:

$$\begin{aligned} \int_K \left[\tilde{\mathbf{u}}_h^{K,i} \big|_{t_i + \Delta t_i} \right]_v [\mathbf{w}_h^K]_v d\mathbf{x} &= \int_K \left[\tilde{\mathbf{u}}_h^{K,i} \big|_{t_i} \right]_v [\mathbf{w}_h^K]_v d\mathbf{x} + \\ &\quad \int_{t_i}^{t_i + \Delta t_i} \int_K \left[F(\tilde{\mathbf{u}}_h^{K,i}) \right]_{vd} \frac{\partial}{\partial x_d} [\mathbf{w}_h^K]_v d\mathbf{x} dt - \\ &\quad \int_{t_i}^{t_i + \Delta t_i} \int_{\partial K} \left[\mathcal{G}(\tilde{\mathbf{u}}_h^{K,i}, \tilde{\mathbf{u}}_h^{K',i}, \mathbf{n}) \right]_v [\mathbf{w}_h^K]_v ds(\mathbf{x}) dt + \\ &\quad \int_{t_i}^{t_i + \Delta t_i} \int_K \left[S(\tilde{\mathbf{u}}_h^{K,i}) \right]_v [\mathbf{w}_h^K]_v d\mathbf{x} dt. \end{aligned} \quad (2.21)$$

Again we require eq. (2.21) to hold for all $v \in \mathcal{V}$, $\mathbf{w}_h \in \mathbb{W}_h$, $K \in \mathcal{K}_h$ and $i \in \mathcal{I}$. Note, however, that the scheme is incomplete, since we only know $\tilde{\mathbf{u}}_h^i|_t$ at the discrete time steps $t \in \{t_i, t_i + \Delta t_i\}$, not within the open interval, i.e. for $t \in (t_i, t_i + \Delta t_i)$. As commonly done in a DG framework we therefore proceed by replacing $\tilde{\mathbf{u}}_h$ on the interval $(t_i, t_i + \Delta t_i)$ by an approximation $\tilde{\mathbf{q}}_h^i \in \tilde{\mathbb{W}}_h^i$ which we call space-time predictor.

2.1.6 Space-time predictor

To derive a procedure to compute the space-time predictor $\tilde{q}_h^i \in \tilde{W}_h^i$ we again start from the original PDE (2.1), but this time we do not use a spatial test function $w_h \in W_h$, but a space-time test function $\tilde{w}_h^i \in \tilde{W}_h^i$. If we furthermore replace the solution u by the space-time predictor $\tilde{q}_h^i \in \tilde{W}_h^i$, integrate over the space-time element $K \times [t_i, t_i + \Delta t_i]$ and apply the divergence theorem analogously to eq. (2.12) we obtain the following relation:

$$\begin{aligned} & \int_{t_i}^{t_i+\Delta t_i} \int_K \frac{\partial}{\partial t} [\tilde{q}_h^{K,i}]_v [\tilde{w}_h^{K,i}]_v dxdt - \\ & \int_{t_i}^{t_i+\Delta t_i} \int_K [F(\tilde{q}_h^{K,i})]_{vd} \frac{\partial}{\partial x_d} [\tilde{w}_h^{K,i}]_v dxdt + \\ & \int_{t_i}^{t_i+\Delta t_i} \int_{\partial K} [\mathcal{G}(\tilde{q}_h^{K,i}, \tilde{q}_h^{K+i}, n)]_v [\tilde{w}_h^{K,i}]_v ds(x)dt = \\ & \int_{t_i}^{t_i+\Delta t_i} \int_K [S(\tilde{q}_h^{K,i})]_v [\tilde{w}_h^{K,i}]_v dxdt. \end{aligned} \quad (2.22)$$

We require eq. (2.22) to hold for all $v \in \mathcal{V}$, $\tilde{w}_h^i \in \tilde{W}_h^i$, $K \in \mathcal{K}_h$ and $i \in \mathcal{I}$.

The assumption that the solution is balanced, i.e. that there is no net inflow or outflow for cells $K \in \mathcal{K}_h$ allows us to drop the third term. Together with integration by parts in time applied to the first term this yields

$$\begin{aligned} & \int_K [\tilde{q}_h^{K,i}]_{t_i+\Delta t_i} [\tilde{w}_h^{K,i}]_{t_i+\Delta t_i} dx - \int_{t_i}^{t_i+\Delta t_i} \int_K [\tilde{q}_h^{K,i}]_v \frac{\partial}{\partial t} [\tilde{w}_h^{K,i}]_v dxdt = \\ & \int_K [\tilde{q}_h^{K,i}]_{t_i} [\tilde{w}_h^{K,i}]_{t_i} dx + \int_{t_i}^{t_i+\Delta t_i} \int_K [F(\tilde{q}_h^{K,i})]_{vd} \frac{\partial}{\partial x_d} [\tilde{w}_h^{K,i}]_v dxdt + \\ & \int_{t_i}^{t_i+\Delta t_i} \int_K [S(\tilde{q}_h^{K,i})]_v [\tilde{w}_h^{K,i}]_v dxdt, \end{aligned} \quad (2.23)$$

which we require to hold for all $v \in \mathcal{V}$, $\tilde{w}_h^i \in \tilde{W}_h^i$, $K \in \mathcal{K}_h$ and $i \in \mathcal{I}$. In conjunction with the initial condition

$$\tilde{q}_h^{K,i} \Big|_{t_i} = \tilde{u}_h^{K,i} \quad (2.24)$$

and an initial guess

$$\tilde{q}_h^{K,i} \Big|_t = \tilde{u}_h^{K,i} \forall t \in (t_i, t_i + \Delta t_i] \quad (2.25)$$

this relation can be used as a fixed-point iteration to find the cell-local space-time predictor $\tilde{q}_h^{K,i}$.

In the following two sections we will introduce mappings from spatial elements K and space-time elements $K \times [t_i, t_i + \Delta t_i]$ to spatial and space-time

2.1. A D -dimensional ADER-DG scheme with MUSCL-Hancock a-posteriori subcell limiting for non-linear hyperbolic conservation laws

reference cells and orthogonal bases for the spaces \mathbb{W}_h and $\tilde{\mathbb{W}}_h^i$. We will then insert these results into eq. (2.23) and derive a fully-discrete iterative method to compute the cell-local space-time predictor $\tilde{\mathbf{q}}_h^{K,i}$.

2.1.7 Reference Elements and Mappings

Let $\hat{K} := [0, 1]^D$ be the spatial reference element and $\boldsymbol{\xi} \in \hat{K}$ be a point therein. Let $[0, 1]$ be the reference time interval and $\tau \in [0, 1]$ be a point reference time. We can then introduce the following mappings:

Spatial mappings: Let $K \in \mathcal{K}_h$ be a cell in global coordinates with extent $\Delta \mathbf{x}^K$ and “lower-left corner” \mathbf{P}_K , more precisely that is

$$[\Delta \mathbf{x}^K]_d = \max_{x \in K} [x]_d - \min_{x \in K} [x]_d \quad (2.26)$$

and

$$[\mathbf{P}_K]_d = \min_{x \in K} [x]_d \quad (2.27)$$

for $d \in \mathcal{D}$. We can then define a mapping

$$\boldsymbol{\mathcal{X}}_K : \hat{K} \rightarrow K, \boldsymbol{\xi} \mapsto \boldsymbol{\mathcal{X}}_K(\boldsymbol{\xi}) = \mathbf{x} \quad (2.28)$$

via the relation

$$[x]_d = [\boldsymbol{\mathcal{X}}_K(\boldsymbol{\xi})]_d = [\mathbf{P}_K]_d + [\Delta \mathbf{x}]_d [\boldsymbol{\xi}]_d \quad (2.29)$$

for $d \in \mathcal{D}$ (i.e. no summation on d) and for all $x \in K$, $\boldsymbol{\xi} \in \hat{K}$ and $K \in \mathcal{K}_h$.

Temporal mappings: Let $[t_i, t_i + \Delta t_i], i \in \mathcal{I}$ be an interval in global time. The mapping

$$\mathcal{T}_i : [0, 1] \rightarrow [t_i, t_i + \Delta t_i], \tau \mapsto \mathcal{T}_i(\tau) = t_i + \Delta t_i \tau = t \quad (2.30)$$

maps a point $\tau \in [0, 1]$ in reference time to a point $t \in [t_i, t_i + \Delta t_i]$ in global time for all $i \in \mathcal{I}$.

The inverse mappings, the Jacobian matrices and the Jacobi determinants of the mappings are given in the following:

Spatial mappings: The inverse spatial mappings

$$\boldsymbol{\mathcal{X}}_K^{-1} : K \rightarrow \hat{K}, \mathbf{x} \mapsto \boldsymbol{\mathcal{X}}_K^{-1}(\mathbf{x}) = \boldsymbol{\xi} \quad (2.31)$$

are defined via the relation

$$[\boldsymbol{\xi}]_d = [\boldsymbol{\mathcal{X}}_K^{-1}(\mathbf{x})]_d = \frac{1}{[\Delta \mathbf{x}^K]_d} ([x]_d - [\mathbf{P}_K]_d) \quad (2.32)$$

for $d \in \mathcal{D}$ and for all $\xi \in \hat{K}$, $x \in K$ and $K \in \mathcal{K}_h$. The Jacobian of \mathcal{X}_K is found to be

$$\left[\frac{\partial \mathcal{X}_K}{\partial \xi} \right]_{dd'} = \frac{\partial [\mathcal{X}_K]_d}{\partial \xi_{d'}} = [\Delta x^K]_d \delta_{dd'}, \quad (2.33)$$

where $d, d' \in \mathcal{D}$ (i.e. no summation on d) and for all $K \in \mathcal{K}_h$. As usual $\delta_{dd'}$ denotes the Kronecker delta defined as

$$\delta_{dd'} = \begin{cases} 0 & \text{if } d \neq d' \\ 1 & \text{if } d = d'. \end{cases} \quad (2.34)$$

The Jacobi determinant of \mathcal{X}_K for $K \in \mathcal{K}_h$ then simply is

$$J_{\mathcal{X}_K} = \left\| \frac{\partial \mathcal{X}_K}{\partial \xi} \right\| = \prod_{d=1}^D [\Delta x^K]_d, \quad (2.35)$$

i.e. the determinant is constant for all $\xi \in \hat{K}$.

Temporal mappings: The inverse temporal mappings are given as

$$\mathcal{T}_i^{-1} : [t_i, t_i + \Delta t_i] \rightarrow [0, 1], t \mapsto \mathcal{T}_i^{-1}(t) = \frac{t - t_i}{\Delta t_i} = \tau \quad (2.36)$$

for all $\tau \in [0, 1]$, $t \in [t_i, t_i + \Delta t_i]$ and $i \in \mathcal{I}$. In the trivial case of a one-dimensional mapping the Jacobian of \mathcal{T}_i is a scalar which in turn is its own determinant. One finds

$$\frac{d\mathcal{T}_i}{d\tau} = \Delta t_i = J_{\mathcal{T}_i} \quad (2.37)$$

which again is constant for all $\tau \in [0, 1]$.

2.1.8 Orthogonal Bases for the Finite-Dimensional Function Spaces

In section 2.1.5 we introduced finite-dimensional, cell-wise polynomial function spaces \mathbb{W}_h and $\tilde{\mathbb{W}}_h^i$ for spatial and space-time ansatz and test functions, respectively. On our way towards a fully discrete version of the relations (2.23) and (2.21) to obtain the space-time predictor and the solution at the next time step, respectively, we will now derive a set of functions that form bases for the two function spaces of interest. Following the approach presented by Dumbars et al. in ??, throughout the thesis we will use the set of Lagrange functions with nodes located at the roots of the Legendre polynomials and tensor products thereof. In the later chapters of this work it will become obvious why this particular choice is highly favorable. For the moment the two major reasons shall be stated as an outlook:

2.1. A D -dimensional ADER-DG scheme with MUSCL-Hancock a-posteriori subcell limiting for non-linear hyperbolic conservation laws

1. Numerical integration using the Gauss-Legendre method is simple and computationally cheap, since the function values at the Gauss-Legendre nodes are directly available as they are equal to the degrees of freedom representing the local polynomial.
2. The resulting bases are orthogonal, which in turn makes sure that the resulting DG-matrices exhibit a sparse block structure allowing computations to be carried out efficiently in a dimension-by-dimension manner.

Lagrange Interpolation

Let $f \in \mathcal{Q}_N([0, 1])$ be a polynomial of degree less or equal than N and for the index set $\mathcal{N} := \{0, 1, \dots, N\}$ let $\{\hat{\xi}_n\}_{n \in \mathcal{N}}$ be a set of distinct nodes in $[0, 1]$. Then the Lagrange interpolation of f ,

$$\hat{f}(\xi) = \sum_{n=0}^N L_n(\xi) f(\xi_n) \quad (2.38)$$

with Lagrange functions

$$L_n(\xi) = \prod_{m=0, m \neq n}^N \frac{\xi - \hat{\xi}_m}{\hat{\xi}_n - \hat{\xi}_m} \quad (2.39)$$

is exact, i.e.

$$f(\xi) = \hat{f}(\xi) \quad \forall \xi \in [0, 1]. \quad (2.40)$$

Since therefore every polynomial $f \in \mathcal{Q}_N([0, 1])$ can be represented as a linear combination of the Legendre polynomials L_n , $n \in \mathcal{N}$, the set of functions $\{L_n\}_{n \in \mathcal{N}}$ is a basis of $\mathcal{Q}_N([0, 1])$.

The following observation is an important property of the Lagrange polynomials:

$$L_n(\hat{\xi}_{n'}) = \delta_{nn'}, \quad (2.41)$$

i.e. at each node $\hat{\xi}_n$ only L_n has value 1 and all other polynomials evaluate to 0.

Legendre Polynomials and Gauss-Legendre Integration

Let $P_0 : [-1, 1] \rightarrow \mathbb{R}, \xi \mapsto 1$ and $P_1 : [-1, 1] \rightarrow \mathbb{R}, \xi \mapsto \xi$ be the zeroth and the first Legendre polynomial, respectively. Then the $N + 1$ -st Legendre polynomial can be defined via the following recurrence relation:

$$P_{N+1}(\xi) = \frac{1}{N+1} ((2N+1)P_N(\xi) - nP_{N-1}(\xi)). \quad (2.42)$$

2. THEORY

Let $\{\tilde{\xi}_n\}_{n \in \mathcal{N}}$ be the roots of the $N + 1$ -st Legendre polynomial L_{N+1} . Then $\{\hat{\xi}_n\}_{n \in \mathcal{N}}$ with

$$\hat{\xi}_n = \frac{1}{2}(\tilde{\xi}_n + 1) \quad (2.43)$$

are the roots of the $N + 1$ -st Legendre polynomial linearly mapped to the interval $(0, 1)$. In conjunction with a set of suitable weights $\{\hat{\omega}_n\}_{n \in \mathcal{N}}$ Gauss-Legendre integration can be used to integrate polynomials of degree up to $2N + 1$ over the interval $[0, 1]$ exactly, i.e.

$$\int_0^1 f(\xi) d\xi = \sum_{n=0}^N \hat{\omega}_n f(\hat{\xi}_n) \quad \forall f \in \mathbb{Q}_{2N+1}([0, 1]). \quad (2.44)$$

A script on how to find the weights $\{\hat{\xi}_n\}_{n \in \mathcal{N}}$ can be found in appendix XXX.

Scalar-valued Basis Functions on the One-dimensional Reference Element

Let $\{\hat{\psi}_n\}_{n \in \mathcal{N}}$ be the set of $N + 1$ Lagrange polynomials with nodes at the roots of the $N + 1$ -st Legendre polynomial linearly mapped to the interval $[0, 1]$, i.e.

$$\hat{\psi}_n(x) = \sum_{n'=0}^N \frac{x - \hat{x}_{n'}}{\hat{x}_n - \hat{x}_{n'}} \quad (2.45)$$

for $n \in \mathcal{N}$. Since $\{\hat{\psi}_n\}_{n \in \mathcal{N}}$ are Lagrange polynomials and the roots $\{\hat{x}_n\}_{n \in \mathcal{N}}$ are distinct the set is a basis of $\mathbb{Q}_N([0, 1])$. Since furthermore

$$\langle \hat{\psi}_n, \hat{\psi}_m \rangle_{L^2([0, 1])} = \int_0^1 \hat{\psi}_n(x) \hat{\psi}_m(x) dx = \sum_{n'=0}^N \hat{\omega}'_n \hat{\psi}_n(\hat{x}_{n'}) \hat{\psi}_m(\hat{x}_{n'}) = \hat{\omega}_n \delta_{mn} \quad (2.46)$$

for all $m, n \in \mathcal{N}$ (i.e. no summation over n), the set is even an orthogonal basis of $\mathbb{Q}_N([0, 1])$ with respect to the L^2 -scalar product as defined above. In this derivation we used the fact that $\hat{\psi}_n \hat{\psi}_m$ has degree $2N$ and that Gauss-Legendre integration with $N + 1$ nodes is exact for polynomials up to degree $2N + 1$.

Scalar-valued Basis Functions on the Spatial Reference Element

For the vector-valued index set $\mathcal{N} := \{0, 1, \dots, N\}^D$ let us define the set of scalar-valued spatial basis functions $\{\hat{\phi}_n\}_{n \in \mathcal{N}}$ on $\hat{K} := [0, 1]^D$ as

$$\hat{\phi}_n(\xi) = \prod_{d=1}^D \hat{\psi}_{[n]_d}([\xi]_d) = \hat{\psi}_{[n]_d}([\xi]_d), \quad (2.47)$$

2.1. A D -dimensional ADER-DG scheme with MUSCL-Hancock a-posteriori subcell limiting for non-linear hyperbolic conservation laws

i.e. $\{\hat{\phi}_n\}_{n \in \mathcal{N}}$ is the tensor product of $\{\hat{\psi}_n\}_{n \in \mathcal{N}}$ and as such it is a basis of $\mathbb{Q}([0, 1]^D) = \mathbb{Q}(\hat{K})$. If we define

$$\left[\hat{\xi}_n\right]_d = \hat{\xi}_{[n]_d} \quad (2.48)$$

and

$$\prod_{d=1}^D \hat{\omega}_{[n]_d}, \quad (2.49)$$

for all $d \in \mathcal{V}$ and $n \in \mathcal{N}$, we furthermore observe that the basis is orthogonal with respect to the L^2 -scalar product, since

$$\begin{aligned} \left\langle \hat{\phi}_n, \hat{\phi}_m \right\rangle_{L^2(\hat{K})} &= \int_{\hat{K}} \hat{\phi}_n(\xi) \hat{\phi}_m(\xi) d\xi = \\ \sum_{n' \in \mathcal{N}} \left(\hat{\omega}_{n'} \hat{\phi}_n(\hat{\xi}_{n'}) \hat{\phi}_m(\hat{\xi}_{n'}) \right) &= \hat{\omega}_n \delta_{nm} \end{aligned} \quad (2.50)$$

for all $n, m \in \mathcal{N}$. The natural extensions of the Kronecker delta for vector-valued indices is defined as follows:

$$\delta_{nm} = \prod_{d=1}^D \delta_{[n]_d [m]_d} = \delta_{[n]_d [m]_d}. \quad (2.51)$$

Scalar-valued Basis Functions on the Space-time Reference Element

Analogously to the procedure illustrated above for the spatial reference element \hat{K} we can define a basis $\{\hat{\theta}_{nl}\}_{n \in \mathcal{N}, l \in \mathcal{N}}$ of $\mathbb{Q}_N(\hat{K} \times [0, 1])$ on the reference space-time element $\hat{K} \times [0, 1]$ as

$$\hat{\theta}_{nl}(\xi, \tau) = \hat{\phi}_n(\xi) \hat{\psi}_l(\tau), \quad (2.52)$$

which again is orthogonal, since

$$\left\langle \hat{\theta}_{nl}, \hat{\theta}_{mk} \right\rangle_{L^2(\hat{K} \times [0, 1])} = \int_0^1 \int_{\hat{K}} \hat{\theta}_{nl} \hat{\theta}_{mk} d\xi d\tau = \hat{\omega}_n \hat{\omega}_l \delta_{nm} \delta_{lk} \quad (2.53)$$

for all $n, m \in \mathcal{N}$ and $l, k \in \mathcal{N}$.

Vector-valued Basis Functions on the Spatial Reference Element

If we define $\{\hat{\phi}_{nv}\}_{n \in \mathcal{N}, v \in \mathcal{V}}$ as

$$\hat{\phi}_{nv} = \hat{\phi}_n e_v, \quad (2.54)$$

where e_v is the v -th unit vector, i.e.

$$[e_v]_{v'} = \delta_{vv'} \quad (2.55)$$

for $v, v' \in \mathcal{V}$. Since

$$\begin{aligned} \langle \hat{\phi}_{nv}, \hat{\phi}_{n'v'} \rangle_{L^2(\hat{K})^V} &= \int_{\hat{K}} [\hat{\phi}_{nv}]_j [\hat{\phi}_{n'v'}]_j d\boldsymbol{\xi} = \\ &= ([e_v]_j [e_{v'}]_j) \int_0^1 \int_{\hat{K}} \hat{\phi}_n \hat{\phi}_{n'} d\boldsymbol{\xi} = \hat{\omega}_n \delta_{nn'} \delta_{vv'} \end{aligned} \quad (2.56)$$

for all $n, n' \in \mathcal{N}$ and $v, v' \in \{1, 2, \dots, V\}$ the set is an orthogonal basis for $Q_N(\hat{K})^V$.

Vector-valued Basis Functions on the Space-time Reference Element

The set $\{\hat{\theta}_{nlv}\}_{n \in \mathcal{N}, l \in \mathcal{N}, v \in \mathcal{V}}$ defined as

$$\hat{\theta}_{nlv}(\boldsymbol{\xi}, \tau) = \hat{\theta}_{nl}(\boldsymbol{\xi}, \tau) e_v = \hat{\phi}_n(\boldsymbol{\xi}) \hat{\psi}_l(\tau) e_v \quad (2.57)$$

is a basis of $Q_N(\hat{K} \times [0, 1])^V$. Since furthermore

$$\langle \hat{\theta}_{nlv}, \hat{\theta}_{n'l'v'} \rangle_{L^2(\hat{K} \times [0, 1])^V} = \int_0^1 \int_{\hat{K}} [\hat{\theta}_{nlv}]_j [\hat{\theta}_{n'l'v'}]_j d\boldsymbol{\xi} d\tau = \hat{\omega}_n \hat{\omega}_l \delta_{nn'} \delta_{ll'} \delta_{vv'}, \quad (2.58)$$

for all $n, n' \in \mathcal{N}$, $l, l' \in \mathcal{N}$ and $v, v' \in \mathcal{V}$, the set is an orthogonal basis with respect to the respective L^2 -scalar product.

2.1.9 Basis Functions in Global Coordinates

We can use the mappings derived in ch. 2.1.7 to map the basis functions to global coordinates. For the vector-valued basis functions on a spatial element K we obtain

$$\boldsymbol{\phi}_{nv}^K(\mathbf{x}) = \begin{cases} \left(\hat{\phi}_{nv} \circ \boldsymbol{\chi}_K^{-1} \right) (\mathbf{x}) & \text{if } \mathbf{x} \in K \\ 0 & \text{otherwise,} \end{cases} \quad (2.59)$$

and for the vector-valued basis functions on a space-time element $K \times [t_i, t_i + \Delta t_i]$ we have

$$\boldsymbol{\theta}_{nlv}^{Ki}(\mathbf{x}, t) = \begin{cases} \left(\hat{\theta}_{nlv} \circ \left(\boldsymbol{\chi}_K^{-1}, \mathcal{T}_i^{-1} \right) \right) (\mathbf{x}, t) & \text{if } \mathbf{x} \in K \text{ and } t \in [t_i, t_i + \Delta t_i] \\ 0 & \text{otherwise} \end{cases} \quad (2.60)$$

for $n \in \mathcal{N}$, $l \in \{0, 1, \dots, N\}$ as well as $v \in \mathcal{V}$ and for all $K \in \mathcal{K}_h$ and $i \in \mathcal{I}$.

2.1.10 A Fully-discrete Iterative Method for the Space-time Predictor

We recall relation (2.25) for the space-time predictor. Plugging in the initial condition (2.24) yields

$$\begin{aligned} & \int_K \left[\tilde{\mathbf{q}}_h^{K,i} \Big|_{t_i+\Delta t_i} \right]_j \left[\tilde{\mathbf{w}}_h^{K,i} \Big|_{t_i+\Delta t_i} \right]_j d\mathbf{x} - \int_{t_i}^{t_i+\Delta t_i} \int_K \left[\tilde{\mathbf{q}}_h^{K,i} \right]_j \frac{\partial}{\partial t} \left[\tilde{\mathbf{w}}_h^{K,i} \right]_j dxdt = \\ & \int_K \left[\tilde{\mathbf{u}}_h^{K,i} \Big|_{t_i} \right]_j \left[\tilde{\mathbf{w}}_h^{K,i} \Big|_{t_i} \right]_j d\mathbf{x} + \int_{t_i}^{t_i+\Delta t_i} \int_K \left[\mathbf{F}(\tilde{\mathbf{q}}_h^{K,i}) \right]_{jk} \frac{\partial}{\partial x_k} \left[\tilde{\mathbf{w}}_h^{K,i} \right]_j dxdt + \\ & \int_{t_i}^{t_i+\Delta t_i} \int_K \left[\mathbf{S}(\tilde{\mathbf{q}}_h^{K,i}) \right]_j \left[\tilde{\mathbf{w}}_h^{K,i} \right]_j dxdt, \end{aligned} \quad (2.61)$$

which we require to hold for all $\tilde{\mathbf{w}}_h \in \tilde{\mathbb{W}}_h$, $K \in \mathcal{K}_h$ and $i \in \mathcal{I}$.

Making use of the bases we derived in the previous section the cell-local space-time predictor $\tilde{\mathbf{q}}_h^{K,i}$ can be represented by a tensor of coefficients $\hat{\mathbf{q}}^{K,i}$ ("degrees of freedom") as follows:

$$\tilde{\mathbf{q}}_h^{K,i} = \left[\hat{\mathbf{q}}^{K,i} \right]_{nlv} \boldsymbol{\theta}_{nlv}^{Ki}. \quad (2.62)$$

The initial condition $\tilde{\mathbf{u}}_h^{K,i} \Big|_{t_i}$ can be represented as

$$\tilde{\mathbf{u}}_h^{K,i} \Big|_{t_i} = \left[\hat{\mathbf{u}}^{K,i} \right]_{nv} \boldsymbol{\phi}_{nv}^K, \quad (2.63)$$

where

$$\left[\hat{\mathbf{u}}^{K,i} \right]_{nv} = \left[\tilde{\mathbf{u}}_h^{K,i} \Big|_{(\mathcal{X}_K(\xi_n), t_i)} \right]_v. \quad (2.64)$$

2. THEORY

Inserting eqs. (2.62) and (2.63) into eq. (2.61) and introduction of the iteration index $r \in \{0, 1, \dots, R\}$ leads to the following iterative scheme for the degrees of freedom of the cell-local space-time predictor:

$$\begin{aligned}
 & \underbrace{\int_K \left[\left[\hat{\mathbf{q}}^{K,i,r+1} \right]_{nlv} \boldsymbol{\theta}_{nlv}^{Ki} \Big|_{t_i+\Delta t_i} \right]_j \left[\boldsymbol{\theta}_{\alpha\beta\gamma}^{Ki} \Big|_{t_i+\Delta t_i} \right]_j dx}_{\text{S-I}} - \\
 & \underbrace{\int_{t_i}^{t_i+\Delta t_i} \int_K \left[\left[\hat{\mathbf{q}}^{K,i,r+1} \right]_{nlv} \boldsymbol{\theta}_{nlv}^{Ki} \right]_j \frac{\partial}{\partial t} \left[\boldsymbol{\theta}_{\alpha\beta\gamma}^{Ki} \right]_j dx dt}_{\text{S-II}} = \\
 & \underbrace{\int_K \left[\left[\hat{\mathbf{u}}^{K,i} \right]_{nv} \boldsymbol{\phi}_{nv}^K \right]_j \left[\boldsymbol{\theta}_{\alpha\beta\gamma}^{Ki} \Big|_{t_i} \right]_j dx}_{\text{S-III}} + \\
 & \underbrace{\int_{t_i}^{t_i+\Delta t_i} \int_K \left[\mathbf{F} \left(\left[\hat{\mathbf{q}}^{K,i,r} \right]_{nlv} \boldsymbol{\theta}_{nlv}^{Ki} \right) \right]_{jk} \frac{\partial}{\partial x_k} \left[\boldsymbol{\theta}_{\alpha\beta\gamma}^{Ki} \right]_j dx dt}_{\text{S-IV}} + \\
 & \underbrace{\int_{t_i}^{t_i+\Delta t_i} \int_K \left[\mathbf{S} \left(\left[\hat{\mathbf{q}}^{K,i,r} \right]_{nlv} \boldsymbol{\theta}_{nlv}^{Ki} \right) \right]_j \left[\boldsymbol{\theta}_{\alpha\beta\gamma}^{Ki} \right]_j dx dt}_{\text{S-V}}. \tag{2.65}
 \end{aligned}$$

We require this relation to hold for all $\alpha \in \mathcal{N}$, $\beta \in \mathcal{N}$ and $\gamma \in \mathcal{V}$.

As initial condition, i.e. for $r = 0$, we use

$$\left[\hat{\mathbf{q}}^{K,i,0} \right]_{nvl} = \left[\hat{\mathbf{u}}^{K,i} \right]_{nv} \tag{2.66}$$

for all time degrees of freedom $l \in \mathcal{N}$.

We will now proceed in a term-by-term fashion to rewrite all integrals with respect to reference coordinates so that we can finally derive a complete rule on how to compute $\hat{\mathbf{q}}^{K,i,r+1}$ that holds for all $K \in \mathcal{K}_h$ and $i \in \mathcal{I}$.

2.1. A D -dimensional ADER-DG scheme with MUSCL-Hancock a-posteriori subcell limiting for non-linear hyperbolic conservation laws

Term S-I

The first term of eq. (2.65) can be rewritten with respect to reference coordinates as follows:

$$\begin{aligned}
& \int_K \left[\left[\hat{\mathbf{q}}^{K,i,r+1} \right]_{nlv} \boldsymbol{\theta}_{nlv}^{Ki} \Big|_{t_i+\Delta t_i} \right]_j \left[\boldsymbol{\theta}_{\alpha\beta\gamma}^{Ki} \right]_{t_i+\Delta t_i} d\mathbf{x} = \\
& \int_K \left[\hat{\mathbf{q}}^{K,i,r+1} \right]_{nlv} \phi_n^K \left(\psi_l^i \Big|_{t_i+\Delta t_i} \right) [\mathbf{e}_v]_j \phi_\alpha^K \left(\psi_\beta^i \Big|_{t_i+\Delta t_i} \right) [\mathbf{e}_\gamma]_j d\mathbf{x} = \\
& J_{\mathcal{X}_K} \int_{\hat{K}} \left[\hat{\mathbf{q}}^{K,i,r+1} \right]_{nlv} \hat{\phi}_n \left(\hat{\psi}_l \Big|_1 \right) [\mathbf{e}_v]_j \hat{\phi}_\alpha \left(\hat{\psi}_\beta \Big|_1 \right) [\mathbf{e}_\gamma]_j d\hat{\mathbf{x}} = \\
& J_{\mathcal{X}_K} \sum_{\alpha' \in \mathcal{N}} \left(\hat{\omega}_{\alpha'} \left[\hat{\mathbf{q}}^{K,i,r+1} \right]_{nlv} \hat{\phi}_n(\hat{\xi}_{\alpha'}) \left(\hat{\psi}_l \Big|_1 \right) [\mathbf{e}_v]_j \hat{\phi}_\alpha(\hat{\xi}_{\alpha'}) \left(\hat{\psi}_\beta \Big|_1 \right) [\mathbf{e}_\gamma]_j \right) = \\
& J_{\mathcal{X}_K} \sum_{\alpha' \in \mathcal{N}} \left(\hat{\omega}_{\alpha'} \left[\hat{\mathbf{q}}^{K,i,r+1} \right]_{nlv} \delta_{n\alpha'} \left(\hat{\psi}_l \Big|_1 \right) \delta_{vj} \delta_{\alpha\alpha'} \left(\hat{\psi}_\beta \Big|_1 \right) \delta_{j\gamma} \right) = \\
& J_{\mathcal{X}_K} \hat{\omega}_\alpha \left[\hat{\psi}_\beta \Big|_1 \hat{\psi}_l \Big|_1 \right] \left[\hat{\mathbf{q}}^{K,i,r+1} \right]_{\alpha l \gamma} = \\
& J_{\mathcal{X}_K} \hat{\omega}_\alpha [\mathbf{R}]_{\beta,l} \left[\hat{\mathbf{q}}^{K,i,r+1} \right]_{\alpha l \gamma},
\end{aligned} \tag{2.67}$$

where we remember from eq. (2.35) that

$$J_{\mathcal{X}_K} = \prod_{d=1}^D [\Delta \mathbf{x}]_d \tag{2.68}$$

and we define the matrix \mathbf{R} representing the Right Reference Element Mass Operator as

$$[\mathbf{R}]_{i,j} := \left[\hat{\psi}_i \Big|_1 \hat{\psi}_j \Big|_1 \right]_{i,j} \tag{2.69}$$

for $i, j \in \mathcal{N}$. A Python script to compute \mathbf{R} can be found in appendix ??.

Term S-II

The second term of eq. (2.65) can be rewritten with respect to reference coordinates as follows:

$$\begin{aligned}
 & \int_{t_i}^{t_i+\Delta t_i} \int_K \left[\hat{\mathbf{q}}^{K,i,r+1} \right]_{nlv} \boldsymbol{\theta}_{nlv}^{Ki} \left[\frac{\partial}{\partial t} \left[\boldsymbol{\theta}_{\alpha\beta\gamma}^{Ki} \right]_j \right] dx dt = \\
 & \int_{t_i}^{t_i+\Delta t_i} \int_K \left[\hat{\mathbf{q}}^{K,i,r+1} \right]_{nlv} \phi_n^K \psi_l^i [e_v]_j \phi_\alpha^K \left(\frac{\partial}{\partial t} \psi_\beta^i \right) [e_\gamma]_j dx dt = \\
 & J_{\mathcal{T}_i} J_{\mathcal{X}_K} \int_0^1 \int_{\hat{K}} \left[\hat{\mathbf{q}}^{K,i,r+1} \right]_{nlv} \hat{\phi}_n \hat{\psi}_l [e_v]_j \hat{\phi}_\alpha \left(\frac{1}{\Delta t_i} \frac{\partial}{\partial \tau} \hat{\psi}_\beta \right) [e_\gamma]_j d\xi d\tau = \\
 & J_{\mathcal{T}_i} J_{\mathcal{X}_K} \sum_{\alpha' \in \mathcal{N}} \sum_{\beta' \in \mathcal{N}} \left(\hat{\omega}_{\alpha'} \hat{\omega}_{\beta'} \left[\hat{\mathbf{q}}^{K,i,r+1} \right]_{nlv} \hat{\phi}_n(\hat{\xi}_{\alpha'}) \hat{\psi}_l(\hat{\tau}_{\beta'}) [e_v]_j \dots \right. \\
 & \quad \left. \dots \hat{\phi}_\alpha(\hat{\xi}_{\alpha'}) \left(\frac{\partial}{\partial \tau} \hat{\psi}_\beta(\hat{\tau}_{\beta'}) \right) [e_\gamma]_j \right) = \\
 & J_{\mathcal{T}_i} J_{\mathcal{X}_K} \sum_{\alpha' \in \mathcal{N}} \sum_{\beta' \in \mathcal{N}} \left(\hat{\omega}_{\alpha'} \hat{\omega}_{\beta'} \left[\hat{\mathbf{q}}^{K,i,r+1} \right]_{nlv} \delta_{n\alpha'} \delta_{l\beta'} \delta_{vj} \dots \right. \\
 & \quad \left. \dots \delta_{\alpha\alpha'} \left(\frac{1}{\Delta t_i} \frac{\partial}{\partial \tau} \hat{\psi}_\beta(\hat{\tau}_{\beta'}) \right) \delta_{\gamma j} \right) = \\
 & J_{\mathcal{T}_i} J_{\mathcal{X}_K} \hat{\omega}_\alpha \frac{1}{\Delta t_i} \sum_{\beta' \in \mathcal{N}} \left(\hat{\omega}_{\beta'} \left[\frac{\partial}{\partial \tau} \hat{\psi}_\beta(\hat{\tau}_{\beta'}) \right] \left[\hat{\mathbf{q}}^{K,i,r+1} \right]_{\alpha\beta'\gamma} \right) = \\
 & J_{\mathcal{T}_i} J_{\mathcal{X}_K} \hat{\omega}_\alpha \frac{1}{\Delta t_i} [\mathbf{K}]_{\beta,\beta'} \left[\hat{\mathbf{q}}^{K,i,r+1} \right]_{\alpha\beta'\gamma} \tag{2.70}
 \end{aligned}$$

where we remember from eq. (2.37) that

$$J_{\mathcal{T}_i} = \Delta t_i, \tag{2.71}$$

so that Δt_i and $1/\Delta t_i$ in eq. (2.70) cancel. In the derivation we made use of the fact that due to the chain rule

$$\frac{\partial}{\partial t} \psi_\beta^i = \frac{\partial}{\partial t} (\hat{\psi}_\beta \circ \mathcal{T}_i^{-1}) = \left(\frac{\partial}{\partial \tau} \hat{\psi}_\beta \right) \left(\frac{\partial}{\partial t} \mathcal{T}_i^{-1} \right) = \frac{1}{\Delta t_i} \frac{\partial}{\partial \tau} \hat{\psi}_\beta. \tag{2.72}$$

We furthermore introduce the matrix \mathbf{K} representing the Reference Element Stiffness Operator given as

$$[\mathbf{K}]_{ij} = \hat{\omega}_j \frac{\partial}{\partial \tau} \hat{\psi}_i(\hat{\tau}_j) \tag{2.73}$$

for $i, j \in \mathcal{N}$. A Python script to compute \mathbf{K} can be found in appendix ??.

Term S-III

The third term of eq. (2.65) can be rewritten with respect to reference coordinates as follows:

$$\begin{aligned}
 & \int_K \left[[\hat{\mathbf{u}}^{K,i}]_{nv} \boldsymbol{\phi}_{nv}^K \right]_j \left[\boldsymbol{\theta}_{\alpha\beta\gamma}^{Ki} \Big|_{t_i} \right]_j d\mathbf{x} = \\
 & \int_K [\hat{\mathbf{u}}^{K,i}]_{nv} \boldsymbol{\phi}_n^K [e_v]_j \boldsymbol{\phi}_\alpha^K \left(\psi_\beta^i \Big|_{t_i} \right) [e_\gamma]_j d\mathbf{x} = \\
 & J\mathcal{X}_K \int_{\hat{K}} [\hat{\mathbf{u}}^{K,i}]_{nv} \hat{\phi}_n [e_v]_j \hat{\phi}_\alpha \left(\hat{\psi}_\beta \Big|_0 \right) [e_\gamma]_j d\hat{\xi} = \\
 & J\mathcal{X}_K \sum_{\alpha' \in \mathcal{N}} \left(\hat{\omega}_{\alpha'} [\hat{\mathbf{u}}^{K,i}]_{nv} \hat{\phi}_n(\xi_{\alpha'}) [e_v]_j \hat{\phi}_\alpha(\xi_{\alpha'}) \left(\hat{\psi}_\beta \Big|_0 \right) [e_\gamma]_j \right) = \\
 & J\mathcal{X}_K \sum_{\alpha' \in \mathcal{N}} \left(\hat{\omega}_{\alpha'} [\hat{\mathbf{u}}^{K,i}]_{nv} \delta_{n\alpha'} \delta_{vj} \delta_{\alpha\alpha'} \left(\hat{\psi}_\beta \Big|_0 \right) \delta_{\gamma j} \right) = \\
 & J\mathcal{X}_K \hat{\omega}_\alpha \left[\hat{\psi}_\beta \Big|_0 \right] [\hat{\mathbf{u}}^{K,i}]_{\alpha\gamma} = \\
 & J\mathcal{X}_K \hat{\omega}_\alpha [l]_\beta [\hat{\mathbf{u}}^{K,i}]_{\alpha\gamma}, \tag{2.74}
 \end{aligned}$$

where we define the vector \mathbf{l} representing the Left Reference Time Flux Operator as

$$[l]_i = \hat{\psi}_i \Big|_0 \tag{2.75}$$

for $i \in \mathcal{N}$. A Python script to compute \mathbf{l} can be found in appendix ??.

Term S-IV

The third term of eq. (2.65) can be rewritten with respect to reference coordinates as follows:

$$\begin{aligned}
& \int_{t_i}^{t_i+\Delta t_i} \int_K \left[\mathbf{F} \left(\left[\hat{\mathbf{q}}^{K,i,r} \right]_{nlv} \boldsymbol{\theta}_{nlv}^{Ki} \right) \right]_{jk} \frac{\partial}{\partial x_k} \left[\boldsymbol{\theta}_{\alpha\beta\gamma}^{Ki} \right]_j d\mathbf{x} dt = \\
& \int_{t_i}^{t_i+\Delta t_i} \int_K \left[\mathbf{F} \left(\left[\hat{\mathbf{q}}^{K,i,r} \right]_{nlv} \phi_n^K \psi_l^i \mathbf{e}_v \right) \right]_{jk} \left(\prod_{d=1, d \neq k}^D \psi_{[\alpha]_d}^K([\mathbf{x}]_d) \right) \psi_\beta^i(t) [\mathbf{e}_\gamma]_j \dots \\
& \dots \left(\frac{\partial}{\partial x_k} \psi_{[\alpha]_k}^K \right) d\mathbf{x} dt = \\
& J_{\mathcal{T}_i} J_{\mathcal{X}_K} \int_0^1 \int_{\hat{K}} \left[\mathbf{F} \left(\left[\hat{\mathbf{q}}^{K,i,r} \right]_{nlv} \hat{\phi}_n \hat{\psi}_l \mathbf{e}_v \right) \right]_{jk} \left(\prod_{d=1, d \neq k}^D \hat{\psi}_{[\alpha]_d}([\boldsymbol{\xi}]_d) \right) \hat{\psi}_\beta(t) [\mathbf{e}_\gamma]_j \dots \\
& \dots \left(\frac{1}{[\Delta \mathbf{x}]_k} \frac{\partial}{\partial \xi_k} \hat{\psi}_{[\alpha]_k}([\boldsymbol{\xi}]_k) \right) d\boldsymbol{\xi} d\tau = \\
& J_{\mathcal{T}_i} J_{\mathcal{X}_K} \sum_{\alpha' \in \mathcal{N}} \sum_{\beta' \in \mathcal{N}} \left(\hat{\omega}_{\alpha'} \hat{\omega}_{\beta'} \left[\mathbf{F} \left(\left[\hat{\mathbf{q}}^{K,i,r} \right]_{nlv} \hat{\phi}_n(\hat{\boldsymbol{\xi}}_{\alpha'}) \hat{\psi}_l(\hat{\tau}_{\beta'}) \mathbf{e}_v \right) \right]_{jk} \dots \right. \\
& \dots \left(\prod_{d=1, d \neq k}^D \hat{\psi}_{[\alpha]_d}([\hat{\boldsymbol{\xi}}_{\alpha'}]_d) \right) \hat{\psi}_\beta(\hat{\tau}_{\beta'}) [\mathbf{e}_\gamma]_j \left(\frac{1}{[\Delta \mathbf{x}]_k} \frac{\partial}{\partial \xi_k} \hat{\psi}_{[\alpha]_k}([\hat{\boldsymbol{\xi}}_{\alpha'}]_k) \right) \left. \right) = \\
& J_{\mathcal{T}_i} J_{\mathcal{X}_K} \sum_{\alpha' \in \mathcal{N}} \sum_{\beta' \in \mathcal{N}} \left(\hat{\omega}_{\alpha'} \hat{\omega}_{\beta'} \left[\mathbf{F} \left(\left[\hat{\mathbf{q}}^{K,i,r} \right]_{nlv} \delta_{n\alpha'} \delta_{l\beta'} \mathbf{e}_v \right) \right]_{jk} \dots \right. \\
& \dots \left(\prod_{d=1, d \neq k}^D \delta_{[\alpha]_d[\alpha']_d} \right) \delta_{\beta\beta'} \delta_{\gamma j} \left(\frac{1}{[\Delta \mathbf{x}]_k} \frac{\partial}{\partial \xi_k} \hat{\psi}_{[\alpha]_k}([\hat{\boldsymbol{\xi}}_{\alpha'}]_k) \right) \left. \right) = \\
& J_{\mathcal{T}_i} J_{\mathcal{X}_K} \hat{\omega}_\beta \sum_{k=1}^D \left(\frac{1}{[\Delta \mathbf{x}]_k} \sum_{\alpha'_k \in \{0,1,\dots,N\}} \left(\prod_{d=0, d \neq k}^D \hat{\omega}_{[\alpha]_d} \dots \right. \right. \\
& \dots \hat{\omega}_{\alpha'_k} \left(\frac{\partial}{\partial \xi_k} \hat{\psi}_{[\alpha]_k}([\hat{\boldsymbol{\xi}}_{\alpha'_k}]) \right) \left[\mathbf{F} \left(\left[\hat{\mathbf{q}}^{K,i,r} \right]_{[\alpha]_{1:k-1}, \alpha'_k, [\alpha]_{k+1:N}] \beta v} \right) \right]_{jk} \left. \right) \left. \right) = \\
& J_{\mathcal{T}_i} J_{\mathcal{X}_K} \hat{\omega}_\beta \sum_{k=1}^D \left(\frac{1}{[\Delta \mathbf{x}]_k} \sum_{\alpha'_k \in \{0,1,\dots,N\}} \left(\prod_{d=0, d \neq k}^D \hat{\omega}_{[\alpha]_d} \dots \right. \right. \\
& \dots [K]_{[\alpha]_k, \alpha'_k} \left[\mathbf{F} \left(\left[\hat{\mathbf{q}}^{K,i,r} \right]_{[\alpha_0, \alpha_1, \dots, \alpha_{k-1}, \alpha'_k, \alpha_{k+1}, \dots, \alpha_N] \beta v} \mathbf{e}_v \right) \right]_{\gamma k} \left. \right) \left. \right),
\end{aligned}$$

2.1. A D -dimensional ADER-DG scheme with MUSCL-Hancock a-posteriori subcell limiting for non-linear hyperbolic conservation laws

where we used that

$$\begin{aligned}
\frac{\partial}{\partial x_k} \theta_{\alpha\beta\gamma}^{Ki}(\mathbf{x}, t) &= \left(\frac{\partial}{\partial x_k} \phi_{\alpha}^K(\mathbf{x}) \right) \psi_{\beta}^i(t) \mathbf{e}_{\gamma} = \left(\frac{\partial}{\partial x_k} \prod_{d=1}^D \psi_{[\alpha]_d}^K([\mathbf{x}]_d) \right) \psi_{\beta}^i(t) \mathbf{e}_{\gamma} = \\
&\left(\prod_{d=1, d \neq k}^D \psi_{[\alpha]_d}^K([\mathbf{x}]_d) \right) \left(\frac{\partial}{\partial x_k} \psi_{[\alpha]_k}^K([\mathbf{x}]_k) \right) \psi_{\beta}^i(t) \mathbf{e}_{\gamma} = \\
&\left(\prod_{d=1, d \neq k}^D \psi_{[\alpha]_d}^K([\mathbf{x}]_d) \right) \left(\frac{\partial}{\partial x_k} \hat{\psi}_{[\alpha]_k} \left([\boldsymbol{\chi}_K^{-1}(\mathbf{x})]_k \right) \right) \psi_{\beta}^i(t) \mathbf{e}_{\gamma} = \\
&\left(\prod_{d=1, d \neq k}^D \psi_{[\alpha]_d}^K([\mathbf{x}]_d) \right) \left(\left(\frac{\partial}{\partial \xi_k} \hat{\psi}_{[\alpha]_k} \left([\boldsymbol{\chi}_K^{-1}(\mathbf{x})]_k \right) \right) \left(\frac{\partial}{\partial x_k} [\boldsymbol{\chi}_K^{-1}(\mathbf{x})]_k \right) \right) \dots \\
&\dots \psi_{\beta}^i(t) \mathbf{e}_{\gamma} = \\
&\left(\prod_{d=1, d \neq k}^D \psi_{[\alpha]_d}^K([\mathbf{x}]_d) \right) \left(\frac{1}{[\Delta \mathbf{x}^K]_k} \frac{\partial}{\partial \xi_k} \hat{\psi}_{[\alpha]_k} \left([\boldsymbol{\chi}_K^{-1}(\mathbf{x})]_k \right) \right) \psi_{\beta}^i(t) \mathbf{e}_{\gamma}.
\end{aligned} \tag{2.77}$$

Term S-V

The fifth term of eq. (2.65) can be rewritten with respect to reference coordinates as follows:

$$\begin{aligned}
&\int_{t_i}^{t_i + \Delta t_i} \int_K \left[S \left([\hat{\mathbf{q}}^{K,i,r}]_{nlv} \theta_{nlv}^{Ki} \right) \right]_j [\theta_{\alpha\beta\gamma}^{Ki}]_j d\mathbf{x} dt = \\
&J_{\mathcal{T}_i} J_{\mathcal{X}_K} \int_0^1 \int_{\hat{K}} \left[S \left([\hat{\mathbf{q}}^{K,i,r}]_{nlv} \hat{\phi}_n \hat{\psi}_l \mathbf{e}_v \right) \right]_j \hat{\phi}_{\alpha} \hat{\psi}_l [\mathbf{e}_{\gamma}]_j d\xi d\tau = \\
&J_{\mathcal{T}_i} J_{\mathcal{X}_K} \sum_{\alpha' \in \mathcal{N}} \sum_{\beta' \in \mathcal{N}} \left(\hat{\omega}_{\alpha'} \hat{\omega}_{\beta'} \left[S \left([\hat{\mathbf{q}}^{K,i,r}]_{nlv} \hat{\phi}_n(\xi_{\alpha'}) \hat{\psi}_l(\hat{\tau}_{\beta'}) \mathbf{e}_v \right) \right]_j \dots \right. \\
&\dots \left. \hat{\phi}_{\alpha}(\xi_{\alpha'}) \hat{\psi}_{\beta}(\hat{\tau}_{\beta'}) [\mathbf{e}_{\gamma}]_j \right) = \\
&J_{\mathcal{T}_i} J_{\mathcal{X}_K} \sum_{\alpha' \in \mathcal{N}} \sum_{\beta' \in \mathcal{N}} \left(\hat{\omega}_{\alpha'} \hat{\omega}_{\beta'} \left[S \left([\hat{\mathbf{q}}^{K,i,r}]_{nlv} \delta_{n\alpha'} \delta_{l\beta'} \mathbf{e}_v \right) \right]_j \delta_{\alpha\alpha'} \delta_{\beta\beta'} \delta_{\gamma j} \right) = \\
&J_{\mathcal{T}_i} J_{\mathcal{X}_K} \hat{\omega}_{\alpha} \hat{\omega}_{\beta} \left[S \left([\hat{\mathbf{q}}^{K,i,r}]_{\alpha\beta v} \mathbf{e}_v \right) \right]_{\gamma}
\end{aligned} \tag{2.78}$$

The Complete Fixed-point Iteration

Now collecting the results from eqs. (2.67), (2.70), (2.74), (2.76) and (2.78) and plugging them back into eq. (2.65) and division by $J_{\mathcal{X}_K}$ and $\hat{\omega}_\alpha$ yields

$$\begin{aligned}
 (\mathbf{R} - \mathbf{K})_{\beta\beta'} [\hat{\mathbf{q}}^{K,i,r+1}]_{\alpha\beta'\gamma} &= [\mathbf{I}]_\beta [\hat{\mathbf{u}}^{K,i}]_{\alpha\gamma} + \\
 J_{\mathcal{T}_i} \frac{\hat{\omega}_\beta}{\hat{\omega}_\alpha} \sum_{k=1}^D &\left(\frac{1}{[\Delta \mathbf{x}]_k} \sum_{\alpha'_k \in \{0,1,\dots,N\}} \left(\prod_{d=0, d \neq k}^D \hat{\omega}_{[\alpha]_d} \dots \right. \right. \\
 \dots [\mathbf{K}]_{[\alpha]_k, \alpha'_k} &\left[F \left([\hat{\mathbf{q}}^{K,i,r}]_{[\alpha_0, \alpha_1, \dots, \alpha_{k-1}, \alpha'_k, \alpha_{k+1}, \dots, \alpha_N] \beta v} \mathbf{e}_v \right) \right]_{\gamma k} \left. \right) \Bigg) + \\
 J_{\mathcal{T}_i} \hat{\omega}_\beta &\left[S \left([\hat{\mathbf{q}}^{K,i,r}]_{\alpha\beta v} \mathbf{e}_v \right) \right]_\gamma, \tag{2.79}
 \end{aligned}$$

which has to hold for all $\alpha \in \mathcal{N}$, $\beta \in \mathcal{N}$ and $\gamma \in \mathcal{V}$. To speed up the computation of the new iterate $\hat{\mathbf{q}}^{K,i,r+1}$ we can compute the inverse $(\mathbf{R} - \mathbf{K})^{-1}$ prior to the simulation. A Python script to compute the matrix can be found in appendix ??.

A possible termination criterion could be $\Delta < \varepsilon$, where $\varepsilon > 0$ is a suitable constant related to the desired accuracy of the iteration, e.g. $\varepsilon = 10^{-7}$ and the squared element-wise residual Δ^2 is defined as follows:

$$\Delta^2 = \sum_{n \in \mathcal{N}} \sum_{l \in \mathcal{N}} \sum_{v \in \mathcal{V}} \left([\hat{\mathbf{q}}^{K,i,r+1}]_{n,l,v} - [\hat{\mathbf{q}}^{K,i,r}]_{n,l,v} \right). \tag{2.80}$$

For linear homogeneous scalar hyperbolic conservation laws and neglecting floating point errors it can be proven that the iteration converges after at most N steps (see ?? for details).

2.1.11 A Fully-discrete Update Scheme for the Time-discrete Solution

Now that we have developed a method to compute the space-time predictor, we can go back to the original one-step, cell-local update scheme given in eq. (2.21). Inserting the local space-time predictor $\tilde{\mathbf{q}}_h^{K,i}$ yields

$$\begin{aligned} \int_K \left[\tilde{\mathbf{u}}_h^{K,i} \Big|_{t_i+\Delta t_i} \right]_v \left[\mathbf{w}_h^K \right]_v dx &= \int_K \left[\tilde{\mathbf{u}}_h^{K,i} \Big|_{t_i} \right]_v \left[\mathbf{w}_h^K \right]_v dx + \\ &\int_{t_i}^{t_i+\Delta t_i} \int_K \left[\mathbf{F}(\tilde{\mathbf{q}}_h^{K,i}) \right]_{vd} \frac{\partial}{\partial x_d} \left[\mathbf{w}_h^K \right]_v dx dt + \\ &\int_{t_i}^{t_i+\Delta t_i} \int_K \left[\mathbf{S}(\tilde{\mathbf{q}}_h^{K,i}) \right]_v \left[\mathbf{w}_h^K \right]_v dx dt - \\ &\int_{t_i}^{t_i+\Delta t_i} \int_{\partial K} \left[\mathcal{G}(\tilde{\mathbf{q}}_h^{K,i}, \tilde{\mathbf{q}}_h^{K+1,i}, \mathbf{n}) \right]_v \left[\mathbf{w}_h^K \right]_v ds(x) dt, \end{aligned} \quad (2.81)$$

which has to hold for all $v \in \mathcal{V}$, $K \in \mathcal{K}_h$, $\mathbf{w}_h \in \mathbb{W}_h$ and $i \in \mathcal{I}$.

Making use of the bases we derived earlier the cell-local solution $\tilde{\mathbf{u}}_h^{K,i}$ at times t_i and $t_i + \Delta t_i$ can be represented by tensors of coefficients $\hat{\mathbf{u}}^{K,i}$ and $\hat{\mathbf{u}}^{K,i+1}$ as

$$\tilde{\mathbf{u}}_h^{K,i} \Big|_{t_i} = \left[\hat{\mathbf{u}}^{K,i} \right]_{n,v} \boldsymbol{\phi}_{n,v}^K \quad (2.82)$$

and

$$\tilde{\mathbf{u}}_h^{K,i} \Big|_{t_i+\Delta t_i} = \left[\hat{\mathbf{u}}^{K,i+1} \right]_{n,v} \boldsymbol{\phi}_{n,v}^K, \quad (2.83)$$

respectively. Inserting eqs. (2.82) and (2.83) and the ansatz for the space-time predictor (2.62) into eq. (2.81) yields

$$\begin{aligned} \underbrace{\int_K \left[\left[\hat{\mathbf{u}}^{K,i+1} \right]_{n,v} \boldsymbol{\phi}_{n,v}^K \right]_j \left[\boldsymbol{\phi}_{\alpha,\gamma}^K \right]_j dx}_{\text{U-I}} &= \underbrace{\int_K \left[\left[\hat{\mathbf{u}}^{K,i} \right]_{n,v} \boldsymbol{\phi}_{n,v}^K \right]_j \left[\boldsymbol{\phi}_{\alpha,\gamma}^K \right]_j dx}_{\text{U-II}} + \\ &\underbrace{\int_{t_i}^{t_i+\Delta t_i} \int_K \left[\mathbf{F} \left(\left[\hat{\mathbf{q}}^{K,i} \right]_{n,l,v} \boldsymbol{\theta}_{n,l,v}^{Ki} \right) \right]_{jk} \frac{\partial}{\partial x_k} \left[\boldsymbol{\phi}_{\alpha,\gamma}^K \right]_j dx dt}_{\text{U-III}} + \\ &\underbrace{\int_{t_i}^{t_i+\Delta t_i} \int_K \left[\mathbf{S} \left(\left[\hat{\mathbf{q}}^{K,i} \right]_{n,l,v} \boldsymbol{\theta}_{n,l,v}^{Ki} \right) \right]_j \left[\boldsymbol{\phi}_{\alpha,\gamma}^K \right]_j dx dt}_{\text{U-IV}} - \\ &\underbrace{\int_{t_i}^{t_i+\Delta t_i} \int_{\partial K} \left[\mathcal{G} \left(\hat{\mathbf{q}}^{K,i}, \hat{\mathbf{q}}^{K+1,i}, \mathbf{n} \right) \right]_j \left[\boldsymbol{\phi}_{\alpha,\gamma}^K \right]_j ds(x) dt}_{\text{U-V}}, \end{aligned} \quad (2.84)$$

which we require to hold for all $\alpha \in \mathcal{N}$, $\gamma \in \mathcal{V}$, $K \in \mathcal{K}_h$ and $i \in \mathcal{I}$. In the following we will again proceed by simplifying each term in reference coordinates separately and then in the end assemble all terms to obtain a complete fully-discrete update scheme.

Term U-I

The first term of eq. (2.84) can be rewritten with respect to reference coordinates as follows:

$$\begin{aligned}
 & \int_K \left[\left[\hat{\mathbf{u}}^{K,i+1} \right]_{n,v} \boldsymbol{\phi}_{n,v}^K \right]_j \left[\boldsymbol{\phi}_{\alpha,\gamma}^K \right]_j d\mathbf{x} = \\
 & \int_K \left[\left[\hat{\mathbf{u}}^{K,i+1} \right]_{n,v} \phi_n^K \mathbf{e}_v \right]_j \left[\phi_\alpha^K \mathbf{e}_\gamma \right]_j d\mathbf{x} = \\
 & J\mathcal{X}_K \int_{\hat{K}} \left[\left[\hat{\mathbf{u}}^{K,i+1} \right]_{n,v} \hat{\phi}_n \mathbf{e}_v \right]_j \left[\hat{\phi}_\alpha \mathbf{e}_\gamma \right]_j d\hat{\boldsymbol{\xi}} = \\
 & J\mathcal{X}_K \sum_{\alpha' \in \mathcal{N}} \left(\hat{\omega}_{\alpha'} \left[\hat{\mathbf{u}}^{K,i+1} \right]_{n,v} \hat{\phi}_n(\hat{\boldsymbol{\xi}}_{\alpha'}) [\mathbf{e}_v]_j \hat{\phi}_\alpha(\hat{\boldsymbol{\xi}}_{\alpha'}) [\mathbf{e}_\gamma]_j \right) = \\
 & J\mathcal{X}_K \sum_{\alpha' \in \mathcal{N}} \left(\hat{\omega}_{\alpha'} \left[\hat{\mathbf{u}}^{K,i+1} \right]_{n,v} \delta_{n\alpha'} \delta_{vj} \delta_{\alpha\alpha'} \delta_{\gamma j} \right) = \\
 & J\mathcal{X}_K \hat{\omega}_\alpha \left[\hat{\mathbf{u}}^{K,i+1} \right]_{\alpha,\gamma}. \tag{2.85}
 \end{aligned}$$

Term U-II

Analogously to the first term of eq. (2.84), the second term can be rewritten as follows:

$$\begin{aligned}
 & \int_K \left[\left[\hat{\mathbf{u}}^{K,i} \right]_{n,v} \boldsymbol{\phi}_{n,v}^K \right]_j \left[\boldsymbol{\phi}_{\alpha,\gamma}^K \right]_j d\mathbf{x} = \\
 & J\mathcal{X}_K \hat{\omega}_\alpha \left[\hat{\mathbf{u}}^{K,i} \right]_{\alpha,\gamma}. \tag{2.86}
 \end{aligned}$$

2.1. A D -dimensional ADER-DG scheme with MUSCL-Hancock a-posteriori subcell limiting for non-linear hyperbolic conservation laws

Term U-III

The third term of eq. (2.84) can be rewritten with respect to reference coordinates as follows:

$$\begin{aligned}
& \int_{t_i}^{t_i+\Delta t_i} \int_K \left[F \left([\hat{q}^{K,i}]_{n,l,v} \theta_{n,l,v}^{Ki} \right) \right]_{jk} \frac{\partial}{\partial x_k} [\phi_{\alpha,\gamma}^K]_j dx dt = \\
& \int_{t_i}^{t_i+\Delta t_i} \int_K \left[F \left([\hat{q}^{K,i}]_{n,l,v} \phi_n^K \psi_l^i e_v \right) \right]_{jk} \frac{\partial}{\partial x_k} \left(\prod_{d=1}^D \psi_{[\alpha]_d}^K([x]_d) \right) [e_\gamma]_j dx dt = \\
& \int_{t_i}^{t_i+\Delta t_i} \int_K \left[F \left([\hat{q}^{K,i}]_{n,l,v} \phi_n^K \psi_l^i e_v \right) \right]_{jk} \left(\prod_{d=1, d \neq k}^D \psi_{[\alpha]_d}^K([x]_d) \right) \frac{1}{[\Delta \mathbf{x}^K]_k} \frac{\partial}{\partial \xi_k} \hat{\psi}_{[\alpha]_k}([\mathbf{x}_K(\mathbf{x})]_k) [e_\gamma]_j dx dt = \\
& J_{\mathcal{T}_i} J_{\mathcal{X}_K} \int_0^1 \int_{\hat{K}} \left[F \left([\hat{q}^{K,i}]_{n,l,v} \hat{\phi}_n \hat{\psi}_l e_v \right) \right]_{kj} \left(\prod_{d=1, d \neq k}^D \hat{\psi}_{[\alpha]_d}([\xi]_d) \right) \frac{1}{[\Delta \mathbf{x}^K]_k} \frac{\partial}{\partial \xi_k} \hat{\psi}_{[\alpha]_k}([\xi]_k) [e_\gamma]_j d\xi d\tau = \\
& J_{\mathcal{T}_i} J_{\mathcal{X}_K} \sum_{\alpha' \in \mathcal{N}} \sum_{\beta' \in \mathcal{N}} \left(\hat{\omega}_{\alpha'} \hat{\omega}_{\beta'} \left[F \left([\hat{q}^{K,i}]_{n,l,v} \hat{\phi}_n(\xi_{\alpha'}) \hat{\psi}(\hat{\tau}_{\beta'}) e_v \right) \right]_{jk} \left(\prod_{d=1, d \neq k}^D \hat{\psi}_{[\alpha]_d}([\xi_{\alpha'}]_d) \right) \frac{1}{[\Delta \mathbf{x}^K]_k} \frac{\partial}{\partial \xi_k} \hat{\psi}_{[\alpha]_k}([\xi_{\alpha'}]_k) \right) = \\
& J_{\mathcal{T}_i} J_{\mathcal{X}_K} \sum_{\alpha' \in \mathcal{N}} \sum_{\beta' \in \mathcal{N}} \left(\hat{\omega}_{\alpha'} \hat{\omega}_{\beta'} \left[F \left([\hat{q}^{K,i}]_{n,l,v} \delta_{n\alpha'} \delta_{l\beta'} e_v \right) \right]_{jk} \left(\prod_{d=1, d \neq k}^D \delta_{[\alpha]_d}[\alpha']_d \right) \frac{1}{[\Delta \mathbf{x}^K]_k} \frac{\partial}{\partial \xi_k} \hat{\psi}_{[\alpha]_k}([\xi_{\alpha'}]_k) \delta_{\gamma j} \right) = \\
& J_{\mathcal{T}_i} J_{\mathcal{X}_K} \hat{\omega}_\alpha \sum_{k=1}^D \left(\sum_{\alpha'_k \in \mathcal{N}} \sum_{\beta' \in \mathcal{N}} \left(\frac{\hat{\omega}_{\beta'}}{\hat{\omega}_{\alpha'_k}} \frac{1}{[\Delta \mathbf{x}^K]_k} \frac{\partial}{\partial \xi_k} \hat{\psi}_{\alpha'_k}([\xi]_{\alpha'_k}) \left[F \left([\hat{q}^{K,i}]_{[\alpha]_1, [\alpha]_2, \dots, [\alpha]_{k-1}, \alpha'_k, [\alpha]_{k+1}, \dots, [\alpha]_D, \beta', v} e_v \right) \right]_{\gamma, k} \right) \right) = \\
& J_{\mathcal{T}_i} J_{\mathcal{X}_K} \hat{\omega}_\alpha \sum_{k=1}^D \left(\sum_{\alpha'_k \in \mathcal{N}} \sum_{\beta' \in \mathcal{N}} \left(\frac{1}{\hat{\omega}_{\alpha'_k}} \frac{1}{[\Delta \mathbf{x}^K]_k} [K]_{\alpha'_k, k} \left[F \left([\hat{q}^{K,i}]_{[\alpha]_1, [\alpha]_2, \dots, [\alpha]_{k-1}, \alpha'_k, [\alpha]_{k+1}, \dots, [\alpha]_D, \beta', v} e_v \right) \right]_{\gamma, k} \right) \right), \tag{2.87}
\end{aligned}$$

where we made use of the fact that du to the chain rule:

$$\begin{aligned}
 \frac{\partial}{\partial x_k} \left(\prod_{d=1}^D \psi_{[\alpha]_d}^K([x]_d) \right) &= \left(\prod_{d=1, d \neq k}^D \psi_{[\alpha]_d}^K([x]_d) \right) \frac{\partial}{\partial x_k} \psi_{[\alpha]_k}^K([x]_k) = \\
 &\left(\prod_{d=1, d \neq k}^D \psi_{[\alpha]_d}^K([x]_d) \right) \frac{\partial}{\partial \xi_j} \hat{\psi}_{[\alpha]_k}([\mathbf{x}_K(\mathbf{x})]_k) \frac{\partial}{\partial x_k} [\mathbf{x}_K(\mathbf{x})]_j = \\
 &\left(\prod_{d=1, d \neq k}^D \psi_{[\alpha]_d}^K([x]_d) \right) \frac{\partial}{\partial \xi_j} \hat{\psi}_{[\alpha]_k}([\mathbf{x}_K(\mathbf{x})]_k) \frac{1}{[\Delta \mathbf{x}^K]_k} \delta_{kj} = \\
 &\left(\prod_{d=1, d \neq k}^D \psi_{[\alpha]_d}^K([x]_d) \right) \frac{1}{[\Delta \mathbf{x}^K]_k} \frac{\partial}{\partial \xi_k} \hat{\psi}_{[\alpha]_k}([\mathbf{x}_K(\mathbf{x})]_k) dx dt. \tag{2.88}
 \end{aligned}$$

Term U-IV

The fourth term of eq. (2.84) can be rewritten with respect to reference coordinates as follows:

$$\begin{aligned}
 &\int_{t_i}^{t_i + \Delta t_i} \int_K \left[S \left([\hat{\mathbf{q}}^{K,i}]_{n,l,v} \boldsymbol{\theta}_{n,l,v}^{Ki} \right) \right]_j [\boldsymbol{\phi}_{\alpha,\gamma}^K]_j dx dt = \\
 &\int_{t_i}^{t_i + \Delta t_i} \int_K \left[S \left([\hat{\mathbf{q}}^{K,i}]_{n,l,v} \boldsymbol{\phi}_n^K \psi_l^i e_v \right) \right]_j \phi_\alpha^K [e_\gamma]_j dx dt = \\
 &J_{\mathcal{T}_i} J_{\mathcal{X}_K} \int_0^1 \int_{\hat{K}} \left[S \left([\hat{\mathbf{q}}^{K,i}]_{n,l,v} \hat{\phi}_n \hat{\psi}_l e_v \right) \right]_j \hat{\phi}_\alpha [e_\gamma]_j d\xi d\tau = \\
 &J_{\mathcal{T}_i} J_{\mathcal{X}_K} \sum_{\alpha' \in \mathcal{N}} \sum_{\beta' \in \mathcal{N}} \left(\hat{\omega}_{\alpha'} \hat{\omega}_{\beta'} \left[S \left([\hat{\mathbf{q}}^{K,i}]_{n,l,v} \hat{\phi}_n(\hat{\xi}_{\alpha'}) \hat{\psi}_l(\hat{\tau}_{\beta'} e_v) \right) \right]_j \hat{\phi}_\alpha(\hat{\xi}_{\alpha'}) [e_\gamma]_j \right) = \\
 &J_{\mathcal{T}_i} J_{\mathcal{X}_K} \sum_{\alpha' \in \mathcal{N}} \sum_{\beta' \in \mathcal{N}} \left(\hat{\omega}_{\alpha'} \hat{\omega}_{\beta'} \left[S \left([\hat{\mathbf{q}}^{K,i}]_{n,l,v} \delta_{n\alpha'} \delta_{l\beta'} e_v \right) \right]_j \delta_{\alpha\alpha'} \delta_{\gamma j} \right) = \\
 &J_{\mathcal{T}_i} J_{\mathcal{X}_K} \hat{\omega}_\alpha \sum_{\beta' \in \mathcal{N}} \left(\hat{\omega}_{\beta'} \left[S \left([\hat{\mathbf{q}}^{K,i}]_{\alpha,\beta',v} e_v \right) \right]_\gamma \right). \tag{2.89}
 \end{aligned}$$

Term U-V

Let $d \in \mathcal{D}$ and $e \in \{0, 1\} := \mathcal{E}$. Then if we define the $D - 1$ -dimensional quadrilateral $\partial \hat{K}_{d,e}$ as

$$\partial \hat{K}_{d,e} = \left\{ \xi \in \hat{K} \mid [\xi]_d = e \right\}, \tag{2.90}$$

2.1. A D -dimensional ADER-DG scheme with MUSCL-Hancock a-posteriori subcell limiting for non-linear hyperbolic conservation laws

the set $\{\partial\hat{K}_{d,e}\}_{d\in\mathcal{D},e\in\mathcal{E}}$ is a partition of the surface $\partial\hat{K}$ of the spatial reference element. By making use of the mappings \mathcal{X}_K that maps points $\xi \in \hat{K}$ to $x \in K$ for all $K \in \mathcal{K}_h$ we can define

$$\partial K_{d,e} = \mathcal{X}_K(\partial\hat{K}_{d,e}), \quad (2.91)$$

where now the set $\{\partial K_{d,e}\}_{d\in\mathcal{D},e\in\mathcal{E}}$ is a quadrilateral partition of the surface ∂K for all cells $K \in \mathcal{K}_h$.

In consequence the surface integral in the fifth term of eq. (2.84) can be rewritten as follows:

$$\begin{aligned} & \int_{t_i}^{t_i+\Delta t_i} \int_{\partial K} \left[\mathcal{G}(\hat{q}^{K,i}, \hat{q}^{K+,i}, n) \right]_j [\phi_{\alpha,\gamma}^K]_j ds(x) dt = \\ & \int_{t_i}^{t_i+\Delta t_i} \sum_{d\in\mathcal{D}} \sum_{e\in\mathcal{E}} \left(\int_{\partial K_{d,e}} \left[\mathcal{G}(\hat{q}^{K,i}, \hat{q}^{K+,i}, e_d) \right]_j \phi_{\alpha}^K [e_{\gamma}]_j ds(x) \right) dt = \\ & J_{\mathcal{T}_i} J_{\mathcal{X}_K} \int_0^1 \sum_{d\in\mathcal{D}} \sum_{e\in\mathcal{E}} \left(\frac{1}{[\Delta x^K]_d} \int_{\partial\hat{K}_{d,e}} \left[\mathcal{G}(\hat{q}^{K,i}, \hat{q}^{K+,i}, (-1)^e e_d) \right]_j \hat{\phi}_{\alpha} [e_d]_j ds(\xi) \right) d\tau = \\ & J_{\mathcal{T}_i} J_{\mathcal{X}_K} \sum_{\beta' \in \mathcal{D}} \hat{\omega}_{\beta'} \sum_{d\in\mathcal{D}} \sum_{e\in\mathcal{E}} \sum_{\alpha' \in \mathcal{N}^-} \left(\hat{\omega}_{\alpha'} \frac{1}{[\Delta x^K]_d} \left[\mathcal{G}(\hat{q}^{K,i}, \hat{q}^{K+,i}, (-1)^e e_d) \right]_j \hat{\phi}_{\alpha^d}(\xi_{\alpha'}) \left(\hat{\psi}_{[\alpha]_d|_e} \right) [e_d]_j \right) = \\ & J_{\mathcal{T}_i} J_{\mathcal{X}_K} \sum_{\beta' \in \mathcal{D}} \hat{\omega}_{\beta'} \sum_{d\in\mathcal{D}} \sum_{e\in\mathcal{E}} \sum_{\alpha' \in \mathcal{N}^-} \left(\hat{\omega}_{\alpha'} \frac{1}{[\Delta x^K]_d} \left[\mathcal{G}(\hat{q}^{K,i}, \hat{q}^{K+,i}, (-1)^e e_d) \right]_j \delta_{\alpha^d \alpha'} \left(\hat{\psi}_{[\alpha]_d|_e} \right) \delta_{\gamma j} \right) = \\ & J_{\mathcal{T}_i} J_{\mathcal{X}_K} \hat{\omega}_{\alpha} \sum_{\beta' \in \mathcal{D}} \sum_{d\in\mathcal{D}} \sum_{e\in\mathcal{E}} \sum_{\alpha'_d \in \mathcal{N}} \left(\frac{\hat{\omega}_{\beta'}}{\hat{\omega}_{\alpha'_d}} \frac{1}{[\Delta x^K]_d} \left[\mathcal{G}(\hat{q}^{K,i}, \hat{q}^{K+,i}, (-1)^e e_d) \right]_{\gamma} \underbrace{\left(\hat{\psi}_{\alpha'_d|_e} \right)}_{\text{F0, F1}} \right). \end{aligned} \quad (2.92)$$

In each term we have to solve a Riemann problem in direction of the unit vector e_d defined as

$$[e_d]_{d'} = \delta_{dd'} \quad (2.93)$$

for $d' \in \mathcal{D}$.

The Complete One-step Update Scheme

Inserting eqs. (2.85) to (2.87), (2.89) and (2.92) into eq. (2.84) and dividing the resulting equation by $\hat{\omega}_\alpha$ and $J_{\mathcal{X}_K}$ yields

$$\begin{aligned}
 [\hat{\mathbf{u}}^{K,i+1}]_{\alpha,\gamma} &= [\hat{\mathbf{u}}^{K,i}]_{\alpha,\gamma} + \\
 &J_{\mathcal{T}_i} \sum_{k=1}^D \left(\sum_{\alpha'_k \in \mathcal{N}} \sum_{\beta' \in \mathcal{N}} \left(\frac{\hat{\omega}_{\beta'}}{\hat{\omega}_{\alpha'_k}} \frac{1}{[\Delta \mathbf{x}^K]_k} \underbrace{\frac{\partial}{\partial \xi_k} \hat{\psi}_{\alpha'_k} \left([\hat{\xi}]_{\alpha'_k} \right)}_{\text{Kxi}_{\alpha'_k}} \right) \left[F \left([\hat{\mathbf{q}}^{K,i}]_{[\alpha]_1, [\alpha]_2, \dots, [\alpha]_{k-1}, \alpha'_k, [\alpha]_{k+1}, \dots, [\alpha]_D}, \beta', v \right) \mathbf{e}_v \right]_{\gamma} \right) \\
 &J_{\mathcal{T}_i} \sum_{\beta' \in \mathcal{N}} \left(\hat{\omega}_{\beta'} \left[S \left([\hat{\mathbf{q}}^{K,i}]_{\alpha, \beta', v} \right) \right]_{\gamma} \right) - \\
 &J_{\mathcal{T}_i} \sum_{\beta' \in \mathcal{D}} \sum_{d \in \mathcal{D}} \sum_{e \in \mathcal{E}} \sum_{\alpha'_d \in \mathcal{N}} \left(\frac{\hat{\omega}_{\beta'}}{\hat{\omega}_{\alpha'_d}} \frac{1}{[\Delta \mathbf{x}^K]_d} \left[\mathcal{G} \left(\hat{\mathbf{q}}^{K,i}, \hat{\mathbf{q}}^{K+,i}, (-1)^e \mathbf{e}_d \right) \right]_{\gamma} \underbrace{\left(\hat{\psi}_{\alpha'_d} \right)_e}_{\text{F0, F1}} \right), \tag{2.94}
 \end{aligned}$$

which we require to hold for $\alpha \in \mathcal{N}$, $\gamma \in \mathcal{V}$, $K \in \mathcal{K}_h$ and $i \in \mathcal{I}$.

Time step restriction

$$\Delta t \leq \frac{1}{D} \frac{1}{(2N+1)} \min_{d \in \mathcal{D}} \left(\frac{[\Delta \mathbf{x}]_d}{\Lambda^d} \right), \tag{2.95}$$

where

$$\Lambda^d = \max_{v \in \mathcal{V}} \text{abs} \left[\lambda^d \right]_v \tag{2.96}$$

and λ^d is a vector containing the V real eigenvalues of the Jacobian

$$\frac{\partial}{\partial x_j} \left[\mathbf{F}(\mathbf{u}(\mathbf{x}, t)) \right]_{id} \tag{2.97}$$

for the respective dimension $d \in \mathcal{D}$.

2.1.12 A Posteriori Subcell Limiting

Motivation:

- Shock = discontinuity

2.1. A D -dimensional ADER-DG scheme with MUSCL-Hancock a-posteriori subcell limiting for non-linear hyperbolic conservation laws

- Discontinuity + high-order DG method leads to Gibbs phenomenon (oscillations)
- Reason: Discon. initial data or spontaneous formation in nonlinear problems
- Problems:
 1. Pointwise first order away from discontinuity
 2. Loss of pointwise convergence at the point of discontinuity
 3. Introduction of artificial and persistent oscillations at the point of discontinuity
- Positive physical quantities such as pressure or density might become negative; simulation might crash
- ADER-DG with a posteriori subcell limiting has very desirable properties (TODO)

Questions:

1. How do we identify cells for which limiting is needed? Troubled cell indicator.
2. How do we achieve high-order accuracy and still ensure non-oscillatory property close to troubled cells? Ideally replace DG solution such that additional numerical viscosity is added only at these cells but nowhere else and preferably without destroying the subcell resolution of the DG method.

Projection and Reconstruction

In order to do FVM we need to project the ADER-DG degrees of freedom $\hat{u}^{K,i}$ to N_S equidistant subcell-averages $\hat{p}^{K_L,i}$ for each cell $K \in \mathcal{K}_h$. We choose $N_S = 2N + 1$, since for explicit Godunov-type finite volume schemes on the subgrid we must satisfy the stability condition

$$\Delta t \leq \frac{1}{d} \frac{1}{N_S} \min_{d \in \mathcal{D}} \left(\frac{[\Delta x]_d}{\Lambda^d} \right). \quad (2.98)$$

Comparing eq. (2.98) to the time step restriction for the ADER-DG scheme given in eq. (2.95) illustrates that the choice $N_S = 2N + 1$ make sure that a) time steps on the ADER-DG grid are also valid on the subgrid and b) that we add the minimum amount of dissipation.

Let L^K be a regular subgrid on cell K consisting of $(N_S)^D = (2N + 1)^D$ subcells denoted by L_j^K , $j \in \{1, 2, \dots, (N_S)^D\} := \mathcal{N}_S$ for $N_S \geq N + 1$. Then we

2. THEORY

can define an alternative representation of the $V(N+1)^D$ degrees of freedom $\hat{\mathbf{u}}^{K,i}$ in terms of $V(N_S)^D$ cell averages $\hat{\mathbf{p}}^{K_L,i}$ using the following relation:

$$\begin{aligned} [\hat{\mathbf{p}}^{K_L,i}]_{\alpha,\gamma} &= \frac{1}{|L_K^\alpha|} \int_{L_K^\alpha} [\hat{\mathbf{u}}_h^{K,i}]_\gamma d\mathbf{x} = \frac{1}{|L_K^\alpha|} \int_{L_K^\alpha} \phi_n^K(\mathbf{x}) [e_v]_\gamma d\mathbf{x} [\hat{\mathbf{u}}^{K,i}]_{n,v} = \\ &= \int_{\hat{K}} \hat{\phi}_n \left(\frac{1}{N_S} \boldsymbol{\alpha} + \frac{1}{N_S} \boldsymbol{\xi} \right) d\boldsymbol{\xi} [\hat{\mathbf{u}}^{K,i}]_{n,\gamma} = \\ &= \sum_{\alpha' \in \mathcal{N}} \left(\hat{\omega}_{\alpha'} \hat{\phi}_n \left(\frac{1}{N_S} \boldsymbol{\alpha} + \frac{1}{N_S} \hat{\boldsymbol{\xi}}_{\alpha'} \right) \right) [\hat{\mathbf{u}}^{K,i}]_{n,\gamma}, \end{aligned} \quad (2.99)$$

which can be carried out efficiently in a dimension-by-dimension manner. See ?? for details.

We can directly derive the following relation:

Reconstruction: If $N_S > N+1$ then we impose additional restrictions:

Identification of Troubled Cells

Candidate solution: $\mathbf{u}_h^*(\mathbf{x}, t^{i+1})$ obtained using unlimited high-order scheme. Apply troubled cell indicator Project, recompute with more robust scheme and restrict.

Physical admissibility detection (PAD): Domain knowledge

$$\pi_k \left(\mathbf{u}_h^*(\mathbf{x}, t^{i+1}) \right) > 0 \quad (2.100)$$

For Euler $\pi_1(\mathbf{u}) = \rho$, $\pi_2(\mathbf{u}) = p$.

Numerical admissibility detection (NAD): Relaxed discrete maximum principle:

$$\min_{\mathbf{x}' \in V(K)} \mathbf{u}_h(\mathbf{x}', t^i) - \delta \leq \mathbf{u}_h^*(\mathbf{x}, t^{i+1}) \leq \max_{\mathbf{x}' \in V(K)} \mathbf{u}_h(\mathbf{x}', t^i) + \delta \quad (2.101)$$

for all Voronoi neighbors K' of $K \in \mathcal{K}_h$.

Proposed relaxation:

$$\delta = \varepsilon \left(\max_{\mathbf{x}' \in V(K)} \left(\mathbf{u}_h^*(\mathbf{x}', t^i) \right) - \min_{\mathbf{x}' \in V(K)} \left(\mathbf{u}_h^*(\mathbf{x}', t^i) \right) \right), \quad (2.102)$$

where $\varepsilon = 10^{-3}$.

Approximation: Evaluate on \mathbf{p}_h^* , i.e. on the projection to linspace. Projection described in the next chapter.

$$\min_{\mathbf{x}' \in V(K)} \mathbf{p}_h(\mathbf{x}', t^i) - \delta \leq \mathbf{p}_h^*(\mathbf{x}, t^{i+1}) \leq \max_{\mathbf{x}' \in V(K)} \mathbf{p}_h(\mathbf{x}', t^i) + \delta \quad (2.103)$$

2.1. A D -dimensional ADER-DG scheme with MUSCL-Hancock a-posteriori subcell limiting for non-linear hyperbolic conservation laws

evaluate in terms of subcell averages.

$$p_h^*(x, t^{i+1}) = \mathcal{P} \left(u_h^*(x, t^{i+1}) \right) \quad (2.104)$$

as defined in the next section.

Important: For troubled cells that have been troubled in the previous step of the simulation we use the FVM result $A(p)$, for newly troubled cells we use $P(u)$.

The MUSCL-Hancock Scheme

second order in both space and time; total variation diminishing, robust, simple, details: ??

Consists of the following steps:

1. Compute slopes:

$$\begin{aligned} [\delta_d^{K,i}]_{\alpha,\gamma} = \minmod \left([\hat{p}^{K_L,i}]_{\alpha+e_d,\gamma} - [\hat{p}^{K_L,i}]_{\alpha,\gamma}, \right. \\ \left. [\hat{p}^{K_L,i}]_{\alpha,\gamma} - [\hat{p}^{K_L,i}]_{\alpha-e_d,\gamma} \right) \end{aligned} \quad (2.105)$$

for all $\alpha \in \{0, 1, \dots, N_S + 1\}^D := \mathcal{N}_{\mathcal{S}}^*$, $\gamma \in \mathcal{V}$, unit vectors e_d , $d \in \mathcal{D}$, subgrid cells $L^K \in K$, troubled grid cells $K \in \mathcal{K}_h^*$ and $i \in \mathcal{I}$. We furthermore use the common definition of the minmod function, namely

$$\minmod(a, b) = \begin{cases} 0 & \text{if } ab \leq 0 \\ a & \text{if } ab > 0 \text{ and } |a| \leq |b| \\ b & \text{if } ab > 0 \text{ and } |b| < |a|. \end{cases} \quad (2.106)$$

2. Evaluate source:

$$[s^{K,i}]_{\alpha,\gamma} = \left[s \left([\hat{q}^{K,i}]_{\alpha} \right) \right]_{\gamma} \quad (2.107)$$

for all $\alpha \in \mathcal{N}_{\mathcal{S}}^*$, $\gamma \in \mathcal{V}$, subgrid cells $L^K \in K$, troubled cells $K \in \mathcal{K}_h^*$ and $i \in \mathcal{I}$.

3. Extrapolate:

$$[w^{K,i}]_{d,e,\alpha,\gamma} = [\hat{u}^{K,i}]_{\alpha,\gamma} + \frac{e}{2} [\delta_d^{K,i}]_{\alpha,\gamma} \quad (2.108)$$

for $\alpha \in \mathcal{N}_{\mathcal{S}}^*$, $\gamma \in \mathcal{V}$, $d \in \mathcal{D}$ and $e \in \{-1, +1\} := \sigma, \dots$

4. Evolve:

$$\begin{aligned} \left[w^{K,i+\frac{1}{2}} \right]_{d,e,\alpha,\gamma} &= \frac{\Delta t_i}{2} \sum_{d' \in \mathcal{D}} \sum_{e' \in \sigma} \left(e' \left[F \left(\left[w^{K,i} \right]_{d',e',\alpha} \right) \right]_{\gamma,d'} / \left[\Delta x^{L^K} \right]_{d'} \right) + \\ &\quad \frac{\Delta t_i}{2} \left[s^{K,i} \right]_{\alpha,\gamma} \end{aligned} \quad (2.109)$$

for all $\alpha \in \mathcal{N}_{\mathcal{S}}^*$, $\gamma \in \mathcal{V}$, $d \in \mathcal{D}$, $e \in \sigma$, ...

5. Solve Riemann problems:

$$\left[f^{K,i} \right]_{d,\alpha,\gamma} = \left[\mathcal{G} \left(\left[w^{K,i+\frac{1}{2}} \right]_{d,+1,\alpha-e_d}, \left[w^{K,i+\frac{1}{2}} \right]_{d,-1,\alpha+e_d}, e_d \right) \right]_{\gamma} \quad (2.110)$$

6. Evolve source

$$\begin{aligned} \left[s^{K,i+\frac{1}{2}} \right]_{\alpha,\gamma} &= \left[s^{K,i} \right]_{\alpha,\gamma} + \\ &\quad \frac{1}{2} \sum_{d' \in \mathcal{D}} \sum_{e' \in \sigma} \left(e' \left[F \left(\left[w^{K,i} \right]_{d',e',\alpha} \right) \right]_{\gamma,d'} / \left[\Delta x^{L^K} \right]_{d'} \right) + \\ &\quad \frac{1}{2} \left[s^{K,i} \right]_{\alpha,\gamma} \end{aligned} \quad (2.111)$$

7. Update solution

$$\begin{aligned} \left[p^{L^K,i+1} \right]_{\alpha,\gamma} &= \\ &\quad \left[p^{L^K,i} \right]_{\alpha,\gamma} - \\ &\quad \Delta t_i \sum_{d \in \mathcal{D}} \left(\left[f^{K,i} \right]_{d,\alpha+e_d,\gamma} - \left[f^{K,i} \right]_{d,\alpha,\gamma} / \left[\Delta x^{L^K} \right]_d \right) + \\ &\quad \Delta t_i \left[s^{K,i+\frac{1}{2}} \right]_{\alpha,\gamma} \end{aligned} \quad (2.112)$$

2.2 Profiling and Energy-aware Computing

A profiling infrastructure for ExaHyPE

- General architecture
- Architecture profiling
- Functionality

Chapter 4

Preliminary profiling results, case studies

- Analytic benchmark: Introduction, derivation
- Pie-chart per kernel
- Case-study: Cache-misses, compile-time (\rightarrow Toolkit philosophy)
- Degree \rightarrow Wallclock, Energy (AMR)
- Static mesh $\Delta x \rightarrow$ Error for polynomials (convergence tables)

Chapter 5

Conclusion and Outlook

- PA is important
- ExaHyPE as an answer to exascale challenges
- Applications

Chapter 6

Acknowledgment
