

Leslie Gonzales*, Margaret Youngman, Kenneth S Henry
University of Rochester Medical Center

leslie_gonzales@urmc.rochester.edu

Cochlear Synaptopathy

Cochlear Synaptopathy (CS) is the loss of synapses between hair cells and auditory-nerve fibers (ANFs).

CS is hypothesized to impair perception of sounds in noise, called hidden hearing loss, but underlying neural mechanisms are unknown.

Previous studies of CS

- Conflicting literature about the changes in the spontaneous rate of ANFs^{3,6}
- Enhancement of ANF onset response^{2,6}


Only a single study has examined the effects of CS on the cochlear nucleus (CN).

Current Study

Quantified effects of CS on neural encoding in the nucleus angularis (NA) of the CN in the budgerigar.

Methods

Animal model: Budgerigar (a parakeet species)




Budgerigar brainstem

Wong et al., 2019


- Anesthesia = ketamine + dexmedetomidine
- Glass electrode (10-80 MΩ impedance)
- Distortion product otoacoustic emissions
- Swept-tone paradigm⁵

- Stimuli
 - Tuning curve, spontaneous rate, rate-level function and post-stimulus time histogram (PSTH)
- Identifying a CN unit based on action potential shape, vector strength (VS) and PSTH shape⁴
- Recordings are conducted at least 6 weeks after KA-exposure


PSTH shapes in the NA




Vector Strength in CN units (PSTH)




Is spontaneous activity of CN units altered by CS?




Does CS affect sensitivity of CN units? (Tuning curve)



Does CS affect frequency tuning of CN units? (Tuning curve)



Are onset responses of CN units altered by CS? (PSTH)



Summary

Neural encoding in the NA of the CN in the budgerigar does not seem to be affected by CS.

Future directions

Study neural encoding of complex sounds in the NA of the CN in the budgerigar. Conduct histology to investigate if molecular changes occur in NA cells of CS-induced animals.

Acknowledgements

This research was supported by R01-DC017519, R01-DC017519-S1, and 1F31DC021889-01A1.

References

- ¹ Budgerigar Brain Atlas. (n.d.). Retrieved January 27, 2026, from <http://www.brauthlab.umd.edu/atlas.htm>
- ² Diuba, A., Grati, P., Jeffers, P. W. C., Nouvian, R., Puel, J.-L., Kujawa, S. G., & Bourien, J. (2025). Phenotypic changes of auditory nerve fibers after excitotoxicity. *Proceedings of the National Academy of Sciences*, 122(14), e2412332122. <https://doi.org/10.1073/pnas.2412332122>
- ³ Furman, A. C., Kujawa, S. G., & Liberman, M. C. (2013). Noise-induced cochlear neuropathy is selective for fibers with low spontaneous rates. *Journal of Neurophysiology*, 110(3), 577–586. <https://doi.org/10.1152/jn.00164.2013>
- ⁴ Köppl, C., & Carr, C. E. (2003). Computational diversity in the cochlear nucleus angularis of the barn owl. *Journal of Neurophysiology*, 89(4), 2313–2329. <https://doi.org/10.1152/jn.00635.2002>
- ⁵ Long, G. R., Talmadge, C. L., & Lee, J. (2008). Measuring distortion product otoacoustic emissions using continuously sweeping primaries. *The Journal of the Acoustical Society of America*, 124(3), 1613–1626. <https://doi.org/10.1121/1.2949505>
- ⁶ Suthakar, K., & Liberman, M. C. (2021). Auditory-nerve responses in mice with noise-induced cochlear synaptopathy. *Journal of Neurophysiology*, 126(6), 2027–2038. <https://doi.org/10.1152/jn.00342.2021>
- ⁷ Wong, S. J., Abrams, K. S., Amburgey, K. N., Wang, Y., & Henry, K. S. (2019). Effects of selective auditory-nerve damage on the behavioral audiogram and temporal integration in the budgerigar. *Hearing Research*, 374, 24–34. <https://doi.org/10.1016/j.heares.2019.01.019>