

# Cooling the Heat: Co-benefits of Opioid-reformulation on Temperature and Intimate Partner Violence

Filippo Pavanello\*

Guglielmo Zappalà<sup>†</sup>

[Most recent version here](#)

## Abstract

Intimate Partner Violence (IPV) is a major public health issue often associated with substance abuse. Building on burgeoning prior work showing that higher temperatures increase crime and violent behavior, we document a strong positive effect of temperatures on IPV cases on female victims in the United States using administrative daily data over thirty years at the jurisdiction level. Combining an exogenous abuse-deterrent reformulation of opioids with random short-run variation in daily temperature, we examine whether reducing prescription opioid misuse exacerbates or mitigates the effect of temperature on IPV. Using a within-county triple difference and an event study design, we document that the reformulation significantly mitigates the temperature-IPV gradient in the short-run in counties with greater initial rates of prescription opioid usage. The effect of the policy is more pronounced in urban and richer counties, explaining disparities in IPV outcomes across racial and ethnic groups.

**Keywords:** Intimate Partner Violence, Temperature, Opioid

**JEL Classification:** I18, J16, K32, K42, L65, Q51, Q54

---

\*University of Bologna, Department of Economics, Italy; Centro Euro-Mediterraneo sui Cambiamenti Climatici, Italy; Ca Foscari University of Venice, Department of Economics, Italy; RFF-CMCC European Institute on Economics and the Environment, Italy. Email: [filippo.pavanello2@unibo.it](mailto:filippo.pavanello2@unibo.it)

<sup>†</sup>UC Santa Barbara, emLab and Bren School of Environmental Science & Management, CA 93117. Email: [gzappala@ucsb.edu](mailto:gzappala@ucsb.edu)

# 1 Introduction

In the United States, intimate partner violence (IPV) is a widespread and major public health issue. According to the [National Coalition Against Domestic Violence \(2020\)](#), more than 10 million adults experience domestic violence annually. The incidence of this phenomenon is strongly gendered: 1 in 4 women and 1 in 10 men experience sexual violence, physical violence and/or stalking by an intimate partner during their lifetime and IPV alone accounts for 18% of all violent crime. The consequences of such experiences can be devastating with the estimated lifetime economic cost associated with IPV equal to \$3.6 trillion, as a result of medical services for IPV-related injuries, lost productivity from paid work, criminal justice and other costs ([Centers for Disease Control and Prevention, 2022](#)).

Understanding the drivers of intimate partner violence is a major priority, although the factors affecting IPV are complex. An extensive body of research has established a link between higher temperature and violent behavior through physiological and psychological mechanisms. Previous research has documented this pattern in the US for criminal activity ([Ranson, 2014](#); [Heilmann et al., 2021](#)), homicides ([Colmer and Doleac, 2023](#)) and child maltreatment ([Evans et al., 2023](#)). An often cited primary risk factor associated with IPV perpetration is substance abuse, which can induce individuals to become aggressive and can intensify impulse control disorders ([Angelucci and Heath, 2020](#); [Schilbach, 2019](#)). In particular, opioid misuse has been indicated as a factor facilitating IPV ([Stone and Rothman, 2019](#); [Radcliffe et al., 2021](#)). Supply-side shocks that restrict access to addictive substances, including prescription opioids, can thus either exacerbate or mitigate violent behavior ([Dave et al., 2023](#); [Evans et al., 2022](#)).

In this paper, we examine whether higher temperatures affect IPV perpetration in the United States and if policies designed to curtail prescription opioid misuse

mitigate or exacerbate this effect. We leverage plausibly exogenous variation in temperature to consider whether increases in daily temperatures are associated with higher IPV rates at the jurisdiction-level. We combine this variation with variation in the reformulation of the main legal opioid - OxyContin - into an abuse-deterrent form in 2010, a supply-side intervention that addresses the overprescription of opioids and mitigates their addictive risks. We combine these two sources of county-level variation in a triple-difference and event-study research designs, where the triple interaction term of temperature, pre-intervention opioid exposure, and post-policy time indicator identifies the moderating role of supply-side shock reducing opioid availability, under the assumption that the temperature-IPV relationship would have stayed the same had the reformulation not occurred.

Using administrative jurisdiction-level data from the FBI's National Incident-Based Reporting System (NIBRS) from 1991 to 2021, we find a strong positive effect of average daily temperature. On average, a one-degree Celsius increase is associated with 0.0058 more daily cases of IPV per 100,000 people, a 0.87% increase compared to the mean. Combining weather variation with county-level variation in exposure to prescription opioids, the triple-difference estimates reveal that one-degree Celsius increase in temperature is associated with 0.008 fewer IPV cases per 100,000 people after the reformulation occurred in 2010, a 7.9% reduction at the mean. The event-study analysis documents that the attenuation factor of the policy does not last more than five years after the policy. Altogether these results suggest a positive unintended co-benefit of the opioid reformulating in mitigating the role of temperature in inducing violent behavior and increases in intimate partner violence.

Our findings contribute to a burgeoning literature that examines the effect of temperature on violent behavior, one of the main channels of the socio-economic impact of climate ([Carleton and Hsiang, 2016](#)). Higher temperature can increase

aggressivity and induce violent behavior through physiological channels (Ranson, 2014; Baylis, 2020; Heilmann et al., 2021; Mukherjee and Sanders, 2021; Behrer and Bolotnyy, 2022). Our contribution stands in the examining the impact of temperature on a specific and widespread type of violent behavior, namely intimate partner violence. We do so at a highly granular spatial and temporal resolution, exploiting daily variation in temperature from hourly information at the jurisdiction-level in the US, holding jurisdiction-month-year, week-of-year and day-of-week factors fixed. Our additional results on the mediating role of the policy environment contributes to the growing literature that evaluates the (un)intended consequences of policies in mediating or exacerbating climate impacts on socio-economic outcomes. Health care policies have been shown to attenuate the temperature-mortality relationship in the United States (Mullins and White, 2020) and in Mexico (Cohen and Dechezleprêtre, 2022). More restrictive gun laws attenuate the temperature-homicide relationship (Colmer and Doleac, 2023). Cash transfers also attenuate the effects of higher temperatures on violent behavior, but only temporarily (Garg et al., 2020). Policies, however, can also have negative externalities, such as the highly subsidize federal crop insurance program that makes farmers more sensitive to extreme heat as a result of moral hazard (Annan and Schlenker, 2015). Our study is the first one that examine the positive externality of a supply-side intervention in opioid availability as a mediator to the temperature-IPV relationship.

Our paper also contributes to the literature that studies the determinants of IPV perpetration. Previous work has focused on economic shocks or other policies that may impact womens bargaining power by documenting the effects of emotional cues (Card and Dahl, 2011), cash transfers (Bobonis et al., 2013; Angelucci and Heath, 2020), family structures (Tur-Prats, 2019), labor market shocks - including gender wage gap (Aizer, 2010) and unemployment (Anderberg et al.,

2016; Tur-Prats, 2021) - education (Erten and Keskin, 2018), divorce laws (Stevenson and Wolfers, 2006), and trade shocks (Erten and Keskin, 2021). Dave et al. (2023) document that the opioid reformulation significantly reduced IPV exposure for women, but induced a notable uptick in heroin-involved IPV. We explore environmental factors as measured by daily temperature as a new determinant, and provide novel evidence on the mediating role of the opioid reformulation. We examine the mechanisms and uncover substantial heterogeneity across sub-populations where the policy was effective, underlining the importance of identifying sub-groups of population where the temperature-IPV gradient is positive and inelastic to policy interventions. In particular, we document that the policy was not effective in marginalised communities, including rural, low-income, high-poverty, non-Hispanic, non-White, Black counties.

Finally, we speak to the literature on the opioid epidemic which has pervaded the United States in the past decades (Arteaga and Barone, 2022; Dave et al., 2023). Evans et al. (2022) document that the reformulation of OxyContin and the implementation of must-access prescription drug monitoring programs increase child physical abuse and neglect. Gihleb et al. (2022) document higher entry into foster care in states with the must-access Prescription Drug Monitoring Programs (PDMPs). Arteaga and Barone (2023) find greater exposure to the opioid epidemic continuously increased the Republican vote share. In this paper, we document the positive externality of the supply-side intervention on opioid availability on the temperature-IPV relationship.

## 2 Background and Data

### 2.1 Potential physiological mechanisms

The primary channels through which increase in temperatures could affect violent behavior are physiological and psychological effects on impulse controls and aggression ([Anderson, 2001](#)). Higher temperatures can deteriorate mental health ([Mullins and White, 2019](#)), increasing anxiety, despair and isolation, which can in turn exacerbate substance use. Recent studies address the “looming confrontation between the world’s complex overdose crisis and its equally intensifying climate emergency” ([Ezell, 2023](#)), documenting that vulnerabilities associated with opioid use disorders are exacerbated by changes in climate, leading to more opioid-related emergency department visits and hospitalisations due to increases in temperature ([Chang et al., 2023](#); [Parks et al., 2023](#)).

Primarily, several prescription drugs, including opioids, have been associated with increases in criminal behavior and violence towards others ([Moore et al., 2010](#); [Sim, 2023](#)), and, in particular, towards intimate partners ([Moore et al., 2011](#)). Often opioid abuse co-moves with alcohol consumption ([Esser et al., 2019, 2021](#)), which is one of the main channels for increases in criminal behavior ([Anderson et al., 2018](#)) and domestic violence ([Klostermann and Fals-Stewart, 2006](#)).<sup>1</sup>

Altogether, there is suggestive evidence of potential interaction between opioids abuse, temperature, and intimate partner violence. Disruptions to opioid access induced by supply-side shocks such as the 2010 OxyContin reformulation that we study in this paper may have an ambiguous effect on the relationship between temperature and intimate partner violence. On the one hand, reduced opioid ac-

---

<sup>1</sup>In this regard, [Cohen and Gonzalez \(2024\)](#) find that 9% of weather-related crimes committed are in fact crimes triggered by an additional use of alcohol determined by weather conditions.

cess may favor substitution into other illicit drugs, inducing increases in violent behavior. On the other hand, supply-side interventions could reduce the overall reduction in the pool of addicts, reducing crime. Which of the two mechanisms prevails is an empirical question that we address in this paper.

## 2.2 Data

We briefly summarize the data (with complementary information provided in the Appendix). First, we retrieve administrative comprehensive data on reported cases of intimate partner violence at the finest temporal and geographical scale (Section 2.2.1). Second, we combine these data with granular weather data to identify the effect of temperature (Section 2.2.2). Third, we need information on the prescriptions of opioids in the United States to test the mediating role of the policy intervention (Section 2.2.3). Last, we combine the resulting data set with a number of additional information at various resolution (individual-, county-, and state-level) to explore mechanisms and channels of the relationship between temperature and intimate partner violence incidence.

### 2.2.1 Intimate Partner Violence Cases

We use data from the FBI’s National Incident-Based Reporting System (NIBRS) from 1991 to 2021, which contains reports of IPV cases to individual law enforcement agencies (*ORIs*, or jurisdictions) including information on the characteristics of the victim (e.g. age, gender), the offender (e.g. gender and relationship to the victim), and the incident date. We construct daily reports of IPV cases at the jurisdiction level. To classify incidents as intimate partner violence cases we follow [Dave et al. \(2023\)](#). We include aggravated assaults, simple assaults, forced sex, and intimidation, experience by female victims, from relationships that consist of

spouses, common-law spouses, boyfriends/girlfriends, homosexual partners, ex-spouses, and ex-boyfriends/girlfriends. Our primary dependent variable is the number of IPV cases per 100,000 people.<sup>2</sup>

Unfortunately, our data also come with drawbacks. First, the number of agencies participating in the NIBRS is increasing over time, moving from 609 in 1991 to 11,384 in 2021. Moreover, departments drop in and out of the sample over time, leading to an unbalanced panel. To attenuate this issue, we construct a panel at the jurisdiction-day level that is balanced at the year level, and we exploit within-year variation to obtain our estimates.<sup>3</sup>

Second, NIBRS are not representative of the whole United States, as only a self-selected sample of agencies report their crime. Nonetheless, despite this geographic coverage gaps, the NIBRS data are considered to be the most consistent and comparable national data available on daily crime rates in the US (US DOJ, 2018) and have been used widely in the literature (Card and Dahl, 2011; Burkhardt et al., 2019; Jones, 2022; Colmer and Doleac, 2023).

### 2.2.2 Weather

We process weather data from the ERA5-Land reanalysis product (Muñoz Sabater, 2019), which provides hourly temperature and precipitation from 1950 to present at a  $0.10^\circ$  spatial resolution ( $\approx 11\text{km}$ ). We combine weather data with 30 arc-

---

<sup>2</sup>We obtain the daily rate of IPV incidents scaling by population at the jurisdiction-level from (1) the [ICPSR website](#) at the University of Michigan and (2) the FBI's [Crime Data Explorer \(CDE\)](#). Jurisdiction-level population is not available for all the agencies. We test that our results are robust to the inclusion of this dropped sample, using the daily count of IPV cases as a dependent variable.

<sup>3</sup>Using the NIBRS data, Colmer and Doleac (2023) construct the panel in a similar way. However, they exclude agencies that did not report 12 months of data for that year to the reporting system. This is because Colmer and Doleac (2023) tests the role of within-year-month changes in concealed carry laws through a staggered design. Hence, they need to be sure how crimes are allocated across the months, avoiding measurement errors in reporting. This is not the case for our work, as we use a pre- and post-treatment policy design. We can so relax their restrictions.



seconds ( $\approx 1\text{km}$ ) population density information (Seirup and Yetman, 2006) to compute the county-level population-weighted average daily temperature and total precipitation for the representative individual in the county.<sup>4</sup> Although the IPV data are originally at the jurisdiction level, geographical coordinates are not available at the jurisdiction level. Therefore, we match jurisdictions to counties and we exploit these variables for the main analysis. We prefer ERA-5 Land weather data over PRISM Climate Group (PRISM, 2024) - another common weather data source for studies in the US (Colmer and Doleac, 2023; Molitor et al., 2023) - to explore the finer original temporal resolution of the ERA-5 Land product (hourly), in subsequent analysis on the mechanisms and robustness of our results.<sup>5</sup> In particular, we compute population-weighted daily maximum, minimum, and nighttime temperature.<sup>6</sup>

### 2.2.3 Opioids

We study a policy that exogenously curtailed access to opioids as a potential moderating or exacerbating factor to the temperature-IPV relationship. Since the 1990s, opioid prescriptions in the United States escalated quickly from 76 million to more than 250 million (Volkow et al., 2014). OxyContin, released by Purdue Pharma, was one of the main catalyst of such an opioid epidemic. OxyContin contained oxycodone - a narcotic analgesic - and was originally used to treat moderate to severe chronic pain. Nevertheless, it also had high risk for addiction and depen-

---

<sup>4</sup>When computing non-linear transformations in temperature and precipitation, we perform them at the grid-cell level before weighing and averaging, in order to preserve non-linearities in the original weather data, as common in climate econometrics (Hsiang, 2016).

<sup>5</sup>In future versions of this work, we aim at replicating the analysis using the PRISM data to check for the robustness of the results.

<sup>6</sup>Nighttime temperature is defined in three different ways: we consider the average temperature i) between 6pm and 6am; ii) between 8pm and 6am, iii) between 8pm and 8am.

dence, with an extensive number of people across the United States that started abusing it. To address the opioid crisis and reduce the misuse of OxyContin, Purdue Pharma developed an abuse deterrent version of the drug, making it more difficult to crush or dissolve. The version was approved by the Food and Drug Administration in 2010, which became effective immediately with the new formulation being distributed and the previous formulation being discontinued (Evans et al., 2019).

The reformulation led to a decrease in OxyContin abuse (Cicero and Ellis, 2015; Sessler et al., 2014). Nevertheless, it emerged a substitution pattern away from OxyContin to other illicit opioids such as heroin and synthetics, documented by an increase in overdoses related to these drugs in the post-reformulation period (Powell and Pacula, 2021). Recent studies document an increase in child physical abuse and neglect after OxyContin's reformulation (Evans et al., 2022) and heroin-involved IPV (Dave et al., 2023).

We exploit the national reformulation of OxyContin in 2010 as a result of an unanticipated, unilateral decision from the producers, Purdue Pharmaceutical. To obtain a measure of pre-intervention exposure to opioid prescription, we use the population-weighted mean number of all Schedule II opioid prescriptions per capita for the period 2006 to 2009 from the Centers for Disease Control (CDC) (Evans et al., 2022). This measure accounts for a broader set of prescription opioids than the intervention which was only targeting OxyContin, but it allows for more granular geographical variation at the county-level than previous studies exploiting state-level variation (Alpert et al., 2022). Appendix Figure A1 shows the spatial distribution of exposure to prescription opioids prior to the reformulation.

#### 2.2.4 Additional data

In additional analysis, we set out to better understand the mechanisms. We use auxiliary data for this purpose, which we briefly summarize here.<sup>7</sup>

**Medical marijuana laws.** We explore the substitution/complementary role of legal marijuana on access restrictions to opioids. Legal marijuana reduce opioid addiction and overdose deaths (Powell et al., 2018). We use information at the state-level with medical marijuana access in place (Evans et al., 2022) to study if its availability as a therapeutic substitute can mediate or exacerbate the differential effects of OxyContin reformulation on the temperature-IPV relationship. If legal access to medical marijuana favors substitution away from opioids, we should observe the opioid reformulation to be effective on attenuating the temperature effect only in states without medical marijuana laws. We test for this hypothesis in Section 5.

**Socio-demographic covariates.** We combine the data set with a plethora of socio-demographic covariates at the county-level. We include the percentage of the county population living in rural areas from the 2010 Census Bureau (Evans et al., 2022).<sup>8</sup> We use two measures to define rich and poor counties, respectively the median household income and the share of population in poverty from the Economic Research Service at the US Department of Agriculture<sup>9</sup>. We obtain county-level population by age, gender and ethnicity groups from the 2010 US Census Bureau,

---

<sup>7</sup>In the future version of this work, which will be available for the conference, we are going to explore further mechanisms on the relationship between IPV and temperature. This includes the role of alcohol consumption and mental health, happiness, and well-being using the Behavioral Risk Factor Surveillance System (BRFSS) survey, and potential temperature-induced changes in time-use as measured in the American Time Use Survey (ATUS).

<sup>8</sup>Data are publicly available [here](#).

<sup>9</sup>Data are publicly available [here](#).

Population Division<sup>10</sup>.

## 2.3 Descriptive Statistics

Our final combined dataset includes 11,176 jurisdictions (in 1,635 counties) across the United States, providing 44,170,732 unique jurisdiction-day observations. Table 1 provides summary statistics on the main variables of interest. On average, in our sample period 0.06 daily cases of IPV per 100,000 are reported. This rate is greater in the period before Oxycontin reformulation, and slightly smaller after the policy. Using data from Centers for Disease Control (CDC), the population-weighted mean number of all Schedule II opioid prescriptions per capita in the county for the period 2006 to 2009 is centered at 0.89 prescriptions. As for weather variables, in the whole sample the average daily temperature and total daily precipitation are 11.09 °C and 0.001 metres, respectively.

## 3 Temperature and Intimate Partner Violence

This section examines the impact of temperature on intimate partner violence (IPV) on females. First, we analyse the base relationship between daily temperature realizations and jurisdiction reports of IPV per 100,000 people. Next, we test the heterogeneity of the relationship across socio-demographic characteristics, climatic areas, and opioid exposure. We conclude providing robustness tests for our empirical analysis.

---

<sup>10</sup>Data are publicly available [here](#).

**Table 1:** Descriptive Statistics

	1991-2021		1991-2009		2010-2021	
	Mean	SD	Mean	SD	Mean	SD
<b>NBRIS</b>						
IPV on female per 100,000 people	0.06	0.59	0.08	0.72	0.05	0.52
<b>Opioid use</b>						
Per capita opioid prescriptions			0.89	0.35		
<b>ERA5-Land</b>						
Average daily temperature (°C)	11.09	10.50	9.46	10.13	11.88	10.58
Total daily precipitation (m)	0.00	0.01	0.00	0.01	0.00	0.01
Observations	44,170,732		26,425,814		17,744,918	

**Notes:** NBRIS descriptive statistics are weighted by jurisdiction-location population.

### 3.1 Econometric Framework

We model the relationship between temperature and intimate partner violence as follows:

$$Y_{idmy} = f(T_{c(i)dmy}, P_{c(i)dmy}) + \mu_{im y} + \phi_{wy} + \delta_{dw} + \varepsilon_{idmy} \quad (1)$$

where  $Y_{idmy}$  is the number of reported cases of intimate partner violence on female per 100,000 by jurisdiction  $i$  in day  $d$  of month  $m$  and year  $y$ ;  $\mu_{im y}$  are jurisdiction-month-year fixed effects;  $\phi_{wy}$  and  $\delta_{dw}$  are respectively week-of-year and day-of-week fixed effects; and, finally,  $\varepsilon_{idmy}$  is the stochastic error term, and it is clustered at the county-level. Equation 1 is estimated with population weights at the jurisdiction level. However, we also present estimates without including any regression weights, since population weights may lead to less precise estimates, as is common with data that represent group-level averages (Solon et al., 2015).

The term  $f(T_{c(i)dmy}, P_{c(i)dmy})$  is a function of average daily temperature (in °C)

and daily precipitation (in m). In the baseline specification, we linearly model the two weather variables:

$$f(T_{c(i)dmy}, P_{c(i)dmy}) = \beta_1 T_{c(i)dmy} + \beta_2 P_{c(i)dmy}$$

The coefficients  $\beta_1$  and  $\beta_2$  capture the linear impact of temperature and precipitation exploiting plausibly exogenous quasi-random variation in daily weather realizations (Deschênes and Greenstone, 2007). We also test for alternative specifications, where we account for potential non-linearities in the relationship between temperature and IPV.

## 3.2 Results

Table 2 reports the results from the estimation of Equation 1. Column (1) displays the unconditional estimate of the relationship between the number of intimate partner violence cases per 100,000 people and daily realizations of temperature and precipitation. We find that, on average, a one-degree Celsius increase in average daily temperature is associated with an increase by 0.0090 more cases of intimate partner violence per 100,000 people. This corresponds to a 1.34% increase with respect to the mean. Moreover, we find that precipitation is also associated with an increase in cases of intimate partner violence.

In Columns (2)-(3), we exploit month-year and county-month-year variation to identify the relationship temperature-intimate partner violence exposure by females. We estimate that, on average, a one-degree Celsius increase in average daily temperature is associated with an increase by 0.0107-0.0221 more cases of intimate partner violence per 100,000 people, corresponding to a 1.60%-3.30% increase with respect to the mean. The effect of precipitation is not robust to the inclusion of

county-month-year fixed effects.

Columns (4)-(5) replace county-month-year variation with, respectively, jurisdiction and jurisdiction-month-year fixed effects. As a result, we absorb much more variation in the relationship between intimate partner violence and temperature. We find that, on average, a one-degree Celsius is associated with 0.0040-0.0058 more cases of intimate partner violence per 100,000 people, a 0.60%-0.87% increase compared to the mean.

Finally, in Column (6) we present the results of our preferred specification. We estimate an identical specification to Column (5), but we weight the regression using jurisdiction-location population weights. Adding weights reduces the accuracy of the estimate. However, the magnitude of the temperature coefficient remains identical to the unweighted estimate. We find that, on average, that a one-degree Celsius increase in average daily temperature is associated with 0.0005 more cases of intimate partner violence per 100,000 people. This corresponds to a 0.87% increase compared to the weighted average.

Our baseline results are robust to various specification tests, including dropping Covid period or controlling for Covid period (Table A3); trimming the sample (Table A4); expressing the dependent variable as count variable; testing alternative fixed-effects (Table A6), and standard errors clustered at the state level (Table A7). We also explore non-linear specifications of temperature, using 5-degree temperature bins (Figure A2 Table A8), share of hours in a day in a temperature bin (Table A9), and a up to 4-degree polynomial (Table A10). The estimates suggest that a linear specification is a good approximation. Finally, we use lags and leads to control for potential displacements of intimate partner violence cases, and anticipatory behaviours (Table A11). We observe displacement effects offsetting nearly 60 percent of the contemporaneous effect after 21 days. As Cohen and Gonzalez (2024), we find no impact of leads, except for the first lead for temperature. This correlation

most likely is due to the correlation during night between the average temperature on day  $d$  and the average temperature on day  $d - 1$ .

Moreover, our estimates align in sign and magnitude with previous literature on the relationship between temperature and crimes in the United States. [Ranson \(2014\)](#) shows that the temperature-violent crime relationship is approximately linear. He finds that an additional day between 90 and 99 °F ( $\approx$  between 32 and 35 °C) is associated with an increase by 0.6%, 0.9%, 0.7% and 0.4% in murder, rape, aggravated and simple assault cases respectively. [Colmer and Doleac \(2023\)](#) estimates that a one-degree Celsius is associated with an increase between 0.8 and 5.7% in the mean murder rate, depending on the specification.

**Table 2:** Temperature and Intimate Partner Violence

	Intimate Partner Violence per 100,000 people					
	(1)	(2)	(3)	(4)	(5)	(6)
Temperature (°C)	0.0090*** (0.0006)	0.0221*** (0.0018)	0.0107*** (0.0010)	0.0040*** (0.0002)	0.0058*** (0.0003)	0.0005** (0.0002)
Precipitation (m)	0.6838*** (0.2552)	1.464*** (0.2831)	0.3675 (0.2368)	-0.3330* (0.1883)	-0.0232 (0.2146)	-0.0126 (0.0101)
Observations	44,170,732	44,170,732	44,170,732	44,170,732	44,170,732	44,170,732
Dependent Variable Mean	0.66975	0.66975	0.66975	0.66975	0.66975	0.05741
Month-Year FE		✓				
Week-of-Year FE		✓	✓	✓	✓	✓
Day-of-Week FE		✓	✓	✓	✓	✓
County-Month-Year FE			✓			
Jurisdiction FE				✓		
Jurisdiction-Month-Year FE					✓	✓
Jurisdiction-Location Population Weights						✓

**Notes:** The dependent variable is the number of intimate partner violence per 100,000 people. Temperature is the average daily temperature measured in degrees Celsius (°C). Precipitation is the total daily precipitation measured in metres (m). Standard errors are clustered at the county level. Significance levels are indicated as follows: \* $p < 0.10$ , \*\* $p < 0.05$ , \*\*\* $p < 0.01$ .

### 3.3 Heterogeneity

We present a heterogeneity analysis to provide additional insights on the relationship between temperature and intimate partner violence.



**County-level socio-demographic characteristics.** We first test the presence of heterogeneous effects of temperature on IPV incidence across socio-demographic characteristics at the county-level.<sup>11</sup> Table A1 reports the coefficients for each socio-demographic feature considered. First, we test the urban-rural divide, defining a jurisdiction as urban if it is within a county whose urban population share is above the median populationweighted urban share in the sample. Our estimates suggest that the effect of temperature on IPV on females is more pronounced in urban areas. This is consistent with the findings from Cohen and Gonzalez (2024), where the authors find that a relationship between temperature and crime only for municipalities with more than 10,000 inhabitants in Mexico. This can be explained through different factors, such as urban heat island effect, changes in social interactions during very hot days.<sup>12</sup> Additionally, we find that counties with above median poverty rate and below level of income drive the relationship

**Climatic conditions.** There are numerous ways in which people and communities adapt to their current climate, including biological acclimatization, infrastructure investments, architectural styles, cooling appliances (e.g. air conditioning). It is beyond the scope of this paper to identify each heterogeneity component of adaptation to the impact of temperature. However, we can explore for the role of long-term climatic conditions as a proxy for behavioral and physiological adap-

---

<sup>11</sup>To do so, we allow the temperature-intimate partner violence relationship to vary cross-sectionally across each group (Carleton et al., 2022). We then specify the regression as follows:

$$Y_{aidmy} = f_a(T_{c(i)dmy}, P_{c(i)dmy}) + \mu_{aimy} + \phi_{wy} + \delta_{dw} + \varepsilon_{aidmy}$$

where the function  $f(T_{c(i)dmy}, P_{c(i)dmy})$  is interacted with a categorical (or dummy) variable identifying the specific heterogeneity group of interest  $a$ , and  $\gamma_{aidy}$  are group-jurisdiction-month-year fixed effects. Notice that the model does not include uninteracted terms for the group  $a$  because they are collinear with  $\gamma_{aidy}$ .

<sup>12</sup>High temperatures may discourage outdoor activities (Graff Zivin and Neidell, 2014) and exacerbate feelings of isolation (Mullins and White, 2019).


tation. If any form of adaptation is taking place, we would expect that the effect of temperature diminishes as we move along the climate distribution, from colder to warmer counties. In Table A2 we test for this hypothesis, reporting the coefficients of the interaction with terciles of a 30-year mean of average daily temperature. Our estimates suggest that average daily temperature increases the number of IPV cases on females especially in the warmer counties of the United States. In magnitude, the effect is about 2 to 5 times what it is estimated in cold and temperature counties. This evidence of no mediating effect coming from adaptation. This is in contrast with the literature on physical health (Heutel et al., 2021), but it is consistent with findings on the relationships between temperature and mental health, and between temperature and violence (Mullins and White, 2019; Evans et al., 2023).

**Levels of Opioid Prescriptions.** To add on previous literature that documents a positive association between warmer temperature and opioid abuse (Chang et al., 2023; Parks et al., 2023), we explore whether the effect of temperature on IPV varies by opioid prescriptions per capita. Figure 1 reports the marginal effect of temperature on IPV cases on females interacted with the population-weighted rate of OxyContin misuse prior to the reformulation. We document a positive effect, i.e., higher temperature increases IPV in counties with higher opioid abuse, suggesting a compounding effect of environmental factors and opioid use disorders.

## 4 Opioid reformulation in temperature-induced IPV

In this section we explore whether OxyContin reformulation in 2010 has amplified or reduced the impact of temperature on intimate partner violence (IPV) on

**Figure 1:** Temperature and Opioid prescription reformulation



*Notes:* The figure plots the marginal effect of temperature on IPV on female victims per 100,000 by population-weighted mean per capita opioid prescriptions for the pre-period policy 2006 to 2009.

females. We also provide several robustness tests that corroborate our estimates.

#### 4.1 A triple difference approach

To assess whether policy interventions during the time period of the analysis mediate or exacerbate the temperature-IPV relationship, we design a triple difference (DiDiD) specification which exploits within-county changes in opioid reformulation alongside with within-county variation in temperature over time. We estimate the following specification:

$$Y_{idmcy} = f(T_{c(i)dmy}, exposure_c, post_y) + g(P_{c(i)dmy}) + \mu_{imy} + \phi_{wy} + \delta_{dw} + \varepsilon_{idmcy} \quad (2)$$

where we interact daily county-level temperature  $T_{c(i)dm_y}$  with county-level pre-2010 exposure to prescription opioids ( $exposure_c$ ) and an indicator variable  $post_y$  which takes value of one for the post-reformulation period, starting from 2010. We also allow for heterogeneous effects of OxyContin’s reformulation and estimate differential impacts by time window after the policy intervention. To do so, we interact temperature and pre-intervention exposure with four dummies that, respectively, take value of one for i) pre-reformulation years (2006 to 2009); ii) years immediately following reformulation (2011-2013); iii) years 2014 to 2016 for medium-run impacts; iv) long-run impacts for several years post-reformulation (2017-onwards).

As in the baseline specification in Equation (1), we control for precipitation an account for jurisdiction-month-year, week-of-year, day-of-week fixed effects. We also include precipitation and year-specific temperature controls, accounting for time-varying changes in the direct effects of temperature and precipitation on homicide. This implies that we allow for the temperature-IPV relationship to vary flexibly over time, and we focus on identifying the relative effect of temperature between high- and low-exposure mean opioid prescriptions per capita before and after the reformulation.

The coefficient on the triple interaction term is identified under the assumption that the temperature-IPV relationship would have stayed constant had the opioid reformulation not occurred. The most conservative specification also accounts for month-year jurisdiction-specific unobserved heterogeneity, which provides further support for the assumption that within-jurisdiction variation in daily temperature is uncorrelated with other unobserved factors that may also affect the probability of intimate partner violence.

To rule out any potential differential pre-intervention trends between high- and low-exposure counties to prescription opioids, we also estimate an event study

specification where the interaction term between daily temperature and pre-2010 exposure to prescription opioids is interacted with a set of event year coefficients for each of the 16 years in the sample. We so identify differences in the temperature-IPV relationship between counties with higher and lower pre-intervention exposure in year  $y$  compared to 2010, the year OxyContin was reformulated.


We estimate Equation (2) using weighted-least squares where the weights are the average population in the jurisdiction during the sample period. Standard errors are clustered at the county-level.

## 4.2 Results

To explore the potential moderating or exacerbating role of the opioid reformulation, Figure 2 shows the regression-adjusted relationship between temperature and intimate partner violence cases separately in a sample after the policy reformulation with low opioid prescriptions in Panel A and high opioid prescriptions in Panel B. We determine each group using the medial opioid prescription (1.02) sample split. The relationship between temperature and IPV is positive in both panels, confirming the previous result that higher temperature increases IPV cases. In counties where opioid prescriptions per capita are high, however, the slope of the positive gradient reduces dramatically and is not statistically different than zero at conventional levels. This result suggests that the opioid reformulation has a mitigating effect on the relationship between temperature and IPV.

To provide further evidence on the role of the opioid reformulation, we present in Figure 3 the results from the triple difference (DiDiD) research design. The graph on the left-hand side reports the coefficient associated with the triple interaction term between daily temperature, pre-reformulation exposure, post-2010 time

**Figure 2:** Temperature and Opioid prescription reformulation



*Notes:* Panels (a) and (b) show binned scatterplots with 50 bins and a linear regression on the underlying data. Each shows the correlation net of jurisdiction-month-year, week-of-year, day-of-week fixed effects between residualized IPV rate and residualized temperature. The panels show the relationship for below- (a) and above-median (b) exposure to prescription opioids in the sample after the opioid reformulation in 2010.

indicator is negative and statistically significant, indicating that a one-degree Celsius increase in temperature is associated with 0.008 fewer IPV cases per 100,000 people after the reformulation occurred in 2010, a 7.9% reduction at the mean. Compared to our baseline estimate of the temperature-IPV relationship ( $0.0005/1^{\circ}\text{C}$ , in column 6, Table 2), the results suggest that the opioid reformulation strongly attenuates the temperature-IPV relationship.


On the right-hand side of Figure 3, we explore the hypothesis that the attenuation role of OxyContin's reformulation in the temperature-IPV relationship varies over time since the implementation of the policy. We consider short-run, medium-run, and long-run impacts of the policy by estimating four different triple interaction terms that also include the pre-policy period. The triple interaction term with the time indicator of the pre-policy period is small in magnitude and not statistically significant at conventional levels, allaying concerns on differential pre-intervention trends between high- and low-exposure counties to prescription opi-

oids. The triple interaction term is negative and statistically significant at the 95% level only for the years immediately following reformulation, 2011 to 2013. The coefficient is similar in magnitude to the baseline triple difference estimate: a one-degree Celsius increase in temperature is associated with 0.007 fewer IPV cases per 100,000 people after the reformulation occurred in 2010. The effect fades away for several years post-reformulation suggesting that the attenuation role of opioid reformulation is only temporary.

Our results are robust to a variety of alternative specifications that: use alternative fixed effects, accounting for jurisdiction-day, jurisdiction-month-year, and date fixed effects (Appendix Figure A3); exclude the period of Covid after 2019 (Appendix Figure A4); account for wyear-specific precipitation slopes (Appendix Figure A5); cluster standard errors at the state level (Appendix Figure A6); restrict the sample to different seasons (Appendix Figure A7) and different climates (Appendix Figure A8).

Figure 4 presents an event study visualization of our results. We report point estimates and 95% confidence intervals on the triple interaction terms between daily temperature, pre-reformulation opioid exposure, and year indicators, with 2010, the year in which OxyContin was reformulated, normalized to zero. The estimates coefficients in the years prior to reformulation are statistically indistinguishable from zero, consistent with no trend temperature differences between higher- and lower-exposure counties during the pre-reformulation period. After the reformulation in 2010, for an increase in daily temperatures, the number of intimate partner violence cases per 100,000 inhabitants decreases in high-exposure counties relative to low-exposure counties. The responsiveness of the temperature-IPV relationship decreases in high-exposure resulting in an average relative decrease

**Figure 3:** Triple difference (DiDiD) of Opioid reformulation policy in 2010 on the temperature-IPV relationship




*Notes:* The figure plots the coefficients from the triple interaction term between average daily county-level temperature, pre-intervention exposure as the population-weighted mean per capita opioid prescription, and in the left graph an indicator variable that takes the value of one for the post-reformulation years, 2010-2021, and in the right graph an indicator variable (short-run) equal to one for the years immediately following reformulation, 2011 to 2013, another one (medium-run) equal to one for the years 2014 to 2016, and another indicator variable (long-run) for the years 2017-2021, several years post-reformulation. The regression also controls for year-specific temperature and precipitation coefficients, jurisdiction-month-year, week-of-year and day-of-week fixed effects. Bins represent the 95% confidence intervals with standard errors clustered at the county-level.

of 0.005 IPV cases per 100,000 people/ $1^{\circ}\text{C}/\text{day}$ . The policy-induced attenuation of the temperature effect is persistent up to 5 years after the policy change and fades away after 2015, consistent with the short-run effect highlighted in the triple difference findings.



**Figure 4:** Event study of the differential effect of Opioid reformulation on the temperature-IPV relationship



*Notes:* The figure plots the coefficients associated with the triple interaction term between daily-temperature, pre-intervention opioid exposure and year dummies in a regression where the outcome variable is the number of IPV cases per 100,000 people. The regression also controls for year-specific temperature and precipitation coefficients, jurisdiction-month-year, week-of-year and day-of-week fixed effects. Shaded area represent the 95% confidence intervals with standard errors clustered at the county-level.

## 5 Mechanisms

In this section, we discuss additional results that provide insights into the channels through which opioid reformulation attenuates the temperature-IPV relationship.

We begin by documenting how attenuation effect of opioid reformulation on the temperature-IPV gradient is heterogeneous across socio-economic characteristics at the county-level. We estimate a quadruple interaction terms that combines temperature, pre-policy opioid exposure, a post-policy time indicator and an in-

indicator that categorizes counties above and below the sample median in terms of urban/rural, unemployment levels, pre-dominant ethnic groups (White, Black, Hispanic), income and poverty measures. Figure 5 displays the coefficient on the quadruple interaction term for each socio-demographic margin considered. Although there is meaningful overlap in the confidence intervals across sub-groups of population studied, we discuss in detail below some suggestive evidence of the heterogeneous impacts of the policy.

**Urban-rural divide.** Greater population density may induce a differentially stronger attenuation effect of opioid reformulation in urban areas if IPV is more responsive to temperature in these areas. Conversely, urban areas may facilitate access to opioid substitutes such as illegal substances which can themselves induce increases in IPV and exacerbate the IPV elasticity to temperature. We test for which of these two mechanisms prevails empirically and show that the policy is effective only in urban areas. Opioid reformulation mitigates the temperature-IPV gradient in urban areas, in contrast with previous temperature-homicide studies that document the effect to be attenuated by gun law restrictions only in rural areas (Colmer and Doleac, 2023). A potential explanation behind this finding is that higher temperatures can halt addiction treatment programs and derail critical medication supply chains, which can be particularly true in rural areas, where fewer health-care providers are available, and patients often need to travel substantial distances to receive care.

**Economic stress.** Higher unemployment rates often lead to increased economic stress and hardship in communities. Individuals facing economic difficulties may be more susceptible to substance abuse as a coping mechanism. We document that the opioid reformulation is effective at mitigating the effect of temperature on IPV only in counties with unemployment rate above the median. This result


seems at first in contrast with additional heterogeneous impacts of the policy that, although imprecisely, suggest that the policy is effective in richer counties with lower poverty rates. We explain this contrast with greater access to alternative coping mechanisms in richer counties, such as mental health services, recreational activities, and community support. This could reduce the reliance on opioids as a coping mechanism for stress and pain. In these counties, the policy might reinforce existing support systems. Moreover, opioid abuse can have different socioeconomic drivers. In certain cases, unemployment might be a significant factor, while in richer counties, it could be related to other factors, including stress, mental health, or over-prescription.

**Environmental justice.** The impact of the opioid prescription reformulation policy exhibits heterogeneity across counties with varying racial demographics, emphasizing the intersectionality of environmental justice concerns ([Banzhaf et al., 2019](#)). In counties with predominantly white populations and below-median black population, the policy is effective at mitigating the effect of temperature due to potentially higher access to healthcare resources. In contrast, counties with larger Hispanic populations may face unique challenges such as language barriers and disparities in healthcare access, influencing the policy's efficacy differently. Understanding these variations is crucial for ensuring environmental justice, as vulnerable communities may experience different levels of protection and support.

**Medical Marijuana Laws.** Last, we test for the extent to which the availability of medical marijuana mediates the effects of opioid reformulation on the temperature-IPV relationship through therapeutic substitution effects. We find that in states where medical marijuana is not accessible, the policy is effective at mitigating the harmful effect of temperature. In areas where medical marijuana is legally accessible, people may choose to substitute opioids with marijuana, potentially reducing

the likelihood of turning to more destructive and uncontrolled substances such as heroin. Although associative, these results suggest that access to marijuana dampens the substitution to more destructive and uncontrolled substances, however, also muting the co-benefits of the opioid reformulation on the effect of temperatures.

**Figure 5:** Heterogeneous effect of the Opioid reformulation on the temperature-IPV relationship



*Notes:* The figure plots the coefficients associated with the quadruple interaction term between daily-temperature, pre-intervention opioid exposure, post-2010 dummy, and each population subgroup/county type in a regression where the outcome variable is the number of IPV cases per 100,000 people. The regression also controls for year-specific temperature and precipitation coefficients, jurisdiction-month-year, week-of-year and day-of-week fixed effects. Bins area represent the 95% confidence intervals with standard errors clustered at the county-level.

**Sleep deprivation.** Healthy sleep is an important aspect of successfully treating opioid use disorders. Opioid abuse damages sleep duration and quality (Bertz et al., 2019). At the same time, increases in nighttime temperatures can amplify nights of insufficient sleep and deteriorate sleep quality (Minor et al., 2022; Obradovich et al., 2017), which in turn, might diminish the ability to cope with stress, leading individuals to respond to aversive stimuli in an aggressive manner (Rauer and El-Sheikh, 2012). We explore whether the reformulation of opioid prescription can help moderate this channel by using a measure of nighttime temperature (Appendix Table A12). We document that, although nighttime temperature has a positive significant effect on IPV, the effect is attenuated after the reformulation in counties with higher exposure to opioid prescription. This result suggests that the negative supply-side shock induced by the policy can help improve sleep duration and quality and moderate the compounding effect of violence induced by opioid abuse and warmer nighttime temperature.

**Social interactions.** Intimate partner violence is a crime that more often occurs at home rather than outside. Increases in temperature could affect criminal activity by increasing the likelihood of social interaction. Our baseline results confirm this intuition since we document a negative effect of rainier days on IPV. We test whether opioid reformulation has a differential effect on the precipitation-IPV relationship (Appendix Figure A9) and document an effect that is not significant at any conventional level. This result suggests that on opioid reformulation does not affect IPV offenses that are sensitive to precipitation, i.e., that arise in contexts with greater social interaction.<sup>13</sup>

---

<sup>13</sup>In the future version of the manuscript, we plan to explore how our estimates vary between locations of crime (at home, outside, other location) to further disentangle the potential social channel.

## 6 Discussion and conclusions

Our study contributes to understanding how warmer temperature may lead to a higher incidence of female victims of intimate partner violence. We address this question in the United States using administrative daily-level data over the past three decades to test for the effect the impact of higher temperatures on IPV perpetration in the United States and examine the moderating effect of policies targeting prescription opioid misuse.

Under these circumstances the policy environment may exacerbate or mitigate the impact of temperature on intimate partner violence. We combine two quasi-experimental research design to show how the 2010 Oxycontin reformulation has unintendedly attenuated the relationship between temperature and intimate partner violence on females. The policy has been particularly effective in urban and richer counties, and where potential substitutes like marijuana are not legally available. Using triple-difference and event-study designs, our findings reveal a positive association between higher temperatures and IPV rates. However, after the opioid reformulation, the relationship weakens, indicating a 7.9% reduction in IPV cases per 100,000 people for every one-degree Celsius temperature increase in high opioid exposure counties. The event-study analysis suggests that the policy's moderating effect diminishes after five years.

Armed with the estimates of our baseline specification, we conduct a back-of-the-envelope calculation to monetize the net social benefit of the opioid reformulation policy on the temperature-IPV relationship. We estimate that, on average, after OxyContin reformulation one-degree Celsius increase is associated with 9,714 fewer female victims of IPV. Using the lifetime cost of an intimate partner violence of \$103,767 (\$2014) from [Peterson et al. \(2018\)](#) we calculate that, on average, the policy has generated in the period 2010-2021 an annual social benefit

of approximately \$308 million for one-degree Celsius increase in the temperature for the interquartile shift in opioid prescriptions.<sup>14</sup> This back-to-the-envelope estimate highlights that attenuating the effect of temperature on IPV has large welfare implications.

---

<sup>14</sup>This is obtained as follows:  $-0.008 \text{ IPV cases per } 100,000 \text{ people} \times 365 \text{ days} \times \text{Population between } 2006\text{-}2009 \times 0.32 \text{ (opioid prescription interquartile range)} \times \$103,767$

## References

- Aizer, A. (2010). The gender wage gap and domestic violence. *American Economic Review* 100(4), 1847–1859.
- Alpert, A., W. N. Evans, E. M. Lieber, and D. Powell (2022). Origins of the opioid crisis and its enduring impacts. *The Quarterly Journal of Economics* 137(2), 1139–1179.
- Anderberg, D., H. Rainer, J. Wadsworth, and T. Wilson (2016). Unemployment and domestic violence: Theory and evidence. *The Economic Journal* 126(597), 1947–1979.
- Anderson, C. A. (2001). Heat and violence. *Current directions in psychological science* 10(1), 33–38.
- Anderson, D. M., B. Crost, and D. I. Rees (2018). Wet laws, drinking establishments and violent crime. *The Economic Journal* 128(611), 1333–1366.
- Angelucci, M. and R. Heath (2020). Women empowerment programs and intimate partner violence. In *AEA Papers and Proceedings*, Volume 110, pp. 610–614.
- Annan, F. and W. Schlenker (2015). Federal crop insurance and the disincentive to adapt to extreme heat. *American Economic Review* 105(5), 262–266.
- Arteaga, C. and V. Barone (2022). A manufactured tragedy: The origins and deep ripples of the opioid epidemic. *Working paper*. [https://viquibarone.github.io/baronevictoria/Opioids\\_ArteagaBarone.pdf](https://viquibarone.github.io/baronevictoria/Opioids_ArteagaBarone.pdf).
- Arteaga, C. and V. Barone (2023). Democracy and the opioid epidemic. *Working paper*. [https://viquibarone.github.io/baronevictoria/OpioidsDemocracy\\_ArteagaBarone.pdf](https://viquibarone.github.io/baronevictoria/OpioidsDemocracy_ArteagaBarone.pdf).



- Banzhaf, S., L. Ma, and C. Timmins (2019). Environmental justice: The economics of race, place, and pollution. *Journal of Economic Perspectives* 33(1), 185–208.
- Baylis, P. (2020). Temperature and temperament: Evidence from Twitter. *Journal of Public Economics* 184, 104161.
- Behrer, A. P. and V. Bolotnyy (2022). Heat, crime, and punishment. Policy Research Working Paper 9909, World Bank, Washington, DC.
- Bertz, J. W., D. H. Epstein, D. Reamer, W. J. Kowalczyk, K. A. Phillips, A. P. Kennedy, M. L. Jobes, G. Ward, B. A. Plitnick, M. G. Figueiro, et al. (2019). Sleep reductions associated with illicit opioid use and clinic-hour changes during opioid agonist treatment for opioid dependence: Measurement by electronic diary and actigraphy. *Journal of Substance Abuse Treatment* 106, 43–57.
- Bobonis, G. J., M. González-Brenes, and R. Castro (2013). Public transfers and domestic violence: The roles of private information and spousal control. *American Economic Journal: Economic Policy* 5(1), 179–205.
- Burkhardt, J., J. Bayham, A. Wilson, E. Carter, J. D. Berman, K. O'Dell, B. Ford, E. V. Fischer, and J. R. Pierce (2019). The effect of pollution on crime: Evidence from data on particulate matter and ozone. *Journal of Environmental Economics and Management* 98, 102267.
- Card, D. and G. B. Dahl (2011). Family violence and football: The effect of unexpected emotional cues on violent behavior. *The Quarterly Journal of Economics* 126(1), 103–143.
- Carleton, T., A. Jina, M. Delgado, M. Greenstone, T. Houser, S. Hsiang, A. Hultgren, R. E. Kopp, K. E. McCusker, I. Nath, et al. (2022). Valuing the global mortality

- consequences of climate change accounting for adaptation costs and benefits. *The Quarterly Journal of Economics* 137(4), 2037–2105.
- Carleton, T. A. and S. M. Hsiang (2016). Social and economic impacts of climate. *Science* 353(6304), aad9837.
- Centers for Disease Control and Prevention (2022). Fast facts: Preventing intimate partner violence. Retrieved from <https://www.cdc.gov/violenceprevention/intimatepartnerviolence/fastfact.html>. Accessed on 01-15-2024.
- Chang, H. H., H. Zhang, A. D. Latimore, B. P. Murray, R. R. D’Souza, N. Scovronick, M. O. Gribble, and S. T. Ebel (2023). Associations between short-term ambient temperature exposure and emergency department visits for amphetamine, cocaine, and opioid use in California from 2005 to 2019. *Environment international* 181, 108233.
- Cicero, T. J. and M. S. Ellis (2015). Abuse-deterrent formulations and the prescription opioid abuse epidemic in the united states: lessons learned from oxycontin. *JAMA psychiatry* 72(5), 424–430.
- Cohen, F. and A. Dechezleprêtre (2022). Mortality, temperature, and public health provision: evidence from mexico. *American Economic Journal: Economic Policy* 14(2), 161–192.
- Cohen, F. and F. Gonzalez (2024). Understanding the link between temperature and crime. *American Economic Journal: Economic Policy*.
- Cohen, F., F. Gonzalez, et al. (2018). Understanding interpersonal violence: the impact of temperatures in mexico. Technical report, Grantham Research Institute on Climate Change and the Environment.

- Colmer, J. and J. L. Doleac (2023). Access to guns in the heat of the moment: more restrictive gun laws mitigate the effect of temperature on violence. *Review of Economics and Statistics*, 1–40.
- Dave, D. M., B. Erten, P. Keskin, and S. Zhang (2023). From addiction to aggression: The spillover effects of opioid policies on intimate partner violence. NBER Working Papers 31609, National Bureau of Economic Research.
- Deschênes, O. and M. Greenstone (2007). The economic impacts of climate change: evidence from agricultural output and random fluctuations in weather. *American Economic Review* 97(1), 354–385.
- Erten, B. and P. Keskin (2018). For better or for worse?: Education and the prevalence of domestic violence in Turkey. *American Economic Journal: Applied Economics* 10(1), 64–105.
- Erten, B. and P. Keskin (2021). Trade-offs? the impact of wto accession on intimate partner violence in cambodia. *Review of Economics and Statistics*, 1–40.
- Esser, M. B., G. P. Guy Jr, K. Zhang, and R. D. Brewer (2019). Binge drinking and prescription opioid misuse in the us, 2012–2014. *American journal of preventive medicine* 57(2), 197–208.
- Esser, M. B., C. M. Pickens, G. P. Guy Jr, and M. E. Evans (2021). Binge drinking, other substance use, and concurrent use in the us, 2016–2018. *American journal of preventive medicine* 60(2), 169–178.
- Evans, M. F., L. Gazze, and J. Schaller (2023). Temperature and maltreatment of young children. NBER Working Papers 31522, National Bureau of Economic Research.

- Evans, M. F., M. C. Harris, and L. M. Kessler (2022). The hazards of unwinding the prescription opioid epidemic: Implications for child maltreatment. *American Economic Journal: Economic Policy* 14(4), 192–231.
- Evans, W. N., E. M. Lieber, and P. Power (2019). How the reformulation of oxycontin ignited the heroin epidemic. *Review of Economics and Statistics* 101(1), 1–15.
- Ezell, J. M. (2023). Climate change and the opioid epidemic. *Journal of addiction medicine* 17(5), 500–502.
- Garg, T., G. C. McCord, and A. Montfort (2020). Can social protection reduce environmental damages? *Available at SSRN* 3465356.
- Gihleb, R., O. Giuntella, and N. Zhang (2022). The effect of mandatory-access prescription drug monitoring programs on foster care admissions. *Journal of Human Resources* 57(1), 217–240.
- Graff Zivin, J. and M. Neidell (2014). Temperature and the allocation of time: Implications for climate change. *Journal of Labor Economics* 32(1), 1–26.
- Heilmann, K., M. E. Kahn, and C. K. Tang (2021). The urban crime and heat gradient in high and low poverty areas. *Journal of Public Economics* 197, 104408.
- Heutel, G., N. H. Miller, and D. Molitor (2021). Adaptation and the mortality effects of temperature across us climate regions. *Review of Economics and Statistics* 103(4), 740–753.
- Hsiang, S. (2016). Climate econometrics. *Annual Review of Resource Economics* 8, 43–75.
- Jones, B. A. (2022). Dust storms and violent crime. *Journal of Environmental Economics and Management* 111, 102590.

- Klostermann, K. C. and W. Fals-Stewart (2006). Intimate partner violence and alcohol use: Exploring the role of drinking in partner violence and its implications for intervention. *Aggression and violent behavior* 11(6), 587–597.
- Minor, K., A. Bjerre-Nielsen, S. S. Jonasdottir, S. Lehmann, and N. Obradovich (2022). Rising temperatures erode human sleep globally. *One Earth* 5(5), 534–549.
- Molitor, D., J. T. Mullins, and C. White (2023). Air pollution and suicide in rural and urban America: Evidence from wildfire smoke. *Proceedings of the National Academy of Sciences* 120(38), e2221621120.
- Moore, B. C., C. J. Easton, and T. J. McMahon (2011). Drug abuse and intimate partner violence: a comparative study of opioid-dependent fathers. *American Journal of Orthopsychiatry* 81(2), 218.
- Moore, T. J., J. Glenmullen, and C. D. Furberg (2010). Prescription drugs associated with reports of violence towards others. *PloS one* 5(12), e15337.
- Mukherjee, A. and N. J. Sanders (2021). The causal effect of heat on violence: Social implications of unmitigated heat among the incarcerated. NBER Working Papers 28987, National Bureau of Economic Research.
- Mullins, J. T. and C. White (2019). Temperature and mental health: Evidence from the spectrum of mental health outcomes. *Journal of health economics* 68, 102240.
- Mullins, J. T. and C. White (2020). Can access to health care mitigate the effects of temperature on mortality? *Journal of Public Economics* 191, 104259.
- Muñoz Sabater, J. (2019). ERA5-Land hourly data on single levels from 1950 to present. Accessed: 11-17-2023.

- National Coalition Against Domestic Violence (2020). Domestic violence. Retrieved from [https://assets.speakcdn.com/assets/2497/domestic\\_violence-2020080709350855.pdf?1596811079991](https://assets.speakcdn.com/assets/2497/domestic_violence-2020080709350855.pdf?1596811079991). Accessed on 01-15-2024.
- Obradovich, N., R. Migliorini, S. C. Mednick, and J. H. Fowler (2017). Nighttime temperature and human sleep loss in a changing climate. *Science advances* 3(5), e1601555.
- Parks, R. M., S. T. Rowland, V. Do, A. K. Boehme, F. Dominici, C. L. Hart, and M.-A. Kioumourtzoglou (2023). The association between temperature and alcohol-and substance-related disorder hospital visits in New York State. *Communications Medicine* 3(1), 118.
- Peterson, C., M. C. Kearns, W. L. McIntosh, L. F. Estefan, C. Nicolaidis, K. E. McCollier, A. Gordon, and C. Florence (2018). Lifetime economic burden of intimate partner violence among US adults. *American journal of preventive medicine* 55(4), 433–444.
- Powell, D. and R. L. Pacula (2021). The evolving consequences of oxycontin reformulation on drug overdoses. *American Journal of Health Economics* 7(1), 41–67.
- Powell, D., R. L. Pacula, and M. Jacobson (2018). Do medical marijuana laws reduce addictions and deaths related to pain killers? *Journal of Health Economics* 58, 29–42.
- PRISM (2024). Prism climate data. <https://prism.oregonstate.edu/>. Accessed 01-16-2024.
- Radcliffe, P., D. Gadd, J. Henderson, B. Love, D. Stephens-Lewis, A. Johnson, E. Gilchrist, and G. Gilchrist (2021). What role does substance use play in intimate partner violence? A narrative analysis of in-depth interviews with men

- in substance use treatment and their current or former female partner. *Journal of Interpersonal Violence* 36(21-22), 10285–10313.
- Ranson, M. (2014). Crime, weather, and climate change. *Journal of Environmental Economics and Management* 67(3), 274–302.
- Rauer, A. J. and M. El-Sheikh (2012). Reciprocal pathways between intimate partner violence and sleep in men and women. *Journal of Family Psychology* 26(3), 470.
- Schilbach, F. (2019). Alcohol and self-control: A field experiment in India. *American Economic Review* 109(4), 1290–1322.
- Seirup, L. and G. Yetman (2006). U.s. census grids (summary file 1), 2000. <https://doi.org/10.7927/H4B85623>. Accessed 11-17-2023.
- Sessler, N. E., J. M. Downing, H. Kale, H. D. Chilcoat, T. F. Baumgartner, and P. M. Coplan (2014). Reductions in reported deaths following the introduction of extended-release oxycodone (oxycontin) with an abuse-deterrent formulation. *Pharmacoepidemiology and drug safety* 23(12), 1238–1246.
- Sim, Y. (2023). The effect of opioids on crime: Evidence from the introduction of oxycontin. *International Review of Law and Economics* 74, 106136.
- Solon, G., S. J. Haider, and J. M. Wooldridge (2015). What are we weighting for? *Journal of Human Resources* 50(2), 301–316.
- Stevenson, B. and J. Wolfers (2006). Bargaining in the shadow of the law: Divorce laws and family distress. *The Quarterly Journal of Economics* 121(1), 267–288.
- Stone, R. and E. F. Rothman (2019). Opioid use and intimate partner violence: A systematic review. *Current Epidemiology Reports* 6, 215–230.


- Tur-Prats, A. (2019). Family types and intimate partner violence: A historical perspective. *Review of Economics and Statistics* 101(5), 878–891.
- Tur-Prats, A. (2021). Unemployment and intimate partner violence: A cultural approach. *Journal of Economic Behavior & Organization* 185, 27–49.
- Volkow, N. D. et al. (2014). Americas addiction to opioids: Heroin and prescription drug abuse. *Senate Caucus on International Narcotics Control* 14, 1–16.



## A Online Appendix

### A.1 Additional Figures

**Figure A1:** Pre-reformulation opioid exposure, 2006-2009



*Notes:* Figure shows the population-weighted average number of opioids prescriptions per capita in the pre-reformulation period from 2006 to 2009 for 341 counties as reported in [Evans et al. \(2022\)](#). Sample mean is 0.89, standard deviation is 0.35.

## **A.2 Additional Results**

### **A.2.1 Temperature and Intimate Partner Violence: Heterogeneity**

**Table A1:** Heterogeneity — County-level Socio-demographics

	Intimate Partner Violence per 100,000 people					
	(1)	(2)	(3)	(4)	(5)	(6)
Temperature × Rural	0.0003 (0.0002)					
Temperature × Urban	0.0010*** (0.0003)					
Temperature × Poverty Rate (Below Median)		0.0003* (0.0002)				
Temperature × Poverty Rate (Above Median)		0.0015** (0.0006)				
Temperature × Income (Above Median)			0.0004* (0.0002)			
Temperature × Income (Below Median)			0.0020** (0.0008)			
Temperature × White % (Below Median)				0.0005 (0.0003)		
Temperature × White % (Above Median)				0.0004**		
Temperature × Black % (Below Median)					(0.0002)	
Temperature × Black % (Above Median)					0.0004** (0.0001)	
Temperature × Hispanic % (Below Median)					0.0005 (0.0003)	
Temperature × Hispanic % (Above Median)						0.0008*** (0.0003)
						0.0004* (0.0003)
Observations	44,160,870	44,168,174	44,168,174	44,160,870	44,160,870	44,160,870
Dependent Variable Mean	0.05742	0.05742	0.05742	0.05742	0.05742	0.05742
Precipitation Controls	✓	✓	✓	✓	✓	✓
Week-of-Year FE	✓	✓	✓	✓	✓	✓
Day-of-Week FE	✓	✓	✓	✓	✓	✓
Jurisdiction-Month-Year × Median Urban FE	✓					
Jurisdiction-Month-Year × Median Poverty Rate FE		✓				
Jurisdiction-Month-Year × Median Income FE			✓			
Jurisdiction-Month-Year × Median White (%) FE				✓		
Jurisdiction-Month-Year × Median Black (%) FE					✓	
Jurisdiction-Month-Year × Median Hispanic (%) FE						✓
Jurisdiction-Location Population Weights	✓	✓	✓	✓	✓	✓

**Notes:** The dependent variable is the number of intimate partner violence per 100,000 people. Temperature is the average daily temperature measured in degrees Celsius (°C). Precipitation is the total daily precipitation measured in metres (m). Standard errors are clustered at the county level. Significance levels are indicated as follows: \*p<0.10, \*\* p<0.05, \*\*\* p<0.01.

**Table A2:** Heterogeneity — Climatic Conditions

	Intimate Partner Violence per 100,000 people	
	(1)	(2)
Temperature × Cold	0.0043*** (0.0003)	0.0003 (0.0002)
Temperature × Mild	0.0058*** (0.0004)	0.0006** (0.0002)
Temperature × Warm	0.0102*** (0.0007)	0.0016** (0.0007)
Observations	44,170,732	44,170,732
Dependent Variable Mean	0.05742	0.05742
Precipitation Controls	✓	✓
Week-of-Year FE	✓	✓
Day-of-Week FE	✓	✓
Jurisdiction-Month-Year × Climate Terciles FE	✓	✓
Jurisdiction-Location Population Weights		✓

**Notes:** The dependent variable is the number of intimate partner violence per 100,000 people. Temperature is the daily average temperature measured in degrees Celsius (°C). Precipitation is the total daily precipitation measured in metres (m). Counties are grouped based on terciles of their 30-year mean of daily average temperature. Standard errors are clustered at the county level. Significance levels are indicated as follows: \*p<0.10, \*\* p<0.05, \*\*\* p<0.01.

## A.2.2 Temperature and Intimate Partner Violence: Robustness

**Table A3:** The Relationship between Temperature and Intimate Partner Violence — Accounting for Covid Period

	Intimate Partner Violence per 100,000 people	
	1991-2019 (1)	1991-2021 (2)
Temperature (°C)	0.0004** (0.0002)	0.0005** (0.0002)
Precipitation (m)	-0.0138 (0.0090)	-0.0126 (0.0101)
Observations	37,687,828	44,170,732
Dependent variable mean	0.04676	0.05742
Week-of-Year FE	✓	✓
Day-of-Week FE	✓	✓
Jurisdiction-Month-Year FE	✓	✓
Covid Dummy		✓
Jurisdiction-Location Population Weights	✓	✓

**Notes:** The dependent variable is the number of intimate partner violence per 100,000 people. In Column (2) we include a dummy (0,1) from 2020 onwards to identify the years of Covid-19 outbreak. Standard errors are clustered at the county level. Significance levels are indicated as follows: \*p<0.10, \*\* p<0.05, \*\*\* p<0.01.

**Table A4:** The Relationship between Temperature and Intimate Partner Violence — Trimming the Sample

	Intimate Partner Violence per 100,000 people		
	IPV < 100 (1)	IPV < 10 (2)	IPV < 1 (3)
Temperature (°C)	0.0005** (0.0002)	0.0004** (0.0002)	1.81e-05* (9.31e-06)
Precipitation (m)	-0.0126 (0.0101)	-0.0097 (0.0089)	-0.0004 (0.0009)
Observations	44,156,376	43,458,777	40,599,988
Dependent variable mean	0.05258	0.04563	0.00452
Week-of-Year FE	✓	✓	✓
Day-of-Week FE	✓	✓	✓
Jurisdiction-Month-Year FE	✓	✓	✓
Jurisdiction-Location Population Weights	✓	✓	✓

**Notes:** The dependent variable is the number of intimate partner violence per 100,000 people. Standard errors are clustered at the county level. Significance levels are indicated as follows: \*p<0.10, \*\* p<0.05, \*\*\* p<0.01.

**Table A5:** Relationship between Temperature and Intimate Partner Violence — Count Dependent Variable

	# Intimate Partner Violence		
	OLS (1)	PPML (2)	PPML (3)
Temperature (°C)	0.0011*** (0.0004)	0.0185*** (0.0035)	0.0104*** (0.0016)
Precipitation (m)	-0.0279 (0.0560)	0.1790 (0.4844)	-0.1017 (0.3538)
Observations	44,170,732	40,613,425	23,638,551
Dependent Variable Mean	0.11771	0.11771	0.11771
Week-of-Year FE	✓	✓	✓
Day-of-Week FE	✓	✓	✓
County-Month-Year FE		✓	
Jurisdiction-Month-Year FE	✓		✓
Jurisdiction-Location Population Weights	✓	✓	✓

**Notes:** The dependent variable is the number of intimate partner violence cases. Standard errors are clustered at the county level. Significance levels are indicated as follows: \*p<0.10, \*\* p<0.05, \*\*\* p<0.01.

**Table A6:** Relationship between Temperature and Intimate Partner Violence — Alternative Fixed-effects

	Intimate Partner Violence per 100,000 people			
	(1)	(2)	(3)	(4)
Temperature (°C)	0.0005** (0.0002)	0.0003* (0.0002)	0.0008*** (0.0003)	0.0007*** (0.0002)
Precipitation (m)	-0.0126 (0.0101)	-0.0425 (0.0264)	-0.0334* (0.0174)	-0.0189 (0.0154)
Cumulative Effect of Temperature				0.0003*** (0.0001)
Cumulative Effect of All Temperature Leads				0.000259** (0.000106)
Cumulative Effect of 2nd to 7th Temperature Leads				-1.71e-05 (4.97e-05)
Observations	44,170,732	44,170,732	44,170,732	43,979,205
Dependent Variable Mean	0.05741	0.05741	0.05741	0.05741
Week-of-Year FE	✓	✓		
Day-of-Week FE	✓	✓		
Jurisdiction-Month-Year FE	✓		✓	✓
Jurisdiction-Day FE		✓	✓	✓
Date FE			✓	✓
Jurisdiction-Location Population Weights	✓	✓	✓	✓

**Notes:** The dependent variable is the number of intimate partner violence per 100,000 people. Column (1) reports the baseline results. In Columns (3) and (4) we use the same fixed effect specification from [Cohen et al. \(2018\)](#). In Column (4) we also control for 7-day lags and leads. In the same column we report: (i) the cumulative effect of temperature, summing the coefficients of contemporaneous temperature and the 7-day lags; (ii) the cumulative effects of the temperature leads. Standard errors are clustered at the county level. Significance levels are indicated as follows: \*p<0.10, \*\* p<0.05, \*\*\* p<0.01.




**Table A7:** Relationship between Temperature and Intimate Partner Violence — Alternative Standard Errors

	Intimate Partner Violence per 100,000 people	
	(1)	(2)
Temperature (°C)	0.0005** (0.0002)	0.0005* (0.0002)
Precipitation (m)	-0.0126 (0.0101)	-0.0126 (0.0116)
Observations	44,170,732	44,170,732
Dependent Variable Mean	0.05741	0.05741
Week-of-Year FE	✓	✓
Day-of-Week FE	✓	✓
Jurisdiction-Month-Year FE	✓	✓
Jurisdiction-Location Population Weights	✓	✓

**Notes:** The dependent variable is the number of intimate partner violence per 100,000 people. In Column (1) standard errors are clustered at the county level. In Column (2) standard errors are clustered at the state level. Significance levels are indicated as follows: \*p<0.10, \*\* p<0.05, \*\*\* p<0.01

**Figure A2:** The Relationship between Temperature and Intimate Partner Violence  
— Temperature Bins



**Notes:** The dependent variable is the number of intimate partner violence per 100,000 people. Temperature is modelled using seven 5-degree Celsius intervals. The coefficients indicate the effect of an additional day in the j-th bin on IPV cases per 100,000 people, relative to the reference interval 15-19 °C. The regression also controls for total daily precipitation measured in metres (m), jurisdiction-month-year, week-of-year and day-of-week fixed effects. The shaded area represents 95% confidence intervals with standard errors clustered at the county-level.

**Table A8:** The Relationship between Temperature and Intimate Partner Violence — Temperature Bins

	Intimate Partner Violence per 100,000 people				
	(1)	(2)	(3)	(4)	(5)
< 5 (°C)	-0.0189** (0.0074)	-0.0958*** (0.0298)	-0.0177** (0.0083)	-0.0046 (0.0032)	-0.0086** (0.0040)
5-10 (°C)	-0.0027 (0.0021)	-0.0582*** (0.0168)	-0.0121** (0.0056)	-0.0051** (0.0025)	-0.0062** (0.0029)
10-15 (°C)	-0.0027 (0.0021)	-0.0312*** (0.0087)	-0.0066** (0.0030)	-0.0037** (0.0016)	-0.0035** (0.0017)
20-25 (°C)	0.0095** (0.0042)	0.0379*** (0.0143)	0.0069* (0.0037)	0.0044*** (0.0017)	0.0036* (0.0019)
25-30 (°C)	0.0556*** (0.0190)	0.0988*** (0.0311)	0.0162** (0.0073)	0.0099*** (0.0029)	0.0074** (0.0032)
≥ 30 (°C)	0.0933* (0.0522)	0.1379*** (0.0409)	0.0211*** (0.0080)	0.0133** (0.0055)	0.0087** (0.0035)
Precipitation (m)	0.1018 (0.1619)	0.0956 (0.1018)	0.0443 (0.0340)	-0.0151 (0.0180)	-0.0002 (0.0103)
Observations	44,170,732	44,170,732	44,170,732	44,170,732	44,170,732
Dependent Variable Mean	0.05741	0.05741	0.05741	0.05741	0.05741
Month-Year FE		✓			
Week-of-Year FE		✓	✓	✓	✓
Day-of-Week FE		✓	✓	✓	✓
County-Month-Year FE			✓		
Jurisdiction FE				✓	
Jurisdiction-Month-Year FE					✓
Jurisdiction-Location Population Weights	✓	✓	✓	✓	✓

**Notes:** The dependent variable is the number of intimate partner violence per 100,000 people. Temperature is modelled using temperature bins, where each bin is a dummy variable equal to 1 if the average temperature on a day falls within the specific bin. The omitted temperature category is 15-20 °C. Standard errors are clustered at the county level. Significance levels are indicated as follows: \*p<0.10, \*\* p<0.05, \*\*\* p<0.01.

**Table A9:** The Relationship between Temperature and Intimate Partner Violence — Share of Hours

	Intimate Partner Violence per 100,000 people				
	(1)	(2)	(3)	(4)	(5)
< 5 (°C)	-0.0053 (0.0065)	-0.0948*** (0.0271)	-0.0194** (0.0086)	-0.0038 (0.0035)	-0.0091** (0.0040)
5-10 (°C)	0.0191** (0.0076)	-0.0413*** (0.0123)	-0.0117** (0.0048)	-0.0050* (0.0027)	-0.0062** (0.0027)
10-15 (°C)	0.0139** (0.0068)	-0.0138* (0.0080)	-0.0061*** (0.0021)	-0.0031* (0.0016)	-0.0035*** (0.0013)
20-25 (°C)	0.0294** (0.0127)	0.0713** (0.0299)	0.0117 (0.0073)	0.0072** (0.0031)	0.0060 (0.0038)
25-30 (°C)	0.0668** (0.0263)	0.1374*** (0.0472)	0.0226* (0.0116)	0.0131*** (0.0041)	0.0106* (0.0054)
≥ 30 (°C)	0.1128** (0.0512)	0.1750*** (0.0576)	0.0315** (0.0137)	0.0186*** (0.0062)	0.0157** (0.0067)
Precipitation (m)	0.1345 (0.1532)	0.1596 (0.1081)	0.0613 (0.0427)	-0.0055 (0.0172)	0.0091 (0.0134)
Observations	44,170,732	44,170,732	44,170,732	44,170,732	44,170,732
Dependent Variable Mean	0.05741	0.05741	0.05741	0.05741	0.05741
Month-Year FE		✓			
Week-of-Year FE		✓	✓	✓	✓
Day-of-Week FE		✓	✓	✓	✓
County-Month-Year FE			✓		
Jurisdiction FE				✓	
Jurisdiction-Month-Year FE					✓
Jurisdiction-Location Population Weights	✓	✓	✓	✓	✓

**Notes:** The dependent variable is the number of intimate partner violence per 100,000 people. Temperature is modelled using temperature bins, where each bin is a share of hours during a day where hourly temperature falls within the specific bin. The omitted temperature category is 15-20 °C. Standard errors are clustered at the county level. Significance levels are indicated as follows: \*p<0.10, \*\* p<0.05, \*\*\* p<0.01.

**Table A10:** Relationship between Temperature and Intimate Partner Violence  
— Polynomials (up to the 4th degree)

	Intimate Partner Violence per 100,000 people			
	(1)	(2)	(3)	(4)
Temperature (°C)	0.0005** (0.0002)	0.0003* (0.0002)	0.0003* (0.0002)	0.0003* (0.0001)
Temperature <sup>2</sup>		1.51e-05** (6.03e-06)	1.45e-05** (6.59e-06)	1.71e-05** (8.61e-06)
Temperature <sup>3</sup>			3.11e-08 (9.65e-08)	2.43e-07 (1.52e-07)
Temperature <sup>4</sup>				-8.85e-09 (7.58e-09)
Precipitation (m)	-0.0126 (0.0101)	0.0058 (0.0129)	0.0060 (0.0126)	0.0064 (0.0127)
Observations	44,170,732	44,170,732	44,170,732	44,170,732
Dependent Variable Mean	0.05741	0.05741	0.05741	0.05741
Week-of-Year FE	✓	✓	✓	✓
Day-of-Week FE	✓	✓	✓	✓
Jurisdiction-Month-Year FE	✓	✓	✓	✓
Jurisdiction-Location Population Weights	✓	✓	✓	✓

**Notes:** The dependent variable is the number of intimate partner violence per 100,000 people. Standard errors are clustered at the county level. Significance levels are indicated as follows: \*p<0.10, \*\* p<0.05, \*\*\* p<0.01


**Table A11:** The Relationship between Temperature and Intimate Partner Violence — Lags and Leads

	Intimate Partner Violence per 100,000 people				
	7-day Lags (1)	14-day Lags (2)	21-day Lags (3)	7-day Leads (4)	7-day Lags and Leads (5)
Contemporaneous Temperature (°C)	0.0005** (0.0003)	0.0005** (0.0003)	0.0005* (0.0003)	0.0004** (0.0002)	0.0004* (0.0002)
Precipitation (m)	-0.0090 (0.0101)	-0.0080 (0.0099)	-0.0069 (0.0103)	-0.0034 (0.0120)	-0.0036 (0.0108)
Cumulative Effect of Temperature	0.0003** (0.0001)	0.0002** (9.41e-05)	0.0002* (0.0001)		0.0002* (8.82e-05)
Cumulative Effect of All Temperature Leads				0.0002* (0.0001)	0.0002* (9.24e-05)
Cumulative Effect of 2nd to 7th Temperature Leads				4.25e-05 (4.84e-05)	-1.23e-05 (2.31e-05)
Observations	44,074,951	43,979,170	43,883,389	44,074,986	43,979,205
Dependent variable mean	0.05741	0.05741	0.05741	0.05741	0.05741
Week-of-Year FE	✓	✓	✓	✓	✓
Day-of-Week FE	✓	✓	✓	✓	✓
Jurisdiction-Month-Year FE	✓	✓	✓	✓	✓
Jurisdiction-Location Population Weights	✓	✓	✓	✓	✓

**Notes:** The dependent variable is the number of intimate partner violence per 100,000 people. The cumulative effect of temperature is the sum of the coefficients of contemporaneous temperature and its lags. Standard errors are clustered at the county level. Significance levels are indicated as follows: \*p<0.10, \*\* p<0.05, \*\*\* p<0.01.


### A.2.3 Opioid reformulation

**Figure A3:** Triple difference (DiDiD) of Opioid reformulation policy on the temperature-IPV relationship with alternative fixed effects



*Notes:* The figure plots the coefficients from the triple interaction term between average daily county-level temperature, pre-intervention exposure as the population-weighted mean per capita opioid prescription, and in the left graph an indicator variable that takes the value of one for the post-reformulation years, 2010-2021, and in the right graph an indicator variable (short-run) equal to one for the years immediately following reformulation, 2011 to 2013, another one (medium-run) equal to one for the years 2014 to 2016, and another indicator variable (long-run) for the years 2017-2021, several years post-reformulation. The regression also controls for jurisdiction-day, jurisdiction-month-year, and date fixed effects. Bins represent the 95% confidence intervals with standard errors clustered at the county-level.


**Figure A4:** Triple difference (DiDiD) of Opioid reformulation policy in 2010 on the temperature-IPV relationship (excluding Covid period)



*Notes:* The figure plots the coefficients from the triple interaction term between average daily county-level temperature, pre-intervention exposure as the population-weighted mean per capita opioid prescription, and in the left graph an indicator variable that takes the value of one for the post-reformulation years, 2010-2019, and in the right graph an indicator variable (short-run) equal to one for the years immediately following reformulation, 2011 to 2013, another one (medium-run) equal to one for the years 2014 to 2016, and another indicator variable (long-run) for the years 2017-2019, several years post-reformulation. We exclude observations in the Covid period defined as years 2020 and 2021. The regression also controls for year-specific temperature and precipitation coefficients, jurisdiction-month-year, week-of-year and day-of-week fixed effects. Bins represent the 95% confidence intervals with standard errors clustered at the county-level.




**Figure A5:** Triple difference (DiDiD) of Opioid reformulation policy in 2010 on the temperature-IPV relationship (accounting for year-specific precipitation effects)




*Notes:* The figure plots the coefficients from the triple interaction term between average daily county-level temperature, pre-intervention exposure as the population-weighted mean per capita opioid prescription, and in the left graph an indicator variable that takes the value of one for the post-reformulation years, 2010-2019, and in the right graph an indicator variable (short-run) equal to one for the years immediately following reformulation, 2011 to 2013, another one (medium-run) equal to one for the years 2014 to 2016, and another indicator variable (long-run) for the years 2017-2019, several years post-reformulation. The regression also controls for year-specific temperature and year-specific precipitation, jurisdiction-month-year, week-of-year and day-of-week fixed effects. Bins represent the 95% confidence intervals with standard errors clustered at the county-level.

**Figure A6:** Triple difference (DiDiD) of Opioid reformulation policy in 2010 on the temperature-IPV relationship (state-level clustered standard errors)




*Notes:* The figure plots the coefficients from the triple interaction term between average daily county-level temperature, pre-intervention exposure as the population-weighted mean per capita opioid prescription, and in the left graph an indicator variable that takes the value of one for the post-reformulation years, 2010-2019, and in the right graph an indicator variable (short-run) equal to one for the years immediately following reformulation, 2011 to 2013, another one (medium-run) equal to one for the years 2014 to 2016, and another indicator variable (long-run) for the years 2017-2019, several years post-reformulation. The regression also controls for year-specific temperature and precipitation, jurisdiction-day, jurisdiction-month-year, week-of-year and day-of-week fixed effects. Bins represent the 95% confidence intervals with standard errors clustered at the state-level.

**Figure A7:** Leave-one-season-out (LOSO) triple difference of Opioid reformulation policy in 2010 on the temperature-IPV relationship



*Notes:* The figure plots the coefficients from the triple interaction term between average daily county-level temperature, pre-intervention exposure as the population-weighted mean per capita opioid prescription in a regression where we restrict the estimation sample leaving one season out at a time. Winter is defined as December, January, and February. Spring is defined as March, April, May. Summer is defined as June, July, August. Fall is defined as September, October, November. The regression also controls for year-specific temperature and precipitation, jurisdiction-month-year, week-of-year and day-of-week fixed effects. Bins represent the 95% confidence intervals with standard errors clustered at the county-level.

**Figure A8:** Leave-one-climate-out (LOCO) triple difference of Opioid reformulation policy in 2010 on the temperature-IPV relationship




*Notes:* The figure plots the coefficients from the triple interaction term between average daily county-level temperature, pre-intervention exposure as the population-weighted mean per capita opioid prescription in a regression where we restrict the estimation sample leaving counties that fall in a tercile of climate (defined from the average temperature in a county) each at a time. The regression also controls for year-specific temperature and precipitation, jurisdiction-month-year, week-of-year and day-of-week fixed effects. Bins represent the 95% confidence intervals with standard errors clustered at the county-level. The coefficient on the restricted sample excluding cold counties is statistically significant at 90% level.

**Table A12:** The Relationship between Temperature and Intimate Partner Violence

	Intimate Partner Violence per 100,000 people		
	(1)	(2)	(3)
<b>Panel A: DiDiD</b>			
Nighttime temperature $\times$ Opioid exposure	0.00980*** (0.00236)	0.00957*** (0.00233)	0.00936*** (0.00232)
Nighttime temperature $\times$ Opioid exposure $\times$ Post-policy	-0.00819*** (0.00226)	-0.00793*** (0.00222)	-0.00771*** (0.00221)
<b>Panel B: DiDiD by time window</b>			
Nighttime temperature $\times$ Opioid exposure	0.00751** (0.00347)	0.00791** (0.00346)	0.00792** (0.00345)
Nighttime temperature $\times$ Opioid exposure $\times$ Pre-policy	0.00228 (0.00287)	0.00166 (0.00281)	0.00144 (0.00276)
Nighttime temperature $\times$ Opioid exposure $\times$ Short-run post-policy	-0.00905** (0.00361)	-0.00938*** (0.00358)	-0.00931*** (0.00356)
Nighttime temperature $\times$ Opioid exposure $\times$ Medium-run post-policy	-0.00552 (0.00362)	-0.00584 (0.00359)	-0.00577 (0.00356)
Nighttime temperature $\times$ Opioid exposure $\times$ Long-run post-policy	-0.00238 (0.00324)	-0.00283 (0.00325)	-0.00296 (0.00327)
Nighttime	6pm-6am	8pm-6am	8pm-8am
Observations	8,494,904	8,494,904	8,494,904
Dependent Variable Mean	0.095	0.095	0.095
Week-of-Year FE	✓	✓	✓
Day-of-Week FE	✓	✓	✓
Jurisdiction-Month-Year FE	✓	✓	✓
Jurisdiction-Location Population Weights	✓	✓	✓

**Notes:** The dependent variable is the number of intimate partner violence per 100,000 people. Nighttime temperature is the average hourly temperature in the period of the day report in the row "Nighttime" for each column, measured in degrees Celsius (°C). Precipitation is the total daily precipitation measured in metres (m). Standard errors are clustered at the county level. Significance levels are indicated as follows: \*p<0.10, \*\* p<0.05, \*\*\* p<0.01.

**Figure A9:** Effect of the Opioid reformulation on the precipitation-IPV relationship



*Notes:* The figure plots the coefficients associated with the triple interaction term between daily-precipitation, pre-intervention opioid exposure and year dummies in a regression where the outcome variable is the number of IPV cases per 100,000 people. The regression also controls for year-specific temperature and precipitation coefficients, jurisdiction-month-year, week-of-year and day-of-week fixed effects. Shaded area represent the 95% confidence intervals with standard errors clustered at the county-level.

**Table A13:** The Relationship between Nighttime Temperature and Intimate Partner Violence

	Intimate Partner Violence per 100,000 people		
	(1)	(2)	(3)
Nighttime Temperature (°C)	0.000521** (0.000250)	0.000505** (0.000242)	0.000495** (0.000238)
Precipitation (m)	0.000142 (0.0128)	-0.00580 (0.0117)	-0.00992 (0.0109)
Nighttime	6pm-6am	8pm-6am	8pm-8am
R <sup>2</sup>	0.252	0.252	0.252
Observations	44,170,732	44,170,732	44,170,732
Dependent variable mean	0.0574	0.0574	0.0574
Nighttime temperature mean (°C)	12.31	11.82	11.28
Jurisdiction-Month-Year FE	✓	✓	✓
Week-of-Year FE	✓	✓	✓
Day-of-Week FE	✓	✓	✓
Jurisdiction-Location Population Weights	✓	✓	✓

**Notes:** The dependent variable is the number of intimate partner violence per 100,000 people. Standard errors are clustered at the county level. Significance levels are indicated as follows: \*p<0.10, \*\* p<0.05, \*\*\* p<0.01