

NOAA Buoy Data

June 2016

Introduction

The National Oceanic and Atmospheric Administration (NOAA) is the American federal agency in charge of collecting information and making decisions related to the oceans and the atmosphere. Throughout North America, they supply weather stations which are located both along the coast as well as in the middle of the ocean (on buoys). Among other variables, the weather stations collect information on wind, humidity, temperature, visibility, and atmospheric pressure. The data is all publicly available on NOAA's website.

Data information & loading data

All the buoys are listed at http://www.ndbc.noaa.gov/to_station.shtml. The Santa Monica buoy information is at http://www.ndbc.noaa.gov/station_page.php?station=46025. The historical data is given at http://www.ndbc.noaa.gov/station_history.php?station=46025.

Always a good idea to look at the data! One thing to notice is that there are some variables coded as 99/999/9999. From user experience, we surmise that those values should be NA. Additionally, if we want to consider only the 2014 data, we should remove any previous data.

```
summary(buoy_data)
```

```
##           #YY           MM           DD           hh
##  Min.      :2013   Length:8611   Length:8611   Length:8611
##  1st Qu.:2014   Class :character Class :character Class :character
##  Median :2014   Mode  :character Mode  :character Mode  :character
##  Mean      :2014
##  3rd Qu.:2014
##  Max.      :2014
##           mm           WDIR           WSPD           GST
##  Min.      :50   Min.      : 1.0   Min.      :0.000   Min.      :0.000
##  1st Qu.:50   1st Qu.:165.0   1st Qu.:1.900   1st Qu.:2.600
##  Median :50   Median :266.0   Median :3.100   Median :3.900
##  Mean      :50   Mean      :231.3   Mean      :3.408   Mean      :4.106
##  3rd Qu.:50   3rd Qu.:298.0   3rd Qu.:4.600   3rd Qu.:5.400
##  Max.      :50   Max.      :360.0   Max.      :9.900   Max.      :9.900
##           WVHT           DPD           APD           MWD
##  Min.      : 0.360   Min.      : 2.74   Min.      : 3.620   Min.      : 1.0
##  1st Qu.: 0.810   1st Qu.:10.00   1st Qu.: 5.420   1st Qu.:204.0
##  Median : 0.990   Median :12.90   Median : 6.090   Median :253.0
##  Mean      : 1.273   Mean      :12.25   Mean      : 6.612   Mean      :236.5
##  3rd Qu.: 1.280   3rd Qu.:14.81   3rd Qu.: 7.220   3rd Qu.:268.0
##  Max.      :99.000   Max.      :99.00   Max.      :99.000   Max.      :999.0
##           PRES           ATMP           WTMP           DEWP
##  Min.      :1003   Min.      :10.10   Min.      :12.90   Min.      : -8.70
##  1st Qu.:1012   1st Qu.:15.20   1st Qu.:15.80   1st Qu.:11.50
##  Median :1014   Median :17.30   Median :18.60   Median :13.50
##  Mean      :1016   Mean      :17.25   Mean      :18.52   Mean      :13.22
##  3rd Qu.:1017   3rd Qu.:19.30   3rd Qu.:21.00   3rd Qu.:15.70
```

```
## Max. :9999 Max. :24.50 Max. :24.50 Max. :99.00
## VIS TIDE
## Min. :99 Min. :99
## 1st Qu.:99 1st Qu.:99
## Median :99 Median :99
## Mean :99 Mean :99
## 3rd Qu.:99 3rd Qu.:99
## Max. :99 Max. :99
```

```
buoy_data <- buoy_data %>%
  mutate(WVHT = ifelse(WVHT==99, NA, WVHT)) %>%
  mutate(DPD = ifelse(DPD==99, NA, DPD)) %>%
  mutate(APD = ifelse(APD==99, NA, APD)) %>%
  mutate(MWD = ifelse(MWD==999, NA, MWD)) %>%
  mutate(PRES = ifelse(PRES==9999, NA, PRES)) %>%
  mutate(DEWP = ifelse(DEWP==99, NA, DEWP)) %>%
  select(-VIS, -TIDE) %>% filter(`#YY`==2014)

dim(buoy_data)
```

```
## [1] 8610 16
```

```
summary(buoy_data)
```

```
##      #YY      MM      DD      hh
## Min. :2014 Length:8610 Length:8610 Length:8610
## 1st Qu.:2014 Class :character Class :character Class :character
## Median :2014 Mode  :character Mode  :character Mode  :character
## Mean   :2014
## 3rd Qu.:2014
## Max.   :2014
##
##      mm      WDIR      WSPD      GST
## Min. :50 Min. : 1.0 Min. :0.000 Min. :0.000
## 1st Qu.:50 1st Qu.:165.0 1st Qu.:1.900 1st Qu.:2.600
## Median :50 Median :266.0 Median :3.100 Median :3.900
## Mean   :50 Mean   :231.3 Mean   :3.408 Mean   :4.106
## 3rd Qu.:50 3rd Qu.:298.0 3rd Qu.:4.600 3rd Qu.:5.400
## Max.   :50 Max.   :360.0 Max.   :9.900 Max.   :9.900
##
##      WVHT      DPD      APD      MWD
## Min. :0.360 Min. : 2.74 Min. : 3.62 Min. : 1.0
## 1st Qu.:0.810 1st Qu.:10.00 1st Qu.: 5.42 1st Qu.:204.0
## Median :0.990 Median :12.90 Median : 6.09 Median :253.0
## Mean   :1.091 Mean   :12.09 Mean   : 6.44 Mean   :235.1
## 3rd Qu.:1.280 3rd Qu.:14.81 3rd Qu.: 7.21 3rd Qu.:268.0
## Max.   :4.800 Max.   :23.53 Max.   :13.52 Max.   :359.0
## NA's :16 NA's :16 NA's :16 NA's :16
##      PRES      ATMP      WTMP      DEWP
## Min. :1003 Min. :10.10 Min. :12.90 Min. : -8.7
## 1st Qu.:1012 1st Qu.:15.20 1st Qu.:15.80 1st Qu.:11.5
## Median :1014 Median :17.30 Median :18.60 Median :13.5
## Mean   :1015 Mean   :17.25 Mean   :18.52 Mean   :13.2
## 3rd Qu.:1017 3rd Qu.:19.30 3rd Qu.:21.00 3rd Qu.:15.7
## Max.   :1029 Max.   :24.50 Max.   :24.50 Max.   :21.0
```


```
## NA's :1
```

```
NA's :2
```

Using dynamic data within a typical classroom

One might be interested in the difference between the wind temperature and the air temperature. Generally, the air temperature is cooler than the wind temperature, but confidence intervals and prediction intervals allow us to quantify the difference. Note that the data lend themselves nicely to ideas of paired observations acting as a univariate sample. As expected, a 95% confidence interval for the true difference in temperatures gives us a value of between 1.25 and 1.31 degrees. However, 95% of the individual observations have a difference in wind and air temperature between -1.5 degrees (air is warmer) and 4.06 degrees (wind is warmer).

```
buoy_data$TempDiff <- buoy_data$WTMP - buoy_data$ATMP  
densityplot(~TempDiff, data=buoy_data)
```



```
tempdiff.mod <- lm(TempDiff ~ 1, data=buoy_data)  
tempdiff.func <- makeFun(tempdiff.mod)  
tempdiff.func()
```

```
##      1  
## 1.276249
```

```
tempdiff.func( interval="prediction")
```

```
##      fit      lwr      upr  
## 1 1.276249 -1.50516 4.057657
```


```
tempdiff.func( interval="confidence")
```

```
##      fit      lwr      upr  
## 1 1.276249 1.246275 1.306222
```

Thinking outside the box

The data are nicely set up to think about analyses in the time domain. Indeed, looking at the autocorrelation function shows clear 24-hour trends for the wind speed variable.


```
acf(buoy_data$WSPD, main="Series: Wind Speed", xlab="Lag (hours)")
```



Although a full analysis of the data would warrant multiple years of data (so as to understand yearly trends), we can estimate the spectral density of the time series using a smoothed periodogram.

```
spec.pgram(buoy_data$WSPD, spans=c(50), xlab="Frequency = 1/period",  
            main="Wind Speed, Smoothed Periodogram")  
abline(v=c(1/12, 1/24), lty=2)
```

Wind Speed, Smoothed Periodogram



In the smoothed periodogram, the x-axis is the frequency (one over the period) and y-axis represents the correlation (normalized) between the cosine wave at that frequency and the time series. We can see that wind speed has strong correlation at period 12 hours and period 24 hours.

Additional ideas for analysis:

A more sophisticated analysis or longer project could include collecting data from multiple buoys, extended years, and/or additional information on storms <https://www.ncdc.noaa.gov/stormevents/>.