

L7.2.X Worked Examples

Zonghua Gu 2021

Recall: TD, Sarsa, Q Learning

- TD solves [BEV] by sampling:
 - $V(S_t) \leftarrow V(S_t) + \alpha(R_{t+1} + \gamma V(S_{t+1}) - V(S_t))$
- Sarsa and Expected Sarsa solve [BEA] by sampling:
 - $Q(S_t, A_t) \leftarrow Q(S_t, A_t) + \alpha(R_{t+1} + \gamma Q(S_{t+1}, A_{t+1}) - Q(S_t, A_t))$
 - $Q(S_t, A_t) \leftarrow Q(S_t, A_t) + \alpha(R_{t+1} + \gamma \sum_{a'} \pi(a'|S_{t+1}) Q(S_{t+1}, a') - Q(S_t, A_t))$
- Q Learning solves [BOA] by sampling:
 - $Q(S_t, A_t) \leftarrow Q(S_t, A_t) + \alpha(R_{t+1} + \gamma \max_{a'} Q(S_{t+1}, a') - Q(S_t, A_t))$

- [BEV] Bellman Expectation Equation for State Value Function:
 - $v_\pi(s) = \sum_a \pi(a|s) \sum_{r,s'} p(r, s'|s, a) [r + \gamma v_\pi(s')]$
- [BEA] Bellman Expectation Equation for Action Value Function
 - $q_\pi(s, a) = \sum_{r,s'} p(r, s'|s, a) [r + \gamma \sum_{a'} \pi(a'|s') q_\pi(s', a')]$
- [BOA] Bellman Optimality Equation for Optimal Action Value Function:
 - $q_*(s, a) = \sum_{r,s'} p(r, s'|s, a) [r + \gamma \max_{a'} q_*(s', a')]$


TD, Sarsa, QL w. $\alpha = 1$

- With learning rate $\alpha = 1$, each $V(S_t)$ or $Q(S_t, A_t)$ is completely overwritten by in each update
- update equations simplify to:
 - MC: $V(S_t) \leftarrow V(S_t) + \alpha(G_t - V(S_t)) = G_t$
 - TD: $V(S_t) \leftarrow V(S_t) + \alpha(R_{t+1} + \gamma V(S_{t+1}) - V(S_t)) = R_{t+1} + \gamma V(S_{t+1})$
 - Sarsa: $Q(S_t, A_t) \leftarrow Q(S_t, A_t) + \alpha(R_{t+1} + \gamma Q(S_{t+1}, A_{t+1}) - Q(S_t, A_t)) = R_{t+1} + \gamma Q(S_{t+1}, A_{t+1})$
 - QL: $Q(S_t, A_t) \leftarrow Q(S_t, A_t) + \alpha(R_{t+1} + \gamma \max_{a'} Q(S_{t+1}, a') - Q(S_t, A_t)) = R_{t+1} + \gamma \max_{a'} Q(S_{t+1}, a')$

Two-Branch Example

Two-Branch Example



- An episodic MDP w. deterministic env, 3 states $\{B, C, D\}$ and 2 actions $\{1, 2\}$ at each state. Discount factor $\gamma = 1$, learning rate $\alpha = 1$. The initial state of each episode is B .



Policy Iteration

1.1 Policy Evaluation of Random Policy


- Bellman Exp Equation: $v_{\pi}(s) = \sum_a \pi(a|s) q_{\pi}(s, a); q_{\pi}(s, a) = R_s^a + \gamma v_{\pi}(s')$
- $v_{\pi}(B) = .5[q_{\pi}(B, a1) + q_{\pi}(B, a2)] = .5[1 + v_{\pi}(C) + v_{\pi}(D)]$
 - $q_{\pi}(B, a1) = 1 + v_{\pi}(C), q_{\pi}(B, a2) = 0 + v_{\pi}(D)$
- $v_{\pi}(C) = .5[q_{\pi}(C, a1) + q_{\pi}(C, a2)] = 2$
 - $q_{\pi}(C, a1) = 1, q_{\pi}(C, a2) = 3$
- $v_{\pi}(D) = .5[q_{\pi}(D, a1) + q_{\pi}(D, a2)] = -48$
 - $q_{\pi}(D, a1) = -100, q_{\pi}(D, a2) = 4$
- Solution: $v_{\pi}(B) = -22.5, v_{\pi}(C) = 2, v_{\pi}(D) = -48$



	$V_{\pi}(B)$	$V_{\pi}(C)$	$V_{\pi}(D)$
Iter1	-22.5	3	-48
Iter2	4	3	4
Iter3	4	3	4



1.2 Policy Improvement

- Plug in values from PE to get new policy
- $\pi'(B) = \operatorname{argmax}_a (q_\pi(B, a1), q_\pi(B, a2)) = a1$
 - $q_\pi(B, a1) = 1 + v_\pi(C) = 3, q_\pi(B, a2) = 0 + v_\pi(D) = -22.5$
- $\pi'(C) = \operatorname{argmax}_a (q_\pi(C, a1), q_\pi(C, a2)) = a2$
 - $q_\pi(C, a1) = 1, q_\pi(C, a2) = 3$
- $\pi'(D) = \operatorname{argmax}_a (q_\pi(D, a1), q_\pi(D, a2)) = a2$
 - $q_\pi(C, a1) = -100, q_\pi(C, a2) = 4$



2.1 Policy Evaluation of Det Policy


- Bellman Exp Equation: $v_{\pi}(s) = \sum_a \pi(a|s)q_{\pi}(s, a); q_{\pi}(s, a) = R_s^a + \gamma v_{\pi}(s')$
- $v_{\pi}(B) = q_{\pi}(B, a1) = 1 + v_{\pi}(C)$
 - $q_{\pi}(B, a1) = 1 + v_{\pi}(C)$
- $v_{\pi}(C) = q_{\pi}(C, a2) = 3$
 - $q_{\pi}(C, a2) = 3$
- $v_{\pi}(D) = q_{\pi}(D, a2) = 4$
 - $q_{\pi}(D, a2) = 4$
- Solution: $v_{\pi}(B) = 4, v_{\pi}(C) = 3, v_{\pi}(D) = 4$



	$V_{\pi}(B)$	$V_{\pi}(C)$	$V_{\pi}(D)$
Iter1	-22.5	3	-48
Iter2	4	3	4
Iter3	4	3	4



2.2 Policy Improvement

- Plug in values from PE to get new policy
- $\pi'(B) = \operatorname{argmax}_a (q_\pi(B, a1), q_\pi(B, a2)) = a1 \text{ or } a2$
 - $q_\pi(B, a1) = 1 + v_\pi(C) = 4, q_\pi(B, a2) = 0 + v_\pi(D) = 4$
- $\pi'(C) = \operatorname{argmax}_a (q_\pi(C, a1), q_\pi(C, a2)) = a2$
 - $q_\pi(C, a1) = 1, q_\pi(C, a2) = 3$
- $\pi'(D) = \operatorname{argmax}_a (q_\pi(D, a1), q_\pi(D, a2)) = a2$
 - $q_\pi(D, a1) = -100, q_\pi(D, a2) = 4$



3.1 Policy Evaluation


- Bellman Exp Equation: $v_{\pi}(s) = \sum_a \pi(a|s) q_{\pi}(s, a); q_{\pi}(s, a) = R_s^a + \gamma v_{\pi}(s')$
- $v_{\pi}(B) = .5[q_{\pi}(B, a1) + q_{\pi}(B, a2)] = .5[1 + v_{\pi}(C) + v_{\pi}(D)]$
 - $q_{\pi}(B, a1) = 1 + v_{\pi}(C), q_{\pi}(B, a2) = 0 + v_{\pi}(D)$
- $v_{\pi}(C) = q_{\pi}(C, a2) = 3$
 - $q_{\pi}(C, a2) = 3$
- $v_{\pi}(D) = q_{\pi}(D, a2) = 4$
 - $q_{\pi}(D, a2) = 4$
- Solution: $v_{\pi}(B) = 4, v_{\pi}(C) = 3, v_{\pi}(D) = 4$



	$V_{\pi}(B)$	$V_{\pi}(C)$	$V_{\pi}(D)$
Iter1	-22.5	3	-48
Iter2	4	3	4
Iter3	4	3	4

3.2 Policy Improvement



- Plug in values from PE to get new policy
- $\pi'(B) = \operatorname{argmax}_a (q_\pi(B, a1), q_\pi(B, a2)) = a1 \text{ or } a2$
 - $q_\pi(B, a1) = 1 + v_\pi(C) = 4, q_\pi(B, a2) = 0 + v_\pi(D) = 4$
- $\pi'(C) = \operatorname{argmax}_a (q_\pi(C, a1), q_\pi(C, a2)) = a2$
 - $q_\pi(C, a1) = 1, q_\pi(C, a2) = 3$
- $\pi'(D) = \operatorname{argmax}_a (q_\pi(D, a1), q_\pi(D, a2)) = a2$
 - $q_\pi(D, a1) = -100, q_\pi(D, a2) = 4$
- Policy has converged



Value Iteration

Value Iteration

- Bellman Opt Equation: $v_*(s) = \max_a q_*(s, a); q_*(s, a) = R_s^a + \gamma v_*(s')$
- $v_*(B) = \max_a [q_*(B, a1), q_*(B, a2)] = \max[1 + v_*(C), v_*(D)]$
 - $q_*(B, a1) = 1 + v_*(C), q_*(B, a2) = 0 + v_*(D)$
- $v_*(C) = \max_a [q_*(C, a1), q_*(C, a2)] = q_*(C, a2) = 3$
 - $q_*(C, a1) = 1, q_*(C, a2) = 3$
- $v_*(D) = \max_a [q_*(D, a1), q_*(D, a2)] = 4$
 - $q_*(D, a1) = -100, q_*(D, a2) = 4$
- We use Value Iteration to solve it. Table shows the iteration process until convergence (not using in-place updates for clarity). Solution: $v_*(1) = -3, v_*(2) = -2, v_*(3) = -1$
- Optimal policy: $\pi_*(B) = \operatorname{argmax}_a q_*(B, a) = a1 \text{ or } a2; \pi_*(C) = \operatorname{argmax}_a q_*(C, a) = a2; \pi_*(D) = \operatorname{argmax}_a q_*(D, a) = a2$




	$V_\pi(B)$	$V_\pi(C)$	$V_\pi(D)$
Init	0	0	0
Iter1	0	3	4
Iter2	4	3	4
Iter3	4	3	4

MC

MC, Episodes $3 \times (B, a2, 0, D, a1, -100, T)$


- MC update equation: $V(S_t) \leftarrow G_t$
- EP1:
 - $G(D) = -100 + V(T) = -100, V(D) = G(D) = -100$
 - $G(B) = 0 + G(D) = -100, V(D) = G(D) = -100$
- EP2:
 - $G(D) = -100 + V(T) = -100, V(D) = G(D) = -100$
 - $G(B) = 0 + G(D) = -100, V(D) = G(D) = -100$
- EP3:
 - $G(D) = -100 + V(T) = -100, V(D) = G(D) = -100$
 - $G(B) = 0 + G(D) = -100, V(D) = G(D) = -100$



	$V(B)$	$V(D)$
Init	0	0
EP1	-100	-100
EP2	-100	-100
EP3	-100	-100

MC, Episodes $n \times (B, a2, 0, D, a2, 4, T)$

- MC update equation: $V(S_t) \leftarrow G_t$
- EP1:
 - $G(D) = -100 + V(T) = \textcolor{red}{4}, V(D) = G(D) = \textcolor{red}{4}$
 - $G(B) = 0 + G(D) = 4, V(D) = G(D) = 4$
- EP2:
 - $G(D) = -100 + V(T) = 4, V(D) = G(D) = 4$
 - $G(B) = 0 + G(D) = 4, V(D) = G(D) = 4$
- EP3:
 - $G(D) = -100 + V(T) = 4, V(D) = G(D) = 4$
 - $G(B) = 0 + G(D) = 4, V(D) = G(D) = 4$




	$V(B)$	$V(D)$
Init	0	0
EP1	4	4
EP2	4	4
EP3	4	4

TD

TD, Episodes $n \times (B, a2, 0, D, a1, -100, T)$


- TD update equation: $V(S_t) \leftarrow R_{t+1} + \gamma V(S_{t+1})$
- EP1:
 - $V(B) \leftarrow R_{t+1} + \gamma V(D) = 0 + 0 = 0$
 - $V(D) \leftarrow R_{t+1} + \gamma V(T) = -100 - 0 = -100$
- EP2:
 - $V(B) \leftarrow R_{t+1} + \gamma V(D) = 0 - 100 = -100$
 - $V(D) \leftarrow R_{t+1} + \gamma V(T) = -100 - 0 = -100$
- EP3:
 - $V(B) \leftarrow R_{t+1} + \gamma V(D) = 0 - 100 = -100$
 - $V(D) \leftarrow R_{t+1} + \gamma V(T) = -100 - 0 = -100$



	$V(B)$	$V(D)$
Init	0	0
EP1	0	-100
EP2	-100	-100
EP3	-100	-100

TD, Episodes $3 \times (B, a2, 0, D, a2, 4, T)$

- TD update equation: $V(S_t) \leftarrow R_{t+1} + \gamma V(S_{t+1})$
- EP1:
 - $V(B) \leftarrow R_{t+1} + \gamma V(D) = 0 + 0 = 0$
 - $V(D) \leftarrow R_{t+1} + \gamma V(T) = 4 - 0 = 4$
- EP2:
 - $V(B) \leftarrow R_{t+1} + \gamma V(D) = 0 + 4 = 4$
 - $V(D) \leftarrow R_{t+1} + \gamma V(T) = 4 - 0 = 4$
- EP3:
 - $V(B) \leftarrow R_{t+1} + \gamma V(D) = 0 + 4 = 4$
 - $V(D) \leftarrow R_{t+1} + \gamma V(T) = 4 - 0 = 4$




	$V(B)$	$V(D)$
Init	0	0
EP1	0	4
EP2	4	4
EP3	4	4

Sarsa

Sarsa, Episodes $3 \times (B, a2, 0, D, a1, -100, T)$


- Sarsa update equation: $Q(S_t, A_t) \leftarrow R_{t+1} + \gamma Q(S_{t+1}, A_{t+1})$
- EP1:
 - $Q(B, a2) \leftarrow R_{t+1} + \gamma Q(D, a1) = 0 - 0 = 0$
 - $Q(D, a1) \leftarrow R_{t+1} + \gamma Q(T, -) = -100 + 0 = -100$
- EP2:
 - $Q(B, a2) \leftarrow R_{t+1} + \gamma Q(D, a1) = 0 - 100 = -100$
 - $Q(D, a1) \leftarrow R_{t+1} + \gamma Q(T, -) = -100 + 0 = -100$
- EP3:
 - $Q(B, a2) \leftarrow R_{t+1} + \gamma Q(D, a1) = 0 - 100 = -100$
 - $Q(D, a1) \leftarrow R_{t+1} + \gamma Q(T, -) = -100 + 0 = -100$



	$Q(B, a1)$	$Q(B, a2)$	$Q(D, a1)$	$Q(D, a2)$
Init	0	0	0	0
EP1	0	0	-100	0
EP2	0	-100	-100	0
EP3	0	-100	-100	0

Sarsa, Episodes $3 \times (B, a2, 0, D, a2, 4, T)$


- Sarsa update equation: $Q(S_t, A_t) \leftarrow R_{t+1} + \gamma Q(S_{t+1}, A_{t+1})$
- EP1:
 - $Q(B, a2) \leftarrow R_{t+1} + \gamma Q(D, a2) = 0 - 0 = 0$
 - $Q(D, a2) \leftarrow R_{t+1} + \gamma Q(T, -) = 4 + 0 = 4$
- EP2:
 - $Q(B, a2) \leftarrow R_{t+1} + \gamma Q(D, a2) = 0 + 4 = 4$
 - $Q(D, a2) \leftarrow R_{t+1} + \gamma Q(T, -) = 4 + 0 = 4$
- EP3:
 - $Q(B, a2) \leftarrow R_{t+1} + \gamma Q(D, a2) = 0 + 4 = 4$
 - $Q(D, a2) \leftarrow R_{t+1} + \gamma Q(T, -) = 4 + 0 = 4$



	$Q(B, a1)$	$Q(B, a2)$	$Q(D, a1)$	$Q(D, a2)$
Init	0	0	0	0
EP1	0	0	0	4
EP2	0	4	0	4
EP3	0	4	0	4

QL, Episodes $3 \times (B, a2, 0, D, a1, -100, T)$

- QL update equation: $Q(S_t, A_t) \leftarrow R_{t+1} + \gamma \max_{a'} Q(S_{t+1}, a')$
- EP1:
 - $Q(B, a2) \leftarrow R_{t+1} + \gamma \max_a Q(D, a) = 0 + \max(0, 0) = 0$
 - $Q(D, a1) \leftarrow R_{t+1} + \gamma Q(T, -) = -100 + 0 = -100$
- EP2:
 - $Q(B, a2) \leftarrow R_{t+1} + \gamma \max_a Q(D, a) = 0 + \max(-100, 0) = 0$
 - $Q(D, a1) \leftarrow R_{t+1} + \gamma Q(T, -) = -100 + 0 = -100$
- EP3:
 - $Q(B, a2) \leftarrow R_{t+1} + \gamma \max_a Q(D, a) = 0 + \max(-100, 0) = 0$
 - $Q(D, a1) \leftarrow R_{t+1} + \gamma Q(T, -) = -100 + 0 = -100$




	$Q(B, a1)$	$Q(B, a2)$	$Q(D, a1)$	$Q(D, a2)$
Init	0	0	0	0
EP1	0	0	-100	0
EP2	0	0	-100	0
EP3	0	0	-100	0

Q Learning

QL, Episodes $3 \times (B, 2, 0, D, 2, 4, T)$


- QL update equation: $Q(S_t, A_t) \leftarrow R_{t+1} + \gamma \max_{a'} Q(S_{t+1}, a')$
- EP1:
- $Q(B, a2) \leftarrow R_{t+1} + \gamma \max_a Q(D, a) = 0 + \max(0, 0) = 0$
- $Q(D, a2) \leftarrow R_{t+1} + \gamma Q(T, -) = 4 + 0 = 4$
- EP2:
- $Q(B, a2) \leftarrow R_{t+1} + \gamma \max_a Q(D, a) = 0 + 4 = 4$
- $Q(D, a2) \leftarrow R_{t+1} + \gamma Q(T, -) = 4 + 0 = 4$
- EP3:
- $Q(B, a2) \leftarrow R_{t+1} + \gamma \max_a Q(D, a) = 0 + 4 = 4$
- $Q(D, a2) \leftarrow R_{t+1} + \gamma Q(T, -) = 4 + 0 = 4$



	$Q(B, a1)$	$Q(B, a2)$	$Q(D, a1)$	$Q(D, a2)$
Init	0	0	0	0
EP1	0	0	0	4
EP2	0	4	0	4
EP3	0	4	0	4


Comparisons

- MC and TD:
 - Transition $(D, a1, -100, T)$ drives $V(D) \rightarrow -100$; $V(D)$ drives $V(B) \rightarrow -100$.
 - Transition $(D, a2, 4, T)$ drives $V(D) \rightarrow 4$; $V(D)$ drives $V(B) \rightarrow 4$.
 - Final values of $V(B), V(D)$ depend on relative execution frequencies of the 2 transitions (e.g., ϵ -greedy).
- Sarsa:
 - Transition $(D, a1, -100, T)$ drives $Q(D, a1) \rightarrow -100$; $Q(D, a1)$ drives $Q(B, a2) \rightarrow -100$.
 - Transition $(D, a2, 4, T)$ drives $Q(D, a2) \rightarrow 4$; $Q(D, a2)$ drives $Q(B, a2) \rightarrow 4$.
 - Final value of $Q(B, 2)$ depends on relative execution frequencies of the 2 transitions (e.g., ϵ -greedy).
- QL:
 - Transition $(D, a1, -100, T)$ drives $Q(D, a1) \rightarrow -100$; $Q(D, a1)$ does not affect $Q(B, a2)$ since
 - $\max_a Q(D, a) = \max(Q(D, a1), Q(D, a2)) = 0$. (assuming $Q(D, a2)$ is initialized to 0 and it never updated)
 - Transition $(D, a2, 4, T)$ drives $Q(D, a2) \rightarrow 4$, which in turn drives $Q(B, a2) \rightarrow 4$.
- Our example here only illustrates the policy evaluation for a given set of episodes, not control. If we consider control, and Sarsa or QL uses ϵ -greedy policy with small ϵ , then the agent will likely avoid action a1 in state D after taking it for the 1st time.



Sarsa w. ϵ -greedy


- Suppose EP1 is $(B, a2, 0, D, a1, -100, T)$. After EP1:
- $Q(B, a2) \leftarrow R_{t+1} + \gamma Q(D, a1) = 0 - 0 = 0$, $Q(D, a1) \leftarrow R_{t+1} + \gamma Q(T, -) = -100 + 0 = -100$
- Suppose EP2 starts with $(B, a2, 0, D)$, then in state D, the agent is likely to select action $\operatorname{argmax}_a \{Q(D, a1) = -100, Q(D, a2) = 0\} = a2$ based on ϵ -greedy, so the episode is $(B, a2, 0, D, a2, 4, T)$. After EP2:
- $Q(B, a2) \leftarrow R_{t+1} + \gamma Q(D, a1) = 0 - 100 = -100$, $Q(D, a2) \leftarrow R_{t+1} + \gamma Q(T, -) = 4 + 0 = 4$
- In EP3, in initial state B, the agent is likely to select action $\operatorname{argmax}_a \{Q(B, a1) = 0, Q(B, a2) = -100\} = a1$. Suppose the episode is $(B, a1, 1, C, a1, 1, T)$
- $Q(B, a1) \leftarrow R_{t+1} + \gamma Q(C, a1) = 1 + 0 = 1$, $Q(C, a1) \leftarrow R_{t+1} + \gamma Q(T, -) = 1 + 0 = 1$
- In EP4, in initial state B, the agent is likely to select action $\operatorname{argmax}_a \{Q(B, a1) = 1, Q(B, a2) = -100\} = a1$. in state D, the agent is likely to select action $\operatorname{argmax}_a \{Q(D, a1) = 1, Q(D, a2) = 0\} = a1$. Suppose the episode is again $(B, a1, 1, C, a1, 1, T)$
- $Q(B, a1) \leftarrow R_{t+1} + \gamma Q(C, a1) = 1 + 1 = 2$, $Q(C, a1) \leftarrow R_{t+1} + \gamma Q(T, -) = 1 + 0 = 1$
- if the agent always follows the greedy action, it will always follow the trajectory $(B, a1, 1, C, a1, 1, T)$ and never learn anything new, e.g., it will never experience the trajectories $(B, a1, 1, C, a2, 3, T)$, $(B, a2, 0, D, a2, 4, T)$. It got scared when $Q(B, a2)$ was updated to -100 after EP2 and never wanted to take action a2 in state B, but if it were more adventurous and tried it, it will likely experience EP5 $(B, a2, 0, D, a2, 4, T)$:
- $Q(B, a2) \leftarrow R_{t+1} + \gamma Q(D, a2) = 0 + 4 = 4$, $Q(D, a2) \leftarrow R_{t+1} + \gamma Q(T, -) = 4 + 0 = 4$
- Now you can see the importance of exploration by selecting the non-greedy action occasionally.



	$Q(B, a1)$	$Q(B, a2)$	$Q(C, a1)$	$Q(C, a2)$	$Q(D, a1)$	$Q(D, a2)$
Init	0	0	0	0	0	0
EP1	0	0	0	0	-100	0
EP2	0	-100	0	0	-100	4
EP3	1	-100	1	0	-100	4
EP4	2	-100	1	0	-100	4
EP5	2	4	1	0	-100	4

QL w. ϵ -greedy

- Suppose EP1 is $(B, a2, 0, D, a1, -100, T)$. After EP1:
- $Q(B, a2) \leftarrow R_{t+1} + \gamma \max_a Q(D, a) = 0 + \max(0, 0) = 0$, $Q(D, a1) \leftarrow R_{t+1} + \gamma Q(T, -) = -100 + 0 = -100$
- Suppose EP2 starts with $(B, a2, 0, D)$, then in state D, the agent is likely to select action $\text{argmax}_a \{Q(D, a1) = -100, Q(D, a2) = 0\} = a2$ based on ϵ -greedy, so the episode is $(B, a2, 0, D, a2, 4, T)$. After EP2:
- $Q(B, a2) \leftarrow R_{t+1} + \gamma \max_a Q(D, a) = 0 + \max(-100, 0) = 0$, $Q(D, a2) \leftarrow R_{t+1} + \gamma Q(T, -) = 4 + 0 = 4$
- In EP3, in initial state B, the agent is equally likely to select action $a1$ and $a2$ since $Q(B, a1) = Q(B, a2) = 0$. Suppose the episode is $(B, a1, 1, C, a1, 1, T)$
- $Q(B, a1) \leftarrow R_{t+1} + \gamma \max_a Q(C, a) = 1 + \max(0, 0) = 1$, $Q(C, a1) \leftarrow R_{t+1} + \gamma Q(T, -) = 1 + 0 = 1$
- In EP4, in initial state B, the agent is likely to select action $\text{argmax}_a \{Q(B, a1) = 1, Q(B, a2) = 0\} = a1$. in state D, the agent is likely to select action $\text{argmax}_a \{Q(D, a1) = 1, Q(D, a2) = 0\} = a1$. Suppose the episode is $(B, a1, 1, C, a1, 1, T)$
- $Q(B, a1) \leftarrow R_{t+1} + \gamma \max_a Q(C, a) = 1 + \max(1, 0) = 2$, $Q(C, a1) \leftarrow R_{t+1} + \gamma Q(T, -) = 1 + 0 = 1$
- The difference from Sarsa lies in $Q(B, a2)$, which stays at 0 until the agent experienced EP4. So it got less scared than Sarsa (where $Q(B, a2)$ was updated to -100 after EP2), so QL agent is more likely to explore unseen states.
- Suppose EP5 is $(B, a2, 0, D, a2, 4, T)$:
- $Q(B, a2) \leftarrow R_{t+1} + \gamma \max_a Q(D, a) = 0 + \max(-100, 4) = 4$, $Q(D, a2) \leftarrow R_{t+1} + \gamma Q(T, -) = 4 + 0 = 4$




	$Q(B, a1)$	$Q(B, a2)$	$Q(C, a1)$	$Q(C, a2)$	$Q(D, a1)$	$Q(D, a2)$
Init	0	0	0	0	0	0
EP1	0	0	0	0	-100	0
EP2	0	0	0	0	-100	4
EP3	1	0	1	0	-100	4
EP4	2	0	1	0	-100	4
EP5	2	4	1	0	-100	4

Linear Chain Example

Linear Chain Example

- Consider the following MDP. Environment is deterministic. In each state, there are two possible actions $a \in \{l, r\}$, where l corresponds to moving left, and r corresponds to moving right. Each movement incurs a reward of $r = -1$. State $s = 4$ is the goal state: taking any action from $s = 4$ results in reward of $r = 0$ and ends the episode by going into the terminal state, hence $V(4) = 0$, $Q(4, a) = 0$ for any action a . (Alternatively, we can view state 4 as the terminal state itself.) Assume $\gamma = 1$, $\alpha = 1$. All value functions are initialized to 0.
- A. Use Policy Iteration, Value Iteration to derive optimal policy.





TD, Sarsa, QL

- B. Consider 8 given consecutive episodes in the form of (s,a,r) (we do not consider ϵ -greedy exploration here):
 1. EP1: (1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)
 2. EP2: (1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)
 3. EP3: (1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)
 4. EP4: (3, l, -1), (2, l, -1), (1, l, -1), (1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)
 5. EP5: (3, l, -1), (2, l, -1), (1, l, -1), (1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)
 6. EP6: (3, l, -1), (2, l, -1), (1, l, -1), (1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)
 7. EP7: (3, l, -1), (2, l, -1), (1, l, -1), (1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)
 8. EP8: (3, l, -1), (2, l, -1), (1, l, -1), (1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)
- Derive the following:
 1. State value functions (V functions) after TD learning.
 2. State-action value functions (Q Functions) after Sarsa, and the resulting policy.
 3. State-action value functions (Q Functions) after Q learning, and the resulting policy.

Policy Iteration

1.1 Policy Evaluation of Random Policy


- Bellman Exp Equation: $v_{\pi}(s) = \sum_a \pi(a|s)q_{\pi}(s, a)$; $q_{\pi}(s, a) = R_s^a + \gamma v_{\pi}(s')$
- $v_{\pi}(1) = .5[q_{\pi}(1, l) + q_{\pi}(1, r)] = -1 + .5[v_{\pi}(1) + v_{\pi}(2)]$
 - $q_{\pi}(1, l) = -1 + v_{\pi}(1), q_{\pi}(1, r) = -1 + v_{\pi}(2)$
- $v_{\pi}(2) = .5[q_{\pi}(2, l) + q_{\pi}(2, r)] = -1 + .5[v_{\pi}(1) + v_{\pi}(3)]$
 - $q_{\pi}(2, l) = -1 + v_{\pi}(1), q_{\pi}(2, r) = -1 + v_{\pi}(3)$
- $v_{\pi}(3) = .5[q_{\pi}(3, l) + q_{\pi}(3, r)] = -1 + .5 v_{\pi}(2)$
 - $q_{\pi}(3, l) = -1 + v_{\pi}(2), q_{\pi}(3, r) = -1 + v_{\pi}(4) = -1$
- Solution: $v_{\pi}(1) = -12, v_{\pi}(2) = -10, v_{\pi}(3) = -6$



	$V_{\pi}(1)$	$V_{\pi}(2)$	$V_{\pi}(3)$
Iter1	-12	-10	-6
Iter2	4	3	4



1.2 Policy Improvement

- Plug in values from PE to get new policy
- $\pi'(1) = \operatorname{argmax}_a (q_\pi(1, l), q_\pi(1, r)) = r$
 - $q_\pi(1, l) = -1 + v_\pi(1) = -13$, $q_\pi(1, r) = -1 + v_\pi(2) = -11$,
- $\pi'(2) = \operatorname{argmax}_a (q_\pi(2, l), q_\pi(2, r)) = r$
 - $q_\pi(2, l) = -1 + v_\pi(1) = -13$, $q_\pi(2, r) = -1 + v_\pi(3) = -7$
- $\pi'(3) = \operatorname{argmax}_a (q_\pi(3, l), q_\pi(3, r)) = r$
 - $q_\pi(3, l) = -1 + v_\pi(2) = -11$, $q_\pi(3, r) = -1$



2.1 Policy Evaluation of Det Policy


- $v_{\pi}(1) = 1.0q_{\pi}(1, r) = -1 + v_{\pi}(2)$
 - $q_{\pi}(1, r) = -1 + v_{\pi}(2)$
- $v_{\pi}(2) = 1.0q_{\pi}(2, r) = -1 + v_{\pi}(3)$
 - $q_{\pi}(2, r) = -1 + v_{pi}(3)$
- $v_{\pi}(3) = 1.0q_{\pi}(3, r) = -1$
 - $q_{\pi}(3, r) = -1$
- Solution: $v_{\pi}(1) = -3, v_{\pi}(2) = -2, v_{\pi}(3) = -1$



	$V_{\pi}(1)$	$V_{\pi}(2)$	$V_{\pi}(3)$
Iter1	-12	-10	-6
Iter2	-3	-2	-1

2.2 Policy Improvement



- Plug in values from PE to get new policy
- $\pi'(1) = \operatorname{argmax}_a(q_\pi(1, l), q_\pi(1, r)) = r$
 - $q_\pi(1, l) = -1 - 3 = -4, q_\pi(1, r) = -1 - 2 = -3$
- $\pi'(2) = \operatorname{argmax}_a(q_\pi(2, l), q_\pi(2, r)) = r$
 - $q_\pi(2, l) = -1 - 3 = -4, q_\pi(2, r) = -1 - 1 = -2$
- $\pi'(3) = \operatorname{argmax}_a(q_\pi(3, l), q_\pi(3, r)) = r$
 - $q_\pi(3, l) = -1 - 2 = -3, q_\pi(3, r) = -1$
- Policy is now stable



Value Iteration

Value Iteration

- Bellman Opt Equation: $v_*(s) = \max_a q_*(s, a); q_*(s, a) = R_s^a + \gamma v_*(s')$
- $v_*(1) = \max_a [q_*(1, l), q_*(1, r)] = \max_a [-1 + v_*(1), -1 + v_*(2)]$
 - $q_*(1, l) = -1 + v_*(1), q_*(1, r) = -1 + v_*(2)$
- $v_*(2) = \max_a [q_*(2, l), q_*(2, r)] = \max_a [-1 + v_*(1), -1 + v_*(3)]$
 - $q_*(2, l) = -1 + v_*(1), q_*(2, r) = -1 + v_*(3)$
- $v_*(3) = \max_a [q_*(3, l), q_*(3, r)] = \max_a [-1 + v_*(2), -1 + v_*(4)] = \max_a [-1 + v_*(2), -1]$
 - $q_*(3, l) = -1 + v_*(2), q_*(3, r) = -1 + v_*(4) = -1$
- We use Value Iteration to solve it. Table shows the iteration process until convergence (not using in-place updates for clarity). Solution: $v_*(1) = -3, v_*(2) = -2, v_*(3) = -1$
- Optimal policy: $\pi_*(1) = \operatorname{argmax}_a q_*(1, a) = r; \pi_*(2) = \operatorname{argmax}_a q_*(2, a) = r; \pi_*(3) = \operatorname{argmax}_a q_*(3, a) = r$



(The $V_*(4)$ column is omitted since it is always 0)

	$V_*(1)$	$V_*(2)$	$V_*(3)$
Init	0	0	0
Iter1	-1	-1	-1
Iter2	-2	-2	-1
Iter3	-3	-2	-1
Iter4	-3	-2	-1

MC

Recall: MC Prediction Details

- State s may be visited multiple times in the same episode; let us call the first time it is visited in an episode the first visit to s .
 - The **first-visit** MC method estimates $v_\pi(s)$ as the average of the returns following first visits to s ,
 - The **every-visit** MC method averages the returns following all visits to s (show below).

MC prediction, for estimating $V \approx v_\pi$

Input: a policy π to be evaluated

Initialize:

$V(s) \in \mathbb{R}$, arbitrarily, for all $s \in \mathcal{S}$

$Returns(s) \leftarrow$ an empty list, for all $s \in \mathcal{S}$

Loop forever (for each episode):

Generate an episode following π : $S_0, A_0, R_1, S_1, A_1, R_2, \dots, S_{T-1}, A_{T-1}, R_T$

$G \leftarrow 0$

Loop for each step of episode, $t = T-1, T-2, \dots, 0$:

$G \leftarrow \gamma G + R_{t+1}$

Append G to $Returns(S_t)$

$V(S_t) \leftarrow \text{average}(Returns(S_t))$



Go backwards from last non-terminal S_{T-1}



Can compute running avg w. incremental update $V(S_t) \leftarrow_\alpha Returns(S_t)$

- TD update equation: $V(S_t) \leftarrow G_t$
- $V(4) \equiv 0$. Initialize $V(1) = V(2) = V(3) = 0$,
- EP1: $(1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)$

1. $G(3) \leftarrow -1, V(3) \leftarrow G(3) = -1$
2. $G(2) \leftarrow -1 + \gamma G(3) = -2, V(2) \leftarrow G(2) = -2$
3. $G(1) \leftarrow -1 + \gamma G(2) = -3, V(1) \leftarrow G(1) = -3$


- EP2: $(1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)$

1. $G(3) \leftarrow -1, V(3) \leftarrow G(3) = -1$
2. $G(2) \leftarrow -1 + \gamma G(3) = -2, V(2) \leftarrow G(2) = -2$
3. $G(1) \leftarrow -1 + \gamma G(2) = -3, V(1) \leftarrow G(1) = -3$

- EP3: $(1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)$

1. $G(3) \leftarrow -1, V(3) \leftarrow G(3) = -1$
2. $G(2) \leftarrow -1 + \gamma G(3) = -2, V(2) \leftarrow G(2) = -2$
3. $G(1) \leftarrow -1 + \gamma G(2) = -3, V(1) \leftarrow G(1) = -3$

MC EP1-3




TD	$V(1)$	$V(2)$	$V(3)$
Init	0	0	0
After EP1	-3	-2	-1
After EP2	-3	-2	-1
After EP3	-3	-2	-1
After EP4	-3.5	-3.5	-3.5
After EP5	-3.5	-3.5	-3.5
After EP6	-3.5	-3.5	-3.5
After EP7	-3.5	-3.5	-3.5
After EP8	-3.5	-3.5	-3.5

- TD update equation: $V(S_t) \leftarrow G_t$
- EP4:
 $(3, l, -1), (2, l, -1), (1, l, -1), (1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)$

MC EP4-8

1. $G(3) \leftarrow -1$ (2nd visit)
2. $G(2) \leftarrow -1 + \gamma G(3) = -2$ (2nd visit)
3. $G(1) \leftarrow -1 + \gamma G(2) = -3$ (2nd visit)
4. $G(1)' \leftarrow -1 + \gamma G(1) = -4$ (1st visit)
5. $G(2)' \leftarrow -1 + \gamma G(1)' = -5$ (1st visit)
6. $G(3)' \leftarrow -1 + \gamma G(2)' = -6$ (1st visit)
7. (Every visit):
8. $V(3) = .5(G(3) + G(3)') = -3.5$
9. $V(2) = .5(G(2) + G(2)') = -3.5$
10. $V(1) = .5(G(1) + G(1)') = -3.5$
11. (First visit):
12. $V(3) = G(3)' = -6$
13. $V(2) = G(2)' = -5$
14. $V(1) = G(1)' = -4$

- EP5-8 omitted



TD	$V(1)$	$V(2)$	$V(3)$
Init	0	0	0
After EP1	-3	-2	-1
After EP2	-3	-2	-1
After EP3	-3	-2	-1
After EP4	-3.5	-3.5	-3.5
After EP5	-3.5	-3.5	-3.5
After EP6	-3.5	-3.5	-3.5
After EP7	-3.5	-3.5	-3.5
After EP8	-3.5	-3.5	-3.5

TD

- TD update equation: $V(S_t) \leftarrow R_{t+1} + V(S_{t+1})$
 - With $\gamma = 1, \alpha = 1$, each $V(s)$ is completely replaced/overwritten by the TD update
- $V(4) \equiv 0$. Initialize $V(1) = V(2) = V(3) = 0$,

• EP1: $(1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)$

1. $V(1) \leftarrow -1 + V(2) = -1 + 0 = -1$

2. $V(2) \leftarrow -1 + V(3) = -1 + 0 = -1$

3. $V(3) \leftarrow -1 + V(4) = -1 + 0 = -1$

• EP2: $(1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)$

1. $V(1) \leftarrow -1 + V(2) = -1 - 1 = -2$

2. $V(2) \leftarrow -1 + V(3) = -1 - 1 = -2$

3. $V(3) \leftarrow -1 + V(4) = -1 + 0 = -1$

• EP3: $(1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)$


1. $V(1) \leftarrow -1 + V(2) = -1 - 2 = -3$

2. $V(2) \leftarrow -1 + V(3) = -1 - 1 = -2$

3. $V(3) \leftarrow -1 + V(4) = -1 + 0 = -1$

- Arrows denote bootstrap dependencies, e.g., $V(1)$ bootstraps off $V(2)$, $V(2)$ bootstraps off $V(3)$, $V(3)$ bootstraps off $V(4)$. They also denote direction of information flow during learning, e.g., $V(4) \equiv 0$ is the external learning signal, and info flows $V(4) \rightarrow V(3) \rightarrow V(2) \rightarrow V(1)$.

TD EP1-3



TD	$V(1)$	$V(2)$	$V(3)$
Init	0	0	0
After EP1	-1	-1	-1
After EP2	-2	-2	-1
After EP3	-3	-2	-1
After EP4	-5	-4	-1
After EP5	-7	-6	-1
After EP6	-9	-8	-1
After EP7	-11	-10	-1
After EP8	-13	-12	-1

- TD update equation: $V(S_t) \leftarrow R_{t+1} + V(S_{t+1})$

1. EP4:

$(3, l, -1), (2, l, -1), (1, l, -1), (1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)$

2. $V(3) \leftarrow -1 + V(2) = -1 - 2 = -3$

3. $V(2) \leftarrow -1 + V(1) = -1 - 3 = -4$

4. $V(1) \leftarrow -1 + V(1) = -1 - 3 = -4$

5. $V(1) \leftarrow -1 + V(2) = -1 - 4 = -5$

6. $V(2) \leftarrow -1 + V(3) = -1 - 3 = -4$

7. $V(3) \leftarrow -1 + V(4) = -1 + 0 = -1$

• EP5:

$(3, l, -1), (2, l, -1), (1, l, -1), (1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)$

1. $V(3) \leftarrow -1 + V(2) = -1 - 4 = -5$

2. $V(2) \leftarrow -1 + V(1) = -1 - 5 = -6$

3. $V(1) \leftarrow -1 + V(1) = -1 - 5 = -6$


4. $V(1) \leftarrow -1 + V(2) = -1 - 6 = -7$

5. $V(2) \leftarrow -1 + V(3) = -1 - 5 = -6$

6. $V(3) \leftarrow -1 + V(4) = -1 + 0 = -1$

- EP6-8 omitted. Some arrows are omitted, since some values are overwritten within an episode.

TD EP4-8




TD	V(1)	V(2)	V(3)
Init	0	0	0
After EP1	-1	-1	-1
After EP2	-2	-2	-1
After EP3	-3	-2	-1
After EP4	-5	-4	-1
After EP5	-7	-6	-1
After EP6	-9	-8	-1
After EP7	-11	-10	-1
After EP8	-13	-12	-1

TD failed to converge

- TD failed to converge for this set of episodes. The sequence of TD updates cause all value functions to be increasingly negative.

- For simplicity, consider the infinite sequence of $(2, l, -1), (1, r, -1), (2, l, -1) \dots$
- The sequence of TD updates: $V(2) = -1 + V(1), V(1) = -1 + V(2), \dots$ So $V(1)$ and $V(2)$ bootstrap off each other and both go to $-\infty$.
- An analogy is that two students $V(1)$ and $V(2)$ are copying from each other, but they never get any true reward feedback from the teacher ($V(4) = 0$)



- $V(3)$ is bootstrapped off $V(2)$ when moving left, and is bootstrapped off $V(4) \equiv 0$ when moving right. Steps 1-5 form a bootstrap dependency cycle $V(3) \leftarrow V(2) \leftarrow V(1) \leftarrow V(2) \leftarrow V(3)$ that causes $V(1), V(2), V(3)$ to blow up. Even though $V(3)$ is updated to $V(3) = -1 + V(4) = -1$ when it moves right to state 4, the episode ends immediately afterwards, so $V(1)$ and $V(2)$ do not have a chance to bootstrap off the correct $V(3)$.

- $V(3) \leftarrow -1 + V(2) = -1 - 4 = -5$
- $V(2) \leftarrow -1 + V(1) = -1 - 5 = -6$
- $V(1) \leftarrow -1 + V(1) = -1 - 5 = -6$
- $V(1) \leftarrow -1 + V(2) = -1 - 6 = -7$
- $V(2) \leftarrow -1 + V(3) = -1 - 5 = -6$
- $V(3) \leftarrow -1 + V(4) = -1 + 0 = -1$


- If the episode does not end immediately, but the agent moves left again, then $V(1)$ and $V(2)$ will have a chance to bootstrap off the new $V(3)$, and they may converge to the correct values.

TD	$V(1)$	$V(2)$	$V(3)$
Init	0	0	0
After EP1	-1	-1	-1
After EP2	-2	-2	-1
After EP3	-3	-2	-1
After EP4	-5	-4	-1
After EP5	-7	-6	-1
After EP6	-9	-8	-1
After EP7	-11	-10	-1
After EP8	-13	-12	-1

Sarsa

- Sarsa update equation: $Q(S_t, A_t) \leftarrow Q(S_t, A_t) + \alpha(R_{t+1} + \gamma Q(S_{t+1}, A_{t+1}) - Q(S_t, A_t)) = R_{t+1} + \gamma Q(S_{t+1}, A_{t+1})$
 - With $\gamma = 1, \alpha = 1$, each $Q(S, A)$ is completely replaced/overwritten by the Sarsa update
- $Q(4, a) \equiv 0$. Initialize $Q(1, *) = Q(2, *) = Q(3, *) = 0$
- After EP1: $(1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)$
 - $Q(1, r) \leftarrow -1 + Q(2, r) = -1 + 0 = -1$
 - $Q(2, r) \leftarrow -1 + Q(3, r) = -1 + 0 = -1$
 - $Q(3, r) \leftarrow -1 + Q(4, r) = -1 + 0 = -1$
- After EP2: $(1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)$
 - $Q(1, r) \leftarrow -1 + Q(2, r) = -1 - 1 = -2$
 - $Q(2, r) \leftarrow -1 + Q(3, r) = -1 - 1 = -2$
 - $Q(3, r) \leftarrow -1 + Q(4, r) = -1 + 0 = -1$
- After EP3: $(1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)$
 - $Q(1, r) \leftarrow -1 + Q(2, r) = -1 - 2 = -3$
 - $Q(2, r) \leftarrow -1 + Q(3, r) = -1 - 1 = -2$
 - $Q(3, r) \leftarrow -1 + Q(4, r) = -1 + 0 = -1$

Sarsa EP1-3



Sarsa	$Q(1, l)$	$Q(1, r)$	$Q(2, l)$	$Q(2, r)$	$Q(3, l)$	$Q(3, r)$
Init	0	0	0	0	0	0
After EP1	0	-1	0	-1	0	-1
After EP2	0	-2	0	-2	0	-1
After EP3	0	-3	0	-2	0	-1
After EP4	-4	-3	-1	-2	-1	-1
After EP5	-4	-3	-5	-2	-2	-1
After EP6	-4	-3	-5	-2	-6	-1
After EP7	-4	-3	-5	-2	-6	-1
After EP8	-4	-3	-5	-2	-6	-1

- Sarsa update equation: $Q(S_t, A_t) \leftarrow R_{t+1} + Q(S_{t+1}, A_{t+1})$
- EP4: (3, l, -1), (2, l, -1), (1, l, -1), (1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)

$$1. \quad Q(3, l) \leftarrow -1 + Q(2, l) = -1 + 0 = -1$$

$$2. \quad Q(2, l) \leftarrow -1 + Q(1, l) = -1 + 0 = -1$$

$$3. \quad Q(1, l) \leftarrow -1 + Q(1, r) = -1 - 3 = -4$$

$$4. \quad Q(1, r) \leftarrow -1 + Q(2, r) = -1 - 2 = -3$$

$$5. \quad Q(2, r) \leftarrow -1 + Q(3, r) = -1 - 1 = -2$$

$$6. \quad Q(3, r) \leftarrow -1 + Q(4, r) = -1 + 0 = -1$$

- EP5: (3, l, -1), (2, l, -1), (1, l, -1), (1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)

$$1. \quad Q(3, l) \leftarrow -1 + Q(2, l) = -1 - 1 = -2$$

$$2. \quad Q(2, l) \leftarrow -1 + Q(1, l) = -1 - 4 = -5$$

$$3. \quad Q(1, l) \leftarrow -1 + Q(1, r) = -1 - 3 = -4$$

$$4. \quad Q(1, r) \leftarrow -1 + Q(2, r) = -1 - 2 = -3$$

$$5. \quad Q(2, r) \leftarrow -1 + Q(3, r) = -1 - 1 = -2$$

$$6. \quad Q(3, r) \leftarrow -1 + Q(4, r) = -1 + 0 = -1$$

- EP6: (3, l, -1), (2, l, -1), (1, l, -1), (1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)

$$1. \quad Q(3, l) \leftarrow -1 + Q(2, l) = -1 - 5 = -6$$

$$2. \quad Q(2, l) \leftarrow -1 + Q(1, l) = -1 - 4 = -5$$

$$3. \quad Q(1, l) \leftarrow -1 + Q(1, r) = -1 - 3 = -4$$

$$4. \quad Q(1, r) \leftarrow -1 + Q(2, r) = -1 - 2 = -3$$

$$5. \quad Q(2, r) \leftarrow -1 + Q(3, r) = -1 - 1 = -2$$

$$6. \quad Q(3, r) \leftarrow -1 + Q(4, r) = -1 + 0 = -1$$

- EP7: (3, l, -1), (2, l, -1), (1, l, -1), (1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)

$$1. \quad Q(3, l) \leftarrow -1 + Q(2, l) = -1 - 5 = -6$$

$$2. \quad Q(2, l) \leftarrow -1 + Q(1, l) = -1 - 4 = -5$$


$$3. \quad Q(1, l) \leftarrow -1 + Q(1, r) = -1 - 3 = -4$$

$$4. \quad Q(1, r) \leftarrow -1 + Q(2, r) = -1 - 2 = -3$$

$$5. \quad Q(2, r) \leftarrow -1 + Q(3, r) = -1 - 1 = -2$$

$$6. \quad Q(3, r) \leftarrow -1 + Q(4, r) = -1 + 0 = -1 \text{ (EP8 omitted)}$$

Sarsa EP4-8




Sarsa	$Q(1, l)$	$Q(1, r)$	$Q(2, l)$	$Q(2, r)$	$Q(3, l)$	$Q(3, r)$
Init	0	0	0	0	0	0
After EP1	0	-1	0	-1	0	-1
After EP2	0	-2	0	-2	0	-1
After EP3	0	-3	0	-2	0	-1
After EP4	-4	-3	-1	-2	-1	-1
After EP5	-4	-3	-5	-2	-2	-1
After EP6	-4	-3	-5	-2	-6	-1
After EP7	-4	-3	-5	-2	-6	-1
After EP8	-4	-3	-5	-2	-6	-1

Q values have converged at EP6. Bootstrap dependency arrows are omitted for EP7-8, since they are the same as EP6. **Red arrows** denote the stable set of dependencies that keep the Q values stable after EP6.

Comments on Sarsa

- State-action value functions for moving right look reasonable: $Q(1, r) = -3, Q(2, r) = -2, Q(3, r) = -1$.
- State-action value functions for moving left look unreasonable: $Q(1, l) = -4, Q(2, l) = -5, Q(3, l) = -6$. This is because the only trajectory with move left actions are $3 \rightarrow 2 \rightarrow 1 \rightarrow 1 \rightarrow 2 \rightarrow 3 \rightarrow 4$, the Q values are updated based on only this episode (on-policy), i.e., from state 3 taking action left, it can only take the above trajectory, and reach the goal in 6 steps, hence $Q(3, l) = -6$. If we had collected more trajectories like $3 \rightarrow 2 \rightarrow 3 \rightarrow 4$, then Sarsa could learn the more accurate Q value $Q(3, l) = -1 + Q(2, r) = -3$.
- Even though the Q values for left actions are inaccurate, the greedy policy is still optimal since right action is always better than left:
 - $\pi_*(1) = \operatorname{argmax}_a (Q(1, l), Q(1, r)) = r$
 - $\pi_*(2) = \operatorname{argmax}_a (Q(2, l), Q(2, r)) = r$
 - $\pi_*(3) = \operatorname{argmax}_a (Q(3, l), Q(3, r)) = r$




Sarsa	$Q(1, l)$	$Q(1, r)$	$Q(2, l)$	$Q(2, r)$	$Q(3, l)$	$Q(3, r)$
Init	0	0	0	0	0	0
After EP1	0	-1	0	-1	0	-1
After EP2	0	-2	0	-2	0	-1
After EP3	0	-3	0	-2	0	-1
After EP4	-4	-3	-1	-2	-1	-1
After EP5	-4	-3	-5	-2	-2	-1
After EP6	-4	-3	-5	-2	-6	-1
After EP7	-4	-3	-5	-2	-6	-1
After EP8	-4	-3	-5	-2	-6	-1

Why Sarsa did not blow up

- TD: $V(s)$ is updated regardless if agent moves left or right. $V(3)$ is bootstrapped off $V(2)$ when moving left, and is bootstrapped off $V(4) \equiv 0$ when moving right. bootstrap dependency cycle $V(3) \leftarrow V(2) \leftarrow V(1) \leftarrow V(2) \leftarrow V(3)$ that causes $V(1), V(2), V(3)$ to blow up
- Sarsa: when agent moves left, $Q(s, l)$ is updated; when agent moves right, $Q(s, r)$ is updated. $Q(3, r)$ is always bootstrapped off $Q(4, r) \equiv 0$. So there is no bootstrap dependency cycle like TD. The linear dependency chain from $Q(4, r)$ to $Q(3, l)$ determines the stable values:

1. $Q(3, l) \leftarrow -1 + Q(2, l) = -1 - 5 = -6$
2. $Q(2, l) \leftarrow -1 + Q(1, l) = -1 - 4 = -5$
3. $Q(1, l) \leftarrow -1 + Q(1, r) = -1 - 3 = -4$
4. $Q(1, r) \leftarrow -1 + Q(2, r) = -1 - 2 = -3$
5. $Q(2, r) \leftarrow -1 + Q(3, r) = -1 - 1 = -2$
6. $Q(3, r) \leftarrow -1 + Q(4, r) = -1 + 0 = -1$



Sarsa	$Q(1, l)$	$Q(1, r)$	$Q(2, l)$	$Q(2, r)$	$Q(3, l)$	$Q(3, r)$
Init	0	0	0	0	0	0
After EP1	0	-1	0	-1	0	-1
After EP2	0	-2	0	-2	0	-1
After EP3	0	-3	0	-2	0	-1
After EP4	-4	-3	-1	-2	-1	-1
After EP5	-4	-3	-5	-2	-2	-1
After EP6	-4	-3	-5	-2	-6	-1
After EP7	-4	-3	-5	-2	-6	-1
After EP8	-4	-3	-5	-2	-6	-1

Q Learning

- QL update equation: $(S_t, A_t) \leftarrow Q(S_t, A_t) + \alpha (R_{t+1} + \gamma \max_{a'} Q(S_{t+1}, a') - Q(S_t, A_t)) = R_{t+1} + \gamma \max_{a'} Q(S_{t+1}, a')$
 - With $\gamma = 1, \alpha = 1$, each $Q(S, A)$ is completely replaced overwritten by the Q update

- After EP1: $(1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)$

$$1. \quad Q(1, r) \leftarrow -1 + \max_{a'} Q(2, a') = -1 + \max(0, 0) = -1$$

$$2. \quad Q(2, r) \leftarrow -1 + \max_{a'} Q(3, a') = -1 + \max(0, 0) = -1$$

$$3. \quad Q(3, r) \leftarrow -1 + \max_{a'} Q(4, a') = -1 + 0 = -1$$

- After EP2: $(1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)$

$$1. \quad Q(1, r) \leftarrow -1 + \max_{a'} Q(2, a') = -1 + \max(-1, 0) = -1$$

$$2. \quad Q(2, r) \leftarrow -1 + \max_{a'} Q(3, a') = -1 + \max(-1, 0) = -1$$

$$3. \quad Q(3, r) \leftarrow -1 + \max_{a'} Q(4, a') = -1 + 0 = -1$$


- After EP3: $(1, r, -1), (2, r, -1), (3, r, -1), (4, r, 0)$

$$1. \quad Q(1, r) \leftarrow -1 + \max_{a'} Q(2, a') = -1 + \max(-1, 0) = -1$$

$$2. \quad Q(2, r) \leftarrow -1 + \max_{a'} Q(3, a') = -1 + \max(-1, 0) = -1$$

$$3. \quad Q(3, r) \leftarrow -1 + \max_{a'} Q(4, a') = -1 + 0 = -1$$

QL EP1-3



QL	$Q(1, l)$	$Q(1, r)$	$Q(2, l)$	$Q(2, r)$	$Q(3, l)$	$Q(3, r)$
Init	0	0	0	0	0	0
After EP1	0	-1	0	-1	0	-1
After EP2	0	-1	0	-1	0	-1
After EP3	0	-1	0	-1	0	-1
After EP4	-1	-2	-1	-2	-1	-1
After EP5	-2	-3	-2	-2	-2	-1
After EP6	-3	-3	-3	-2	-3	-1
After EP7	-4	-3	-4	-2	-3	-1
After EP8	-4	-3	-4	-2	-3	-1

- EP4: (3,l,-1), (2,l,-1), (1,l,-1), (1,r,-1), (2,r,-1), (3,r,-1), (4,r,0)

1. $Q(3,l) \leftarrow -1 + \max_{a'} Q(2,a') = -1 + \max(0, -1) = -1$
2. $Q(2,l) \leftarrow -1 + \max_{a'} Q(1,a') = -1 + \max(0, -1) = -1$
3. $Q(1,l) \leftarrow -1 + \max_{a'} Q(1,a') = -1 + \max(0, -1) = -1$
4. $Q(1,r) \leftarrow -1 + \max_{a'} Q(2,a') = -1 + \max(-1, -1) = -2$
5. $Q(2,r) \leftarrow -1 + \max_{a'} Q(3,a') = -1 + \max(-1, -1) = -2$
6. $Q(3,r) \leftarrow -1 + \max_{a'} Q(4,a') = -1 + 0 = -1$


- EP5: (3,l,-1), (2,l,-1), (1,l,-1), (1,r,-1), (2,r,-1), (3,r,-1), (4,r,0)

1. $Q(3,l) \leftarrow -1 + \max_{a'} Q(2,a') = -1 + \max(-1, -2) = -2$
2. $Q(2,l) \leftarrow -1 + \max_{a'} Q(1,a') = -1 + \max(-1, -2) = -2$
3. $Q(1,l) \leftarrow -1 + \max_{a'} Q(1,a') = -1 + \max(-1, -2) = -2$
4. $Q(1,r) \leftarrow -1 + \max_{a'} Q(2,a') = -1 + \max(-2, -2) = -3$
5. $Q(2,r) \leftarrow -1 + \max_{a'} Q(3,a') = -1 + \max(-2, -1) = -2$
6. $Q(3,r) \leftarrow -1 + \max_{a'} Q(4,a') = -1 + 0 = -1$

- EP6: (3,l,-1), (2,l,-1), (1,l,-1), (1,r,-1), (2,r,-1), (3,r,-1), (4,r,0)

1. $Q(3,l) \leftarrow -1 + \max_{a'} Q(2,a') = -1 + \max(-2, -2) = -3$
2. $Q(2,l) \leftarrow -1 + \max_{a'} Q(1,a') = -1 + \max(-2, -3) = -3$
3. $Q(1,l) \leftarrow -1 + \max_{a'} Q(1,a') = -1 + \max(-2, -3) = -3$
4. $Q(1,r) \leftarrow -1 + \max_{a'} Q(2,a') = -1 + \max(-3, -2) = -3$
5. $Q(2,r) \leftarrow -1 + \max_{a'} Q(3,a') = -1 + \max(-3, -1) = -2$
6. $Q(3,r) \leftarrow -1 + \max_{a'} Q(4,a') = -1 + 0 = -1$

QL EP4-6



QL	$Q(1,l)$	$Q(1,r)$	$Q(2,l)$	$Q(2,r)$	$Q(3,l)$	$Q(3,r)$
Init	0	0	0	0	0	0
After EP1	0	-1	0	-1	0	-1
After EP2	0	-1	0	-1	0	-1
After EP3	0	-1	0	-1	0	-1
After EP4	-1	-2	-1	-2	-1	-1
After EP5	-2	-3	-2	-2	-2	-1
After EP6	-3	-3	-3	-2	-3	-1
After EP7	-4	-3	-4	-2	-3	-1
After EP8	-4	-3	-4	-2	-3	-1

- EP7: (3,l,-1), (2,l,-1), (1,l,-1), (1,r,-1), (2,r,-1), (3,r,-1), (4,r,0)

$$1. \quad Q(3,l) \leftarrow -1 + \max_{a'} Q(2,a') = -1 + \max(-3,-2) = -3$$

$$2. \quad Q(2,l) \leftarrow -1 + \max_{a'} Q(1,a') = -1 + \max(-3,-3) = -4$$

$$3. \quad Q(1,l) \leftarrow -1 + \max_{a'} Q(1,a') = -1 + \max(-3,-3) = -4$$

$$4. \quad Q(1,r) \leftarrow -1 + \max_{a'} Q(2,a') = -1 + \max(-4,-2) = -3$$

$$5. \quad Q(2,r) \leftarrow -1 + \max_{a'} Q(3,a') = -1 + \max(-3,-1) = -2$$

$$6. \quad Q(3,r) \leftarrow -1 + \max_{a'} Q(4,a') = -1 + 0 = -1$$

- EP8: (3,l,-1), (2,l,-1), (1,l,-1), (1,r,-1), (2,r,-1), (3,r,-1), (4,r,0)

$$1. \quad Q(3,l) \leftarrow -1 + \max_{a'} Q(2,a') = -1 + \max(-4,-2) = -3$$

$$2. \quad Q(2,l) \leftarrow -1 + \max_{a'} Q(1,a') = -1 + \max(-4,-3) = -4$$

$$3. \quad Q(1,l) \leftarrow -1 + \max_{a'} Q(1,a') = -1 + \max(-4,-3) = -4$$


$$4. \quad Q(1,r) \leftarrow -1 + \max_{a'} Q(2,a') = -1 + \max(-4,-2) = -3$$

$$5. \quad Q(2,r) \leftarrow -1 + \max_{a'} Q(3,a') = -1 + \max(-3,-1) = -2$$

$$6. \quad Q(3,r) \leftarrow -1 + \max_{a'} Q(4,a') = -1 + 0 = -1$$


- Q values have converged at EP7. Bootstrap dependency arrows are omitted for EP8, since they are the same as EP7. **Red arrows** denote the stable set of dependencies that keep the Q values stable after EP7.

QL EP7-8



QL	$Q(1,l)$	$Q(1,r)$	$Q(2,l)$	$Q(2,r)$	$Q(3,l)$	$Q(3,r)$
Init	0	0	0	0	0	0
After EP1	0	-1	0	-1	0	-1
After EP2	0	-1	0	-1	0	-1
After EP3	0	-1	0	-1	0	-1
After EP4	-1	-2	-1	-2	-1	-1
After EP5	-2	-3	-2	-2	-2	-1
After EP6	-3	-3	-3	-2	-3	-1
After EP7	-4	-3	-4	-2	-3	-1
After EP8	-4	-3	-4	-2	-3	-1

Comments on QL



QL	$Q(1, l)$	$Q(1, r)$	$Q(2, l)$	$Q(2, r)$	$Q(3, l)$	$Q(3, r)$
Init	0	0	0	0	0	0
After EP1	0	-1	0	-1	0	-1
After EP2	0	-1	0	-1	0	-1
After EP3	0	-1	0	-1	0	-1
After EP4	-1	-2	-1	-2	-1	-1
After EP5	-2	-3	-2	-2	-2	-1
After EP6	-3	-3	-3	-2	-3	-1
After EP7	-4	-3	-4	-2	-3	-1
After EP8	-4	-3	-4	-2	-3	-1

- QL converges. All state-action value functions look reasonable.
- $Q(1, r) = -3, Q(2, r) = -2, Q(3, r) = -1$. The optimal path can be derived from bootstrap dependencies, e.g., dependency chain $Q(1, r) \leftarrow Q(2, r) \leftarrow Q(3, r)$ corresponds to the optimal path $1 \rightarrow 2 \rightarrow 3 \rightarrow 4$.
- $Q(1, l) = -4$: If agent moves left in state 1, dependency chain $Q(1, l) \leftarrow Q(1, r) \leftarrow Q(2, r) \leftarrow Q(3, r)$ corresponds to the optimal path $1 \rightarrow 1 \rightarrow 2 \rightarrow 3 \rightarrow 4$ w. 4 steps to reach goal state 4.
- $Q(2, l) = -4$: If agent moves left in state 2, dependency chain $Q(2, l) \leftarrow Q(1, r) \leftarrow Q(2, r) \leftarrow Q(3, r)$ corresponds to the optimal path $2 \rightarrow 1 \rightarrow 2 \rightarrow 3 \rightarrow 4$ w. 4 steps to reach goal state 4.
- $Q(3, l) = -3$: If agent moves left in state 3, dependency chain $Q(3, l) \leftarrow Q(2, r) \leftarrow Q(3, r)$ corresponds to the optimal path $3 \rightarrow 2 \rightarrow 3 \rightarrow 4$ w. 3 steps to reach goal state 4.
- QL is smarter than Sarsa: since it is off-policy, agent can learn the correct Q value functions that correspond to trajectories that it has never actually experienced, e.g., if If agent moves left in state 3, even though it has never experienced the trajectory $3 \rightarrow 2 \rightarrow 3 \rightarrow 4$, the bootstrap dependency $Q(3, l) \leftarrow Q(2, r)$ lead to that trajectory instead of the experienced trajectory $3 \rightarrow 2 \rightarrow 1 \rightarrow 1 \rightarrow 2 \rightarrow 3 \rightarrow 4$.
- The intermediate Q values before convergence may not correspond to a valid policy, e.g., before EP7, $\arg\max_a Q(1, a) = l$, so the agent would be stuck in state 1 trying to go left forever.
- Q values learned by QL are accurate, and the greedy policy is optimal:
- $\pi_*(1) = \arg\max_a (Q(1, l), Q(1, r)) = r$
- $\pi_*(2) = \arg\max_a (Q(2, l), Q(2, r)) = r$
- $\pi_*(3) = \arg\max_a (Q(3, l), Q(3, r)) = r$