



Characteristic Boolean algebras in classical realizability: a completeness result


Guillaume Geoffroy


IRIF, Université Paris Cité



















Variant $\frac{1}{2}$




Variant $\frac{1}{2}$




Variant $1/2$




Variant ¹/₂




Variant ¹/₂




Variant ¹/₂




Variant $\frac{1}{2}$



Variant $\frac{1}{2}$







Variant $\frac{1}{2}$
Win if $r > 0$




Variant ²/₃




Variant ²/₃




Variant ²/₃




Variant ²/₃





Variant ²/₃





Variant 3/5





Variant k/n

Win if $r > ???$



Puzzle for you!



Classical realizability

Formulas: (A, B, \dots)

- ▶ first-order
- ▶ language of Boolean algebras

Programs: $t, u ::= x \mid \lambda x. t \mid tu$


Realizability relation: “ t realizes A ”

Classical realizability

Formulas: (A, B, \dots)

- ▶ first-order
- ▶ language of Boolean algebras

Programs: $t, u ::= x \mid \lambda x. t \mid tu$

Realizability relation: “ t realizes A ”

e.g. $\lambda x. \lambda y. xyy$
realizes $(A \rightarrow A \rightarrow B) \rightarrow A \rightarrow B$
for all A, B

A completeness result

Adequacy (standard): if A is true in all Boolean algebras,
then there exists a λ -term t that realizes A

A completeness result

Adequacy (standard): if A is true in all Boolean algebras,
then there exists a λ -term t that realizes A

Theorem: A is true in all Boolean algebras
iif there exists a λ -term t that realizes A

Back to the game

Allowed questions
(variant k/n)



Formula $A_{k,n}$

Initial situation
($N_{\text{red}}, N_{\text{balls}}$)



Formula $A_{N_{\text{red}}, N_{\text{balls}}}$

Back to the game

Allowed questions
(variant k/n)



Formula $A_{k,n}$

Initial situation
($N_{\text{red}}, N_{\text{balls}}$)



Formula $A_{N_{\text{red}}, N_{\text{balls}}}$

Winning strategy



λ -term that realizes
 $A_{k,n} \rightarrow A_{N_{\text{red}}, N_{\text{balls}}}$

Back to the game

Allowed questions
(variant k/n)



Formula $A_{k,n}$

Initial situation
($N_{\text{red}}, N_{\text{balls}}$)



Formula $A_{N_{\text{red}}, N_{\text{balls}}}$

\exists Winning strategy



In all Boolean algebras,
 $A_{k,n} \rightarrow A_{N_{\text{red}}, N_{\text{balls}}}$ holds

Back to the game

Allowed questions
(variant k/n)

Formula $A_{k,n}$
“ \exists at most $f(k/n)$ elements”

Initial situation
($N_{\text{red}}, N_{\text{balls}}$)

Formula $A_{N_{\text{red}}, N_{\text{balls}}}$
“ \exists at most $f(N_{\text{red}}/N_{\text{balls}})$ elements”

\exists Winning strategy

In all Boolean algebras,
 $A_{k,n} \rightarrow A_{N_{\text{red}}, N_{\text{balls}}}$ holds

Back to the game

Allowed questions
(variant k/n)

Formula $A_{k,n}$
“ \exists at most $f(k/n)$ elements”

Initial situation
($N_{\text{red}}, N_{\text{balls}}$)

Formula $A_{N_{\text{red}}, N_{\text{balls}}}$
“ \exists at most $f(N_{\text{red}}/N_{\text{balls}})$ elements”

\exists Winning strategy

In all Boolean algebras,
 $A_{k,n} \rightarrow A_{N_{\text{red}}, N_{\text{balls}}}$ holds
 $f(k/n) \leq f(N_{\text{red}}/N_{\text{balls}})$

Translating variant ^{2/3}

► The key: $\bullet \Leftrightarrow \perp$ $\bullet \Leftrightarrow \top$









Translating variant ^{2/3}

- The key:
-  $\Leftrightarrow \perp$
 -  $\Leftrightarrow \top$
 - universal quantification \Leftrightarrow intersection type

Translating variant $2/3$

- ▶ The key: $\bullet \text{ (red)} \Leftrightarrow \perp$ $\bullet \text{ (blue)} \Leftrightarrow \top$
 - universal quantification \Leftrightarrow intersection type

- ▶ Variant $2/3$ is the intersection of:



Translating variant $2/3$

- ▶ The key: $\bullet \text{ (red)} \Leftrightarrow \perp$ $\bullet \text{ (blue)} \Leftrightarrow \top$
 - universal quantification \Leftrightarrow intersection type
- ▶ Variant $2/3$ is the intersection of:

$\perp \rightarrow \perp \rightarrow \perp \rightarrow \perp$

$\perp \rightarrow \perp \rightarrow \top \rightarrow \perp$

$\perp \rightarrow \top \rightarrow \perp \rightarrow \perp$

$\top \rightarrow \perp \rightarrow \perp \rightarrow \perp$

$\top \rightarrow \top \rightarrow \top \rightarrow \top$

$\top \rightarrow \top \rightarrow \perp \rightarrow \top$

$\top \rightarrow \perp \rightarrow \top \rightarrow \top$

$\perp \rightarrow \top \rightarrow \top \rightarrow \top$

Translating variant $2/3$

- The key: $\bullet \Leftrightarrow \perp$ $\bullet \Leftrightarrow \top$
• universal quantification \Leftrightarrow intersection type

- Variant $2/3$ is the intersection of:

$\perp \rightarrow \perp \rightarrow \perp \rightarrow \perp$
 $\perp \rightarrow \perp \rightarrow \top \rightarrow \perp$
 $\perp \rightarrow \top \rightarrow \perp \rightarrow \perp$
 $\top \rightarrow \perp \rightarrow \perp \rightarrow \perp$

$\top \rightarrow \top \rightarrow \top \rightarrow \top$
 $\top \rightarrow \top \rightarrow \perp \rightarrow \top$
 $\top \rightarrow \perp \rightarrow \top \rightarrow \top$
 $\perp \rightarrow \top \rightarrow \top \rightarrow \top$

- $A_{2,3}$ is the formula:

$$\forall x \forall y \forall z (x \neq 0 \rightarrow y \neq 0 \rightarrow z \neq 0 \rightarrow (x \wedge y \vee y \wedge z \vee z \wedge x) \neq 0)$$

Translating variant $2/3$

- ▶ The key: $\bullet \Leftrightarrow \perp$ $\bullet \Leftrightarrow \top$
 - universal quantification \Leftrightarrow intersection type

- ▶ Variant $2/3$ is the intersection of:

$$\begin{array}{l} \perp \rightarrow \perp \rightarrow \perp \rightarrow \perp \\ \perp \rightarrow \perp \rightarrow \top \rightarrow \perp \\ \perp \rightarrow \top \rightarrow \perp \rightarrow \perp \\ \top \rightarrow \perp \rightarrow \perp \rightarrow \perp \end{array}$$
$$\begin{array}{l} \top \rightarrow \top \rightarrow \top \rightarrow \top \\ \top \rightarrow \top \rightarrow \perp \rightarrow \top \\ \top \rightarrow \perp \rightarrow \top \rightarrow \top \\ \perp \rightarrow \top \rightarrow \top \rightarrow \top \end{array}$$

- ▶ $A_{2,3}$ is the formula:

$$\forall x \forall y \forall z (x \neq 0 \rightarrow y \neq 0 \rightarrow z \neq 0 \rightarrow (x \wedge y \vee y \wedge z \vee z \wedge x) \neq 0)$$

