

Differential program semantics now with real bi-orthogonality pieces

Guillaume Geoffroy

DIAPASoN, Unibo

11 December 2019

λ -terms – syntax

$$t, u ::= x \mid \lambda x. t \mid tu$$

λ -terms – syntax

$$t, u ::= x \mid \lambda x. t \mid tu \\ \mid \langle t, u \rangle \mid \rho_L(t) \mid \rho_R(t)$$

λ -terms – syntax

$$\begin{aligned} t, u \quad ::= & \quad x \mid \lambda x. t \mid tu \\ & \mid \langle t, u \rangle \mid \rho_L(t) \mid \rho_R(t) \\ & \mid f(t_1, \dots, t_n) \quad (f : \mathbb{R}^n \rightarrow \mathbb{R}, n \in \mathbb{N}) \end{aligned}$$

λ -terms – reduction

► $(\lambda x. t)u \rightarrow_{\beta} t[x := u]$

λ -terms – reduction

- ▶ $(\lambda x. t)u \rightarrow_{\beta} t[x := u]$
- ▶ $f(t_1, \dots, t_j, g(u_1, \dots, u_k), v_1, \dots, v_l) \rightarrow_{\beta}$
 $(f \circ_j g)(t_1, \dots, t_j, u_1, \dots, u_k, v_1, \dots, v_l)$

λ -terms – reduction

- ▶ $(\lambda x. t)u \rightarrow_{\beta} t[x := u]$
- ▶ $f(t_1, \dots, t_j, g(u_1, \dots, u_k), v_1, \dots, v_l) \rightarrow_{\beta}$
 $(f \circ_j g)(t_1, \dots, t_j, u_1, \dots, u_k, v_1, \dots, v_l)$
 $\text{add}(1, \text{add}(2, 3))$

λ -terms – reduction

- ▶ $(\lambda x. t)u \rightarrow_{\beta} t[x := u]$
- ▶ $f(t_1, \dots, t_j, g(u_1, \dots, u_k), v_1, \dots, v_l) \rightarrow_{\beta}$
 $(f \circ_j g)(t_1, \dots, t_j, u_1, \dots, u_k, v_1, \dots, v_l)$
 $\text{add}(1, \text{add}(2, 3)) \rightarrow_{\beta} \text{add}_3(1, 2, 3)$

λ -terms – reduction

- ▶ $(\lambda x. t)u \rightarrow_{\beta} t[x := u]$
- ▶ $f(t_1, \dots, t_j, g(u_1, \dots, u_k), v_1, \dots, v_l) \rightarrow_{\beta}$
 $(f \circ_j g)(t_1, \dots, t_j, u_1, \dots, u_k, v_1, \dots, v_l)$
 $\text{add}(1, \text{add}(2, 3)) \rightarrow_{\beta} \text{add}_3(1, 2, 3) \rightarrow_{\beta}^3 6$

λ -terms – reduction

- ▶ $(\lambda x. t)u \rightarrow_{\beta} t[x := u]$
- ▶ $f(t_1, \dots, t_j, g(u_1, \dots, u_k), v_1, \dots, v_l) \rightarrow_{\beta}$
 $(f \circ_j g)(t_1, \dots, t_j, u_1, \dots, u_k, v_1, \dots, v_l)$
 $\text{add}(1, \text{add}(2, 3)) \rightarrow_{\beta} \text{add}_3(1, 2, 3) \rightarrow_{\beta}^3 6$
- ▶ $\rho_L(\langle t, u \rangle) \rightarrow_{\beta} t$
- ▶ $\rho_R(\langle t, u \rangle) \rightarrow_{\beta} u$

λ -terms – types

$$A, B ::= \mathbb{R}$$

λ -terms – types

$$\begin{array}{l} A, B ::= \mathbb{R} \\ \quad | \quad A \rightarrow B \end{array}$$

λ -terms – types

$$\begin{array}{lcl} A, B & ::= & \mathbb{R} \\ & | & A \rightarrow B \\ & | & A \times B \end{array}$$

λ -terms – types

$$\begin{array}{lcl} A, B & ::= & \mathbb{R} \\ & | & A \rightarrow B \\ & | & A \times B \end{array}$$

Expect: confluence + strong normalization

λ -terms – typing

$$\frac{\Gamma, x : A \vdash t : B}{\Gamma \vdash \lambda x. t : A \rightarrow B}$$

etc.

λ -terms – typing

$$\frac{\Gamma, x : A \vdash t : B}{\Gamma \vdash \lambda x. t : A \rightarrow B} \quad \text{etc.}$$

$$\frac{(f : \mathbb{R}^n \rightarrow \mathbb{R}) \quad \Gamma \vdash t_1 : \mathbb{R} \quad \dots \quad \Gamma \vdash t_n : \mathbb{R}}{\Gamma \vdash f(t_1, \dots, t_n) : \mathbb{R}}$$

λ -terms – typing

$$\frac{\Gamma, x : A \vdash t : B}{\Gamma \vdash \lambda x. t : A \rightarrow B} \quad \text{etc.}$$

$$\frac{(f : \mathbb{R}^n \rightarrow \mathbb{R}) \quad \Gamma \vdash t_1 : \mathbb{R} \quad \dots \quad \Gamma \vdash t_n : \mathbb{R}}{\Gamma \vdash f(t_1, \dots, t_n) : \mathbb{R}}$$

$$\Lambda_A := \{\text{closed terms of type } A\}$$

Stacks (*i.e.* tests) – syntax

$\pi ::= \quad / \text{ (closed interval)}$

Stacks (*i.e.* tests) – syntax

$$\pi ::= \quad I \text{ (closed interval)} \\ \quad \mid t \cdot \pi \text{ (} t \text{ closed term)}$$

Stacks (*i.e.* tests) – syntax

$$\pi ::= \begin{array}{l} I \text{ (closed interval)} \\ | \quad t \cdot \pi \text{ (} t \text{ closed term)} \\ | \quad L \cdot \pi \mid R \cdot \pi \end{array}$$

Stacks – typing

$$\frac{(I \text{ closed interval})}{\vdash I : \mathbb{R}}$$

Stacks – typing

$$\frac{(I \text{ closed interval})}{\vdash I : \mathbb{R}}$$

$$\frac{\vdash t : A \quad \vdash \pi : B}{\vdash t \cdot \pi \vdash A \rightarrow B}$$

Stacks – typing

$$\frac{(I \text{ closed interval})}{\vdash I : \mathbb{R}}$$

$$\frac{\vdash t : A \quad \vdash \pi : B}{\vdash t \cdot \pi \vdash A \rightarrow B}$$

$$\frac{\vdash \pi : A}{\vdash L \cdot \pi : A \times B}$$

Stacks – typing

$$\frac{(I \text{ closed interval})}{\vdash I : \mathbb{R}}$$

$$\frac{\vdash \pi : A}{\vdash L \cdot \pi : A \times B}$$

$$\frac{\vdash t : A \quad \vdash \pi : B}{\vdash t \cdot \pi \vdash A \rightarrow B}$$

$$\frac{\vdash \pi : B}{\vdash R \cdot \pi : A \times B}$$

Stacks – typing

$$\frac{(I \text{ closed interval})}{\vdash I : \mathbb{R}}$$

$$\frac{\vdash t : A \quad \vdash \pi : B}{\vdash t \cdot \pi \vdash A \rightarrow B}$$

$$\frac{\vdash \pi : A}{\vdash L \cdot \pi : A \times B}$$

$$\frac{\vdash \pi : B}{\vdash R \cdot \pi : A \times B}$$

$$\Pi_A := \{\text{stacks of type } A\}$$

Orthogonality : relation on $\Lambda_A \times \Pi_A$

► $t \perp\!\!\!\perp I$ iff $t \rightarrow_{\beta}^* r$ for some real $r \in I$

Orthogonality : relation on $\Lambda_A \times \Pi_A$

- ▶ $t \perp\!\!\!\perp I$ iff $t \rightarrow_{\beta}^* r$ for some real $r \in I$
- ▶ $t \perp\!\!\!\perp u \cdot \pi$ iff $tu \perp\!\!\!\perp \pi$

Orthogonality : relation on $\Lambda_A \times \Pi_A$

- ▶ $t \perp\!\!\!\perp I$ iff $t \rightarrow_{\beta}^* r$ for some real $r \in I$
- ▶ $t \perp\!\!\!\perp u \cdot \pi$ iff $tu \perp\!\!\!\perp \pi$
- ▶ $t \perp\!\!\!\perp L \cdot \pi$ iff $\rho_L(t) \perp\!\!\!\perp \pi$
- ▶ $t \perp\!\!\!\perp R \cdot \pi$ iff $\rho_R(t) \perp\!\!\!\perp \pi$

Orthogonality : relation on $\Lambda_A \times \Pi_A$

- ▶ $t \perp\!\!\!\perp I$ iff $t \rightarrow_{\beta}^* r$ for some real $r \in I$
- ▶ $t \perp\!\!\!\perp u \cdot \pi$ iff $tu \perp\!\!\!\perp \pi$
- ▶ $t \perp\!\!\!\perp L \cdot \pi$ iff $\rho_L(t) \perp\!\!\!\perp \pi$
- ▶ $t \perp\!\!\!\perp R \cdot \pi$ iff $\rho_R(t) \perp\!\!\!\perp \pi$

For $Y \subseteq \Pi_A$, $Y^{\perp\!\!\!\perp} := \{t \in \Lambda_A : \forall \pi \in Y \ t \perp\!\!\!\perp \pi\}$

Orthogonality : relation on $\Lambda_A \times \Pi_A$

- ▶ $t \perp\!\!\!\perp I$ iff $t \rightarrow_{\beta}^* r$ for some real $r \in I$
- ▶ $t \perp\!\!\!\perp u \cdot \pi$ iff $tu \perp\!\!\!\perp \pi$
- ▶ $t \perp\!\!\!\perp L \cdot \pi$ iff $\rho_L(t) \perp\!\!\!\perp \pi$
- ▶ $t \perp\!\!\!\perp R \cdot \pi$ iff $\rho_R(t) \perp\!\!\!\perp \pi$

For $Y \subseteq \Pi_A$, $Y^{\perp\!\!\!\perp} := \{t \in \Lambda_A : \forall \pi \in Y \ t \perp\!\!\!\perp \pi\}$

For $X \subseteq \Lambda_A$, $X^{\perp\!\!\!\perp} := \{\pi \in \Pi_A : \forall t \in X \ t \perp\!\!\!\perp \pi\}$

Approximate programs

$$\llbracket A \rrbracket := \{X \subseteq \Lambda_A; X^{\perp\perp} = X\}$$

Approximate programs

$$\begin{aligned} \llbracket A \rrbracket &:= \{X \subseteq \Lambda_A; X^{\perp\perp} = X\} \\ &= \{Y^{\perp}; Y \subseteq \Pi_A\} \end{aligned}$$

Approximate programs

$$\begin{aligned} \llbracket A \rrbracket &:= \{X \subseteq \Lambda_A; X^{\perp\perp} = X\} \\ &= \{Y^{\perp}; Y \subseteq \Pi_A\} \end{aligned} \quad \text{Complete lattice}$$

Approximate programs

$$\begin{aligned} \llbracket A \rrbracket &:= \{X \subseteq \Lambda_A; X^{\perp\perp} = X\} \\ &= \{Y^{\perp}; Y \subseteq \Pi_A\} \end{aligned} \quad \text{Complete lattice}$$

Example (Approximate reals)

- For all I , let $|I| := \{I\}^{\perp\perp}$

Approximate programs

$$\begin{aligned} \llbracket A \rrbracket &:= \{X \subseteq \Lambda_A; X^{\perp\perp} = X\} \\ &= \{Y^{\perp}; Y \subseteq \Pi_A\} \end{aligned} \quad \text{Complete lattice}$$

Example (Approximate reals)

- ▶ For all I , let $|I| := \{I\}^{\perp\perp}$
- ▶ Then $\{I_k; k \in K\}^{\perp\perp} = |\bigcap_{k \in K} I_k|$

Approximate programs

$$\begin{aligned} \llbracket A \rrbracket &:= \{X \subseteq \Lambda_A; X^{\perp\perp} = X\} \\ &= \{Y^{\perp}; Y \subseteq \Pi_A\} \end{aligned} \quad \text{Complete lattice}$$

Example (Approximate reals)

- ▶ For all I , let $|I| := \{I\}^{\perp\perp}$
- ▶ Then $\{I_k; k \in K\}^{\perp\perp} = |\bigcap_{k \in K} I_k|$
- ▶ So $\llbracket \mathbb{R} \rrbracket = \{|I|; I \text{ closed interval}\}$

Approximate programs – examples

$$\llbracket A \rrbracket^* := \llbracket A \rrbracket \setminus \{\emptyset\}$$

Approximate programs – examples

$$\llbracket A \rrbracket^* := \llbracket A \rrbracket \setminus \{\emptyset\}$$

Example (Approximate functions)

- For all $F : \mathbb{R} \times \dots \times \mathbb{R} \rightarrow \{I \neq \emptyset\}$, let
$$|F| := \{r_1 \cdot \dots \cdot r_n \cdot F(r_1, \dots, r_n); r_1, \dots, r_n \in \mathbb{R}\}^{\perp}$$


Approximate programs – examples

$$\llbracket A \rrbracket^* := \llbracket A \rrbracket \setminus \{\emptyset\}$$


Example (Approximate functions)

- ▶ For all $F : \mathbb{R} \times \dots \times \mathbb{R} \rightarrow \{I \neq \emptyset\}$, let
$$|F| := \{r_1 \cdot \dots \cdot r_n \cdot F(r_1, \dots, r_n); r_1, \dots, r_n \in \mathbb{R}\}^{\perp\!\!\!\perp}$$
- ▶ Then $\llbracket \mathbb{R} \rightarrow \dots \rightarrow \mathbb{R} \rightarrow \mathbb{R} \rrbracket^* = \{|F|; F \dots\}$


Approximate programs – examples



Approximate programs – examples



Approximate programs – examples



Approximate programs – examples

Example (Approximate pairs)

- For all $a \in \llbracket A \rrbracket$, $b \in \llbracket B \rrbracket$, let
$$|a \times b| := \left\{ t; t \xrightarrow[\beta]^* \langle u, v \rangle, u \in a, v \in b \right\}$$

Approximate programs – examples

Example (Approximate pairs)

- ▶ For all $a \in \llbracket A \rrbracket$, $b \in \llbracket B \rrbracket$, let
$$|a \times b| := \{t; t \rightarrow_{\beta}^* \langle u, v \rangle, u \in a, v \in b\}$$
- ▶ Then $\llbracket A \times B \rrbracket^* = \{|a \times b|; a \in \llbracket A \rrbracket^*, b \in \llbracket B \rrbracket^*\}$

Substitution

$$\blacktriangleright t[x_1 : A_1, \dots, x_n : A_n] : B$$

Substitution

- ▶ $t[x_1 : A_1, \dots, x_n : A_n] : B$
- ▶ $a_1 \in \llbracket A_1 \rrbracket, \dots, a_n \in \llbracket A_n \rrbracket$

Substitution

► $t[x_1 : A_1, \dots, x_n : A_n] : B$

► $a_1 \in \llbracket A_1 \rrbracket, \dots, a_n \in \llbracket A_n \rrbracket$

► Then let

$$\begin{aligned} & t[x_1 := a_1, \dots, x_n := a_n] \\ & := \left\{ t[x_1 := u_1, \dots, x_n := u_n]; \right. \\ & \quad \left. u_1 \in a_1, \dots, u_n \in a_n \right\}^{\perp\perp\perp} \in \llbracket B \rrbracket \end{aligned}$$

Substitution – sub-functoriality

$$\blacktriangleright t[y_1 : B_1, \dots, y_n : B_n] : C$$

Substitution – sub-functoriality

- ▶ $t[y_1 : B_1, \dots, y_n : B_n] : C$
- ▶ $u_1[x_1 : A_1, \dots, x_m : A_m] : B_1, \dots, u_n[x_1 : A_1, \dots, x_m : A_m] : B_n$

Substitution – sub-functoriality

- ▶ $t[y_1 : B_1, \dots, y_n : B_n] : C$
- ▶ $u_1[x_1 : A_1, \dots, x_m : A_m] : B_1, \dots, u_n[x_1 : A_1, \dots, x_m : A_m] : B_n$
- ▶ $a_1 \in \llbracket A_1 \rrbracket, \dots, a_m \in \llbracket A_m \rrbracket$

Substitution – sub-functoriality

- ▶ $t[y_1 : B_1, \dots, y_n : B_n] : C$
- ▶ $u_1[x_1 : A_1, \dots, x_m : A_m] : B_1, \dots, u_n[x_1 : A_1, \dots, x_m : A_m] : B_n$
- ▶ $a_1 \in \llbracket A_1 \rrbracket, \dots, a_m \in \llbracket A_m \rrbracket$
- ▶ $\rightsquigarrow \subseteq \frac{t[u_1, \dots, u_n][a_1, \dots, a_m]}{t[u_1[a_1, \dots, a_m], \dots, u_n[a_1, \dots, a_m]]}$

Distances – distance spaces

- ▶ $(\mathbb{R}) := \mathbb{R}_+^\infty$
- ▶ $(A \times B) := (A) \times (B)$
- ▶ $(A \rightarrow B) := \llbracket A \rrbracket \rightarrow (B)$

Distances – diameter function

$\delta_A : \llbracket A \rrbracket \rightarrow \langle A \rangle :$

- ▶ $\delta_{\mathbb{R}}(|I|) := \text{length}(I)$
- ▶ $\delta_{A \times B}(p) := (\delta_A(\rho_L(p)), \delta_B(\rho_R(p)))$
- ▶ $\delta_{A \rightarrow B}(f)(a) := \delta_B(fa)$

Distances – diameter function

$$\delta_A : \llbracket A \rrbracket \rightarrow \langle A \rangle:$$

$$\blacktriangleright \delta_{\mathbb{R}}(|I|) := \text{length}(I)$$

$$\blacktriangleright \delta_{A \times B}(p) := (\delta_A(\rho_L(p)), \delta_B(\rho_R(p)))$$

$$\blacktriangleright \delta_{A \rightarrow B}(f)(a) := \delta_B(fa)$$

$$d_A(a, b) := \delta_A(a \vee b)$$

Distances – sub-modularity

Proposition

If $a \wedge b \neq \emptyset$ then

$$\delta_A(a \vee b) + \delta_A(a \wedge b) \leq \delta_A(a) + \delta_A(b)$$

Distances – sub-modularity

Proposition

If $a \wedge b \neq \emptyset$ then

$$\delta_A(a \vee b) + \delta_A(a \wedge b) \leq \delta_A(a) + \delta_A(b)$$

Corollary

For all $a, b, c \in \llbracket A \rrbracket^$,*

$$\delta_A(a \vee c) + \delta_A(b) \leq \delta_A(a \vee b) + \delta_A(b \vee c)$$

Distances – sub-modularity

Proposition

If $a \wedge b \neq \emptyset$ then

$$\delta_A(a \vee b) + \delta_A(a \wedge b) \leq \delta_A(a) + \delta_A(b)$$

Corollary

For all $a, b, c \in \llbracket A \rrbracket^$,*

$$d_A(a, c) + d_A(b, b) \leq d_A(a, b) + d_A(b, c)$$

Distances – partial metric ?

Proposition?

$\llbracket A \rrbracket^*$ is a partial metric space:

- ▶ $d_A(a, a) \leq d_A(a, b)$
- ▶ $d_A(a, b) = d_A(b, a)$
- ▶ $d_A(a, c) + d_A(b, b) \leq d_A(a, b) + d_A(b, c)$
- ▶ if $d_A(a, a) = d_A(a, b) = d_A(b, b)$, then $a = b$

Distances – partial metric ?

Proposition?

$\llbracket A \rrbracket^*$ is a partial metric space:

- ▶ $d_A(a, a) \leq d_A(a, b)$
- ▶ $d_A(a, b) = d_A(b, a)$
- ▶ $d_A(a, c) + d_A(b, b) \leq d_A(a, b) + d_A(b, c)$
- ▶ ~~if $d_A(a, a) = d_A(a, b) = d_A(b, b)$, then $a = b$~~

Distances – locally finite distances

► $(\mathbb{R})_f := \mathbb{R}_+$

Distances – locally finite distances

► $(\mathbb{R})_f := \mathbb{R}_+$

► $(A \times B)_f := (A)_f \times (B)_f$

Distances – locally finite distances

- ▶ $(\mathbb{R})_f := \mathbb{R}_+$
- ▶ $(A \times B)_f := (A)_f \times (B)_f$
- ▶ $(A \rightarrow B)_f :=$
 $\{\varepsilon : \llbracket A \rrbracket \rightarrow (B); \forall u : A, \varepsilon(\{u\}) \in (B)_f\}$

Distances – locally finite distances

- ▶ $(\mathbb{R})_f := \mathbb{R}_+$
- ▶ $(A \times B)_f := (A)_f \times (B)_f$
- ▶ $(A \rightarrow B)_f :=$
 $\{\varepsilon : \llbracket A \rrbracket \rightarrow (B); \forall u : A, \varepsilon(\{u\}) \in (B)_f\}$

$$\llbracket A \rrbracket_f = \{a \in \llbracket A \rrbracket; \delta_A(a) \in (A)_f\}$$

Distances – partial metric

Proposition

$\llbracket A \rrbracket^*$ is almost a partial metric space:

- ▶ $d_A(a, a) \leq d_A(a, b)$
- ▶ $d_A(a, b) = d_A(b, a)$
- ▶ $d_A(a, c) + d_A(b, b) \leq d_A(a, b) + d_A(b, c)$
- ▶ if $d_A(a, a) = d_A(a, b) = d_A(b, b) \in \llbracket A \rrbracket_f$, then $a = b$