## CrossMark

## ERRATUM

## Erratum to: Housing price prediction: parametric versus semi-parametric spatial hedonic models

José-María Montero<sup>1</sup> · Román Mínguez<sup>2</sup> · Gema Fernández-Avilés<sup>1</sup>

Published online: 19 September 2017 © Springer-Verlag GmbH Germany 2017

Erratum to: J Geogr Syst DOI 10.1007/s10109-017-0257-y

In the original publication, in Table 1, there is a misprint in equations under models SDEM and SEM. The specification for the spatially correlated error term,  $\mathbf{u}$ , must be  $\mathbf{u} = \lambda \mathbf{W} \mathbf{u} + \boldsymbol{\epsilon}$ . The correct versions of these equations are given below. The original version of the article has been updated (Table 1).

Department of Statistics, Faculty of Social Sciences, University of Castilla-La Mancha, 16071 Cuenca, Spain



The online version of the original article can be found under doi:10.1007/s10109-017-0257-y.

<sup>☑</sup> José-María Montero jose.mlorenzo@uclm.es

Department of Statistics, Faculty of Law and Social Sciences, University of Castilla-La Mancha, 45071 Toledo, Spain

J.-M. Montero et al.

Table 1 Competing hedonic house price models: parametric and semi-parametric model specifications

Model	Specification	Spatial lag		Spatial	
		Response	Covariates	Error	or covariates
Parametric models					
(i) Conventional a-	(i) Conventional a-spatial hedonic model				
HM	$\mathbf{y} = \alpha \mathbf{\dot{n}}_n + \mathbf{X} \mathbf{\beta} + \mathbf{\epsilon}$				
(iia) SAR models					
SLM	$\mathbf{y} = \rho \mathbf{W} \mathbf{y} + \alpha \mathbf{i}_n + \mathbf{X} \mathbf{\beta} + \mathbf{\epsilon}$	×			
SDM	$\mathbf{y} = \rho \mathbf{W} \mathbf{y} + \alpha \mathbf{i}_n + \mathbf{X} \mathbf{\beta} + \mathbf{W} \mathbf{X} \mathbf{\theta} + \mathbf{\epsilon}$	X	×		
SDEM	$\mathbf{y} = \alpha \mathbf{i}_n + \mathbf{X}\mathbf{\beta} + \mathbf{W}\mathbf{X}\mathbf{\theta} + \mathbf{u},  \mathbf{u} = \lambda \mathbf{W}\mathbf{u} + \mathbf{\epsilon}$		×	×	
(iib) SEM model					
SEM	$\mathbf{y} = \alpha \mathbf{i}_n + \mathbf{X} \mathbf{\beta} + \mathbf{u},  \mathbf{u} = \lambda \mathbf{W} \mathbf{u} + \mathbf{\epsilon}$			×	
Semi-parametric models	odels				
(iii) PSSD-HM and	(iii) PSSD-HM and PSSD-SAR models				
PSSD-HM	$\mathbf{y} = f(s_1, s_2) + \mathbf{X}\mathbf{\beta} + \mathbf{\epsilon}$			×	
PSSD-SLM	$\mathbf{y} = f(s_1, s_2) + \rho \mathbf{W} \mathbf{y} + \mathbf{X} \mathbf{\beta} + \mathbf{\epsilon}$	×		×	
PSSD-SDM	$\mathbf{y} = f(s_1, s_2) + \rho \mathbf{W} \mathbf{y} + \mathbf{X} \mathbf{\beta} + \mathbf{W} \mathbf{X} \mathbf{\theta} + \mathbf{\epsilon}$	×	×	×	
(iv) GAM model					
GAM	$\mathbf{y} = \sum_{r=1}^r f_{1,r}(x_r^+) + lpha \mathbf{i}_n + \mathbf{X}^* \mathbf{\beta} + \mathbf{\epsilon}$				×
(v) GAM-SAR models	dels				
GAM-SLM	$\mathbf{y} = \sum_{r=1}^{l} f_{1,r}(x_r^+) + \alpha \mathbf{i}_n + \rho \mathbf{W} \mathbf{y} + \mathbf{X}^* \mathbf{\beta} + \mathbf{\epsilon}$	×			×
GAM-SDM	$\mathbf{y} = \sum_{r=1}^{l} f_{1,r}(\mathbf{x}_{r}^{+}) + \sum_{r=1}^{l} f_{2,r}(\mathbf{W}\mathbf{x}_{r}^{+}) + \alpha \mathbf{i}_{n} + \rho \mathbf{W}\mathbf{y} + \mathbf{X}^{*} \mathbf{\beta} + \mathbf{W}\mathbf{X}^{*} \mathbf{\theta} + \mathbf{\epsilon}$	×	×		×



Table 1 continued						
Model	Specification	Spatial lag			Spatial	Spatial Nonparametric functions
		Response Covariates Error	Covariates	Error	arin	or covariates
(vi) PSSD-GAM model PSSD-GAM	(e) $\mathbf{y}=f(s_1,s_2)+\sum_{r=1}^l f_{l,r}(x_r^+)+\mathbf{X}^*\mathbf{\beta}+\mathbf{\epsilon}$				×	×
(vii) PSSD-GAM-SAR models	models					
PSSD-GAM-SLM	PSSD-GAM-SLM $\mathbf{y} = f(s_1, s_2) + \sum_{r=1} f_{1,r}(x_r^+) + \rho \mathbf{W} \mathbf{y} + \mathbf{X}^* \mathbf{\beta} + \mathbf{\epsilon}$	×			×	×
PSSD-GAM-SDM	PSSD-GAM-SDM $\mathbf{y} = f(s_1, s_2) + \sum_{r=1}^{l} f_{1,r}(x_r^+) + \sum_{r=1}^{l} f_{2,r}(\mathbf{W} x_r^+) + \rho \mathbf{W} \mathbf{y} + \mathbf{X}^* \mathbf{\beta} + \mathbf{W} \mathbf{X}^* 0 + \mathbf{\epsilon}  \mathbf{x}$		×		×	×

