

Higgs bundles twisted by a vector bundle

Guillermo Gallego

Universidad Complutense de Madrid

26 September 2019

Holomorphic vector bundles over Riemann surfaces

Let X be a compact Riemann surface of genus g .

Holomorphic vector bundles over Riemann surfaces

Let X be a compact Riemann surface of genus g .

- Smooth classification: rank and degree (1st Chern class).

Holomorphic vector bundles over Riemann surfaces

Let X be a compact Riemann surface of genus g .

- Smooth classification: rank and degree (1st Chern class).
- Line bundles: Picard group, $\text{Pic}(X) = H^1(X, \mathcal{O}_X^*)$.

Holomorphic vector bundles over Riemann surfaces

Let X be a compact Riemann surface of genus g .

- Smooth classification: rank and degree (1st Chern class).
- Line bundles: Picard group, $\text{Pic}(X) = H^1(X, \mathcal{O}_X^*)$.
Components of fixed degree $\text{Pic}^d(X)$ are isomorphic to the **Jacobian**

$$J(X) = \frac{H^1(X, \mathcal{O}_X)}{H^1(X, \mathbb{Z})}.$$

Holomorphic vector bundles over Riemann surfaces

Let X be a compact Riemann surface of genus g .

- Smooth classification: rank and degree (1st Chern class).
- Line bundles: Picard group, $\text{Pic}(X) = H^1(X, \mathcal{O}_X^*)$.
Components of fixed degree $\text{Pic}^d(X)$ are isomorphic to the **Jacobian**

$$J(X) = \frac{H^1(X, \mathcal{O}_X)}{H^1(X, \mathbb{Z})}.$$

- $g = 0$, Grothendieck (1957).

Holomorphic vector bundles over Riemann surfaces

Let X be a compact Riemann surface of genus g .

- Smooth classification: rank and degree (1st Chern class).
- Line bundles: Picard group, $\text{Pic}(X) = H^1(X, \mathcal{O}_X^*)$.
Components of fixed degree $\text{Pic}^d(X)$ are isomorphic to the **Jacobian**

$$J(X) = \frac{H^1(X, \mathcal{O}_X)}{H^1(X, \mathbb{Z})}.$$

- $g = 0$, Grothendieck (1957).
- $g = 1$, Atiyah (1957).

Holomorphic vector bundles over Riemann surfaces

Let X be a compact Riemann surface of genus g .

- Smooth classification: rank and degree (1st Chern class).
- Line bundles: Picard group, $\text{Pic}(X) = H^1(X, \mathcal{O}_X^*)$.
Components of fixed degree $\text{Pic}^d(X)$ are isomorphic to the **Jacobian**

$$J(X) = \frac{H^1(X, \mathcal{O}_X)}{H^1(X, \mathbb{Z})}.$$

- $g = 0$, Grothendieck (1957).
- $g = 1$, Atiyah (1957).
- $g \geq 2$?

Holomorphic vector bundles over Riemann surfaces

Let X be a compact Riemann surface of genus g .

- Smooth classification: rank and degree (1st Chern class).
- Line bundles: Picard group, $\text{Pic}(X) = H^1(X, \mathcal{O}_X^*)$.
Components of fixed degree $\text{Pic}^d(X)$ are isomorphic to the **Jacobian**

$$J(X) = \frac{H^1(X, \mathcal{O}_X)}{H^1(X, \mathbb{Z})}.$$

- $g = 0$, Grothendieck (1957).
- $g = 1$, Atiyah (1957).
- $g \geq 2$?

For now on, assume $g \geq 2$.

Definition

A holomorphic vector bundle $E \rightarrow X$ is **stable** if for every proper holomorphic subbundle $E' \subset E$,

$$\frac{\deg E'}{\operatorname{rk} E'} < \frac{\deg E}{\operatorname{rk} E}.$$

Definition

A holomorphic vector bundle $E \rightarrow X$ is **stable** if for every proper holomorphic subbundle $E' \subset E$,

$$\frac{\deg E'}{\operatorname{rk} E'} < \frac{\deg E}{\operatorname{rk} E}.$$

- Notion of stability arising from Mumford's GIT.

Definition

A holomorphic vector bundle $E \rightarrow X$ is **stable** if for every proper holomorphic subbundle $E' \subset E$,

$$\frac{\deg E'}{\operatorname{rk} E'} < \frac{\deg E}{\operatorname{rk} E}.$$

- Notion of stability arising from Mumford's GIT.
- **Moduli space of stable vector bundles:** $\mathcal{N}(n, d)$

Definition

Let \mathbb{E} be a smooth vector bundle of rank n and degree d over X .

Definition

Let \mathbb{E} be a smooth vector bundle of rank n and degree d over X . A **holomorphic structure on \mathbb{E}**

Definition

Let \mathbb{E} be a smooth vector bundle of rank n and degree d over X . A **holomorphic structure on \mathbb{E}** is an operator

$$\bar{\partial}_E : \Omega^{p,q}(X, \mathbb{E}) \longrightarrow \Omega^{p,q+1}(X, \mathbb{E})$$

Definition

Let \mathbb{E} be a smooth vector bundle of rank n and degree d over X . A **holomorphic structure on \mathbb{E}** is an operator

$$\bar{\partial}_E : \Omega^{p,q}(X, \mathbb{E}) \longrightarrow \Omega^{p,q+1}(X, \mathbb{E})$$

which satisfies

$$\bar{\partial}_E(\alpha\psi) = (\bar{\partial}\alpha)\psi + (-1)^p\alpha \wedge \bar{\partial}_E\psi.$$

Definition

Let \mathbb{E} be a smooth vector bundle of rank n and degree d over X . A **holomorphic structure on \mathbb{E}** is an operator

$$\bar{\partial}_E : \Omega^{p,q}(X, \mathbb{E}) \longrightarrow \Omega^{p,q+1}(X, \mathbb{E})$$

which satisfies

$$\bar{\partial}_E(\alpha\psi) = (\bar{\partial}\alpha)\psi + (-1)^p\alpha \wedge \bar{\partial}_E\psi.$$

- $\bar{\partial}_E^2 = 0$ is the integrability condition for the PDE $\bar{\partial}_E s = 0$.

Definition

Let \mathbb{E} be a smooth vector bundle of rank n and degree d over X . A **holomorphic structure on \mathbb{E}** is an operator

$$\bar{\partial}_E : \Omega^{p,q}(X, \mathbb{E}) \longrightarrow \Omega^{p,q+1}(X, \mathbb{E})$$

which satisfies

$$\bar{\partial}_E(\alpha\psi) = (\bar{\partial}\alpha)\psi + (-1)^p\alpha \wedge \bar{\partial}_E\psi.$$

- $\bar{\partial}_E^2 = 0$ is the integrability condition for the PDE $\bar{\partial}_E s = 0$.
- The solutions of $\bar{\partial}_E s = 0$ are holomorphic vector functions (locally free sheaf over \mathcal{O}_X).

Definition

Let \mathbb{E} be a smooth vector bundle of rank n and degree d over X . A **holomorphic structure on \mathbb{E}** is an operator

$$\bar{\partial}_E : \Omega^{p,q}(X, \mathbb{E}) \longrightarrow \Omega^{p,q+1}(X, \mathbb{E})$$

which satisfies

$$\bar{\partial}_E(\alpha\psi) = (\bar{\partial}\alpha)\psi + (-1)^p\alpha \wedge \bar{\partial}_E\psi.$$

- $\bar{\partial}_E^2 = 0$ is the integrability condition for the PDE $\bar{\partial}_E s = 0$.
- The solutions of $\bar{\partial}_E s = 0$ are holomorphic vector functions (locally free sheaf over \mathcal{O}_X).
- $E = (\mathbb{E}, \bar{\partial}_E)$ is a holomorphic vector bundle.

The gauge-theoretic point of view

(\mathbb{E}, h) smooth complex Hermitian vector bundle of rank n and degree d over X .

The gauge-theoretic point of view

(\mathbb{E}, h) smooth complex Hermitian vector bundle of rank n and degree d over X .

h -unitary connections \longleftrightarrow Holomorphic structures on \mathbb{E}

$$\nabla \longmapsto \nabla^{0,1}$$

The gauge-theoretic point of view

(\mathbb{E}, h) smooth complex Hermitian vector bundle of rank n and degree d over X .

h -unitary connections \longleftrightarrow Holomorphic structures on \mathbb{E}

$$\nabla \longmapsto \nabla^{0,1}$$

- \mathcal{A} set of irreducible h -unitary connections on (\mathbb{E}, h) .

The gauge-theoretic point of view

(\mathbb{E}, h) smooth complex Hermitian vector bundle of rank n and degree d over X .

h -unitary connections \longleftrightarrow Holomorphic structures on \mathbb{E}

$$\nabla \longmapsto \nabla^{0,1}$$

- \mathcal{A} set of irreducible h -unitary connections on (\mathbb{E}, h) .
- \mathcal{G} group of unitary gauge transformations of \mathbb{E} .

The gauge-theoretic point of view

(\mathbb{E}, h) smooth complex Hermitian vector bundle of rank n and degree d over X .

h -unitary connections \longleftrightarrow Holomorphic structures on \mathbb{E}

$$\nabla \longmapsto \nabla^{0,1}$$

- \mathcal{A} set of irreducible h -unitary connections on (\mathbb{E}, h) .
- \mathcal{G} group of unitary gauge transformations of \mathbb{E} .
- \mathcal{G} acts on \mathcal{A} by conjugation with momentum map

$$\mu : \mathcal{A} \longrightarrow \text{Lie}(\mathcal{G})$$

$$\nabla \longmapsto F_{\nabla} - 2\pi i \frac{d}{n} \mathbf{1}_{\mathbb{E}} \omega_X.$$

The gauge-theoretic point of view

(\mathbb{E}, h) smooth complex Hermitian vector bundle of rank n and degree d over X .

h -unitary connections \longleftrightarrow Holomorphic structures on \mathbb{E}

$$\nabla \longmapsto \nabla^{0,1}$$

- \mathcal{A} set of irreducible h -unitary connections on (\mathbb{E}, h) .
- \mathcal{G} group of unitary gauge transformations of \mathbb{E} .
- \mathcal{G} acts on \mathcal{A} by conjugation with momentum map

$$\mu : \mathcal{A} \longrightarrow \text{Lie}(\mathcal{G})$$

$$\nabla \longmapsto F_{\nabla} - 2\pi i \frac{d}{n} \mathbf{1}_{\mathbb{E}} \omega_X.$$

- Symplectic quotient: $\mathcal{A} // \mathcal{G} = \mu^{-1}(0) / \mathcal{G}$.

The gauge-theoretic point of view

Theorem (Donaldson's version of Narasimhan–Seshadri)

An irreducible h -unitary connection ∇ satisfies

$$F_{\nabla} = -2\pi i \frac{d}{n} \mathbf{1}_{\mathbb{E}} \omega_X$$

if and only if $(\mathbb{E}, \nabla^{0,1})$ is stable.

The gauge-theoretic point of view

Theorem (Donaldson's version of Narasimhan–Seshadri)

An irreducible h -unitary connection ∇ satisfies

$$F_{\nabla} = -2\pi i \frac{d}{n} \mathbf{1}_{\mathbb{E}} \omega_X$$

if and only if $(\mathbb{E}, \nabla^{0,1})$ is stable.

- Consequence:

$$\mathcal{N}(n, d) \longleftrightarrow \mathcal{A} // \mathcal{G}.$$

The Hitchin system

- $T_{[E]}^* \mathcal{N}(n, d) \cong H^0(X, \text{End} E \otimes K_X).$

The Hitchin system

- $T_{[E]}^*\mathcal{N}(n, d) \cong H^0(X, \text{End}E \otimes K_X)$.
- $(E, \varphi) \in T^*\mathcal{N}(n, d)$, $\varphi : E \rightarrow E \otimes K_X$, characteristic polynomial:

$$P_\varphi(T) = \det(T - \varphi) = T^n + \sum_{i=1}^n \sigma_i(\varphi) T^{n-i}.$$

The Hitchin system

- $T_{[E]}^*\mathcal{N}(n, d) \cong H^0(X, \text{End}E \otimes K_X)$.
- $(E, \varphi) \in T^*\mathcal{N}(n, d)$, $\varphi : E \rightarrow E \otimes K_X$, characteristic polynomial:

$$P_\varphi(T) = \det(T - \varphi) = T^n + \sum_{i=1}^n \sigma_i(\varphi) T^{n-i}.$$

- **Hitchin fibration:**

$$\begin{aligned} H : T^*\mathcal{N}(n, d) &\longrightarrow \mathcal{B} := \bigoplus_{i=1}^n H^0(X, K_X^i) \\ (E, \varphi) &\longmapsto (\sigma_1(\varphi), \dots, \sigma_n(\varphi)) \end{aligned}$$

The Hitchin system

- $T_{[E]}^*\mathcal{N}(n, d) \cong H^0(X, \text{End}E \otimes K_X)$.
- $(E, \varphi) \in T^*\mathcal{N}(n, d)$, $\varphi : E \rightarrow E \otimes K_X$, characteristic polynomial:

$$P_\varphi(T) = \det(T - \varphi) = T^n + \sum_{i=1}^n \sigma_i(\varphi) T^{n-i}.$$

- **Hitchin fibration:**

$$\begin{aligned} H : T^*\mathcal{N}(n, d) &\longrightarrow \mathcal{B} := \bigoplus_{i=1}^n H^0(X, K_X^i) \\ (E, \varphi) &\longmapsto (\sigma_1(\varphi), \dots, \sigma_n(\varphi)) \end{aligned}$$

- **Fibres?**

The spectral correspondence

$\varphi : \mathbb{C}^n \rightarrow \mathbb{C}^n$ endomorphism

The spectral correspondence

$\varphi : \mathbb{C}^n \rightarrow \mathbb{C}^n$ endomorphism :

- $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ eigenvalues of φ ,

The spectral correspondence


$\varphi : \mathbb{C}^n \rightarrow \mathbb{C}^n$ endomorphism :

- $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ eigenvalues of φ ,
- $v_1, \dots, v_n \in \mathbb{C}^n$ eigenvectors of φ .

The spectral correspondence

$\varphi : \mathbb{C}^n \rightarrow \mathbb{C}^n$ endomorphism :

- $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ eigenvalues of φ ,
- $v_1, \dots, v_n \in \mathbb{C}^n$ eigenvectors of φ .



The spectral correspondence

$E \rightarrow X$ holomorphic vector bundle.

The spectral correspondence

$E \rightarrow X$ holomorphic vector bundle.

$\varphi : E \rightarrow E$ vector bundle

endomorphism

The spectral correspondence

$E \rightarrow X$ holomorphic vector bundle.

$\varphi : E \rightarrow E$ vector bundle

endomorphism :

- $\lambda_1(x), \dots, \lambda_n(x) \in \mathbb{C}$
eigenvalues of $\varphi_x : E_x \rightarrow E_x$,

The spectral correspondence

$E \rightarrow X$ holomorphic vector bundle.

$\varphi : E \rightarrow E$ vector bundle

endomorphism :

- $\lambda_1(x), \dots, \lambda_n(x) \in \mathbb{C}$
eigenvalues of $\varphi_x : E_x \rightarrow E_x$,
- $v_1(x), \dots, v_n(x) \in \mathbb{C}^n$
eigenvectors of $\varphi_x : E_x \rightarrow E_x$.

The spectral correspondence

$E \rightarrow X$ holomorphic vector bundle.

$\varphi : E \rightarrow E$ vector bundle

endomorphism :

- $\lambda_1(x), \dots, \lambda_n(x) \in \mathbb{C}$
eigenvalues of $\varphi_x : E_x \rightarrow E_x$,
- $v_1(x), \dots, v_n(x) \in \mathbb{C}^n$
eigenvectors of $\varphi_x : E_x \rightarrow E_x$.
- **Spectral curve:**

$$S_\varphi = \{(x, \lambda_i(x)), i = 1, \dots, n\} \subset X \times \mathbb{C}.$$

The spectral correspondence

$E \rightarrow X$ holomorphic vector bundle.

$\varphi : E \rightarrow E$ vector bundle

endomorphism :

- $\lambda_1(x), \dots, \lambda_n(x) \in \mathbb{C}$
eigenvalues of $\varphi_x : E_x \rightarrow E_x$,
- $v_1(x), \dots, v_n(x) \in \mathbb{C}^n$
eigenvectors of $\varphi_x : E_x \rightarrow E_x$.
- **Spectral curve:**

$$S_\varphi = \{(x, \lambda_i(x)), i = 1, \dots, n\} \subset X \times \mathbb{C}.$$

- $S_\varphi \xrightarrow{\pi} X$ ramified cover.

The spectral correspondence

$E \rightarrow X$ holomorphic vector bundle.

$\varphi : E \rightarrow E$ vector bundle


endomorphism :

- $\lambda_1(x), \dots, \lambda_n(x) \in \mathbb{C}$
eigenvalues of $\varphi_x : E_x \rightarrow E_x$,
- $v_1(x), \dots, v_n(x) \in \mathbb{C}^n$
eigenvectors of $\varphi_x : E_x \rightarrow E_x$.

- **Spectral curve:**

$$S_\varphi = \{(x, \lambda_i(x)), i = 1, \dots, n\} \subset X \times \mathbb{C}.$$

- $S_\varphi \xrightarrow{\pi} X$ ramified cover.



The spectral correspondence

$E \rightarrow X$ holomorphic vector bundle.

$\varphi : E \rightarrow E$ vector bundle


endomorphism :

- $\lambda_1(x), \dots, \lambda_n(x) \in \mathbb{C}$
eigenvalues of $\varphi_x : E_x \rightarrow E_x$,
- $v_1(x), \dots, v_n(x) \in \mathbb{C}^n$
eigenvectors of $\varphi_x : E_x \rightarrow E_x$.

- **Spectral curve:**

$$S_\varphi = \{(x, \lambda_i(x)), i = 1, \dots, n\} \subset X \times \mathbb{C}.$$

- $S_\varphi \xrightarrow{\pi} X$ ramified cover.



The spectral correspondence

$E \rightarrow X$ holomorphic vector bundle.

$\varphi : E \rightarrow E \otimes L$ **twisted** vector
bundle endomorphism

The spectral correspondence

$E \rightarrow X$ holomorphic vector bundle.

$\varphi : E \rightarrow E \otimes L$ **twisted** vector
bundle endomorphism :

- $\lambda_1(x), \dots, \lambda_n(x) \in L_x$
eigenvalues of
 $\varphi_x : E_x \rightarrow E_x \otimes L_x,$

The spectral correspondence

$E \rightarrow X$ holomorphic vector bundle.

$\varphi : E \rightarrow E \otimes L$ **twisted** vector
bundle endomorphism :

- $\lambda_1(x), \dots, \lambda_n(x) \in L_x$
eigenvalues of
 $\varphi_x : E_x \rightarrow E_x \otimes L_x,$
- $v_1(x), \dots, v_n(x) \in E_x$
eigenvectors of
 $\varphi_x : E_x \rightarrow E_x \otimes L_x.$

The spectral correspondence

$E \rightarrow X$ holomorphic vector bundle.

$\varphi : E \rightarrow E \otimes L$ **twisted** vector
bundle endomorphism :

- $\lambda_1(x), \dots, \lambda_n(x) \in L_x$

eigenvalues of

$$\varphi_x : E_x \rightarrow E_x \otimes L_x,$$

- $v_1(x), \dots, v_n(x) \in E_x$

eigenvectors of

$$\varphi_x : E_x \rightarrow E_x \otimes L_x.$$

- **Spectral curve** (locally):

$$S_\varphi = \{(x, \lambda_i(x)), i = 1, \dots, n\} \subset L.$$

The spectral correspondence

$E \rightarrow X$ holomorphic vector bundle.

$\varphi : E \rightarrow E \otimes L$ **twisted** vector
bundle endomorphism :

- $\lambda_1(x), \dots, \lambda_n(x) \in L_x$

eigenvalues of

$$\varphi_x : E_x \rightarrow E_x \otimes L_x,$$

- $v_1(x), \dots, v_n(x) \in E_x$

eigenvectors of

$$\varphi_x : E_x \rightarrow E_x \otimes L_x.$$

- **Spectral curve** (locally):

$$S_\varphi = \{(x, \lambda_i(x)), i = 1, \dots, n\} \subset L.$$

- $S_\varphi \xrightarrow{\pi} X$ ramified cover.

The spectral correspondence

$E \rightarrow X$ holomorphic vector bundle.


$\varphi : E \rightarrow E \otimes L$ **twisted** vector bundle endomorphism :

- $\lambda_1(x), \dots, \lambda_n(x) \in L_x$
eigenvalues of
 $\varphi_x : E_x \rightarrow E_x \otimes L_x$,
- $v_1(x), \dots, v_n(x) \in E_x$
eigenvectors of
 $\varphi_x : E_x \rightarrow E_x \otimes L_x$.

- **Spectral curve** (locally):

$$S_\varphi = \{(x, \lambda_i(x)), i = 1, \dots, n\} \subset L.$$

- $S_\varphi \xrightarrow{\pi} X$ ramified cover.



The spectral correspondence


$E \rightarrow X$ holomorphic vector bundle.

$\varphi : E \rightarrow E \otimes L$ **twisted** vector bundle endomorphism :

- $\lambda_1(x), \dots, \lambda_n(x) \in L_x$
eigenvalues of
 $\varphi_x : E_x \rightarrow E_x \otimes L_x$,
- $v_1(x), \dots, v_n(x) \in E_x$
eigenvectors of
 $\varphi_x : E_x \rightarrow E_x \otimes L_x$.
- **Spectral curve** (locally):

$$S_\varphi = \{(x, \lambda_i(x)), i = 1, \dots, n\} \subset L.$$

- $S_\varphi \xrightarrow{\pi} X$ ramified cover.



The spectral correspondence

Theorem

Let $b \in \bigoplus_{i=1}^n H^0(X, L^i)$ such that the spectral curve S_b is irreducible and smooth.

The spectral correspondence

Theorem

Let $b \in \bigoplus_{i=1}^n H^0(X, L^i)$ such that the spectral curve S_b is irreducible and smooth. There is a bijective correspondence

$$\begin{aligned} \mathrm{Pic}^\delta(S_b) &\longleftrightarrow \{[(E, \varphi)] \mid P_\varphi(T) = P_b(T), \mathrm{rk} E = n, \deg E = d\} \\ [L] &\longmapsto [(\pi_* L, \pi_* \lambda)]. \end{aligned}$$

The spectral correspondence

Theorem

Let $b \in \bigoplus_{i=1}^n H^0(X, L^i)$ such that the spectral curve S_b is irreducible and smooth. There is a bijective correspondence

$$\begin{aligned} \mathrm{Pic}^\delta(S_b) &\longleftrightarrow \{[(E, \varphi)] \mid P_\varphi(T) = P_b(T), \mathrm{rk} E = n, \deg E = d\} \\ [L] &\longmapsto [(\pi_* L, \pi_* \lambda)]. \end{aligned}$$

with

$$\delta = d + \frac{n(n-1)}{2}d.$$

The spectral correspondence

Theorem

Let $b \in \bigoplus_{i=1}^n H^0(X, L^i)$ such that the spectral curve S_b is irreducible and smooth. There is a bijective correspondence

$$\begin{aligned} \mathrm{Pic}^\delta(S_b) &\longleftrightarrow \{[(E, \varphi)] \mid P_\varphi(T) = P_b(T), \mathrm{rk} E = n, \deg E = d\} \\ [L] &\longmapsto [(\pi_* L, \pi_* \lambda)]. \end{aligned}$$

with

$$\delta = d + \frac{n(n-1)}{2}d.$$

- Hitchin (1987) for $L = K_X$.

The spectral correspondence

Theorem

Let $b \in \bigoplus_{i=1}^n H^0(X, L^i)$ such that the spectral curve S_b is irreducible and smooth. There is a bijective correspondence

$$\begin{aligned} \mathrm{Pic}^\delta(S_b) &\longleftrightarrow \{[(E, \varphi)] \mid P_\varphi(T) = P_b(T), \mathrm{rk} E = n, \deg E = d\} \\ [L] &\longmapsto [(\pi_* L, \pi_* \lambda)]. \end{aligned}$$

with

$$\delta = d + \frac{n(n-1)}{2}d.$$

- Hitchin (1987) for $L = K_X$.
- Beauville, Narasimhan and Ramanan (1989) for general L .

The Hitchin system

- $(E, \varphi) \in T^*\mathcal{N}(n, d)$, characteristic polynomial:

$$P_\varphi(T) = \det(T - \varphi) = T^n + \sum_{i=1}^n \sigma_i(\varphi) T^{n-i}.$$

- **Hitchin fibration:**

$$\begin{aligned} H : T^*\mathcal{N}(n, d) &\longrightarrow \mathcal{B} := \bigoplus_{i=1}^n H^0(X, K_X^i) \\ (E, \varphi) &\longmapsto (\sigma_1(\varphi), \dots, \sigma_n(\varphi)) \end{aligned}$$

The Hitchin system

- $(E, \varphi) \in T^*\mathcal{N}(n, d)$, characteristic polynomial:

$$P_\varphi(T) = \det(T - \varphi) = T^n + \sum_{i=1}^n \sigma_i(\varphi) T^{n-i}.$$

- **Hitchin fibration:**

$$\begin{aligned} H : T^*\mathcal{N}(n, d) &\longrightarrow \mathcal{B} := \bigoplus_{i=1}^n H^0(X, K_X^i) \\ (E, \varphi) &\longmapsto (\sigma_1(\varphi), \dots, \sigma_n(\varphi)) \end{aligned}$$

- The fibres are open (stability) subsets of tori, $H^{-1}(b) \subset \text{Pic}^\delta(S_b) \cong J(S_b)$.

The Hitchin system

- $(E, \varphi) \in T^*\mathcal{N}(n, d)$, characteristic polynomial:

$$P_\varphi(T) = \det(T - \varphi) = T^n + \sum_{i=1}^n \sigma_i(\varphi) T^{n-i}.$$

- **Hitchin fibration:**

$$H : T^*\mathcal{N}(n, d) \longrightarrow \mathcal{B} := \bigoplus_{i=1}^n H^0(X, K_X^i)$$
$$(E, \varphi) \longmapsto (\sigma_1(\varphi), \dots, \sigma_n(\varphi))$$

- The fibres are open (stability) subsets of tori, $H^{-1}(b) \subset \text{Pic}^\delta(S_b) \cong J(S_b)$.
- What about the remaining points of the Jacobian?

The Hitchin system

- $(E, \varphi) \in T^*\mathcal{N}(n, d)$, characteristic polynomial:

$$P_\varphi(T) = \det(T - \varphi) = T^n + \sum_{i=1}^n \sigma_i(\varphi) T^{n-i}.$$

- **Hitchin fibration:**

$$\begin{aligned} H : T^*\mathcal{N}(n, d) &\longrightarrow \mathcal{B} := \bigoplus_{i=1}^n H^0(X, K_X^i) \\ (E, \varphi) &\longmapsto (\sigma_1(\varphi), \dots, \sigma_n(\varphi)) \end{aligned}$$

- The fibres are open (stability) subsets of tori, $H^{-1}(b) \subset \text{Pic}^\delta(S_b) \cong J(S_b)$.
- What about the remaining points of the Jacobian?
Wider notion of stability for pairs (E, φ) .

Definition

A **Higgs bundle** is a pair (E, φ) where $E \rightarrow X$ is a holomorphic vector bundle and the **Higgs field** $\varphi : E \rightarrow E \otimes K_X$ is a K_X -twisted endomorphism.

Definition

A **Higgs bundle** is a pair (E, φ) where $E \rightarrow X$ is a holomorphic vector bundle and the **Higgs field** $\varphi : E \rightarrow E \otimes K_X$ is a K_X -twisted endomorphism.

A Higgs bundle (E, φ) is **stable** if for every proper holomorphic subbundle $E' \subset E$ such that $\varphi(E') \subset E' \otimes K_X$

$$\frac{\deg E'}{\operatorname{rk} E'} < \frac{\deg E}{\operatorname{rk} E}.$$

Definition

A **Higgs bundle** is a pair (E, φ) where $E \rightarrow X$ is a holomorphic vector bundle and the **Higgs field** $\varphi : E \rightarrow E \otimes K_X$ is a K_X -twisted endomorphism.

A Higgs bundle (E, φ) is **stable** if for every proper holomorphic subbundle $E' \subset E$ such that $\varphi(E') \subset E' \otimes K_X$

$$\frac{\deg E'}{\operatorname{rk} E'} < \frac{\deg E}{\operatorname{rk} E}.$$

- **Moduli space of stable Higgs bundles:** $\mathcal{M}(n, d)$

Definition

A **Higgs bundle** is a pair (E, φ) where $E \rightarrow X$ is a holomorphic vector bundle and the **Higgs field** $\varphi : E \rightarrow E \otimes K_X$ is a K_X -twisted endomorphism.

A Higgs bundle (E, φ) is **stable** if for every proper holomorphic subbundle $E' \subset E$ such that $\varphi(E') \subset E' \otimes K_X$

$$\frac{\deg E'}{\operatorname{rk} E'} < \frac{\deg E}{\operatorname{rk} E}.$$

- **Moduli space of stable Higgs bundles:** $\mathcal{M}(n, d)$
- Hitchin (1987), Simpson (1988), Nitsure (1991)

Definition

A **Higgs bundle** is a pair (E, φ) where $E \rightarrow X$ is a holomorphic vector bundle and the **Higgs field** $\varphi : E \rightarrow E \otimes K_X$ is a K_X -twisted endomorphism.

A Higgs bundle (E, φ) is **stable** if for every proper holomorphic subbundle $E' \subset E$ such that $\varphi(E') \subset E' \otimes K_X$

$$\frac{\deg E'}{\operatorname{rk} E'} < \frac{\deg E}{\operatorname{rk} E}.$$

- **Moduli space of stable Higgs bundles:** $\mathcal{M}(n, d)$
- Hitchin (1987), Simpson (1988), Nitsure (1991)
- The Hitchin fibration extends to a proper map $H : \mathcal{M}(n, d) \rightarrow \mathcal{B}$.

The gauge-theoretic point of view

(\mathbb{E}, h) smooth complex Hermitian vector bundle of rank n and degree d over X .

The gauge-theoretic point of view

(\mathbb{E}, h) smooth complex Hermitian vector bundle of rank n and degree d over X .

- **Hitchin's equations:** $(\nabla, \varphi) \in \mathcal{A} \times \Omega^{1,0}(X, \text{End } \mathbb{E})$.

$$\begin{cases} F_{\nabla} + [\varphi, \varphi^{\dagger}] = -2\pi i \frac{d}{n} \mathbf{1}_{\mathbb{E}} \omega_X \\ \nabla^{0,1} \varphi = 0. \end{cases}$$

The gauge-theoretic point of view

(\mathbb{E}, h) smooth complex Hermitian vector bundle of rank n and degree d over X .

- **Hitchin's equations:** $(\nabla, \varphi) \in \mathcal{A} \times \Omega^{1,0}(X, \text{End } \mathbb{E})$.

$$\begin{cases} F_{\nabla} + [\varphi, \varphi^{\dagger}] = -2\pi i \frac{d}{n} \mathbf{1}_{\mathbb{E}} \omega_X \\ \nabla^{0,1} \varphi = 0. \end{cases}$$

- **Hitchin–Kobayashi correspondence:**

Irreducible solutions of Hitchin's equations \longleftrightarrow Stable Higgs bundles

Definition

Let $V \rightarrow X$ be a rank r holomorphic vector bundle. A **V -twisted Higgs bundle** is a pair (E, φ) , where $E \rightarrow X$ is a holomorphic vector bundle and $\varphi : E \rightarrow E \otimes V$

Definition

Let $V \rightarrow X$ be a rank r holomorphic vector bundle. A **V -twisted Higgs bundle** is a pair (E, φ) , where $E \rightarrow X$ is a holomorphic vector bundle and $\varphi : E \rightarrow E \otimes V$ with

$$\varphi \wedge \varphi = 0 \text{ in } \text{End}E \otimes \Lambda^2 V.$$

Definition

Let $V \rightarrow X$ be a rank r holomorphic vector bundle. A **V -twisted Higgs bundle** is a pair (E, φ) , where $E \rightarrow X$ is a holomorphic vector bundle and $\varphi : E \rightarrow E \otimes V$ with

$$\varphi \wedge \varphi = 0 \text{ in } \text{End}E \otimes \Lambda^2 V.$$

- The condition $\varphi \wedge \varphi = 0$ means that the components of φ commute, $[\varphi_i, \varphi_j] = 0$.

Higgs bundles twisted by a vector bundle

Definition

Let $V \rightarrow X$ be a rank r holomorphic vector bundle. A **V -twisted Higgs bundle** is a pair (E, φ) , where $E \rightarrow X$ is a holomorphic vector bundle and $\varphi : E \rightarrow E \otimes V$ with

$$\varphi \wedge \varphi = 0 \text{ in } \text{End}E \otimes \Lambda^2 V.$$

- The condition $\varphi \wedge \varphi = 0$ means that the components of φ commute, $[\varphi_i, \varphi_j] = 0$.
- Same stability condition than that for (classical) Higgs bundles.

Definition

Let $V \rightarrow X$ be a rank r holomorphic vector bundle. A **V -twisted Higgs bundle** is a pair (E, φ) , where $E \rightarrow X$ is a holomorphic vector bundle and $\varphi : E \rightarrow E \otimes V$ with

$$\varphi \wedge \varphi = 0 \text{ in } \text{End}E \otimes \Lambda^2 V.$$

- The condition $\varphi \wedge \varphi = 0$ means that the components of φ commute, $[\varphi_i, \varphi_j] = 0$.
- Same stability condition than that for (classical) Higgs bundles.
- Moduli space: $\mathcal{M}_V(n, d)$.

Higgs bundles twisted by a vector bundle

Definition

Let $V \rightarrow X$ be a rank r holomorphic vector bundle. A **V -twisted Higgs bundle** is a pair (E, φ) , where $E \rightarrow X$ is a holomorphic vector bundle and $\varphi : E \rightarrow E \otimes V$ with

$$\varphi \wedge \varphi = 0 \text{ in } \text{End}E \otimes \Lambda^2 V.$$

- The condition $\varphi \wedge \varphi = 0$ means that the components of φ commute, $[\varphi_i, \varphi_j] = 0$.
- Same stability condition than that for (classical) Higgs bundles.
- Moduli space: $\mathcal{M}_V(n, d)$.
- Studied by D. Xie and K. Yonekura in the context of $\mathcal{N} = 1$ supersymmetric gauge theories.

The gauge-theoretic point of view

(\mathbb{E}, h) smooth complex Hermitian vector bundle of rank n and degree d over X .

Fix h_V Hermitian metric on the holomorphic vector bundle $V \rightarrow X$.

The gauge-theoretic point of view

(\mathbb{E}, h) smooth complex Hermitian vector bundle of rank n and degree d over X .

Fix h_V Hermitian metric on the holomorphic vector bundle $V \rightarrow X$.

- **Generalized Hitchin's equations:** $(\nabla, \varphi) \in \mathcal{A} \times \Omega^0(X, \text{End } \mathbb{E} \otimes V)$

$$\begin{cases} F_{\nabla} + [\varphi, \varphi^{\dagger}]_{h, h_V} \omega_X = -2\pi i \frac{d}{n} \mathbf{1}_{\mathbb{E}} \omega_X \\ (\nabla^{0,1} \otimes \mathbf{1}_V + \mathbf{1}_{\text{End } \mathbb{E}} \otimes \nabla_V^{0,1}) \varphi = 0 \\ \varphi \wedge \varphi = 0. \end{cases}$$

The gauge-theoretic point of view

(\mathbb{E}, h) smooth complex Hermitian vector bundle of rank n and degree d over X .

Fix h_V Hermitian metric on the holomorphic vector bundle $V \rightarrow X$.

- **Generalized Hitchin's equations:** $(\nabla, \varphi) \in \mathcal{A} \times \Omega^0(X, \text{End } \mathbb{E} \otimes V)$

$$\begin{cases} F_{\nabla} + [\varphi, \varphi^{\dagger}]_{h, h_V} \omega_X = -2\pi i \frac{d}{n} \mathbf{1}_{\mathbb{E}} \omega_X \\ (\nabla^{0,1} \otimes \mathbf{1}_V + \mathbf{1}_{\text{End } \mathbb{E}} \otimes \nabla_V^{0,1}) \varphi = 0 \\ \varphi \wedge \varphi = 0. \end{cases}$$

- **Hitchin–Kobayashi correspondence:**

Irreducible solutions of generalized Hitchin's equations \longleftrightarrow Stable V -twisted Higgs bundles

The generalized Hitchin system

- $(E, \varphi) \in \mathcal{M}_V(n, d)$, characteristic polynomial:

$$P_\varphi(T) = \det(T - \varphi) = T^n + \sum_{i=1}^n \sigma_i(\varphi) T^{n-i}.$$

The generalized Hitchin system

- $(E, \varphi) \in \mathcal{M}_V(n, d)$, characteristic polynomial:

$$P_\varphi(T) = \det(T - \varphi) = T^n + \sum_{i=1}^n \sigma_i(\varphi) T^{n-i}.$$

- **Hitchin fibration:**

$$\begin{aligned} H_V : \mathcal{M}_V(n, d) &\longrightarrow \bigoplus_{i=1}^n H^0(X, \operatorname{Sym}^i V) \\ (E, \varphi) &\longmapsto (\sigma_1(\varphi), \dots, \sigma_n(\varphi)) \end{aligned}$$

The generalized Hitchin system

- $(E, \varphi) \in \mathcal{M}_V(n, d)$, characteristic polynomial:

$$P_\varphi(T) = \det(T - \varphi) = T^n + \sum_{i=1}^n \sigma_i(\varphi) T^{n-i}.$$

- **Hitchin fibration:**

$$\begin{aligned} H_V : \mathcal{M}_V(n, d) &\longrightarrow \bigoplus_{i=1}^n H^0(X, \operatorname{Sym}^i V) \\ (E, \varphi) &\longmapsto (\sigma_1(\varphi), \dots, \sigma_n(\varphi)) \end{aligned}$$

- Not surjective!!

The generalized Hitchin system

- $(E, \varphi) \in \mathcal{M}_V(n, d)$, characteristic polynomial:

$$P_\varphi(T) = \det(T - \varphi) = T^n + \sum_{i=1}^n \sigma_i(\varphi) T^{n-i}.$$

- **Hitchin fibration:**

$$\begin{aligned} H_V : \mathcal{M}_V(n, d) &\longrightarrow \bigoplus_{i=1}^n H^0(X, \operatorname{Sym}^i V) \\ (E, \varphi) &\longmapsto (\sigma_1(\varphi), \dots, \sigma_n(\varphi)) \end{aligned}$$

- Not surjective!! $\mathcal{B}_V = \operatorname{im} H_V$

The generalized spectral correspondence

- The spectral curve can be defined analogously in V , as the solutions of $P_\varphi(T)$.

The generalized spectral correspondence

- The spectral curve can be defined analogously in V , as the solutions of $P_\varphi(T)$.
- The number of equations is bigger than the dimension.

The generalized spectral correspondence

- The spectral curve can be defined analogously in V , as the solutions of $P_\varphi(T)$.
- The number of equations is bigger than the dimension. However...

The generalized spectral correspondence

- The spectral curve can be defined analogously in V , as the solutions of $P_\varphi(T)$.
- The number of equations is bigger than the dimension. However...
- **Main idea:** *Since the components of φ commute, generically they can be diagonalized simultaneously and can be written as polynomials of one of them.*

The generalized spectral correspondence

- The spectral curve can be defined analogously in V , as the solutions of $P_\varphi(T)$.
- The number of equations is bigger than the dimension. However...
- **Main idea:** *Since the components of φ commute, generically they can be diagonalized simultaneously and can be written as polynomials of one of them.*

Theorem

Take $b \in \mathcal{B}_V$ such that the spectral curve S_b is integral and smooth.

The generalized spectral correspondence

- The spectral curve can be defined analogously in V , as the solutions of $P_\varphi(T)$.
- The number of equations is bigger than the dimension. However...
- **Main idea:** *Since the components of φ commute, generically they can be diagonalized simultaneously and can be written as polynomials of one of them.*

Theorem

Take $b \in \mathcal{B}_V$ such that the spectral curve S_b is integral and smooth. Then there is a bijective correspondence

$$\mathrm{Pic}(S_b) \longleftrightarrow \{[(E, \varphi)] \mid S_\varphi = S_b\}.$$

Open questions and further directions

Open questions:

Open questions and further directions

Open questions:

- Complete the Hitchin-Kobayashi correspondence.

Open questions and further directions

Open questions:

- Complete the Hitchin-Kobayashi correspondence.
- GIT construction of the moduli space.

Open questions and further directions

Open questions:

- Complete the Hitchin-Kobayashi correspondence.
- GIT construction of the moduli space.
- Computation of the dimension of $\mathcal{M}_V(n, d)$ using deformation theory.

Open questions and further directions

Open questions:

- Complete the Hitchin-Kobayashi correspondence.
- GIT construction of the moduli space.
- Computation of the dimension of $\mathcal{M}_V(n, d)$ using deformation theory.
- Genus of the spectral curve and relationship between the degrees.

Open questions and further directions

Open questions:

- Complete the Hitchin-Kobayashi correspondence.
- GIT construction of the moduli space.
- Computation of the dimension of $\mathcal{M}_V(n, d)$ using deformation theory.
- Genus of the spectral curve and relationship between the degrees.
- Conditions for the curve to be smooth.

Open questions and further directions

Open questions:

- Complete the Hitchin-Kobayashi correspondence.
- GIT construction of the moduli space.
- Computation of the dimension of $\mathcal{M}_V(n, d)$ using deformation theory.
- Genus of the spectral curve and relationship between the degrees.
- Conditions for the curve to be smooth.
- Study the ramification divisor.

Open questions and further directions

Open questions:

- Complete the Hitchin-Kobayashi correspondence.
- GIT construction of the moduli space.
- Computation of the dimension of $\mathcal{M}_V(n, d)$ using deformation theory.
- Genus of the spectral curve and relationship between the degrees.
- Conditions for the curve to be smooth.
- Study the ramification divisor.

Further directions:

Open questions and further directions

Open questions:

- Complete the Hitchin-Kobayashi correspondence.
- GIT construction of the moduli space.
- Computation of the dimension of $\mathcal{M}_V(n, d)$ using deformation theory.
- Genus of the spectral curve and relationship between the degrees.
- Conditions for the curve to be smooth.
- Study the ramification divisor.

Further directions:

- Study the case $V = L_1 \oplus L_2$ with $L_1 \otimes L_2 = K_X$ (V is Calabi–Yau).

Open questions and further directions


Open questions:


- Complete the Hitchin-Kobayashi correspondence.
- GIT construction of the moduli space.
- Computation of the dimension of $\mathcal{M}_V(n, d)$ using deformation theory.
- Genus of the spectral curve and relationship between the degrees.
- Conditions for the curve to be smooth.
- Study the ramification divisor.


Further directions:


- Study the case $V = L_1 \oplus L_2$ with $L_1 \otimes L_2 = K_X$ (V is Calabi–Yau).
- Donaldson–Thomas invariants.

Bibliography

 A. Beauville, M. S. Narasimhan, and S. Ramanan.
Spectral curves and the generalised theta divisor.
J. reine angew. Math, 398(169-179):3, 1989.

 N. Hitchin.
The self-duality equations on a Riemann surface.
Proceedings of the London Mathematical Society, 3(1):59–126, 1987.

 N. Hitchin.
Stable bundles and integrable systems.
Duke mathematical journal, 54(1):91–114, 1987.

 D. Xie and K. Yonekura.
Generalized Hitchin system, spectral curve and $\mathcal{N} = 1$ dynamics.
Journal of High Energy Physics, 2014(1):1, 2014.