

Numerical Methods in Soft Matter Exercises

Guglielmo Bordin

University of Padua, 2024

Contents

1	Sampling via hit-and-miss and inversion	1
1.1	Sampling in d -dimensional domains by hit-and-miss	1
1.2	Sampling from a distribution with the inversion method	1
1.3	Another sampling with the inversion method	1
1.4	Additional exercises on the inversion method	3
2	Sampling via transformation of coordinates	7
2.1	Sampling uniformly within a unit radius disk	7
2.2	Box–Muller transform	7
2.3	Rejection method	8

Sampling via hit-and-miss and inversion

1.1 Sampling in d -dimensional domains by hit-and-miss

Source codes `src/011a_rect_hit_miss.c, src/011b_disk_hit_miss.c`

Analysis scripts `src/011_hit_miss.R`

Executables `exe/011a_rect_hit_miss, exe/011b_disk_hit_miss`

The error as a function of the number of throws is shown in Fig. 1.1.1. In the disk case, it goes under 1 % with as little as 10 000 ‘throws’.

1.2 Sampling from a distribution with the inversion method

Source codes `src/012_inversion_power34.c`

Analysis scripts `src/012-4_inversion_method.R`

Executables `exe/012_inversion_power34`

The idea is to sample from the probability distribution $\rho_n(x) = cx^n$ in $[0, 1]$. First, using the normalization condition we can find out what c should be:

$$1 = \int_0^1 cx^n dx = \frac{c}{n+1} \implies c = n+1. \quad (1.2.1)$$

Then, we find the expression of the associated cumulative density function:

$$F_n(x) = (n+1) \int_0^x y^n dy = x^{n+1}, \quad (1.2.2)$$

and invert it:

$$u = x^{n+1} \implies x = u^{1/(n+1)}. \quad (1.2.3)$$

So, we just have to sample a random floating point number between 0 and 1 and raise it to the power of $1/(n+1)$ to get $x \sim (n+1)x^n$. Results from a sampling of 50 000 iterations for $n = 3, 4$ are shown in Fig. 1.2.1.

1.3 Another sampling with the inversion method

Source codes `src/012_inversion_power2.c`

Analysis scripts `src/012-4_inversion_method.R`

Executables `exe/012_inversion_power2`




Figure 1.1.1: percentage error as a function of the number of ‘throws’ for Ex. 1.1. The black line shows the average over 100 realizations, with the standard deviation shaded in grey.




Figure 1.2.1: histogram of 50 000 samples from $\rho_n(x) = cx^n$, $x \in [0, 1]$, for $n = 3$ (above) and $n = 4$ (below), obtained with the inversion method. The theoretical distribution is overlaid with a solid line.




Figure 1.3.1: histogram of 50 000 samples from $\rho(x) = 3x^2/8$, $x \in [0, 2]$, obtained with the inversion method. The theoretical distribution is overlaid with a solid line.

I slightly modified the code from the previous exercise, since c is different now. The distribution is $\rho(x) = cx^2$ with $x \in [0, 2]$, so:

$$1 = \int_0^2 cx^2 dx = \frac{8}{3}c \implies c = \frac{3}{8}. \quad (1.3.1)$$

The cumulative is then

$$F(x) = \frac{3}{8} \int_0^x y^2 dy = \frac{x^3}{8} \implies x = 2u^{1/3}. \quad (1.3.2)$$

Once again, you can see the comparison between a 50 000-points histogram and the theoretical curve in Fig. 1.3.1.

1.4 Additional exercises on the inversion method

Source codes `src/014a_inversion_exp.c`, `src/014b_inversion_exp2.c`,
 `src/014c_inversion_powerlaw.c`

Analysis scripts `src/012-4_inversion_method.R`

Executables `exe/014_inversion_exp`, `exe/014b_inversion_exp2`,
 `exe/014c_inversion_powerlaw`

The first exercise here asked to sample from $\rho_1(x) = \mu e^{-\mu x}$, with $x \in [0, \infty)$. The cumulative and the sampling rule are then

$$F_1(x) = \mu \int_0^x e^{-\mu y} dy = \mu(1 - e^{-\mu x}) \implies x = -\log(1 - u)/\mu. \quad (1.4.1)$$

I ran 50 000 samples as before, choosing $\mu = 2$; see Fig. 1.4.1.

The second distribution was $\rho_2(x) = 2xe^{-x^2}$, $x \in [0, \infty)$. Cumulative and sampling rule:

$$F_2(x) = 2 \int_0^x e^{-y^2} dy = 1 - e^{-x^2} \implies x = \sqrt{-\log(1 - u)}. \quad (1.4.2)$$

A histogram of 50 000 samples is shown in Fig. 1.4.2.




Figure 1.4.1: histogram of 50 000 samples from $\rho(x) = 2e^{-2x}$, $x \in [0, \infty)$, obtained with the inversion method. The theoretical distribution is overlaid with a solid line.




Figure 1.4.2: histogram of 50 000 samples from $\rho(x) = 2xe^{-x^2}$, $x \in [0, \infty)$, obtained with the inversion method. The theoretical distribution is overlaid with a solid line.




Figure 1.4.3: histogram of 50 000 samples from $\rho(x) = 375/(5+x)^4$, $x \in [0, \infty)$, obtained with the inversion method. The theoretical distribution is overlaid with a solid line.

The last distribution was $\rho_3(x) = c(a+bx)^{-n}$, with $x \in [0, \infty)$ and $n > 1$. The normalization constant is given by

$$1 = \int_0^\infty c(a+bx)^{-n} dx = \frac{a^{1-n}c}{b(n-1)} \implies c = b(n-1)a^{n-1}. \quad (1.4.3)$$

By integrating we obtain

$$F_3(x) = c \int_0^x (a+by)^{-n} dy = \frac{c}{b(1-n)} [(a+bx)^{1-n} - a^{1-n}] = 1 - a^{n-1}(a+bx)^{1-n}, \quad (1.4.4)$$

from which we can sample with

$$u = 1 - a^{n-1}(a+bx)^{1-n} \implies x = \frac{a}{b} [(1-u)^{1/(1-n)} - 1]. \quad (1.4.5)$$

I chose $a = 1/2$, $b = 1/10$ and $n = 4$ – resulting in $\rho_3(x) = 375/(5+x)^4$ – for a run of 50 000 samples, shown in Fig. 1.4.3.

Sampling via transformation of coordinates

2.1 Sampling uniformly within a unit radius disk

Source codes `src/021_disk_sampling.c`

Analysis scripts `src/021_disk_sampling.R`

Executables `exe/021_disk_sampling`

If we want to sample points uniformly distributed over the unit disk, we cannot simply generate $r \sim \mathcal{U}_{[0,1]}$ and $\vartheta \sim \mathcal{U}_{[0,2\pi]}$. The result of doing that is shown in the left graph of Fig. 2.1.1. While the angular distribution poses no problem, there is an undesired higher density of points close to the centre of the disk.

The reason for this becomes clear when you consider the number of points falling inside a ring of given thickness δr . Consider a ring with inner radius r_1 and outer radius $r_1 + \delta r$, with $\delta r \ll r_1$, and another with radii $r_2, r_2 + \delta r$. Despite the ratio of their areas being

$$\frac{\pi[(r_1 + \delta r)^2 - r_1^2]}{\pi[(r_2 + \delta r)^2 - r_2^2]} = \frac{2r_1\delta r + \mathcal{O}(\delta r^2)}{2r_2\delta r + \mathcal{O}(\delta r^2)} = \frac{r_1}{r_2} + \mathcal{O}(\delta r), \quad (2.1.1)$$

the fraction of points within each ring is in both cases proportional to δr , if r is sampled uniformly. Thus, the number of points falling at a distance r should be proportional to r :

$$\rho_r(r) = 2r, \quad (2.1.2)$$

with the 2 in front to ensure normalization over $[0, 1]$. By computing the cumulative and inverting we get

$$F_r(r) = 2 \int_0^r x \, dx = r^2 \implies u = r^2 \implies r = \sqrt{u}. \quad (2.1.3)$$

The graph on the right in Fig. 2.1.1 shows how the sampling becomes visibly uniform with this adjustment.

2.2 Box–Muller transform

Source codes `src/022_box_muller.c`

Analysis scripts `src/022_box_muller.R`

Executables `exe/022_box_muller`

I started from the factorization of $\rho_{x,y}(x, y)$:

$$\rho_{x,y}(x, y) = \frac{1}{2\pi} \exp\left(-\frac{x^2 + y^2}{2}\right) = \rho_x(x)\rho_y(y), \quad \text{with } \rho_x(x) = \frac{e^{-x^2/2}}{\sqrt{2\pi}}. \quad (2.2.1)$$




Figure 2.1.1: visualization of 50 000 samples on the unit disk, with the ‘naïve’ method of uniform sampling both on r and ϑ (left), and with the correct method described in the text (right).

If we switch to polar coordinates this becomes, remembering the factor r coming from the Jacobian,

$$\rho_{r,\vartheta}(r, \vartheta) = \frac{r}{2\pi} e^{-r^2/2}, \quad (2.2.2)$$

with $\rho_r(r) = r e^{-r^2/2}$ and $\rho_\vartheta(\vartheta) = 1/2\pi$.

Then, we can marginalize over ϑ to get $\rho_r(r)$:

$$\rho_r(r) = \frac{1}{2\pi} \int_0^{2\pi} r e^{-r^2/2} d\vartheta = r e^{-r^2/2}. \quad (2.2.3)$$

As usual, we compute the cumulative and invert it to generate a random $r \sim \rho_r(r)$:

$$F_r(r) = \int_0^r x e^{-x^2/2} dx = 1 - e^{-r^2/2} \implies e^{-r^2/2} = 1 - u. \quad (2.2.4)$$

Since u is distributed uniformly, we can redefine it as $1 - u$ for simplicity:

$$-\frac{r^2}{2} = \log u \implies r = \sqrt{-2 \log u}. \quad (2.2.5)$$

Then, to get the angle ϑ we recover the *conditional* distribution $\rho_{\vartheta|r}(\vartheta | r)$:

$$\rho_{\vartheta|r}(\vartheta | r) = \frac{\rho_{r,\vartheta}(r, \vartheta)}{\rho_r(r)} = \frac{1}{2\pi}. \quad (2.2.6)$$

Thus, for ϑ we have simply to sample from $\mathcal{U}_{[0,2\pi]}$.

The idea then is to sample two numbers u_1, u_2 from $\mathcal{U}_{[0,1]}$ at each iteration; at that point we can calculate

$$x = \sqrt{-2 \log u_1} \cos(2\pi u_2), \quad y = \sqrt{-2 \log u_1} \sin(2\pi u_2), \quad (2.2.7)$$

to get two numbers x, y distributed according to a standard Gaussian $\mathcal{N}(0, 1)$. We can also get a $z \sim \mathcal{N}(\mu, \sigma^2)$ afterwards by multiplying x or y by σ and summing the mean: $z = \mu + \sigma x$. I ran a simulation with 100 000 samples choosing $\mu = 1$ and $\sigma = 2$, and you can see the results in Fig. 2.2.1.

2.3 Rejection method

Source codes `src/023a_rejection_sampling.c,`
 `src/023b_rejection_accratio.c`




Figure 2.2.1: histograms of 100 000 samples – half in the upper panel, half in the lower one – from a $N(1, 4)$ distribution, obtained with the Box–Muller method. The theoretical distribution is overlaid with a solid line.

Analysis scripts `src/023_rejection_method.R`

Executables `exe/023a_rejection_sampling,`
 `exe/023b_rejection_accratio`

Here I implemented the sampling of random numbers from the distribution

$$f(x) = \frac{2}{\sqrt{\pi}} e^{-x^2} \quad (2.3.1)$$

over $[0, \infty)$. We start from a function $g(x)$ that could serve as an upper limit to $f(x)$,

$$g(x) = \begin{cases} A & \text{for } 0 \leq x \leq p, \\ \frac{A}{p} x e^{p^2 - x^2} & \text{for } x > p. \end{cases} \quad (2.3.2)$$

First we normalize it to turn it into a proper probability distribution:

$$1 = \int_0^\infty g(x) dx = Ap + \frac{A}{p} e^{p^2} \int_p^\infty x e^{-x^2} dx = A \left(p + \frac{1}{2p} \right) \implies A = \frac{2p}{1 + 2p^2}. \quad (2.3.3)$$

Then we need a constant c such that $cg(x) \geq f(x)$ everywhere. The maximum of $f(x)$ is in $x = 0$, where $f(0) = 2/\sqrt{\pi}$, so we set $cA = 2/\sqrt{\pi}$. Another thing to consider is that $g(x)$ can have a maximum larger than A in $[p, \infty)$:

$$g'(x) = \begin{cases} 0 & \text{for } 0 \leq x \leq p, \\ \frac{A}{p} (1 - 2x^2) e^{p^2 - x^2} & \text{for } x > p. \end{cases} \quad (2.3.4)$$

Thus in the region $[p, \infty)$ the derivative is zero in $x = 1/\sqrt{2}$. To avoid this we have to choose $p \geq 1/\sqrt{2} \approx 0.707$. An example of sampling with $p = 1/\sqrt{2}$ is given by the histogram in




Figure 2.3.1: histogram of 50 000 samples from the distribution $f(x) = 2e^{-x^2}/\sqrt{\pi}$ in $[0, \infty)$ obtained with the rejection method. The theoretical distribution is overlaid with a solid line.

Fig. 2.3.1. A plot of the acceptance ratio as a function of p , starting from the minimum value $1/\sqrt{2}$, is shown in Fig. 2.3.2; obviously the acceptance ratio decreases as p increases, as the gap between $cg(x)$ and $f(x)$ grows wider:

$$\begin{aligned}
 \int_0^\infty (cg(x) - f(x)) dx &= \frac{2}{\sqrt{\pi}} \int_0^p dx + \frac{2}{p\sqrt{\pi}} \int_p^\infty xe^{p^2-x^2} dx - 1 \\
 &= \frac{2p}{\sqrt{\pi}} + \frac{2}{p\sqrt{\pi}} \frac{1}{2} - 1 \\
 &= \frac{2}{\sqrt{\pi}} \left(p + \frac{1}{2p} \right) - 1 \xrightarrow{p \rightarrow \infty} \infty.
 \end{aligned} \tag{2.3.5}$$




Figure 2.3.2: acceptance ratio over 100 000 samples as a function of p in Ex. 2.3.