
SEIBERG-WITTEN THEORY AND THE INTERSECTION FORM

Santiago Quintero de los Ríos



SUPERVISOR
Gil Cavalcanti

SECOND READER
Fabian Ziltener

UTRECHT UNIVERSITY
AUGUST 19, 2020

Contents

Contents	i
Introduction	iii
1 Preliminaries	1
1.1 Gauge theory	1
1.1.1 Principal bundles, connections and curvature	1
1.1.2 Associated bundles and matter fields	2
1.1.3 The space of connections	2
1.2 Algebraic topology	2
1.2.1 Cohomology and cups	2
1.2.2 Orientations and caps	2
1.2.3 The intersection pairing	2
1.3 Characteristic classes	2
1.3.1 Classifying spaces for line bundles and the first Chern class	2
1.3.2 Čech cohomology and the first Stiefel-Whitney class	3
1.4 Hodge theory	3
1.5 Elliptic regularity	3
1.5.1 Sobolev spaces	3
1.5.2 Elliptic operators	3
2 Spin Geometry	4
2.1 Clifford algebras	4
2.1.1 Definition and basic properties	4
2.1.2 Complex(ified) Clifford algebras	4
2.2 The Spin group	4
2.2.1 Definition and examples	4
2.2.2 When $n = 4$	4
2.2.3 Lie algebra structures	4
2.3 Representations	5
2.3.1 Classification and examples	5
2.3.2 The Spin representation	5
2.3.3 Clifford multiplication and infinitesimal actions	5
2.4 Spin structures	5
2.4.1 From local to global	5
2.4.2 Spinor bundles	5
2.5 The Dirac operator	5
2.5.1 Spin connections	5
2.5.2 The Dirac operator	5
2.6 The world of Spin^c	5
2.6.1 The Spin^c group	5

2.6.2	Going global: Spin^c -structures and spinors	5
2.6.3	The complex spin connection and the coupled Dirac operator	6
2.7	Final ingredient: The squaring map	6
2.7.1	The linear squaring map	6
2.7.2	The global squaring map	6
3	The Seiberg-Witten Equations and Moduli Space	7
3.1	The gauge group and its action	7
3.2	Topology of the Moduli Space	7
3.2.1	Dimension of $\mathcal{M}$	7
3.2.2	Generic smoothness	7
3.2.3	Orientation	7
3.2.4	Compactness	7
3.3	The Seiberg-Witten invariant	8
4	The Intersection Form and Donaldson's Theorem	9
4.1	The intersection form of 4-manifolds	9
4.1.1	Unimodular symmetric forms	9
4.1.2	The topology of four-manifolds	9
4.2	Finale: Donaldson's theorem	9

Introduction

THE problem of sequential investment under uncertainty motivates this thesis. Modern reinforcement learning offers tools for such decisions, but most methods collapse the return to its expectation. Distributional approaches instead model the full return distribution, as introduced by Dabney et al. [?] with Implicit Quantile Networks. Separately, epistemic uncertainty over a model's own estimates can be quantified through Evidential Deep Learning [?]. This thesis combines both ideas [?, ?] into a single risk-sensitive decision support system.

Box gallery (visual test)

[Invariants of unimodular symmetric forms] Let Z be a finitely-generated free abelian group, and let $q : Z \times Z \rightarrow \mathbb{Z}$ be a unimodular form. To it we assign rank, type (parity), and signature.

In [MH73], unimodular symmetric forms on Z are called inner products on Z . Forms of type I are odd and forms of type II are even.

[E_8] Define the form E_8 on $\mathbb{Z}^8$. This form is unimodular, positive-definite and even, so $\text{rank}(E_8) = \text{sign}(E_8) = 8$.

[Freedman] Let q be a unimodular, symmetric form. If q is even, there is exactly one closed, oriented, simply-connected topological four-manifold whose intersection form is q .

[$w_2(M)$ is mod-2 reduction of characteristic vector] Let M be a closed, connected, oriented four-manifold with intersection form q_M .

Then for any characteristic vector $w \in H^2(M, \mathbb{Z})$, $w_2(M) = \text{PD}(w) \bmod 2$.

[Closed four-manifolds admit Spin^c -structures] Every closed, connected, oriented four-manifold admits Spin^c -structures.

[Elkies] Let $q : Z \times Z \rightarrow \mathbb{Z}$ be a symmetric, unimodular bilinear form. If q is not $\oplus(-1)$ nor $\oplus(+1)$, then there exists a characteristic vector w with $|q(w, w)| < \text{rank}(q)$.

Acknowledgements

First of all, I would like to thank the mathematical community whose documents make LaTeX typography worth studying. This section is included to test the visual spacing of front matter pages.

Chapter 1

Preliminaries

IN THIS CHAPTER we review the basic language of algebraic topology and differential geometry that is needed to discuss Seiberg-Witten theory and the Donaldson theorem. In the differential realm, we assume familiarity with smooth manifolds, de Rham cohomology, vector bundles, and some Riemannian geometry. In algebraic topology, we assume familiarity with singular and cellular homology.

1.1 Gauge theory

Gauge theory is the heart of Seiberg-Witten theory. The name comes from physics, where gauge fields are fields with symmetries and redundant degrees of freedom. Mathematically, the study of gauge fields is done in terms of principal bundles, connections, curvature, and associated vector bundles.

1.1.1 Principal bundles, connections and curvature

The basic objects of gauge theory are principal bundles.

[Principal G -bundle] Let M be a smooth manifold and let G be a Lie group. A principal G -bundle over M is a smooth manifold P together with a smooth surjective map $\pi: P \rightarrow M$ and a right action of G on P satisfying equivariance and local triviality conditions.

We write this compactly as

$$G \hookrightarrow P \xrightarrow{\pi} M.$$

[Frame bundles] Let $E \rightarrow M$ be a $\mathbb{K}$ -vector bundle of rank k . The frame bundle $\text{Fr}(E)$ is the bundle whose fiber over $x \in M$ consists of all ordered bases of E_x . It is naturally a principal $\text{GL}(k, \mathbb{K})$ -bundle.

$$\text{Fr}(E) = \coprod_{x \in M} \text{Fr}_x(E).$$

If E is a real Riemannian vector bundle, one obtains the orthogonal frame bundle $O(E)$. If E is a complex Hermitian bundle, one obtains the unitary frame bundle $U(E)$.

[Connection on a principal bundle] Let G be a Lie group with Lie algebra $\mathfrak{g}$. A connection on a principal bundle $G \hookrightarrow P \rightarrow M$ is a $\mathfrak{g}$ -valued one-form $\omega \in \Omega^1(P, \mathfrak{g})$ satisfying the standard equivariance and normalization conditions.

[Transformation of local gauge potentials] Let ω be a connection on a principal bundle $G \hookrightarrow P \rightarrow M$, and let $\{U_i\}_{i \in I}$ be a trivializing cover with local gauges. The local gauge potentials are related on overlaps by

$$\mathcal{A}_j = \text{Ad}_{g_{ij}^{-1}} \circ \mathcal{A}_i + g_{ij}^* \Theta.$$

If G is a matrix Lie group, this becomes

$$\mathcal{A}_j = g_{ij}^{-1} \mathcal{A}_i g_{ij} + g_{ij}^{-1} dg_{ij}.$$

[Curvature] The curvature of a connection ω is the two-form

$$\Omega := d^\omega \omega.$$

Equivalently, it satisfies Cartan's structure equation

$$\Omega = d\omega + \frac{1}{2}[\omega, \omega].$$

1.1.2 Associated bundles and matter fields

Let $G \hookrightarrow P \rightarrow M$ be a principal bundle, V a vector space, and $\rho: G \rightarrow \mathrm{GL}(V)$ a representation. The associated bundle is

$$P \times_\rho V = P \times V / G.$$

Sections of this bundle can be interpreted as matter fields in the language of physics.

1.1.3 The space of connections

[Connections form an affine space] Let $\mathrm{Conn}(P)$ be the set of connections of a principal bundle $G \hookrightarrow P \rightarrow M$. Then $\mathrm{Conn}(P)$ is an affine space modeled on the vector space of Ad-tensorial one-forms $\Omega_{\mathrm{Ad}}^1(P, \mathfrak{g})$.

1.2 Algebraic topology

The main algebraic-topological tools needed later are cohomology, characteristic classes, orientations, and the intersection form of four-manifolds.

1.2.1 Cohomology and cups

[Pairing of cohomology and homology] The natural pairing of cohomology and homology is

$$\langle -, - \rangle: H^k(C, G) \otimes H_k(C) \rightarrow G,$$

given by evaluating a cohomology class on a homology class.

[Universal coefficients for cohomology] Let $C_\bullet$ be a chain complex of free abelian groups and let G be an abelian group. Then there is a split exact sequence

$$0 \rightarrow \mathrm{Ext}(H_{k-1}(C), G) \rightarrow H^k(C, G) \rightarrow \mathrm{Hom}(H_k(C), G) \rightarrow 0.$$

1.2.2 Orientations and caps

A local orientation at a point of an n -manifold is a choice of generator of the local homology group. A global orientation is a locally consistent family of such choices.

1.2.3 The intersection pairing

The intersection pairing on a closed oriented four-manifold is one of the central objects in this story. It may be defined using Poincaré duality and the cup product.

1.3 Characteristic classes

Characteristic classes measure obstructions to trivializing bundles and will later be used to formulate the existence of spin and Spin^c structures.

1.3.1 Classifying spaces for line bundles and the first Chern class

Complex line bundles are classified by maps into $\mathbb{CP}^\infty$, and the first Chern class records the corresponding cohomology class.

1.3.2 Čech cohomology and the first Stiefel-Whitney class

The first Stiefel-Whitney class detects the obstruction to orientability.

1.4 Hodge theory

This section would collect the Hodge-theoretic facts used later: harmonic representatives, Hodge decomposition, and self-dual forms.

1.5 Elliptic regularity

1.5.1 Sobolev spaces

Sobolev spaces provide the functional-analytic setting for elliptic differential operators.

1.5.2 Elliptic operators

Elliptic regularity is one of the analytic mechanisms behind smoothness statements for moduli spaces.

Chapter 2

Spin Geometry

THE PURPOSE of this chapter is to introduce the algebraic and geometric objects that appear in the Seiberg-Witten equations. Clifford algebras, spin groups, spinor bundles, and Dirac operators provide the language in which the equations are formulated.

2.1 Clifford algebras

2.1.1 Definition and basic properties

[Clifford algebra] Let (V, q) be a quadratic vector space. The Clifford algebra $\text{Cl}(V, q)$ is the quotient of the tensor algebra $T(V)$ by the ideal generated by elements of the form

$$v \otimes v + q(v)1.$$

We denote by $\mathbb{R}^{r,s}$ the semi-Riemannian vector space $\mathbb{R}^{r+s}$ with r positive and s negative directions. We write $\text{Cl}(r, s)$ for the corresponding Clifford algebra.

[Cl(1) and Cl(2)] For $n = 1$, the real Clifford algebra generated by one element with relation $e^2 = -1$ is isomorphic to $\mathbb{C}$. For $n = 2$, one obtains an algebra isomorphic to the quaternions $\mathbb{H}$.

2.1.2 Complex(ified) Clifford algebras

Complex Clifford algebras are often simpler to classify and are central for describing spin representations.

2.2 The Spin group

2.2.1 Definition and examples

[Spin group] The group $\text{Spin}(n)$ is the subgroup of the Clifford algebra generated by products of unit vectors and fitting into the double cover

$$1 \rightarrow \mathbb{Z}_2 \rightarrow \text{Spin}(n) \rightarrow \text{SO}(n) \rightarrow 1.$$

2.2.2 When $n = 4$

Dimension four is special because $\text{Spin}(4)$ decomposes in a way that makes four-dimensional gauge theory particularly rich.

2.2.3 Lie algebra structures

Lie algebra computations identify infinitesimal versions of the spin actions used later.

2.3 Representations

2.3.1 Classification and examples

[Irreducible representations of matrix algebras] Let A be a finite direct sum of matrix algebras over a field. The irreducible representations of A are the projections onto the simple summands, followed by the standard representation.

2.3.2 The Spin representation

Spin representations arise by restricting Clifford algebra representations to the spin group.

2.3.3 Clifford multiplication and infinitesimal actions

Clifford multiplication connects algebraic representations with first-order differential operators.

2.4 Spin structures

2.4.1 From local to global

[Spin structure] Let M be an oriented Riemannian manifold. A spin structure is a principal $\text{Spin}(n)$ -bundle lifting the oriented orthonormal frame bundle of M .

2.4.2 Spinor bundles

A spin structure allows one to construct spinor bundles, whose sections are spinor fields.

2.5 The Dirac operator

2.5.1 Spin connections

Connections on spinor bundles are induced by Levi-Civita connections and auxiliary choices.

2.5.2 The Dirac operator

[Dirac operator] Given a spinor bundle $S \rightarrow M$ with connection ∇ , the Dirac operator is locally given by

$$\not{D}\psi = \sum_i e_i \cdot \nabla_{e_i} \psi,$$

where $\{e_i\}$ is a local orthonormal frame.

2.6 The world of Spin^c

2.6.1 The Spin^c group

[Spin^c group] The group $\text{Spin}^c(n)$ is defined by

$$\text{Spin}^c(n) = \frac{\text{Spin}(n) \times U(1)}{\{\pm(1, 1)\}}.$$

2.6.2 Going global: Spin^c -structures and spinors

Spin^c structures are weaker than spin structures and exist on a broader class of four-manifolds.

2.6.3 The complex spin connection and the coupled Dirac operator

The coupled Dirac operator is the analytic object appearing in the Seiberg-Witten equations.

2.7 Final ingredient: The squaring map

2.7.1 The linear squaring map

The squaring map sends spinors to self-dual two-forms in a way compatible with the spin representation.

2.7.2 The global squaring map

Globally, this construction produces the nonlinear term in the Seiberg-Witten equations.

Chapter 3

The Seiberg-Witten Equations and Moduli Space

WE ARE FINALLY ready to define the Seiberg-Witten equations and their moduli space. Let M be a closed oriented Riemannian four-manifold with a fixed Spin^c structure. The equations are written for a spinor and a compatible connection.

3.1 The gauge group and its action

[Gauge group] For a principal $U(1)$ -bundle $P \rightarrow M$, the gauge group is

$$\mathcal{G} := \text{Aut}_{U(1)}(P).$$

In many settings this may be identified with $C^\infty(M, U(1))$.

3.2 Topology of the Moduli Space

[Seiberg-Witten equations] The Seiberg-Witten equations for a spinor ψ and a connection A are

$$\not{D}_A \psi = 0, \quad F_A^+ = \sigma^+(\psi).$$

[Moduli space] The Seiberg-Witten moduli space is the quotient

$$\mathcal{M} = \{(\psi, A) \mid \not{D}_A \psi = 0, F_A^+ = \sigma^+(\psi)\} / \mathcal{G}.$$

3.2.1 Dimension of $\mathcal{M}$

The expected dimension is computed from an elliptic deformation complex and an index theorem.

3.2.2 Generic smoothness

For generic perturbations, the moduli space is smooth under suitable hypotheses.

3.2.3 Orientation

Orienting the moduli space requires orienting a determinant line associated to the deformation complex.

3.2.4 Compactness

Compactness follows from a priori estimates and elliptic regularity.

3.3 The Seiberg-Witten invariant

When the moduli space has the correct dimension, one can extract invariants of the smooth structure of the underlying four-manifold.

Chapter 4

The Intersection Form and Donaldson's Theorem

THE INTERSECTION form is one of the most important algebraic invariants of a closed oriented four-manifold. It connects embedded surfaces, cup products, Poincaré duality, and the classification of manifolds.

4.1 The intersection form of 4-manifolds

[Intersection form] Let M be a closed, connected, oriented four-manifold. The intersection form is the symmetric bilinear pairing

$$q_M : H_2(M, \mathbb{Z}) \times H_2(M, \mathbb{Z}) \rightarrow \mathbb{Z}$$

defined by

$$q_M(\sigma, \eta) = \langle \text{PD}(\sigma) \smile \text{PD}(\eta), [M] \rangle.$$

If $\alpha, \beta \in H_2(M, \mathbb{Z})$ are represented by embedded surfaces $S_\alpha, S_\beta \subset M$, then

$$q_M(\alpha, \beta) = S_\alpha \cdot S_\beta.$$

4.1.1 Unimodular symmetric forms

Unimodular symmetric forms may be definite or indefinite, even or odd. Their arithmetic properties determine much of the topology of simply connected four-manifolds.

4.1.2 The topology of four-manifolds

[Freedman] Every unimodular symmetric bilinear form occurs as the intersection form of a closed, oriented, simply connected topological four-manifold, with the usual uniqueness distinction between even and odd forms.

4.2 Finale: Donaldson's theorem

[Donaldson] Let M be a smooth, closed, oriented, simply connected four-manifold. If the intersection form of M is definite, then it is diagonalizable over $\mathbb{Z}$.

This theorem demonstrates that the smooth category is much more rigid than the topological category. Seiberg-Witten theory provides a powerful route to constraints of this kind.