

## Assignment

August 27, 2024

1. The position vectors of points  $P$  and  $Q$  are  $\vec{p}$  and  $\vec{q}$  respectively. The point  $R$  divides the line segment  $PQ$  in the ratio  $3 : 1$  and  $\vec{S}$  is the midpoint of the line segment  $PR$ . The position vector of  $\vec{S}$  is:

- (a)  $\frac{\vec{p}+3\vec{q}}{4}$
- (b)  $\frac{\vec{p}+3\vec{q}}{8}$
- (c)  $\frac{5\vec{p}+3\vec{q}}{4}$
- (d)  $\frac{5\vec{p}+3\vec{q}}{8}$

2. For the matrix  $A = \begin{bmatrix} 2 & -1 & 1 \\ \lambda & 2 & 0 \\ 1 & -2 & 3 \end{bmatrix}$  to be invertible, the value of  $\lambda$  is:

- (a) 0
- (b) 10
- (c)  $\mathbb{R} - \{10\}$
- (d)  $\mathbb{R} - \{-10\}$

3. The angle which the line  $\frac{x}{1} = \frac{y}{-1} = \frac{z}{2}$  makes with the positive direction of the  $y$ -axis is:

- (a)  $\frac{5\pi}{6}$
- (b)  $\frac{3\pi}{4}$
- (c)  $\frac{5\pi}{4}$
- (d)  $\frac{7\pi}{4}$

4. The Cartesian equation of the line passing through the point  $(1, -3, 2)$  and parallel to the line:

$$\vec{r} = (2 + \lambda)\hat{i} + \lambda\hat{j} + (-1 + \lambda)\hat{k} \quad (1)$$

is:

- (a)  $\frac{x-1}{2} = \frac{y+3}{0} = \frac{z-2}{-1}$
- (b)  $\frac{x+1}{1} = \frac{y-3}{1} = \frac{z+2}{2}$
- (c)  $\frac{x+1}{2} = \frac{y-3}{0} = \frac{z+2}{-1}$
- (d)  $\frac{x-1}{1} = \frac{y+3}{1} = \frac{z-2}{2}$

5. If  $A = \begin{bmatrix} x & 0 \\ 1 & 1 \end{bmatrix}$  and  $B = \begin{bmatrix} 4 & 0 \\ -1 & 1 \end{bmatrix}$  then the value of  $x$  for which  $A^2 = B$  is:

- (a)  $-2$
  - (b)  $2$
  - (c)
  - (d)  $2$  or  $-2$
  - (e)  $1$
6. Given a curve  $y = 7x - x^3$  and  $x$  increases at the rate of 2 units per second, the rate at which the slope of the curve is changing when  $x = 5$  is:
- (a)  $-60$  units/sec
  - (b)  $60$  units/sec
  - (c)  $-70$  units/sec
  - (d)  $-140$  units/sec
7. Let  $f(x) = \begin{bmatrix} x^2 & \sin x \\ p & -1 \end{bmatrix}$  where  $p$  is a constant. The value of  $p$  for which  $f'(0) = 1$  is:
- (a)  $R$
  - (b)  $1$
  - (c)  $0$
  - (d)  $-1$
8. If  $A$  and  $B$  are events such that  $P(A/B) = P(B/A) \neq 0$ , then:
- (a)  $A \subset B$ , but  $A \neq B$
  - (b)  $A = B$
  - (c)  $A \cap B = \emptyset$
  - (d)  $P(A) = P(B)$
9. A function  $f : \mathbb{R} \rightarrow \mathbb{R}$  defined as  $f(x) = x^2 - 4x + 5$  is:
- (a) injective but not surjective
  - (b) surjective but not injective
  - (c) both injective and surjective
  - (d) neither injective nor surjective
10. If  $A$  is a square matrix of order 3 such that the value of  $|\text{adj} A| = 8$ , then the value of  $|A^T|$  is:
- (a)  $\sqrt{2}$
  - (b)  $-\sqrt{2}$
  - (c)  $8$

- (d)  $2\sqrt{2}$
11. If  $\int_{-2}^3 x^2 dx = k \int_0^3 x^2 dx + \int_2^3 x^2 dx$ , then the value of  $k$  is:
- (a) 2  
(b) 1  
(c) 0  
(d)  $\frac{1}{2}$
12. The value of  $\int_1^0 \log x dx$  is:
- (a) 0  
(b) 1  
(c)  $e$   
(d)  $e \log e$
13. The area bounded by the curve  $y = \sqrt{x}$ , the  $y$ -axis, and between the lines  $y = 0$  and  $y = 3$  is:
- (a)  $2\sqrt{3}$   
(b) 27  
(c) 9  
(d) 3
14. The order of the differential equation:

$$\frac{d^3 y}{dx^3} + x \left( \frac{dy}{dx} \right)^5 = 4 \log \left( \frac{d^4 y}{dx^4} \right)$$

is:

- (a) not defined  
(b) 3  
(c) 4  
(d) 5
15. If the inverse of the matrix  $\begin{bmatrix} 7 & -3 & -3 \\ -1 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}$  is the matrix  $\begin{bmatrix} 1 & 3 & 3 \\ 1 & \lambda & 3 \\ 1 & 3 & 4 \end{bmatrix}$ , then the value of  $\lambda$  is:
- (a) -4  
(b) 1  
(c) 3  
(d) 4

16. Find the matrix  $A^2$ , where  $A = [a_{ij}]$  is a  $2 \times 2$  matrix whose elements are given by  $a_{ij} = \max(i, j) - \min(i, j)$ :
- (a)  $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$
- (b)  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$
- (c)  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$
- (d)  $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$
17. Derivative of  $e^{\sin^2 x}$  with respect to  $\cos x$  is:
- (a)  $\sin x e^{a \sin^2 x}$
- (b)  $\cos x e^{\sin^2 x}$
- (c)  $2 \cos x e^{\sin^2 x}$
- (d)  $-2 \sin^2 x \cos x e^{\sin^2 x}$
18. The function  $f(x) = \frac{x}{2} + \frac{2}{x}$  has a local minimum at  $x$  equal to:
- (a) 2
- (b) 1
- (c) 0
- (d) -2

## Assertion - Reason Based Questions

**Direction:** In questions numbers 19 and 20, two statements are given: one labeled Assertion (A) and the other labeled Reason (R). Select the correct answer from the following options:

- (A) Both Assertion (A) and Reason (R) are true and the Reason (R) is the correct explanation of the Assertion (A).
- (B) Both Assertion (A) and Reason (R) are true and Reason (R) is not the correct explanation of the Assertion (A).
- (C) Assertion (A) is true, but Reason (R) is false.
- (D) Assertion (A) is false, but Reason (R) is true.
19. **Assertion (A):** Domain of  $y = \cos^{-1}(x)$  is  $[-1, 1]$ .  
**Reason (R):** The range of the principal value branch of  $y = \cos^{-1}(x)$  is  $[0, \pi] - \left\{\frac{\pi}{2}\right\}$ .