

Title: Pharmacy Dosing Service – Intravenous to Enteral Therapy Conversion Program

CARE BRAVELY

Department: Pharmacy

Document Owner: Name

Approver(s): Lisa Polinsky, AVP Pharmacy Services

Original: 01/11/2022

Updated: 04/02/2024

Expiration Date: 01/31/2025

Site(s):

- | | |
|--|--|
| ☒ Sinai Hospital of Baltimore | ☐ Practice Dynamics |
| ☒ Northwest Hospital Center | ☐ Baltimore Child Abuse Center |
| ☒ Carroll Hospital Center | ☐ BridgingLife |
| ☒ Levindale Hebrew Geriatric Center and Hospital | ☐ William E. Kahlert Regional Cancer Center |
| ☒ Grace Medical Center | ☐ LifeBridge Health & Fitness |
| | ☐ Carroll Health Group |

Appendix A: Medications Approved for Pharmacist IV to Enteral Conversion

Medication	IV Medication	Enteral Medication+
Azithromycin	250 – 500 mg IV Q24H	250 – 500 mg PO Q24H
Ciprofloxacin	400 mg IV Q12H 400 mg IV Q8H	500 mg PO Q12H 750 mg PO Q12H
Clindamycin	600 mg IV Q8H	300 to 450 mg PO Q6-8H 600 mg PO Q8H* (*dosing conversion for following indications: anthrax, barbesiosis, pneumonia due to MRSA, septic arthritis)
Dexamethasone	Total daily dose and frequency are the same (1:1 conversion)	
Doxycycline	100 mg IV Q12H	100 mg PO Q12H
Famotidine	20 mg IV Q12H 40 mg IV Q24H	20 mg PO Q12H 40 mg PO Q24H
Fluconazole	200 – 400 mg IV Q24H	200 – 400 mg PO Q24H
Folic Acid	1mg IV Q24H	1mg PO 24H
Lacosamide	Total daily dose and frequency are the same (1:1 conversion)	
Levetiracetam	Total daily dose and frequency are the same (1:1 conversion)	
Levofloxacin	250 – 750 mg IV, frequency	250 – 750 mg PO, same frequency
Levothyroxine	IV Q24H	PO Q24H *IV dose is half the enteral dose E.g.: 0.1 mg PO equals 0.05 mg IV
Linezolid	600 mg IV Q12H	600 mg PO Q12H
Metronidazole	500 mg IV Q8H	500 mg PO Q8H
Pantoprazole <i>(*Sinai/Northwest/ Carroll/Levindale)</i>	40 mg IV Q24H	40 mg <u>PO</u> Q24H Esomeprazole 40 mg capsule <u>NG/G(PEG)-tube</u> Q24H *Lansoprazole SoluTab 30 mg <u>J-tube</u> Q24H
Thiamine	100mg IV	100mg PO
Valproic Acid	Total daily dose conversion for oral and IV is 1:1. Oral doses are generally given twice daily [^]	

+ medication may be given through oral or enteral tube route.

* see procedure 3a and 3b in the policy for inclusion and exclusion criteria

[^] Valproic acid: If converting from oral to IV dosing, total IV daily dose should be divided for Q 6-8hr administration. Review home medication history and Micromedex for dosing interval consideration.