



A review of chromite mining in Sukinda Valley of India: impact and potential remediation measures

Suman Nayak, Rangabhashiyam S, Balasubramanian P & Paresh Kale

To cite this article: Suman Nayak, Rangabhashiyam S, Balasubramanian P & Paresh Kale (2020): A review of chromite mining in Sukinda Valley of India: impact and potential remediation measures, International Journal of Phytoremediation, DOI: [10.1080/15226514.2020.1717432](https://doi.org/10.1080/15226514.2020.1717432)

To link to this article: <https://doi.org/10.1080/15226514.2020.1717432>



Published online: 07 Feb 2020.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

A review of chromite mining in Sukinda Valley of India: impact and potential remediation measures

Suman Nayak^a, Rangabhashiyam S.^b, Balasubramanian P.^a, and Paresh Kale^c

^aDepartment of Biotechnology and Medical Engineering, National Institute of Technology Rourkela, Rourkela, India; ^bSchool of Biotechnology, Sastra University, Thanjavur, India; ^cDepartment of Electrical Engineering, National Institute of Technology Rourkela, Rourkela, India

ABSTRACT

Sukinda Valley, one of the highly polluted areas of the world is generating tons of mining waste and causing serious health and environmental issues in its surroundings. Several reports are available reporting the severity of hexavalent chromium, yet little efforts have been made to address the pollution and its remediation due to a lack of proper remedial measures. The review highlights the pros and cons of various physical, chemical and biological techniques used worldwide for the treatment of chromium waste and also suggests better and reliable bioremediation measures. Microbes such as *Acidophilium* and *Acidithiobacillus caldus* (Bioleaching), *Pseudomonas*, *Micrococcus* and *Bacillus* (Bioreduction), *Aerobacterium* and *Saccharomyces* (Biosorption), are widely used for bioremediation of hexavalent chromium owing to their unique metabolic activities, ionic movement through an extracellular membrane, and other cellular adsorptions and reduction properties. The use of native and hybrid combinations of microbes supported by organic supplements is projected as a fast and efficient technique that not only reduces chromium quantity but also maintains the integrity of the microbial sources. Innovation and emphasis on nano-based products like nanocomposite, nano adsorbent, nanoscale zerovalent iron (nZVI) particles and multifunctional plant-growth-promoting bacteria (PGPB) will serve as the next generation environmental remediation technologies in the near future.

KEYWORDS

Bioremediation; chromite mining; nanotechnology; nanosized zerovalent iron; phytoremediation; Sukinda Valley; treatment technologies

Introduction

Mining is one of the most ancient practices carried out across different parts of the world since the growth of human civilization (Hartman and Mutmansky 2002). With ever-increasing demands for minerals and advances in ore-extraction techniques, it had only escalated with times. Currently, mining activities cover more than 2% of the earth's surface in the form of opencast and underground mining (Coelho *et al.* 2011). Mining can be considered as a "Necessary Evil" as on one side it provides several benefits like employment, raw materials for industries, petroleum, natural gas, petrochemicals, and precious metals/stones, but is also a significant source of environmental pollutions, adverse health effects and biodiversity loss (Jaishankar *et al.* 2014). The use of nonscientific mining activities has further added to its severity leading to degradation of land, disturbance of water table, topographic disorder, and alteration to land use patterns in and around mining sites (Bridge 2004; Saviour 2012).

Chromium is a vital metal occurs significantly in the earth's crust, and in trace quantities ($<1,000 \mu\text{g g}^{-1}$) on rocks and soils surface. Naturally, chromium occurs in water as a result of weathering and erosion processes or even due to windblown dust particles. Chromite, the core ore of the

element chromium is an oxide of chromium (Cr) and Iron (Fe) containing about 68% of Cr_2O_3 and 32% of FeO with Cr to Fe ratio of about 8:1. Thus, Cr_2O_3 is the major ore mineral of chromium and is an accessory phase in many ultramafic rocks (Matrosova *et al.* 2020). As per the World Mining Data, the world's chromite production is about 13 billion metric tonnes, enough for human usage for hundreds of years (Reichl *et al.* 2016). Currently, mining activities cover more than 2% of the earth's surface as opencast and underground mining, of which 90% of the resources are in South Africa, Kazakhstan, Turkey, India, and Finland as shown in Figure 1a (Coelho *et al.* 2011). India, with a geographical area of about 328 million hectares has 0.14% of its land use for mining activities (Ministry of Mines 2017). India ranked 4th in terms of volume of production from mining sources and stands 3rd for the production of chromite ores in the world (Soni and Vyas 2015).

Sukinda Valley in Odisha of India produces about 99% of the total chromite ores of the country. Karnataka contributes 0.6%, while states like Nagaland, Manipur, Maharashtra, Jharkhand, Tamil Nadu, and Andhra Pradesh cumulatively for the rest of 0.4% (Indian Bureau of Mines 2016) (Figure 1b). The Sukinda Valley spread in over 200 km^2 holds an estimated 183 million tons of raw chromium deposits (Das *et al.*




Figure 1. (a) Country-wise chromium production (%) during 2016–17 (U.S. Geological Survey, 2018), (b) Chromium production (%) in India during 2016–17 (Indian Bureau of Mines 2016).

Table 1. Occurrence of chromium and its industrial applications.

Forms of Cr	Sources	Applications	Ref.
Cr(0)	Does not occur naturally	Primarily used in making steel and other alloys	(De Oliveira Andrade and Mattje 2012)
Cr(II)	CrBr ₂ , CrCl ₂ , CrF ₂ , CrSe, Cr ₂ Si	Used in chromizing and in the preparation of chromium metal and internal medicine	(Choppala <i>et al.</i> 2013)
Cr(III)	CrB, CrB ₂ , CrBr ₃ , CrCl ₃ ·6H ₂ O, CrCl ₃ , CrF ₃ , CrN, KCr(SO ₄) ₂ ·12H ₂ O, FeCr ₂ O ₄	Widely applied in tanning, green paints, inks, text dyes, and ceramics	(Zayed and Terry 2003)
Cr(IV)	CrO ₂ , CrF ₄	Data tape applications for enterprise-class storage systems and magnetic tape emulsion	(De Oliveira Andrade and Mattje 2012)
Cr(V)	CrO ₄ ³⁻ , KCr ₃ O ₈	Mutagenic and carcinogenic agent	(Zayed and Terry 2003)
Cr(VI)	(NH ₄) ₂ CrO ₄ , BaCrO ₄ , CaCrO ₄ , K ₂ CrO ₄ , K ₂ Cr ₂ O ₇ , PbCrO ₄ . Wolman compounds from NaF and C ₆ H ₄ N ₂ O ₅ with Na ₂ Cr ₂ O ₇ /K ₂ Cr ₂ O ₇ , CrCuO ₄ , copper–chromium–arsenic (CCA) formed by combining CuSO ₄ , Na ₂ Cr ₂ O ₇ and As ₂ O ₃ , composites of copper–chromium–boron (CCB), copper–chromium–fluoride (CCF) and copper–chromium–phosphate (CCP)	Used as preservatives for chrome plating, wood, leather and as coolant	(Saha and Orvig 2010)

2013). Currently, 20 opencast and two underground mines are operational in the valley generating about 160 million tons (MT) of overburden. It also releases 11.73 tons of Cr(VI) to the environment every year, making it the fourth-worst polluted places in the world. Chromium mining in Sukinda Valley had, therefore, resulted in major biodiversity loss and severe health impact for locals human population inhabiting around the belt (Dhal *et al.* 2011; Mishra and Sahu 2013; Biswas *et al.* 2017; Das and Singh 2011; U.S. Geological Survey (USGS) 2015).


Chromium is an industrial metal used in the manufacturing of stainless steel and metallurgy, chemical processing and several other commercial products (Table 1). However, considering its large-scale release to the environment and the fact that it is also a hazardous material causing numerous health disorders, an adaptation of proper remediation measures is the need of the hour. Hence, the study is an attempt to underline the impact of chromium on human health around Sukinda Valley, in India and highlights some of the potential bioremediation techniques.

Chromite pollution

There are about 35 metals found on Earth, of which 23 are heavy metals. Chromium can be found in six different forms and is readily released to the environment. It is of immense concern for human health due to their residential or occupational exposure (Jaishankar *et al.* 2014). The most significant and hazardous impact of chromium is from the mining projects. It affects the soil, air and water resources within the mining sites through acid drainage, erosion of soils and mine wastes, and contaminant leaching (Sharma *et al.* 2012). Chromium enters the ecosystem through anthropogenic activities such as untreated industrial discharge, dumping of solid waste, electroplating, dye, and pigment manufacture, wood preservations, refractory wastes solar cells, and leather tanning industries, etc. (Sharma *et al.* 2012). The generation of chromium and its presence in different ecological niches (Air, water and soil) around Sukinda along with their permissible limits set by various health agencies across the globe are listed in Table 2. The potential hazardous impacts

Table 2. Chromium concentration in Sukinda Valley and their permissible limits.

Components	Maximum permissible limits of Cr	Observed Chromium conc. in Sukinda Valley	Units	References
Air	100 as Cr and 10 in an insoluble form	130	$\mu\text{g m}^{-3}$	(Ministry of Mines, Government of India 2013)
Dust (Work zone)	120	20000		(Ministry of Mines, Government of India 2013)
Workplace	10	52–187		(WHO 2012; Mishra <i>et al.</i> 2016)
Fresh water	0.0001–0.05	5–11	mg l^{-1}	(WHO 2012)
Ground water	<0.001	0.023–1.35		(WHO 2012)
Surface water	0.001–0.01	0.012–3.4		(Mishra and Sahu 2013; Naz <i>et al.</i> 2016)
River water-sediment	200	30–104		(Ministry of Mines, Government of India 2013)
Soil	10–50	47	mg kg^{-1}	(Ertani <i>et al.</i> 2017)
Paddy fields soil segments	350	6–190		(Ministry of Mines, Government of India 2013)
Paddy	1	0.001–142		(Ministry of Mines, Government of India 2013; Qu <i>et al.</i> 2015)
Fish	0.15	3–37.93		(WHO 2012)
Milk	0.01	<0.001–0.69		(Ministry of Mines, Government of India 2013)
Fruits and vegetables	2.3 (dry weight)	5–28		(Eslami <i>et al.</i> 2007; Ministry of Mines, Government of India 2013)

**Figure 2.** Various environmental impacts of chromite mining.

of toxic Cr forms on all aspects of biodiversity are described in Figure 2.

Chromium in all of its forms is highly toxic and is associated with several diseases in humans and plants. Cr(VI) causes respiratory and mucous irritant related diseases and disorders that include irritation of nose and throat, asthma, ulcer in lungs, and perforation (holes) in the septum. The minimal exposures about $0.09\text{--}2\ \mu\text{g m}^{-3}$ of Cr can cause ulcers, septal perforations, and reversible impairment of lung function (WHO 2012; Ertani *et al.* 2017).

Workers in mines and Cr-based industries are exposed to airborne Cr(VI) with high exposures frequently occurring *via* inhalation and dermal contact (Ashley *et al.* 2003). The Odisha Voluntary Health Association reported that 86.42% of deaths in villages surrounding Sukinda Valley are results of excess exposure to Cr(VI) (Das and Singh 2011). While the presence of Cr in the atmosphere is a serious issue,

pollutants contaminating the water sources around Sukinda Valley, *i.e.*, the Dharamsala and the Brahmani rivers regarded as more severe (Das and Singh 2011; Mishra and Sahu 2013).

The soil quality around the Sukinda Valley is slightly better than the other environmental variables (air and water). Chromium concentration in soil and segments of crop fields (paddy) in the valley was lower than their legitimate values of $47\ \text{mg kg}^{-1}$ and $6\text{--}190\ \text{mg kg}^{-1}$ (Ministry of Mines, Government of India 2013; Ertani *et al.* 2017). However, an elevated level of soluble Cr, bioavailable of Cr (III) and other forms can dent the soil quality and its infertility (Jaishankar *et al.* 2014; Reijonen and Hartikainen 2016).

Paddy grains from contaminated sites in the Sukinda region were reckoned to exceed the WHO recommended limit ($1\ \text{mg kg}^{-1}$) by a margin of $16\text{--}740\ \text{mg kg}^{-1}$. Similarly, milk, fruits, and other vegetables were found to contain up




Figure 3. Flowchart representation of some of the most commonly used remediation techniques.

to 5 times higher amount of chromium than their consumable permits (<0.001 – 0.69 mg/kg for milk, 10 – 112 mg kg^{-1} for fruits and vegetables) (Eslami *et al.* 2007; Ministry of Mines, Government of India 2013). In plants, excess quantity of Cr is known to have an impact on plant's metabolism and development resulting in dwarf growth, chlorosis, low crop yield, deferred germination period, senescence, premature leaf fall, abrasion, alter enzymatic and biosynthesis production of growth hormones (Zayed and Terry 2003; Mohanty *et al.* 2009). Cr(VI) is also responsible for carcinogenic effects in human leading to ulcers in stomach and lung, and other health ailments like diarrhea, bronchospasm, pneumonia, skin irritation and gastrointestinal diseases (Dubey *et al.* 2001; Beaumont *et al.* 2008; Das and Singh 2011)

Treatment methods for Cr (VI) pollution


To date, several methods or techniques had been developed to curb pollution caused by chromium. Some of the commonly employed techniques for treatment of Cr waste includes reduction precipitation (Enniya *et al.* 2018), membrane filtration (Yang *et al.* 2003), membrane ion exchange (Yang *et al.* 2014) biological adsorption and reduction (Sahinkaya *et al.* 2012) (Figure 3). Factors like the mobility, distribution, concentration, and speciation of chromium in the sources are known to govern the choice of the best-suited method for remediation of Cr (Zayed and Terry 2003).

Physical and chemical remediation processes though were once primarily used for reduction of Cr heavy metals, problems associated with it such as high capital investment and generation of large volume of secondary waste had restricted their use in recent times (Dhal *et al.* 2013; Pradhan *et al.*

2017; Xia *et al.* 2019). Physicochemical treatment technology such as adsorption is quite implemented in many instances. However, various factors such as adsorbent characteristics and experimental conditions (equilibration time, solution pH, initial Cr(VI) concentration, ionic strength) play key role in determining the overall treatment efficiency (Islam *et al.* 2019). Biological remediation measures are time taking although; considering its eco-friendly approach and easy bioavailability of host sources, it is the first choice for chromium treatment. However, bioremediation is still a challenging technique and most studies remain at the laboratory stage to date (Xia *et al.* 2019). Some of the commonly employed biological approaches for remediation of Cr heavy metals are discussed in this section.

Biobleaching

Biobleaching is a biochemical process that uses both chemical substances and environmental microbes for the extraction of metal ions (Figure 4a). Conversely, with some modification, the process now involves the pretreatment of waste materials with acid and subsequently with iron- and sulfur-oxidizing microbes native to the sludge. Biobleaching can be achieved through direct leaching and indirect leaching. For direct leaching, non-soluble metal sulfides are oxidized to their water-soluble forms by chemolithotrophic bacteria while in the indirect route, elemental sulfur (S) is oxidized first to sulfuric acid by sulfur-oxidizing bacteria followed by the dissolution of metals by strong acid. The equations are as follows:







Figure 4. (a) Bioleaching of heavy metals by using microorganisms, (b) Microbial reduction of chromium.



Parameters that influence the leaching efficiency of Cr are contact time, slag/sludge composition, the concentration of competing ions, and type and concentration of leaching agents (Rastegar *et al.* 2014). Bioleaching of Cr heavy metal via *Thiobacillus neapolitanus*, *Alicyclobacillus ferripilum*, *A. pomorum*, *A. ferrooxydans*, *A. aeris* (Bharadwaj and Ting 2013), *Acidithiobacillus thiooxidans*, *Bacillus sp.* (Chen and Lin 2010), *Clostridium sp.*, and *Thiobacillus thioparus* (Zeng *et al.* 2016) have successfully been reported for different forms of industrial waste.

Recent development has highlighted the use of mixture or hybrid combination of native microbial strains supported by organic supplementary as most competent for bioleaching processes (Pathak *et al.* 2008a; Dhal *et al.* 2013). These efficiencies by mixed or hybrid combinations are credited to the availability of diverse functional groups present in their cell walls and other metabolic cycles. Microbial bioleaching using *Thiobacillus sp.* and *Acidithiobacillus thiooxidans* had shown leaching efficiencies as high as 95% in a reducible fraction, and 60.9–97.9% in oxidizable and residual fractions (Zeng *et al.* 2016). Biosurfactant from plant sources like aescin and saponin in the aqueous medium is reported to have a catalytic effect in leaching of heavy metal from contaminated soil (Hong *et al.* 2002).

Bioreduction

Hexavalent chromium is regarded as more toxic compared to trivalent chromium as the former has a labile center and bears structural similarity with sulfate, thus can easily penetrate into the living cell. Further in solution trivalent chromium undergoes precipitation as hydroxide, but the hexavalent chromium due to its inert nature remains stable (Saha *et al.* 2011). Thus, the reduction of Cr(VI), which is highly water-soluble and mobile into a less toxic form is essential for reducing its harmful effects. Bioreduction of chromium can be defined as the process of breaking down of chromium molecules from its hexavalent Cr(VI) forms into less toxic trivalent (Cr III) state (Figure 4b). The traditional method was the treatment of Cr(VI) with acids followed by precipitation with alkaline ferrous sulfate, sodium

sulfite, sulfur dioxide gas or sodium bisulfite and are still in use in some studies (Barrera-Díaz *et al.* 2012; Lu *et al.* 2019). The reduction mechanism for chromium, (*i.e.*,) traditional, electrocoagulation and microbial are summarized in Table 3.

Although traditional and electrocoagulation reduction are quite successful in the reduction of Cr, primary constraints associated with the methods are a large amount of residual sludge, secondary pollutants, etc. Microbial bio-reduction is preferred over other reduction measures as they are saved to the environment, easy and requires less effort. It is carried out mostly by the enzyme chromate reductases ChrR and YieF commonly found in *Aeromonas*, *Escherichia*, *Pseudomonas*, *Bacillus*, and *Enterobacter* (Chen *et al.* 1998). The alkalic amino acids present at the loop of the enzyme provide sites for binding of NAD(P)H and chromate ions. Under aerobic condition Cr(VI) reduces to Cr(III) through a soluble reductase enzyme that serves as an electron donor for NAD(P)H. Whereas in anaerobic conditions, the reduction of Cr(VI) takes place during the electron transport pathway of cytochrome (Cyt B & C) along the respiratory chains inside the intracellular membrane of the microbial cells (Ahemad 2014).

In aerobic conditions, ChrR catalyzes the reduction of Cr(VI) in two or three stages. Initially, a one-electron shuttle reduces Cr(VI) to a short-lived intermediate of Cr(V) or Cr(IV) followed by a two-electron transfer to reduce it to the thermodynamically stable product Cr(III). The reductase YieF from *E. coli* reduces Cr(VI) directly to its trivalent form by transferring four electrons, wherein three electrons are used up in reducing Cr(VI), while one electron is added to oxygen to produce hydrogen peroxide through hydrogen bonding (Chen *et al.* 1998; Ackerley *et al.* 2004; Zou *et al.* 2013).

The anaerobic reduction of Cr is carried out by Sulfate-reducing bacteria (SRB) *Desulfovibrio vulgaris*. In this conditions, Cr(VI) serves as a terminal electron acceptor in the respiratory chain for a vast array of electron donors, including nitroreductase, flavoprotein, glutaredoxin, NAD(P)H and endogenous electron reserves (Ibrahim *et al.* 2012). Sulfide produced by cytochromes of SRB almost entirely reduced Cr(VI) to Cr(III) and precipitated other metal ions (Losi and Frankenberger 1994). SRB's ability to reduce

Table 3. Different Bio-reduction methods of Cr (VI) and their mechanism.

Methods of Cr(VI) reduction	pH	Multi-step mechanism	Ref.
Traditional	2.5	Step-I: 3 kg of sodium bisulfite added with 2–3 kg of sulfuric acid, reduces 1 kg of Cr(VI) to its trivalent form $4\text{CrO}_4^{2-}(\text{aq}) + 6\text{NaHSO}_3(\text{aq}) + 3\text{H}_2\text{SO}_4(\text{aq}) 8\text{H}^+(\text{aq}) \rightarrow 2\text{Cr}_2(\text{SO}_4)_3(\text{aq}) + 3\text{Na}_2\text{SO}_4(\text{aq}) + 10\text{H}_2\text{O}$ Step-II: Further addition of sodium hydroxide precipitates Cr (III) as indicated as $\text{Cr}_2(\text{SO}_4)_3(\text{aq}) + 6\text{NaOH}(\text{aq}) \rightarrow 2\text{Cr}(\text{OH})_3(\text{s}) + 3\text{Na}_2\text{SO}_4(\text{aq})$ Step-III: After precipitation of Cr (III) hydroxide, the resulting effluent will contain little or no residual chromium.	(Barrera-Díaz <i>et al.</i> 2012)
Electro coagulation	6–8	$4\text{CrO}_4^{2-}(\text{aq}) + 3\text{SO}_2(\text{g}) + 4\text{H}^+(\text{aq}) \rightarrow \text{Cr}_2(\text{SO}_4)_3(\text{aq}) + 2\text{H}_2\text{O}$ Step-I: Anodic reaction: $\text{Fe}(\text{s}) \rightarrow \text{Fe}^{2+}(\text{aq}) + 2\text{e}^-$ Step-II: Cathodic reaction: $2\text{H}_2\text{O} + 2\text{e}^- \rightarrow 2\text{OH}^-(\text{aq}) + \text{H}_2(\text{g})$ Step-III: Overall reactions in bulk solution $\text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 14\text{H}^+(\text{aq}) + 6\text{Fe}^{2+}(\text{aq}) \rightarrow 2\text{Cr}^{3+}(\text{aq}) + 6\text{Fe}^{3+}(\text{aq}) + 7\text{H}_2\text{O}$	(Barrera-Díaz <i>et al.</i> 2012)
Aerobic Bacterial reduction	2–4	Step – I : $\text{Cr}^{\text{VI}} + \text{e}^- \xrightarrow{\text{Bacteria}} \text{Cr}^{\text{V}}$ Step – II : $\text{Cr}^{\text{V}} + 2\text{e}^- \xrightarrow{\text{Bacteria}} \text{Cr}^{\text{III}}$	(Singh <i>et al.</i> 2011)
Anaerobic Bacterial reduction	7–9	Step – I : $\text{CrO}_4^{2-}(\text{aq}) + 8\text{H}^+(\text{aq}) + 3\text{e}^- \xrightarrow{\text{Bacteria}} \text{Cr}^{3+}(\text{aq}) + 4\text{H}_2\text{O}$ Step – II : $\text{Cr}^{3+}(\text{aq}) + 4\text{H}_2\text{O} \xrightarrow{\text{Bacteria}} \text{Cr}(\text{OH})_3(\text{s}) + 3\text{H}^+(\text{aq}) + \text{H}_2\text{O}$ Step – III : $\text{C}_6\text{H}_{12}\text{O}_6 + 8\text{CrO}_4^{2-}(\text{aq}) + 14\text{H}_2\text{O} \xrightarrow{\text{Bacteria}} 8\text{Cr}(\text{OH})_3(\text{s}) + 10\text{OH}^-(\text{aq}) + 6\text{HCO}_3^-(\text{aq})$	(Cheung and Gu 2007; Singh <i>et al.</i> 2011)


chromate is due to its structural similarities with chromate. The reaction kinetics of bioreduction as purposed by Kim *et al.* (2001), involves the formation of chromate sulfide intermediate complex $\{\text{H}_2\text{O}_4\text{Cr}^{\text{VI}}\text{S}\}^{2-}$, followed by intramolecular electron transfer to form Cr(IV) species, and subsequently fast reactions with Cr(III) as the final product. The energy kinetic required by ChrR reductase to reduce each molecule of Cr is given as K_m of 374 μM and V_{max} of 1.72 $\mu\text{mol}/\text{min}/\text{mg}$.

Step I:	$\text{H}_2\text{S} + \text{CrO}_4^{2-} \leftrightarrow \{\text{H}_2\text{O}_4\text{Cr}^{\text{VI}}\text{S}\}^{2-}$	(1)
	$\text{HS}^- + \text{CrO}_4^{2-} + \text{H}^+ \rightarrow \{\text{H}_2\text{O}_4\text{Cr}^{\text{VI}}\text{S}\}^{2-}$	(2)
Step II:	$\{\text{H}_2\text{O}_4\text{Cr}^{\text{VI}}\text{S}\}^{2-} \leftrightarrow \text{Cr}^{\text{V}}\text{O}_3^- + \text{S}^0 + \text{H}_2\text{O}$	(3)
	$\{\text{H}_2\text{O}_4\text{Cr}^{\text{VI}}\text{S}\}^{2-} + \text{H}_2\text{S} \leftrightarrow \text{Cr}^{\text{V}}\text{O}_3^- + \text{HSSH} + \text{H}_2\text{O}$	(4)
Step III:	$\text{Cr}^{\text{IV}} + \text{Cr}^{\text{VI}} \leftrightarrow 2\text{Cr}^{\text{V}}$	(5)
	$2\text{Cr}^{\text{IV}} \leftrightarrow \text{Cr}^{\text{III}} + \text{Cr}^{\text{V}}$	(6)
	$\text{Cr}^{\text{V}} + \text{S}^{0} \leftrightarrow \text{Cr}^{\text{III}} + \text{S}^0$	(7)

Nitroreductase, an inducible chromate reductase dimer, isolated from *Shewanella oneidensis* poses the potential for chromate reduction activity under both aerobic and anaerobic conditions. It uses a two-electron transfer mechanism and is induced by nitrate instead of chromate on being cultivated in a nitrite containing medium (Ackerley *et al.* 2004). Another novel enzyme oxidoreductase isolated from *Thermus scotoductus* is a flavoprotein capable of reduces chromate under similar conditions as nitroreductase (Opperman *et al.* 2010). However, its reduction efficiency increases to 180 times and 50 times than quinone reductase and nitroreductase respectively in the supply of excess oxygen.

In thermophilic methanogens, the extracellular reduction of Cr(VI) is attributed to the presence of ferredoxins-dependent reactions in the electron transport chain. While for intracellular it occurs through NADPH-dependent FMN reductase. The reduction process occurs as a result of diversion of electrons from H_2 to Cr(VI) instead of CO_2 to make CH_4 . Stoichiometric consumption of H_2 in the production

of Cr(III) supports the Electron diversion hypothesis. The enzyme exhibits a V_{max} value of 59.89 $\mu\text{M}/\text{min}/\text{mg}$ and K_m of 86.5 μM , signifying its feasibility in laboratory conditions when $\text{K}_2\text{Cr}_2\text{O}_7$ is used as a substrate (Mala *et al.* 2014; Singh *et al.* 2015). Mass balance studies by Singh *et al.*, (2015) on *Methanogens thermautotrophicus* suggest the Cr(VI) to cell ratio is also responsible for the reduction process. The calculation of stoichiometric ratio shows that for the reduction of every mole of Cr(VI), three moles of H_2 are required that produce two moles of stable Cr(III).



Incorporation of biogenic iron (Fe) (oxy-hydro)oxides (BIOS) with organic carboxylate complex was recently reported to enhance Cr(VI) adsorption and reduction (Whitaker *et al.* 2018). The study discovers that BIOS would substantially reduce Cr(VI) to Cr(III) as evident from X-ray absorption near edge structure (XANES) images. However, the lack of a 3:1 reaction stoichiometry between Fe(II) and Cr(III) suggests the roles of organic matter for auto reaction with Cr(V) and production of organo-Cr(III) complexes (Gong *et al.* 2018). The mass to product ratio indicates a higher accumulation of reduced Cr complexes on cell walls with time rather than inside of the cells. The organo-Cr(III) complexes formed by *Geobacter sulfur reducens* were also found to be more stable than the aqueous Cr(III) or $\text{Cr}(\text{OH})_3$ solution (Gong *et al.* 2018).

Biosorption

Biosorption is a passive and irreversible independent metabolic process performed by microorganisms (refer Figure 5). Adsorption of heavy metals by microbes can be ascribed to the interaction of metals ions with functional groups such as carboxyl, hydroxyl, sulfhydryl, phenolic and extracellular polymeric substances (EPS) that are present on the cell surface of different microbes (Dehghani *et al.* 2016; Enniya *et al.* 2018; Pakade *et al.* 2019). These anionic ligands along




Figure 5. Mechanisms of biosorption of heavy metals by microorganism: metabolic dependent and independent metabolic methods.

with D-galactosamine polymers and polyuronides are greatly considered as heavy metal-binding sites contributing to the biosorption processes in microbes (Vendruscolo *et al.* 2016). The amino groups present in the microalgal species such as *Chlorella miniata* and *Spirulina platensis*, are responsible for their biosorption potential (Han, Shan, *et al.* 2007).

In fungi, chitosan, a compound derived from deacetylated chitin is responsible for synergistic or antagonistic effects resulting in the absorption of Cr heavy metal. Species including but not limited to *Candida sp.* (Vendruscolo *et al.* 2016), *Paecilomyces lilacinus* (Sharma and Adholeya 2011), *Aspergillus sp.* (Rezaei 2016), *Rhizopus rhizus* (Arunbabu *et al.* 2017), *Penicillium sp.* (Zhang *et al.* 2017) are prominent in biosorption of Cr(VI).

Yeasts are a complex group of microbes capable of growing in various environments conditions. The species *Saccharomyces* has been an excellent absorbent agent for biosorption of Cr (Rezaei 2016). However, *Cyberlindnera fabianii* isolated from industrial effluent is by far the best-known yeast species to effectively remove 100% of chromium from after 48 h of treatment. Factors that directly impact biosorption are sources of energy (electron donors), nutrients, pH, temperature, and inhibitory substrates or metabolites concentration, etc. (Vendruscolo *et al.* 2016). According to the Box- Behnken design, the ideal parameters for biosorption is a biosorbent dose of 2 g l^{-1} , Cr(VI) concentration of 67.8 mg l^{-1} and a reaction time of 37.5 min (Vendruscolo *et al.* 2016). The optimum pH for biosorption in fungal species is pH 3.0, while for bacterial species it usually ranges from 7 to 9.

Other miscellaneous sources used as low-cost adsorbents include waste newspapers (Dehghani *et al.* 2016), sugarcane bagasse (Garg *et al.* 2007), activated carbon of apricot kernel (Demirbas *et al.* 2004), longan seed (Yang *et al.* 2015), immobilized sugarcane bagasse (Ullah *et al.* 2013), and animal charcoal (Hyder *et al.* 2015). Activated carbon is an emerging entrant for adsorption of Cr(VI) considering its high specific surface to spore area (Yang *et al.* 2015).

Phytoremediation

Phytoremediation, often referred to as “green remediation” is a comparatively new technique where plant species or materials are applied for *in-situ* reduction, detoxification and stability of pollutants from different sources (Figure 6). Phytoremediation is advantageous over conventional

remediation techniques because of cost efficiency, comfort, and large-scale applicability against a variety of contaminants while keeping the biota intact (Zayed and Terry 2003; Diwan *et al.* 2008; Shams *et al.* 2010).

Academically and applied researches on phytoremediation techniques has unambiguously confirmed that particular plant species on suitable conditions possess the genetic prospective to reduce, metabolize or immobilize an array of hazardous heavy metals from soil and water sources (Shams *et al.* 2010; Ahemad 2015). A model species for Cr (VI) remediation must be fast-growing, have higher biomass, easy to cultivate and harvest and amass high concentrations of Cr in aerial component (Diwan *et al.* 2008). Discovery of hyperaccumulators (ability to uptake bulk quantities of toxic metal pollutants) in plants such as *Spartina argentinensis* (Redondo-Gómez *et al.* 2011), *Leersia hexandra* (Zhang *et al.* 2007) and *Prosopis laevigata* (Buendía-González *et al.* 2010) have further boost remediation technique. The use of rooted and submerged plant species as phytoremediation tools have additional benefits as it helps in facilitate colonization of benthic algae, invertebrates, and other microbes (Zhu *et al.* 1999). Apart from aquatic plant species terrestrial trees, grasses, herbs, and shrubs are actively used in phytoremediation as listed in Table 4.

Pennisetum purpureum, *Brachiaria decumbens*, and *Phragmites australis* growing hydroponically over gravel beds are the most efficient plant confiscating 97–99.6% of Cr within 24 h (Mant *et al.* 2006). Stinging nettle (*Urtica dioica*), Rapeseed (*Brassica napus*), Indian mustard (*Brassica juncea*) and Maize (*Zea mays*) due to their versatile nature and unexplored possibilities against several toxic heavy metals had also been incorporated to the pot experiments to explore their sensitivity toward chromium (Shams *et al.* 2010).

Vallisneria spiralis growing on chromium enriched nutrient medium accumulated a significant amount of chromium in different plant parts like dry weight ($1,050 \mu\text{g g}^{-1}$), leaves ($697 \mu\text{g g}^{-1}$), and rhizomes ($437 \mu\text{g g}^{-1}$) in solution containing as many as $10.0 \mu\text{g ml}^{-1}$ Cr after 72 h of treatment (Vajpayee *et al.* 2001). Tumbleweed (*Salsola kali*) when provided with Cr (VI), was recorded to have a varied amount of Cr distribution in different parts as for roots ($2,900 \text{ mg/kg}$) stems 790 (mg/kg) and leaves (600 mg/kg) (Gardea-Torresdey *et al.* 2005).

Aquatic plants like *Eichhornia crassipes*, *Pistia stratiotes*, and *Salvinianatans* has grown on hexavalent Cr contaminated water have also confirmed experimentally to accumulate reduced Cr (III) in their leaves and roots (Sundaramoorthy *et al.* 2010). *Eichhornia crassipes* (water hyacinth) mount up the huge quantity of Cr in roots ($5,000$ to $6,000 \text{ mg/kg}$) on being supplied with 10 mg l^{-1} Cr for 14 days (Lytle *et al.* 1998). Poor translocation of Cr in shoots section in plants indicates that the obstruction in passes of Cr to any aerial part is not outwitted by supplying Cr as organic complexes or any other forms but is dependent on the concentration and length of exposure period (Diwan *et al.* 2008).




Figure 6. Phytoremediation of heavy metals.

Presence of Potassium (K) as an ingredient in salt solution ($\text{KCr}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$), was reported to have an unexpected positive response in growth of plants and uptake of chromium (Shams *et al.* 2010), while EDTA was found to serve as a hindrance agent to metal intake in hydroponic plants (Sundance sunflowers) (January *et al.* 2008). Genetic studies in willow species like *Salix alba*, *Salix eleagnos*, *Salix fragilis* and *Salix matsudana* had identified the gene “Expressed Sequence Tag” as an essential regulator in plants toward chromium exposure (Quaggiotti *et al.* 2007). Other reports suggest that an interrelated network of physiological biochemical and molecular processes can serve as an integral part, affecting the ability of a plant to accumulate heavy metals (Diwan *et al.* 2008).

Phytoremediation process, although are time-consuming but had proven to disintegrate and degrade chromium present in the soil. Bioinoculants (e.g., multifunctional plant-growth promoting bacteria (PGPB) along with plants) exhibit chromium resistance and reducing traits, providing a better approach for phytoremediation. PGPB improve Cr uptake ability of plants by modifying root architecture, secreting metal sequestering molecules in rhizomes, and lessen chromium-induced phytotoxicity (Ahemad 2015). The use of vermicompost prepared with cow dung using earthworm species *Eurdilus euginae* along with sorghum (*Sorghum bicolor*) was suggested as a potential source of bioremediation in Cd and Cr contaminated soil by Revathi *et al.* (2011).

Nanoremediation

Nanotechnology is a trending technology that had found its application in various fields including medical science, fabrics engineering, defense, space technology and many more.

Nanotechnology is also explored as possible remediation techniques with prominence in absorbance and reduction of Cr(VI) from industrial effluents when used as nano-adsorbents, nano-catalysts and nano-membranes (Lv *et al.* 2012; Mitra *et al.* 2017). Nanomaterials are an ideal candidate for environmental remediation following the characteristics features, i.e., small size, better solubility, large surface area, charges on surface area and varied surface composition (Lv *et al.* 2012; Kim and Roh 2019). Maitlo *et al.* (2019) reviewed the various nanomaterials for the sorptive removal of chromium from water.

The removal of Cr from contaminated water sources by nanoscale zerovalent iron (nZVI) particles employed on herb-residue biochar extracted from *Astragalus membranaceus* was credited to the presence of numerous porous structure and oxygen-rich functional groups present on its surface (Shang *et al.* 2017). ZVI (nZVI) has an advanced and better surface area ($182.97 \text{ m}^2 \text{ g}^{-1}$) enabling them to remove several pollutants like PCBs, TCE, and DDT, Pb(II) through reduction, adsorption and precipitation (Guan *et al.* 2015). Zerovalent iron nanoparticles, when support with resin showed Cr removal efficiency of 84.4% under optimal conditions (Fu *et al.* 2013). Further, the addition of carboxymethyl cellulose (CMC) as a stabilizer for nZVI improves the efficiency to 100% for solutions containing 10 mg/l of Cr (Wang *et al.* 2010). Polymer-nZVI composite film composed of multiple layers of chitosan, polyethylene glycol (PEG) blend, poly (sodium 4-styrene sulphonate) solution (PSS) and live biomass of Sulfate-reducing bacteria (SRB) had a Cr deduction rate of 394 mg/g under anaerobic conditions (Argulwar *et al.* 2018).

The novel nanocomposite of acid-treated rock wool zero-valent iron (ARWZ), when used as a membrane, is reported

Table 4. Bioremediation with their sources and other catalytic factors.

Techniques	Source of remediation	Type	Catalytic factors				Ref.
			pH	Temp. (°C)	Time (Days)	Others	
Bioleaching	<i>Acidithiobacillus thiooxidans</i>	Bacteria	2	30	–	–	(Nareshkumar et al. 2007)
	<i>Acidithiobacillus ferrooxidans</i>		1–2.5	–	–	Fe ³⁺	(Rastegar et al. 2014)
	<i>Geotrichum</i> sp.		–	–	7	Magnetic field and Aerobic condition	(Qu et al. 2016; Kim and Roh 2019)
	<i>Bacillus</i> sp.		(Qu et al. 2016)	–	–	–	–
Bioreduction	<i>Antridiavallantii</i>	Fungus	3	–	28	Citric acid, Oxalic acid	(Sierra-alvarez 2009)
	<i>Brettanomyces</i> sp.	Yeast	–	–	5	–	(Pathak et al. 2009)
	<i>Cellulosimicrobium</i> sp.	Bacteria	–	–	–	–	(Bharagava and Mishra 2018)
	<i>Bacillus</i> sp.		7	35	–	–	(Dhal et al. 2013)
	<i>Planococcusmaritimus</i>		7	35	1–1.16	–	–
	<i>Arthrobaacter</i> sp.		7	35	–	Glycerol and glucose	(Dey and Paul 2015)
	<i>Bacillus sphaericus</i>		–	32	–	120rpm	(Pal et al. 2013)
	<i>Bacillus</i> sp.		8	30	–	–	(Xiao et al. 2017)
	<i>Arthrobaacter</i> sp.		7	35	–	–	(Xiao et al. 2017)
	<i>Escherichia coli</i> Acineto bacteriumii		7.5	–	1.125	–	(Samuel et al. 2012; Talib et al. 2019)
	<i>Bacillus subtilis</i>		–	–	–	–	–
	<i>Enterobacter cloacae</i>		–	30	1	180 rpm	(Harish et al. 2012)
	<i>Pseudomonas</i> sp. (<i>P. aeruginosa</i> and <i>P. stutzeri</i>)		–	37	2	Acetate minimal medium	(Ozturk et al. 2012)
	<i>Bacillus</i> sp.		10	35	–	–	(Ibrahim et al. 2011)
Biosorption	<i>Pseudomonas aeruginosa</i>		–	–	3.5	Immobilized with polyvinyl alcohol, sodium alginate, and multivalled carbon nanotubes	(Pang et al. 2011)
	<i>Halomonas</i> sp.	Proteobacteria	–	37	2.25	–	(Murugavelh 2013)
	<i>Chlorella miniata</i>	Algae	2	–	2.41	–	(Han et al. 2017)
	<i>Aspergillus niger</i> , <i>Aspergillus flavus</i>	Fungus	2 to 9	22–40	1	–	(Park et al. 2005; Kumar and Dwivedi 2019)
	<i>Pichia guilliermondii</i>	Yeast	–	30	3–5	–	(Ksheminska et al. 2008)
	<i>Saccharomyces cerevisiae</i>		–	38	2	–	(Ksheminska et al. 2006)
	Activated carbon	Apple peels	2 to 7	10–40	0.16	–	(Enniya et al. 2018)
	<i>Cyberlindnera bialanii</i>	Yeast	–	–	2	–	(Joutey et al. 2015)
	<i>Aspergillus oryzae</i>	Fungus	5	30–35	–	–	(Kongeevaram et al. 2007)
	<i>Aspergillus niger</i> , <i>Aspergillus sydoni</i> and <i>Penicilliumjanthinellum</i>		2	–	–	–	(Kumar et al. 2008)
	<i>Aspergillus fumigatus</i> <i>Fresenius</i>		5.5	35	5	–	(Dhal et al. 2018)
	<i>Trichoderma</i> sp.		5.5	–	–	–	(Vankar and Bajpai 2008)
	<i>Micrococcus</i> sp.	Bacteria	5	30–35	–	–	(Congeevaram et al. 2007)
	<i>Pantoea</i> sp.		3	–	–	–	(Ozdemir et al. 2004)
Phytoremediation	<i>Spirulina</i> sp.	Algae	5	25	–	Agitation time – 120 min	(Rezaei 2016)
	<i>Cladophora</i> sp.		2	–	–	120 rpm	(Rangabhashiyam and Balasubramanian 2018a)
	<i>Spirogyra</i> species		2	18	0.083	–	(Gupta et al. 2001)
	<i>Chlorella miniata</i>		2	–	0.083	–	(Han, Mao, et al. 2007)
	<i>Sargassumwightii</i> and <i>Caulerparacemosa</i>		5	–	–	–	(Tamilselvan et al. 2012)
	<i>Portulacaoleracea</i>	Biomass	2	–	45min	–	(Shinghal 2015)
	<i>Salsola kali</i>	Herb	5.8	–	–	Agar-based medium	(Gardea-Torresdey et al. 2005)
	<i>Medicago sativa</i> and <i>Sorghum bicolor</i>		–	–	50	–	(Karimi 2013)
	<i>Helianthus annuus</i>		–	–	–	EDTA, Citric Acid	(Turgut et al. 2004)
	<i>Urtica dioica</i>		5.5 to 6.5	–	–	50– 300 mg/l N, 40 to 300 mg/l P ₂ O ₅ and 100–500 mg/l K ₂ O	(Shams et al. 2010)
	<i>Polygonumtuberosa</i>		–	–	–	–	(Ramana et al. 2012)
	<i>Brassica juncea</i>		7 to 7.8	–	125	–	(Diwan et al. 2008)
	<i>Pterocarpus indicus</i> and <i>Jatropha curcas</i>	Tree	–	–	–	–	(Mangkoedhardjo et al. 2008)
	<i>Pterosper mumacrerifolium</i> (shells)		2–10	29–49	–	phosphoric acid, Sulfuric acid	(Rangabhashiyam and Balasubramanian 2018b)
	<i>Pinus sylvestris</i>		7 to 1	25	–	–	(Uzun et al. 2002)
	<i>Penisetum purpureum</i> , <i>Brachiaria decumbens</i> and <i>Phragmites australis</i>	Grass	–	–	1	Gravel beds	(Mant et al. 2006)

to have an unexpectedly high removal capacity of Cr(VI) (197.69 mg/g) in 30 min from industrial effluents and soil through adsorption and reduction. The property is accredited to its vast face area and superior magnetic property (Zhou *et al.* 2018; Hou *et al.* 2019). Carbon nano-onions (CNOs) are materials used as substitute adsorbent to confiscate Cr from the water treatment industry. CNOs are highly sensitive to pH and are therefore dependent on the pH of the waste solution (Sakulthaew *et al.* 2017). More recently, 3D urchin-shaped monometallic palladium along with formic acid was successfully used as nanocatalyst in detoxification Cr containing industrial effluents (Mitra *et al.* 2017).

While the usage of nanomaterials and nano-products are limited for remediation of Cr(VI) pollutants, it is widely in use for treatments of such as Co(II), Ni(II), Cu(II), Cd(II), Pb(II) and As(III) (Anjum *et al.* 2016). Manganese oxides (MnO) nanoparticles are proficient for Cd (II) and Zn (II) while Graphene Nanosheets (GNS)/ δ -MnO₂, Nanocrystalline titanium dioxide, and ZnS nanocrystals are capable for reduction of Ni(II), As(III) Hg(II) (Varma *et al.* 2013; Qu *et al.* 2014). Nanotechnology being a relatively new concept and a rapidly progressing technology it is only a matter of time before it will be able to address the hazardous issues in the environment including that of Cr(VI) pollution.

Others remediation techniques

Although microbial and plant sources have great potential for bioremediation of Cr(VI), modern developments in environmental research and waste management technology have underlined new techniques for the treatment of heavy metals. Some of them are Bioaugmentation, Electrocoagulation, Biosurfactants, Vittrification, Bio-hydrometallurgy, and photocatalytic technology.

Bio-augmentation or seeding involves the deposition of highly specialized and engineered microbial populations to any contaminated site with recalcitrant toxic compounds to foster the degradation rates (Tyagi *et al.* 2011). The technique, however, results in the loss of the indigenous bacterial population while enhancing the growth of protozoans.

Electrocoagulation (EC) is a technique where electrolytic oxidation is introduced to an aqueous medium resulting in the destabilization of suspended or dissolved contaminants and generating coagulant. Electrocoagulation has modestly thrived against Cr(VI) ions from groundwater and needs more technical improvements (Hamdan and El-Naas 2014).

Bio-surfactants are nontoxic washing agents with features of low toxicity, eco-friendly, simple fabrication, and re-usable. Bio-surfactants eliminate the production of secondary pollutants while degrading a significant amount of pollutants. The use of emulsified vegetable oil (EVO) with acetate and Mg(OH)₂ has led to the complete removal of Cr from simulated groundwater and also induced accumulation of microbes related to organic fermentation and iron reduction (Fu *et al.* 2017). The use of biosurfactants from the aqueous extracts of plant biomass in association with anionic and neutral agents has gained tremendous popularity over the years (Mukherjee *et al.* 2016). Nandi *et al.* (2017) reported a

reduction efficiency of 96% for Cr (VI) concentration using biosurfactant from blue-green algae (1 g) in combination with 3×10^{-2} M sodium dodecyl sulfate (SDS). Several studies have also reported the use of aqueous extracts from different plant biomass like *Moringa oliefera*, sugarcane bagasse, siris sawdust containing amino acids and sugars which has been found to reduce Cr (VI) to Cr (III) by about 99% in the presence of anionic and neutral surfactants (Mukherjee, Ghosh, *et al.* 2014; Mukherjee, Saha, *et al.* 2014; Mukherjee *et al.* 2015).

Vitrification is the recovery of heavy metal from electroplating sludge with a reported recovery rate of 95% (Huang *et al.* 2013). Chemical additives (limestone and cullet) followed by heat treatment (1,450 °C) were used for the extraction of Ni and Cu from industrial sludge.

Bio-hydrometallurgy is a relatively new concept that employs various microorganisms along with acids, alkalis or organic solvents to the recovery of valuable metal (Mishra and Rhee 2010).

Photocatalytic processes are another useful technique where solar energy is used to reduce Cr(VI) to Cr(III). The use of Tin(IV) Sulfide/Graphene Oxide photocatalyst demonstrated an increased degradation rate of hexavalent chromium in comparison to normally visible light irradiation (Han *et al.* 2017). The photocatalytic reduction may be applied to organic matters and acids (iron fulvic acids and humic acid) to reduce heavy metals. The photocatalytic reduction is, however, have a high cost and a complicated installation procedure. The key factors influencing the photocatalytic reduction efficiency of Cr (VI) such as pH, activity of the catalyst, adsorption of Cr on the catalyst and other additives were briefly discussed in the recent review article outlined by Zhao *et al.* (2019).

Conclusion

Rapid industrialization makes production and disposing of heavy metal to the surrounding environment inevitable. Chromium pollution in Sukinda Valley had adversely affected the environmental scenario, causing several diseases in humans, animals, and plants. Chromium-containing waste needs special treatment for environmental security and recovery, as available remediation measures (physical, chemical and biological) are restricted in one way or the other (Table 5).

Despite different and distinct from each other in principle, various complementary techniques (chemical, microbial and biological) are in use for the remediation of heavy metal. Detailed knowledge of the sources and chemistry along with information on its potentially toxic effects at the contaminated site are necessary for the selection of the appropriate remedial method. Bioremediation though is a successful technique for the reduction of heavy metal pollutants, may not be useful for all heavy metals in every condition.

Microbial degradation and deterioration is a fast, gentle, energy-efficient method that generates fewer byproducts, yet, time-consuming and needs expert management.

Table 5. Advantages and limitations of some of the commonly used remediation techniques.

Remediation techniques	Advantages	Limitations	Ref.
<i>Physiochemical methods</i>			
Precipitation	Relatively inexpensive and straightforward to operate. Easy availability, moreover, low-cost	A large volume of sludge effectiveness reduced in the presence of multiple contaminants	(Agrawal <i>et al.</i> 2006; Bvrith and Reddy 2013)
Ion exchange	High treatment capacity, high removal efficiency, and fast kinetics	A time-consuming generates significant secondary waste	(Kang <i>et al.</i> 2004; Agrawal <i>et al.</i> 2006)
Membrane separation	Purifies heavy metals, suspended solids, and other associated organic compounds	High cost, membrane fouling, limited flow rates	(Agrawal <i>et al.</i> 2006; Bvrith and Reddy 2013)
Electrochemical	The rapid process, require fewer chemicals, provide high selectivity and produce less sludge	Involve high initial capital cost, production of the dendrite, lose or spongy deposits	(Rana <i>et al.</i> 2004; Barrera-Díaz <i>et al.</i> 2012)
<i>Bioremediation methods</i>			
Bioleaching	Easy to operate, less secondary waste	Very slow, leached toxic materials may run-off to nearby water sources	(Pathak <i>et al.</i> 2008b; Zeng <i>et al.</i> 2016)
Adsorption	Easily available, profitable, easy to operate and cost-efficient	Metal-specific and requires chemical derivatization to improve its sorption capacity	(Bvrith and Reddy 2013; Vendruscolo <i>et al.</i> 2016; Enniya <i>et al.</i> 2018)
Bioreduction	Eco-friendly, require even fewer reagents, easy and inexpensive to operate	Need Isolation, identification, and characterization of microbes from their native soil or aqueous environment before their application, difficult to maintain	(Barrera-Díaz <i>et al.</i> 2012; Harish <i>et al.</i> 2012)
<i>Phytoremediation methods</i>			
	Eco-friendly, high accumulation rate	Long duration time, phytotoxic at high concentrations, the probable route for vegetation contamination in the food chain	(Bvrith and Reddy 2013)
<i>Nanoremediation</i>			
	Eco-friendly, effective in small quantity, large surface area, high sensitivity	Need expertise for synthesis, expensive, difficult to maintain	(Mitra <i>et al.</i> 2017)

Although the success of a mixed or hybrid combination of native bacterial strains are being advocated, physiochemical parameter and culturing medium are to be optimized for increasing the recovery percentage of Cr and other heavy metal. The development of engineered microbes and their genetic manipulation can further boost the technological advances in the treatment of Cr(VI) waste and other toxic metals. Hence, a combined approach involving nanotechnology, microbiology and biotechnology, which are currently under active research, is much needed as the next-generation technologies for environmental pollution management.

Acknowledgments

The authors thank the Department of Biotechnology and Medical Engineering and the Department of Electrical Engineering of National Institute of Technology Rourkela, India for providing the research facility.

Funding

The authors gratefully acknowledge the Department of Science and Technology (DST), India for funding the research under DST-WTI (Water Technology Initiative) [WTI/2015/113].

References

Ackerley DF, Gonzalez CF, Park CH, Ii RB, Next-Generation A, Ackerley DF, Gonzalez CF, Park CH, Ii RB, Keyhan M, *et al.* 2004.

- Chromate-reducing properties of soluble flavoproteins from *Pseudomonas putida* and *Escherichia coli*. Appl Environ Microbiol. 70(2):873–882. doi:10.1128/AEM.70.2.873.
- Agrawal A, Kumar V, Pandey BD. 2006. Remediation options for the treatment of electroplating and leather tanning effluent containing chromium – a review. Miner Process Extr Metall Rev. 27(2):99–130. doi:10.1080/08827500600563319.
- Ahemad M. 2014. Bacterial mechanisms for Cr (VI) resistance and reduction: an overview and recent advances. Folia Microbiol. 59(4): 321–332. doi:10.1007/s12223-014-0304-8.
- Ahemad M. 2015. Enhancing phytoremediation of chromium-stressed soils through plant-growth-promoting bacteria. J Genet Eng Biotechnol. 13(1):51–58. doi:10.1016/j.jgeb.2015.02.001.
- Anjum M, Miandad R, Waqas M, Gehany F, Barakat MA. 2016. Remediation of wastewater using various nano-materials. Arab J Chem. 12(8):4897–4919. doi:10.1016/j.arabjc.2016.10.004.
- Argulwar KVGR, Sudakaran S, Pulimi SV, Chandrasekaran M, Mukherjee NA. 2018. Nano-bio sequential removal of hexavalent chromium using polymer-nZVI composite film and sulfate reducing bacteria under anaerobic condition. Environ Technol Innov. 9: 122–133. doi:10.1016/j.eti.2017.11.006.
- Arunbabu V, Indu KS, Ramasamy EV. 2017. Leachate pollution index as an effective tool in determining the phytotoxicity of municipal solid waste leachate. Waste Manag. 68:329–336. doi:10.1016/j.wasman.2017.07.012.
- Ashley K, Howe AM, Demange M, Nygren O. 2003. Sampling and analysis considerations for the determination of hexavalent chromium in workplace air. J Environ Monitor. 5(5):707. doi:10.1039/b306105c.
- Barrera-Díaz CE, Lugo-Lugo V, Bilyeu B. 2012. A review of chemical, electrochemical and biological methods for aqueous Cr (VI) reduction. J Hazard Mater. 224:1–12. doi:10.1016/j.jhazmat.2012.04.054.
- Beaumont JJ, Sedman RM, Reynolds SD, Sherman CD, Li LH, Howd RA, Sandy MS, Zeise L, Alexeff GV. 2008. Cancer mortality in a Chinese population exposed to hexavalent chromium in drinking water. Epidemiology 19(1):12–23. doi:10.1097/EDE.0b013e31815cea4c.

- Bharadwaj A, Ting YP. 2013. Bioleaching of spent hydrotreating catalyst by acidophilic thermophile *Acidianus brierleyi*: leaching mechanism and effect of decoking. *Bioresour Technol.* 130:673–680. doi:10.1016/j.biortech.2012.12.047.
- Bharagava RN, Mishra S. 2018. Hexavalent chromium reduction potential of *Cellulosimicrobium* sp. isolated from common effluent treatment plant of tannery industries. *Ecotoxicol Environ Saf.* 147: 102–109. doi:10.1016/j.ecoenv.2017.08.040.
- Biswas P, Karn AK, Balasubramanian P, Kale PG. 2017. Biosensor for detection of dissolved chromium in potable water: a review. *Biosens Bioelectron.* 94:589–604. doi:10.1016/j.bios.2017.03.043.
- Bridge G. 2004. Contested terrain: mining and the environment. *Annu Rev Environ Resour.* 29(1):205–259. doi:10.1146/annurev.energy.28.011503.163434.
- Buendía-González L, Orozco-Villafuerte J, Cruz-Sosa F, Barrera-Díaz CE, Vernon-Carter EJ. 2010. *Prosopis laevigata* a potential chromium (VI) and cadmium (II) hyperaccumulator desert plant. *Bioresour Technol.* 101(15):5862–5867. doi:10.1016/j.biortech.2010.03.027.
- Bvrith MV, Reddy VB. 2013. An overview on research trends in remediation of chromium. *Res J Recent Sci.* 2:71–83.
- Chen JM, Hao OJ, Chen JM, Hao OJ. 1998. Microbial chromium (VI) reduction microbial chromium (VI) reduction. *Crit Rev Environ Sci Technol.* 28(3):219–251. doi:10.1080/10643389891254214.
- Chen SY, Lin PL. 2010. Optimization of operating parameters for the metal bioleaching process of contaminated soil. *Sep Purif Technol.* 71(2):178–185. doi:10.1016/j.seppur.2009.11.018.
- Cheung KH, Gu JD. 2007. Mechanism of hexavalent chromium detoxification by microorganisms and bioremediation application potential: a review. *Int Biodeterior Biodegrad.* 59(1):8–15. doi:10.1016/j.ibiod.2006.05.002.
- Choppala G, Bolan N, Park JH. 2013. Chapter two – chromium contamination and its risk management in complex environmental settings. *Adv Agron.* 120:129–172. doi:10.1016/B978-0-12-407686-0.00002-6.
- Coelho PCS, Teixeira JPF, Gonçalves ONBSM. 2011. Mining activities: health impacts. *Encycl Environ Heal.* 2011:788–802. doi:10.1016/B978-0-444-52272-6.00488-8.
- Congeevaram S, Dhanarani S, Park J, Dexilin M, Thamaraiselvi K. 2007. Biosorption of chromium and nickel by heavy metal resistant fungal and bacterial isolates. *J Hazard Mater.* 146(1–2):270–277. doi:10.1016/j.jhazmat.2006.12.017.
- Das A, Singh S. 2011. Occupational health assessment of chromite toxicity among Indian miners. *Indian J Occup Environ Med.* 15(1):6. doi:10.4103/0019-5278.82998.
- Das S, Ram SS, Sahu HK, Rao DS, Chakraborty A, Sudarshan M, Thatoi HN. 2013. A study on soil physico-chemical, microbial and metal content in Sukinda chromite mine of Odisha, India. *Environ Earth Sci.* 69(8):2487–2497. doi:10.1007/s12665-012-2074-4.
- De Oliveira Andrade JJ, Mattje V. 2012. Incorporation of chromium-tanned leather residue in mortars. *Proc Inst Civ Eng - Constr Mater.* 165(2):73–87. doi:10.1680/coma.10.00026.
- Dehghani MH, Sanaei D, Ali I, Bhatnagar A. 2016. Removal of chromium(VI) from aqueous solution using treated waste newspaper as a low-cost adsorbent: Kinetic modeling and isotherm studies. *J Mol Liq.* 215:671–679. doi:10.1016/j.molliq.2015.12.057.
- Demirbas E, Kobya M, Senturk E, Ozkan T. 2004. Adsorption kinetics for the removal of chromium (VI) from aqueous solutions on the activated carbons prepared from agricultural wastes. *WSA.* 30: 533–539. doi:10.4314/wsa.v30i4.5106.
- Dey S, Paul AK. 2015. Hexavalent chromate reduction during growth and by immobilized cells of *Arthrobacter* sp. SUK 1205. *Sci Technol Dev.* 34:158–168. doi:10.3923/std.2015.158.168.
- Dhal B, Abhilash, Pandey BD. 2018. Mechanism elucidation and adsorbent characterization for removal of Cr(VI) by native fungal adsorbent. *Sustain Environ Res.* 28:289–297. doi:10.1016/j.serj.2018.05.002.
- Dhal B, Das NN, Pandey BD, Thatoi HN. 2011. Environmental quality of the Boula-Nuasahi Chromite mine area in India. *Mine Water Environ.* 30(3):191–196. doi:10.1007/s10230-010-0134.
- Dhal B, Das NN, Thatoi HN, Pandey BD. 2013. Characterizing toxic Cr (VI) contamination in chromite mine overburden dump and its bacterial remediation. *J Hazard Mater.* 260:141–149. doi:10.1016/j.jhazmat.2013.04.050.
- Diwan H, Ahmad A, Iqbal M. 2008. Genotypic variation in the phytoremediation potential of Indian mustard for chromium. *Environ Manage.* 41(5):734–741. doi:10.1007/s00267-007-9020-3.
- Dubey CS, Sahoo BK, Nayak NR. 2001. Chromium (VI) in waters in parts of Sukinda chromite valley and health hazards, Orissa, India. *Bull Environ Contam Toxicol.* 67(4):541–548. doi:10.1007/s001280157.
- Enniya I, Rghioui L, Jourani A. 2018. Adsorption of hexavalent chromium in aqueous solution on activated carbon prepared from apple peels. *Sustain Chem Pharm.* 7:9–16. doi:10.1016/j.scp.2017.11.003.
- Ertani A, Mietto A, Borin M, Nardi S. 2017. Chromium in agricultural soils and crops: a review. *Water Air Soil Pollut.* 228:190. doi:10.1007/s11270-017-3356-y.
- Eslami J, Khaniki G, Nurani M, Mehrasbi M, Peyda M, Azimi R. 2007. Heavy metals in edible green vegetables grown along the sites. *J Biol Sci.* 6:943–948. doi:10.3923/jbs.2007.943.948.
- Fu F, Ma J, Xie L, Tang B, Han W, Lin S. 2013. Chromium removal using resin supported nanoscale zero-valent iron. *J Environ Manage.* 128:822–827. doi:10.1016/j.jenvman.2013.06.044.
- Fu R, Wen D, Xia X, Zhang W, Gu Y. 2017. Electrokinetic remediation of chromium (Cr)-contaminated soil with citric acid (CA) and polyaspartic acid (PASP) as electrolytes. *Chem Eng J.* 316:601–608. doi:10.1016/j.cej.2017.01.092.
- Gardea-Torresdey JL, De La Rosa G, Peralta-Videa JR, Montes M, Cruz-Jimenez G, Cano-Aguilera I. 2005. Differential uptake and transport of trivalent and hexavalent chromium by tumbleweed (*Salsola kali*). *Arch Environ Contam Toxicol.* 48(2):225–232. doi:10.1007/s00244-003-0162-x.
- Garg UK, Kaur MP, Garg VK, Sud D. 2007. Removal of hexavalent chromium from aqueous solution by agricultural waste biomass. *J Hazard Mater.* 140(1–2):60–68. doi:10.1016/j.jhazmat.2006.06.056.
- Gong Y, Werth CJ, He Y, Su Y, Zhang Y, Zhou X. 2018. Intracellular versus extracellular accumulation of Hexavalent chromium reduction products by *Geobacter sulfurreducens* PCA. *Environ Pollut.* 240: 485–492. doi:10.1016/j.envpol.2018.04.046.
- Guan X, Sun Y, Qin H, Li J, Lo IMC, He D, Dong H. 2015. The limitations of applying zero-valent iron technology in contaminants sequestration and the corresponding countermeasures: the development in zero-valent iron technology in the last two decades (1994–2014). *Water Res.* 75:224–248. doi:10.1016/j.watres.2015.02.034.
- Gupta VK, Shrivastava AK, Jain N. 2001. Biosorption of chromium(VI) from aqueous solutions by green algae *Spirogyra species*. *Water Res.* 35(17):4079–4085. doi:10.1016/S0043-1354(01)00138-5.
- Hamdan SS, El-Naas MH. 2014. Characterization of the removal of Chromium(VI) from groundwater by electrocoagulation. *J Ind Eng Chem.* 20(5):2775–2781. doi:10.1016/j.jiec.2013.11.006.
- Han L, Mao D, Huang Y, Zheng L, Su Y, Sun S. 2017. Fabrication of unique Tin(IV) sulfide/graphene oxide for photocatalytically treating chromium(VI)-containing wastewater. *J Clean Prod.* 168:519–525. doi:10.1016/j.jclepro.2017.09.027.
- Han X, Shan Y, Hung M, Fung N, Tam Y. 2007. Biosorption and bio-reduction of Cr (VI) by a microalgal isolate, *Chlorella miniata*. *J Hazard Mater.* 146(1–2):65–72. doi:10.1016/j.jhazmat.2006.11.053.
- Hartman HL, Mutmansky JM. 2002. Introduction to mining. In: *Introductory Mining Engineering*. New York: Wiley. p. 1–24.
- Harish R, Samuel J, Mishra R, Chandrasekaran N, Mukherjee A. 2012. Bio-reduction of Cr (VI) by exopolysaccharides (EPS) from indigenous bacterial species of Sukinda chromite mine, India. *Biodegradation.* 23:487–496. doi:10.1007/s10532-011-9527-4.
- Hong K, Tokunaga S, Kajluchi T. 2002. Evaluation of remediation process with plant-derived biosurfactant for recovery of heavy metals from contaminated soils. *Chemosphere.* 43:379–387. doi:10.1016/S0045-6535(02)00321-1.
- Hou S, Wu B, Peng D, Wang Z, Wang Y, Xu H. 2019. Remediation performance and mechanism of hexavalent chromium in alkaline

- soil using multi-layer loaded nano-zero-valent iron. *Env Pollu*. 252: 553–561. doi:10.1016/j.envpol.2019.05.083.
- Huang R, Huang K, Lin Z, Wang J, Lin C, Kuo Y. 2013. Recovery of valuable metals from electroplating sludge with reducing additives via vitrification. *J Environ Manage*. 129:586–592. doi:10.1016/j.jenvman.2013.08.019.
- Hyder AHMG, Begum SA, Egiebor NO. 2015. Adsorption isotherm and kinetic studies of hexavalent chromium removal from aqueous solution onto bone char. *J Environ Chem Eng*. 3(2):1329–1336. doi:10.1016/j.jece.2014.12.005.
- Ibrahim ASS, Elbadawi YB, El-Tayeb MA, Al-Salamah AA. 2012. Hexavalent chromium reduction by novel chromate resistant alkaliphilic *Bacillus* sp. strain KSUCr9a. *Afr J Biotechnol*. 11:3832–3841. doi:10.5897/AJB11.3026.
- Ibrahim ASS, El-Tayeb MA, Elbadawi YB, Al-Salamah AA. 2011. Bioreduction of Cr (VI) by potent novel chromate resistant alkaliphilic *Bacillus* sp. strain KSUCr5 isolated from hypersaline Soda lakes. *Afr J Biotechnol*. 10:7207–7218. doi:10.5897/AJB11.1030.
- Indian Bureau of Mines. 2016. Indian Minerals Yearbook 2015. Indian Miner. Yearb. 2015 (Part-III Miner). Rev. 54th Editi, 1–9.
- Islam MA, Angove MJ, Morton DW. 2019. Recent innovative research on chromium (VI) adsorption mechanism. *Environ Nanotechnol Monit Manag*. 12:100267.
- Jaishankar M, Tseten T, Anbalagan N, Mathew BB, Beeregowda KN. 2014. Toxicity, mechanism and health effects of some heavy metals. *Interdiscip Toxicol*. 7(2):60–72. doi:10.2478/intox-2014-0009.
- January MC, Cutright TJ, Keulen H, Van Wei R. 2008. Hydroponic phytoremediation of Cd, Cr, Ni, As, and Fe: can *Helianthus annuus* hyperaccumulate multiple heavy metals? *Chemosphere*. 70(3): 531–537. doi:10.1016/j.chemosphere.2007.06.066.
- Joutey NT, Sayel H, Bahafid W, Ghachtouli E. 2015. Mechanisms of hexavalent chromium resistance and removal by microorganisms. *Rev Environ Contam Toxicol*. 233:45–70. doi:10.1007/978-3-319-10479-9.
- Kang SY, Lee JU, Moon SH, Kim KW. 2004. Competitive adsorption characteristics of Co^{2+} , Ni^{2+} , and Cr^{3+} by IRN-77 cation exchange resin in synthesized wastewater. *Chemosphere*. 56(2):141–147. doi:10.1016/j.chemosphere.2004.02.004.
- Karimi N. 2013. Comparative phytoremediation of chromium-contaminated soils by Alfalfa (*Medicago sativa*) and Sorghum bicolor (L) Moench. *IJSRES*. 1:44–49. doi:10.12983/ijres-2013-p044-049.
- Kim C, Zhou Q, Deng B, Thornton EC, Xu H. 2001. Chromium (VI) reduction by hydrogen sulfide in aqueous media: stoichiometry and kinetics. *Environ Sci Technol*. 35(11):2219–2225. doi:10.1021/es0017007.
- Kim Y, Roh Y. 2019. Environmental application of biogenic magnetite nanoparticles to remediate chromium (III/VI) contaminated water. *Minerals*. 9(5):260. doi:10.3390/min9050260.
- Ksheminska H, Fedorovych D, Honchar T, Ivash M, Gonchar M. 2008. Yeast tolerance to chromium depends on extracellular chromate reduction and Cr (III) Chelation. *Food Technol Biotechnol*. 46: 419–426.
- Ksheminska HP, Honchar TM, Gayda GZ, Gonchar MV. 2006. Extracellular chromate-reducing activity of the yeast cultures. *Cental Eur J Biol*. 1:137–149. doi:10.2478/s11535-006-0009-3.
- Kumar R, Bishnoi NR, Bishnoi K. 2008. Biosorption of chromium (VI) from aqueous solution and electroplating wastewater using fungal biomass. *Chem Eng J*. 135:202–208. doi:10.1016/j.cej.2007.03.004.
- Kumar V, Dwivedi SK. 2019. Hexavalent chromium reduction ability and bioremediation potential of *Aspergillus flavus* CR500 isolated from electroplating wastewater. *Chemosphere*. 237:124567. doi:10.1016/j.chemosphere.2019.124567.
- Losi ME, Frankenberger WT. 1994. Chromium-resistant microorganisms isolated from evaporation ponds of a metal processing plant. *Water Air Soil Pollut*. 74:405–413.
- Lu X, Li B, Guo L, Zhang G, Wang P. 2019. Reduction and stabilization remediation of hexavalent Chromium (Cr^{6+})-contaminated soil. *Ekoloji*. 28(107):973–980.
- Lv X, Xu J, Jiang G, Tang J, Xu X. 2012. Highly active nanoscale zero-valent iron (nZVI)- Fe_3O_4 nanocomposites for the removal of chromium(VI) from aqueous solutions. *J Colloid Interface Sci*. 369(1):460–469. doi:10.1016/j.jcis.2011.11.049.
- Lytle CM, Lytle FW, Yang N, Qian JH, Hansen D, Zayed A, Terry N. 1998. Reduction of Cr (VI) to Cr (III) by wetland plants: potential for in situ heavy metal detoxification. *Environ Sci Technol*. 32(20): 3087–3093. doi:10.1021/es980089x.
- Maitlo HA, Kim KH, Kumar V, Kim S, Park JW. 2019. Nanomaterials-based treatment options for chromium in aqueous environments. *Environ Int*. 130:104748. doi:10.1016/j.envint.2019.04.020.
- Mala J, Dhanasingh S, Rose C. 2014. Inducible chromate reductase exhibiting extracellular activity in *Bacillus methylotrophicus* for chromium bioremediation. *Microbiol Res*. 170:235–241.
- Mangkoedihardjo S, Ratnawati R, Alfianti N. 2008. Phytoremediation of hexavalent chromium polluted soil using *Pterocarpus indicus* and *Jatropha curcas* L. *World Appl Sci J*. 4:338–342.
- Mant C, Costa S, Williams J, Tambourgi E. 2006. Phytoremediation of chromium by model constructed wetland. *Bioresour Technol*. 97(15):1767–1772. doi:10.1016/j.biortech.2005.09.010.
- Matrosova EA, Bobrov AV, Bindi L. 2020. Basics of geochemistry and mineralogy of chromium. In: *Geochemistry of chromium in the earth's mantle*. Cham: Springer. p. 5–36.
- Ministry of Mines, Government of India. 2013. Monograph on chromite. https://doi.org/https://ibm.gov.in/writereaddata/files/09162014114959Monograph_Chromite.pdf.
- Ministry of Mines, India. 2017. Annual report. https://www.mines.gov.in/writereaddata/UploadFile/Mines_AR_2016-17_English.pdf.
- Mishra D, Rhee Y. 2010. Current research trends of microbiological leaching for metal recovery from industrial wastes. *Curr Res Technol Educ Top Appl Microbiol Microb Biotechnol*. 2:1289–1296.
- Mishra H, Sahu HB. 2013. Environmental scenario of chromite mining at Sukinda Valley – a review. *Int J Environ Eng Manag*. 4:287–292.
- Mishra SR, Pradhan RP, Prusty BAK, Sahu SK. 2016. Meteorology drives ambient air quality in a valley: a case of Sukinda chromite mine, one among the ten most polluted areas in the world. *Environ Monit Assess*. 188:1–17. doi:10.1007/s10661-016-5393-1.
- Mitra S, Sarkar A, Sen S. 2017. Removal of chromium from industrial effluents using nanotechnology: a review. *Nanotechnol Environ Eng*. 2(1):11. doi:10.1007/s41204-017-0022-y.
- Mohanty JK, Rao DS, Paul AK, Khaoash S. 2009. Characterization of high magnesium rocks for suitability as flux in iron and steel industry. *J Geol Min Res*. 1:149–155.
- Mukherjee K, Ghosh D, Saha B. 2014. Surfactant assisted enhancement of bioremediation rate for hexavalent chromium by water extract of Siris (*Albizia lebbek*) sawdust. *TSD*. 51(6):521–527. doi:10.3139/113.110338.
- Mukherjee K, Nandi R, Saha D, Saha B. 2015. Surfactant-assisted enhancement of bioremediation rate for hexavalent chromium by water extract of Sajina (*Moringa oleifera*) flower. *Desalin Water Treat*. 54(2):525–532. doi:10.1080/19443994.2014.884477.
- Mukherjee K, Nandi R, Saha D, Saha B. 2016. Surfactant-assisted bioremediation of hexavalent chromium from contaminated water. *Desalin Water Treat*. 53(3):746–751. doi:10.1080/19443994.2013.842503.
- Mukherjee K, Saha R, Ghosh A, Ghosh SK, Maji PK, Saha B. 2014. Surfactant-assisted bioremediation of hexavalent chromium by use of an aqueous extract of sugarcane bagasse. *Res Chem Intermed*. 40(4):1727–1734. doi:10.1007/s11164-013-1077-4.
- Murugavelh KM. 2013. Bioreduction of chromate by immobilized cells of *Halomonas* sp. *Int J Energy Environ*. 4:349–356.
- Nandi R, Laskar S, Saha B. 2017. Surfactant-promoted enhancement in bioremediation of hexavalent chromium to trivalent chromium by naturally occurring wall algae. *Res Chem Intermed*. 43(3): 1619–1634. doi:10.1007/s11164-016-2719-0.
- Nareshkumar R, Nagendran R, Parvathi K. 2007. Microbial solubilization of heavy metals from soil using indigenous sulfur oxidizing bacterium: effects of sulfur/soil ratio. *J Sci Ind Res*. 66:680–683.
- Naz A, Chowdhury A, Mishra BK, Gupta SK. 2016. Metal pollution in water environment and the associated human health risk from drinking water: a case study of Sukinda chromite mine, India. *Hum Ecol Risk Assess An Int J*. 22(7):1433–1455. doi:10.1080/10807039.2016.1185355.

- Opperman DJ, Sewell BT, Litthauer D, Isupov MN, Littlechild JA, Van Heerden E. 2010. Biochemical and biophysical research communications crystal structure of a thermostable Old Yellow enzyme from *Thermus scotoductus* SA-01. *Biochem Biophys Res Commun.* 393(3): 426–431. doi:10.1016/j.bbrc.2010.02.011.
- Ozdemir G, Ceyhan N, Ozturk T, Akirmak F, Cosar T. 2004. Biosorption of chromium (VI), cadmium (II) and copper (II) by *Pantoea* sp. TEM18. *Chem Eng J.* 102(3):249–253. doi:10.1016/j.cej.2004.01.032.
- Ozturk S, Kaya T, Aslim B, Tan S. 2012. Removal and reduction of chromium by *Pseudomonas* sp. and their correlation to rhamnolipid production. *J Hazard Mater.* 232:64–69. doi:10.1016/j.jhazmat.2012.06.038.
- Pakade VE, Tavengwa NT, Madikizela LM. 2019. Recent advances in hexavalent chromium removal from aqueous solutions by adsorptive methods. *RSC Adv.* 9(45):26142–26164. doi:10.1039/C9RA05188K.
- Pal A, Datta S, Paul AK. 2013. Hexavalent chromium reduction by immobilized cells of *Bacillus sphaericus* AND 303. *Braz Arch Biol Technol.* 56(3):505–512. doi:10.1590/S1516-89132013000300019.
- Pang Y, Zeng G, Tang L, Zhang Y, Liu Y, Lei X, Wu M, Li Z, Liu C. 2011. Cr (VI) reduction by *Pseudomonas aeruginosa* immobilized in a polyvinyl alcohol/Sodium alginate matrix containing multi-walled carbon nanotubes. *Bioresour Technol.* 102(22):10733–10736. doi:10.1016/j.biortech.2011.08.078.
- Park D, Yun Y, Hye J, Moon J. 2005. Mechanism of hexavalent chromium removal by dead fungal biomass of *Aspergillus niger*. *Water Res.* 39(4):533–540. doi:10.1016/j.watres.2004.11.002.
- Pathak A, Dastidar MG, Sreekrishnan TR. 2008a. Bioleaching of heavy metals from anaerobically digested sewage sludge Bioleaching of heavy metals from anaerobically digested sewage sludge. *J Environ Sci Heal A.* 4529:402–411. doi:10.1080/10934520701795624.
- Pathak A, Dastidar MG, Sreekrishnan TR. 2008b. Bioleaching of heavy metals from anaerobically digested sewage sludge. *J Environ Sci Heal A.* 43(4):402–411. doi:10.1080/10934520701795624.
- Pathak A, Dastidar MG, Sreekrishnan TR. 2009. Bioleaching of heavy metals from sewage sludge: a review. *J Environ Manage.* 90(8): 2343–2353. doi:10.1016/j.jenvman.2008.11.005.
- Pradhan D, Behari L, Sawyer M, Rahman PKSM. 2017. Recent bioreduction of hexavalent chromium in wastewater treatment: a review. *J Ind Eng Chem.* 55:1–20. doi:10.1016/j.jiec.2017.06.040.
- Qu M, Chen J, Huang Q, Chen J, Xu Y, Luo J, Wang K, Gao W, Zheng Y. 2016. Bioremediation of hexavalent chromium contaminated soil by a bioleaching system with weak magnetic fields. *Int Biodeterior Biodegrad.* 128:41–47. doi:10.1016/j.ibiod.2016.08.022.
- Qu M, Li W, Zhang C, Huang B, Zhao Y. 2015. Assessing the pollution risk of soil chromium based on loading capacity of paddy soil at a regional scale. *Sci Rep.* 5:1–8. doi:10.1038/srep18451.
- Qu Z, Yan L, Li L, Xu J, Liu M, Li Z, Yan N. 2014. Ultra effective ZnS nanocrystals sorbent for mercury (II) removal based on size-dependent cation exchange. *ACS Appl Mater Interfaces.* 6(20):18026–18032. doi:10.1021/am504896w.
- Quaggiotti S, Barcaccia G, Schiavon M, Nicolé S, Galla G, Rossignolo V, Soattin M, Malagoli M. 2007. Phytoremediation of chromium using *Salix* species: cloning ESTs and candidate genes involved in the Cr response. *Gene.* 402(1–2):68–80. doi:10.1016/j.gene.2007.07.021.
- Ramana S, Biswas AK, Singh AB, Ahirwar NK. 2012. Phytoremediation of chromium by tuberose. *Natl Acad Sci Lett.* 35:71–73. doi:10.1007/s40009-012-0016-z.
- Rana P, Mohan N, Rajagopal C. 2004. Electrochemical removal of chromium from wastewater by using carbon aerogel electrodes. *Water Res.* 38(12):2811–2820. doi:10.1016/j.watres.2004.02.029.
- Rangabhashiyam S, Balasubramanian P. 2018a. Biosorption of hexavalent chromium and malachite green from aqueous effluents, using *Cladophora* sp. *Chem Ecol.* 34(4):371–390. doi:10.1080/02757540.2018.1427232.
- Rangabhashiyam S, Balasubramanian P. 2018b. Adsorption behaviors of hazardous methylene blue and hexavalent chromium on novel materials derived from *Pterospermum acerifolium* shells. *J Mol Liq.* 254:433–445. doi:10.1016/j.molliq.2018.01.131.
- Rastegar SO, Mousavi SM, Shojasadati SA. 2014. Cr and Ni recovery during bioleaching of dewatered metal-plating sludge using *Acidithiobacillus ferrooxidans*. *Bioresour Technol.* 167:61–68. doi:10.1016/j.biortech.2014.05.107.
- Redondo-Gómez S, Mateos-Naranjo E, Vecino-Bueno I, Feldman SR. 2011. Accumulation and tolerance characteristics of chromium in a cordgrass Cr-hyperaccumulator, *Spartina argentinensis*. *J Hazard Mater.* 185(2–3):862–869. doi:10.1016/j.jhazmat.2010.09.101.
- Reichl C, Schatz M, Zsak G. 2016. World mining data. *Fed Minist Sci Res Econ Austria.* 31:1–255.
- Reijonen I, Hartikainen H. 2016. Oxidation mechanisms and chemical bioavailability of chromium in agricultural soil – pH as the master variable. *Appl Geochemistry.* 74:84–93. doi:10.1016/j.apgeochem.2016.08.017.
- Revathi K, Haribabu TE, Sudha PN. 2011. Phytoremediation of chromium contaminated soil using sorghum plant. *Int J Environ Sci.* 2(2):417.
- Rezaei H. 2016. Biosorption of chromium by using *Spirulina* sp. *Arab J Chem.* 9(6):846–853. doi:10.1016/j.arabjc.2013.11.008.
- Saha B, Orvig C. 2010. Biosorbents for hexavalent chromium elimination from industrial and municipal effluents. *Coord Chem Rev.* 254(23–24):2959–2972. https://doi:10.1016/j.ccr.2010.06.005.
- Saha R, Nandi R, Saha B. 2011. Sources and toxicity of hexavalent chromium. *J Coord Chem.* 64(10):1782–1806. doi:10.1080/00958972.2011.583646.
- Sahinkaya E, Altun M, Bektas S, Komnitsas K. 2012. Bioreduction of Cr(VI) from acidic wastewaters in a sulfidogenic ABR. *Miner Eng.* 32:38–44. doi:10.1016/j.mineng.2012.03.014.
- Sakulthaew C, Chokejaroenrat C, Poapolatthep A, Satapanajaru T, Poapolatthep S. 2017. Chemosphere hexavalent chromium adsorption from aqueous solution using carbon. *Chemosphere.* 184:1168–1174. doi:10.1016/j.chemosphere.2017.06.094.
- Samuel J, Paul ML, Pulimi M, Nirmala MJ, Chandrasekaran N, Mukherjee A. 2012. Hexavalent chromium bioremoval through adaptation and consortia development from Sukinda chromite mine isolates. *Ind Eng Chem Res.* 51(9):3740–3749. doi:10.1021/ie201796s.
- Saviour MN. 2012. Environmental impact of soil and sand mining: a review. *Int J Sci Environ Technol.* 1:125–134.
- Shams KM, Tichy G, Fischer A, Sager M, Peer T, Bashar A, Filip K. 2010. Aspects of phytoremediation for chromium contaminated sites using common plants *Urtica dioica*, *Brassica napus* and *Zea mays*. *Plant Soil.* 328(1–2):175–189. doi:10.1007/s11104-009-0095-x.
- Shang J, Zong M, Yu Y, Kong X, Du Q, Liao Q. 2017. Removal of chromium (VI) from water using nanoscale zerovalent iron particles supported on herb-residue biochar. *J Environ Manage.* 197:331–337. doi:10.1016/j.jenvman.2017.03.085.
- Sharma S, Adholeya A. 2011. Detoxification and accumulation of chromium from tannery effluent and spent chrome effluent by *Paecilomyces lilacinus* fungi. *Int Biodeterior Biodegrad.* 65(2): 309–317. doi:10.1016/j.ibiod.2010.12.003.
- Sharma P, Bihari V, Agarwal SK, Verma V, Kesavachandran CN, Pangtey BS, Mathur N, Singh KP, Srivastava M, Goel SK. 2012. Groundwater Contaminated with hexavalent Chromium [Cr (VI)]: a health survey and clinical examination of community inhabitants (Kanpur, India). *PLoS One.* 7(10):e47877–9. doi:10.1371/journal.pone.0047877.
- Shinghal AMADS. 2015. Biosorption of chromium (VI) from aqueous solutions using waste plant biomass. *Int J Environ Sci Technol.* 12: 1415–1426. doi:10.1007/s13762-014-0516-0.
- Sierra-Alvarez R. 2009. Removal of copper, chromium and arsenic from preservative-treated wood by chemical extraction-fungal bioleaching. *Waste Manag.* 29(6):1885–1891. doi:10.1016/j.wasman.2008.12.015.
- Singh R, Dong H, Liu D, Zhao L, Marts AR, Farquhar E, Tierney DL, Almquist CB, Briggs BR. 2015. Reduction of hexavalent chromium by the thermophilic methanogen *Methanothermobacter thermautotrophicus*. *Geochim Cosmochim Acta.* 148:442–456. doi:10.1016/j.gca.2014.10.012.
- Singh R, Misra V, Singh RP. 2011. Synthesis, characterization and role of zero-valent iron nanoparticle in removal of hexavalent chromium

- from chromium-spiked soil. *J Nanopart Res.* 13(9):4063–4073. doi:10.1007/s11051-011-0350-y.
- Soni K, Vyas N, 2015. A Study of occupational health and safety related practices in mining companies of southern Rajasthan: a systematic review. *Int J Adv Res Innov Ideas Educ.* 1:92–103.
- Sundaramoorthy P, Chidambaram A, Ganesh KS, Unnikannan P, Baskaran L, 2010. Chromium stress in paddy: (i) Nutrient status of paddy under chromium stress; (ii) Phytoremediation of chromium by aquatic and terrestrial weeds. *C R Biol.* 333(8):597–607. doi:10.1016/j.crv.2010.03.002.
- Talib NSR, Halmi MIE, Ghani SSA, Zaidan UH, Shukor MYA, 2019. Artificial Neural Networks (ANNs) and Response Surface Methodology (RSM) approach for modelling the optimization of Chromium (VI) reduction by newly isolated *Acinetobacter radiore-sistens* Strain NS-MIE from agricultural soil. *BioMed Res Int.* 2019: 1–14. doi:10.1155/2019/5785387.
- Tamilselvan N, Saurav K, Kannabiran K, 2012. Biosorption of Cr (VI), Cr (III), Pb (II) and Cd (II) from aqueous solutions by *Sargassum wightii* and *Caulerpa racemosa* algal biomass. *J Ocean Univ China* . 11(1):52–58. doi:10.1007/s11802-012-1843-8.
- Turgut C, Pepe MK, Cutright TJ, 2004. The effect of EDTA and citric acid on phytoremediation of Cd, Cr, and Ni from soil using *Helianthus annuus*. *Environ Pollut.* 131(1):147–154. doi:10.1016/j.envpol.2004.01.017.
- Tyagi M, da Fonseca MMR, de Carvalho CCCR, 2011. Bioaugmentation and biostimulation strategies to improve the effectiveness of bioremediation processes. *Biodegradation* 22(2):231–241. doi:10.1007/s10532-010-9394-4.
- U.S. Geological Survey (USGS). 2015. Mineral Commodity Summaries 2015. p. 1–196. doi:10.3133/70140094.
- U.S. Geological Survey (USGS) 2018. Mineral Commodity Summaries 2018. U.S. Geological Survey. doi:10.3133/70194932.
- Ucun H, Bayhan YK, Kaya Y, Cakici A, Algur OF, 2002. Biosorption of chromium (VI) from aqueous solution by cone biomass of *Pinus sylvestris*. *Bioresour Technol.* 85:155–158. doi:10.1016/S0960-8524(02)00086-X.
- Ullah I, Nadeem R, Iqbal M, Manzoor Q, 2013. Biosorption of chromium onto native and immobilized sugarcane bagasse waste biomass. *Ecol Eng.* 60:99–107. doi:10.1016/j.ecoleng.2013.07.028.
- Vajpayee P, Rai UN, Ali MB, Tripathi RD, Yadav V, Sinha S, Singh SN, 2001. Chromium-induced physiologic changes in *Vallisneria spiralis* L. and its role in phytoremediation of tannery. *Bull Environ Contam Toxicol.* 67:246–256. doi:10.1007/s00128-001-0117-8.
- Vankar PS, Bajpai D, 2008. Phyto-remediation of chrome-VI of tannery effluent by *Trichoderma species*. *Desalination* 222(1–3):255–262. doi:10.1016/j.desal.2007.01.168.
- Varma S, Sarode D, Wakale S, Bhanvase BA, Deosarkar MP, 2013. Removal of Nickel from waste water using Graphene Nanocomposite. *Int J Chem Phys Sci.* 2:132–139.
- Vendruscolo F, Luiz G, Roberto N, Filho A, 2016. Biosorption of hexavalent chromium by microorganisms. *Int Biodeterior Biodegrad.* 119:87–95.
- Wang Q, Qian H, Yang Y, Zhang Z, Naman C, Xu X, 2010. Reduction of hexavalent chromium by carboxymethyl cellulose-stabilized zero-valent iron nanoparticles. *J Contam Hydrol.* 114(1–4):35–42. doi:10.1016/j.jconhyd.2010.02.006.
- Whitaker AH, Pena J, Amor M, Duckworth OW, 2018. Cr(VI) uptake and reduction by biogenic iron (oxyhydr)oxides. *Environ Sci Processes Impacts.* 20:1056–1068. doi:10.1039/C8EM00149A.
- WHO. 2012. Guidelines for drinking-water quality. 4th ed. Geneva: World Health Organization.
- Xia S, Song Z, Jeyakumar P, Shaheen SM, Rinklebe J, Ok YS, Bolan N, Wang H. 2019. A critical review on bioremediation technologies for Cr (VI)-contaminated soils and wastewater. *Crit Rev Environ Sci Technol.* 49(12):1027–1078.
- Xiao W, Ye X, Yang X, Zhu Z, Sun C, Zhang Q, 2017. Isolation and characterization of chromium (VI) – reducing *Bacillus sp.* FY1 and *Arthrobacter sp.* WZ2 and their bioremediation potential. *Bioremediat J.* 21:100–108. doi:10.1080/10889868.2017.1282939.
- Yang H, Feng S, Xin Y, Wang W, 2014. Community dynamics of attached and free cells and the effects of attached cells on chalcopryrite bioleaching by *Acidithiobacillus sp.* *Bioresour Technol.* 154: 185–191. doi:10.1016/j.biortech.2013.12.036.
- Yang J, Yu M, Chen W, 2015. Adsorption of hexavalent chromium from aqueous solution by activated carbon prepared from longan seed: kinetics, equilibrium and thermodynamics. *J Ind Eng Chem.* 21:414–422. doi:10.1016/j.jiec.2014.02.054.
- Yang WP, Zhang ZJ, Deng W, 2003. Simultaneous, sensitive and selective on-line chemiluminescence determination of Cr(III) and Cr(VI) by capillary electrophoresis. *Anal Chim Acta.* 485(2):169–177. doi:10.1016/S0003-2670(03)00421-5.
- Zayed AM, Terry N, 2003. Chromium in the environment: factors affecting biological remediation. *Plant Soil.* 249(1):139–156.
- Zeng J, Gou M, Tang Y, Li G, Sun Z, Kida K, 2016. Effective bioleaching of chromium in tannery sludge with an enriched sulfur-oxidizing bacterial community. *Bioresour Technol.* 218:859–866. doi:10.1016/j.biortech.2016.07.051.
- Zhang J, Han Z, Teng B, Chen W, 2017. Biodeterioration process of chromium tanned leather with *Penicillium sp.* *Int Biodeterior Biodegrad.* 116:104–111. doi:10.1016/j.ibiod.2016.10.019.
- Zhang XH, Liu J, Huang HT, Chen J, Zhu YN, Wang DQ, 2007. Chromium accumulation by the hyperaccumulator plant *Leersia hexandra* Swartz. *Chemosphere.* 67(6):1138–1143. doi:10.1016/j.chemosphere.2006.11.014.
- Zhao Z, An H, Lin J, Feng M, Murugadoss V, Ding T, Liu H, Shao Q, Mai X, Wang N, et al. 2019. Progress on the photocatalytic reduction removal of chromium contamination. *Chem Rec.* 19(5): 873–882.
- Zhou L, Li R, Zhang G, Wang D, Cai D, Wu Z, Li R, Zhang G, Wang D, Cai D, et al. 2018. Zero-valent iron nanoparticles supported by functionalized waste rock wool for efficient removal of hexavalent chromium. *Chem Eng J.* 339:85–96. doi:10.1016/j.cej.2018.01.132.
- Zhu YL, Zayed AM, Qian J-H, Souza M, Terry N, 1999. Phytoaccumulation of trace elements by wetland plants: II. Water hyacinth. *J Environ Qual.* 28(1):339–344. doi:10.2134/jeq1999.00472425002800010042x.
- Zou L, Liu P, Li X, 2013. New advances in molecular mechanism of microbial hexavalent chromium reduction. *Int J Biotechnol Food Sci.* 1:46–55.