

# PS4

April 23, 2017

## 0.1 Estimation Of Variance For Single Segment

Let's try to estimate  $\sigma_I$  analytically for one segment.

$$\sigma_I^2 = \left\langle \frac{f^2}{w^2} \right\rangle - \left\langle \frac{f}{w} \right\rangle^2$$

Let  $g = \frac{f}{w}$ , then,

$$\sigma_{f/w}^2 = \langle g^2 \rangle - \langle g \rangle^2$$

We can expand  $g$  using Taylor Series around  $\frac{b+a}{2}$ ,

$$g(x) = g\left(\frac{b+a}{2}\right) + g'\left(\frac{b+a}{2}\right)\left(x - \frac{b+a}{2}\right) + \frac{1}{2}g''\left(\frac{b+a}{2}\right)\left(x - \frac{b+a}{2}\right)^2$$

We can also approximate first and second derivative very accurately for a small region,

$$g'\left(\frac{b+a}{2}\right) = \frac{g(b) - g(a)}{b - a}$$

$$g''\left(\frac{b+a}{2}\right) = \frac{g(b) - 2g\left(\frac{b+a}{2}\right) + g(a)}{\left(\frac{b-a}{2}\right)^2}$$

To simplify equations we can define the following variables,

$$p = g\left(\frac{b+a}{2}\right)$$

$$q = \frac{g(b) - g(a)}{b - a}$$

$$r = \frac{g(b) - 2g\left(\frac{b+a}{2}\right) + g(a)}{2\left(\frac{b-a}{2}\right)^2}$$

$$u = \left(x - \frac{b+a}{2}\right)$$

Then,

$$g(x) = ru^2 + qu + p$$

Since we forced  $w$  to be equal to  $f$  at the boundaries,  $q$  is 0.

Our job reduces to calculate

$$\langle (ru^2 + p)^2 \rangle - \langle ru^2 + p \rangle^2$$

Expanding the terms,

$$\begin{aligned}
\sigma_{f/w}^2 &= \langle r^2 u^4 + 2rp u^2 + p^2 \rangle - \langle r u^2 + p \rangle^2 \\
&= (r^2 \langle u^4 \rangle + 2rp \langle u^2 \rangle + p^2) - (r^2 \langle u^2 \rangle^2 + 2rp \langle u^2 \rangle + p^2) \\
&= r^2 \langle u^4 \rangle - r^2 \langle u^2 \rangle^2 \\
&= r^2 (\langle u^4 \rangle - \langle u^2 \rangle^2)
\end{aligned}$$

This could further be simplified as,

$$\sigma_{f/w}^2 = r^2 \sigma_{u^2}^2$$

Finally  $\sigma_I$  can be estimated as,

$$\begin{aligned}
\sigma_I &= \frac{(b-a)}{\sqrt{N}} r \sigma_{u^2} = \frac{(b-a)}{\sqrt{N}} \frac{g(b) - 2g(\frac{b+a}{2}) + g(a)}{2(\frac{b-a}{2})^2} \sqrt{\left( \frac{2(b-a)^4}{5 \cdot 2^5} - \frac{4(b-a)^4}{9 \cdot 2^6} \right)} \\
\sigma_I &= \sqrt{\left( \frac{2(b-a)^2}{5 \cdot 2^5} - \frac{4(b-a)^2}{9 \cdot 2^6} \right)} \frac{2}{\sqrt{N}} (g(b) - 2g(\frac{b+a}{2}) + g(a))
\end{aligned}$$

or

$$\sigma_I = \sqrt{\left( \frac{(b-a)^6}{320N} - \frac{(b-a)^6}{576N} \right)} g'' \left( \frac{a+b}{2} \right)$$

```

In [11]: import numpy as np
          from numpy.polynomial import Polynomial as P
          #import plotly
          #import plotly.plotly as py
          #import plotly.figure_factory as ff
          import matplotlib.pyplot as plt
          #Integrand function
          def f(x,H):
              return (x-5)*np.exp(-(x/2-3))+H

          #Calculates the coefficients of linear weight function.
          def findw(f,H,lower,upper,normalize):
              #Find the linear function.
              slope=(f(upper,H)-f(lower,H))/(upper-lower)
              a=slope
              b=-slope*upper+f(upper,H)
              #Normalization.
              A=(a/2)*(upper**2)+b*upper-(a/2)*(lower**2)-b*lower
              if normalize:
                  a/=A
                  b/=A
              return [a,b]

```

```

#Performs integration.
def integrate(f, lower, upper, N, C):
    H=C
    w=findw(f, H, lower, upper, True)
    #Generate uniform random inputs.
    inputs=np.random.rand(N)
    a=w[0]/2
    b=w[1]
    c=-(a*lower**2+b*lower)

    SUM=0
    SUM2=0

    inverse_inputs=[]
    for i in inputs:
        p=[(-b-np.sqrt(b**2-4*a*(c-i)))/(2*a), (-b+np.sqrt(b**2-4*a*(c-i)))]
        if p[0]>=lower and p[0]<=upper:
            inverse_inputs.append(p[0])
        else :
            inverse_inputs.append(p[1])

    inverse_inputs=np.array(inverse_inputs)
    #Calculate f(inverse(x))/w(inverse(x)).
    outputsF=f(inverse_inputs, H)
    outputsW=w[0]*(inverse_inputs)+w[1]
    outputs=outputsF/outputsW
    SUM=outputs.sum()
    SUM2=(outputs*outputs).sum()
    var=SUM2/N-(SUM/N)**2
    var=var/N
    #Store generated points for variance calculation.
    Vsum=outputs.sum()
    return Vsum/N-H*(upper-lower), (upper-lower)**2*var


def theoretical_sigma(f, lower, upper, N, C):

    w=findw(f, C, lower, upper, True)

    flower=f(lower, C)
    fmiddle=f((lower+upper)/2, C)
    fupper=f(upper, C)

    wlower=w[0]*lower+w[1]
    wmiddle=w[0]*(lower+upper)/2+w[1]
    wupper=w[0]*upper+w[1]

```

```

gupper=fupper/wupper
gmiddle=fmiddle/wmiddle
glower=flower/wlower

sigma=np.abs(np.sqrt((upper-lower)**6/(320*N)-(upper-lower)**6/(576*N))

    #return (upper-lower)*np.sqrt(var)
    return sigma

low=4.6
up=5.2
#Divide the region into 10 pieces.
l=[low,up]
#Real value of the integral
I_real=-.12002
#N values
N=[100,1000,10000,100000,1000000]

#Integration results.
results=[]
#Standart deviation values
sigmas=[]
for k in N:
    I=0
    sigma=0
    S=1000
    for i in range(0,len(l)-1):
        temp,temp2=integrate(f,l[i],l[i+1],k,S)
        I+=temp
        sigma+=temp2
    results.append(I)
    sigmas.append(np.sqrt(sigma))
    print(k,I,I-I_real,np.sqrt(sigma),theoretical_sigma(f,low,up,k,S)/np.s

100 -0.119462876295 0.000557123704812 0.000893254933342 0.964176912628 0.6237006749
1000 -0.119637669659 0.000382330340745 0.000265470016018 1.0259275099 1.44020159595
10000 -0.120042301541 -2.23015408205e-05 8.58932633227e-05 1.00270469476 -0.2596424
100000 -0.120035753352 -1.57533515444e-05 2.72967667204e-05 0.997748177565 -0.57711
1000000 -0.120018938155 1.06184534208e-06 8.63053525367e-06 0.99791699878 0.1230335

```

In [ ]: