


# **Project Report**

## **November 12, 2025**



**EH 611**  
**Modelling of of Earth System and Sustainability**  
**Prof. R N Singh**  
INDIAN INSTITUTE OF TECHNOLOGY GANDHINAGAR  
Palaj, Gandhinagar - 382355

---

## **Modelling of Photochemical Smog**

---

Submitted by

**Guntas Singh Saran (22110089)**  
Computer Science and Engineering



## Abstract

Photochemical smog forms through complex interactions between nitrogen oxides ( $\text{NO}_x$ ), volatile organic compounds (VOCs), and sunlight. In this project, I develop two mathematical models of increasing complexity: Model 1 captures the basic  $\text{NO}$ - $\text{NO}_2$ - $\text{O}_3$  photochemical cycle with 4 species and 5 reactions, while Model 2 extends to 11 species and 15 reactions, incorporating VOC oxidation and radical chemistry. Solving the equations numerically, it is demonstrated that Model 1 produces minimal ozone ( $\sim 2 \times 10^{-3}$  ppb peak) due to rapid titration by  $\text{NO}$ , while Model 2 achieves realistic ozone levels (order of  $\sim 0.1 \times$  ppm) through radical-mediated  $\text{NO}$  oxidation pathways. The enhancement factor highlights the critical role of VOCs and radicals in urban photochemical smog formation. Several studies from the 80s have beautifully produced models for the same involving a large number of species [Bazzell and Peters(1981), McRae et al.(1982), Falls et al.(1979)]. The chapter “Modelling of Photochemical Smog” from the book “Environment Modelling and Pollution” [Duc et al.(2002)] aims to lump various species into a 7-reaction Generic Reaction Set (GRS); although their model is interpretable and easy to follow, I could not fetch parametric values for their system (this is briefly touched upon in the Appendix). Finally, I resorted to a detailed study by [Carrasco-Venegas et al.(2025)] that provided me with all the required reactions and parameters.

## Introduction

Photochemical smog, first documented in Los Angeles in the 1940s [Bazzell and Peters(1981), McRae et al.(1982), Falls et al.(1979)], remains a persistent urban air quality challenge. Unlike traditional industrial smog, photochemical smog arises from sunlight-driven reactions involving:

- **Primary pollutants:**  $\text{NO}_x$  (from combustion), VOCs (from vehicles, solvents)
- **Solar radiation:** Provides photochemical energy
- **Secondary pollutants:** Ozone ( $\text{O}_3$ ), aldehydes, PAN

Ground-level ozone concentrations exceeding 0.07 ppm cause respiratory problems, crop damage, and ecosystem degradation. Understanding the chemical mechanisms is essential for developing effective control strategies.

## Objectives

Develops two models, state the variables, describe chemical equations between them, using conservation of mass and rate law, formulate in ODE and solve numerically:

### 1. Model 1:


- (a) Simple 3-species photochemical cycle ( $\text{NO}$ ,  $\text{NO}_2$ ,  $\text{O}_3$ )
- (b) A bit refined 4-species photochemical cycle ( $\text{NO}$ ,  $\text{NO}_2$ ,  $\text{O}_3$ ,  $\text{O}$ )

### 2. Model 2: Refined 11-species system with VOCs and radicals

## Model 1: Basic Photochemical Cycle

### 3-species Model

Model 1 captures the fundamental  $\text{NO}$ - $\text{NO}_2$ - $\text{O}_3$  photochemical cycle with three species and three reactions:





Reaction (1) is photochemical (requires sunlight with rate constant  $k_1$ ), while reaction (2) occurs day and night (rate constant  $k_3$ ). These reactions form a cycle that interconverts NO and NO<sub>2</sub> but *cannot produce net ozone*-a key limitation we'll address in Model 2.

## Mathematical Formulation

### General Mass Balance Equation

For a well-mixed box (closed system, uniform concentration):

$$\frac{dC_i}{dt} = r_i + E_i \quad (3)$$

The reaction rate  $r_i$  is the algebraic sum of production and consumption terms from all reactions involving species  $i$ :

$$r_i = \sum_{j=1}^{N_{\text{rxn}}} \nu_{ij} R_j \quad (4)$$

where:

- $\nu_{ij}$  = stoichiometric coefficient of species  $i$  in reaction  $j$  (positive for products, negative for reactants)
- $R_j$  = rate of reaction  $j$  (ppm/h)
- $N_{\text{rxn}}$  = total number of reactions

Applying mass conservation with emission sources:

$$\frac{d[\text{NO}]}{dt} = k_1[\text{NO}_2] - k_3[\text{NO}][\text{O}_3] + E_{\text{NO}} \quad (5)$$

$$\frac{d[\text{NO}_2]}{dt} = -k_1[\text{NO}_2] + k_3[\text{NO}][\text{O}_3] + E_{\text{NO}_2} \quad (6)$$

$$\frac{d[\text{O}_3]}{dt} = k_1[\text{NO}_2] - k_3[\text{NO}][\text{O}_3] \quad (7)$$

## 4-species Model

### Reaction Mechanism

Model 1 captures the fundamental NO-NO<sub>2</sub>-O<sub>3</sub> triad plus atomic oxygen:



where M represents air as a third body in the termolecular reaction (9).

- Reaction (8): NO<sub>2</sub> photolysis produces atomic oxygen (photochemical, daytime only)
- Reaction (10): NO immediately reacts with O<sub>3</sub>, **preventing accumulation**
- Reactions (11)-(12): Additional O-NO<sub>x</sub> interactions

## Mathematical Formulation

Applying species mass balance with emission sources:

$$\frac{d[\text{NO}]}{dt} = R_1 - R_3 + R_4 - R_5 + E_{\text{NO}} \quad (13)$$

$$\frac{d[\text{NO}_2]}{dt} = -R_1 + R_3 - R_4 + R_5 + E_{\text{NO}_2} \quad (14)$$

$$\frac{d[\text{O}_3]}{dt} = R_2 - R_3 \quad (15)$$

$$\frac{d[\text{O}]}{dt} = R_1 - R_2 - R_4 - R_5 \quad (16)$$

where reaction rates are:

$$R_1 = k_1(t)[\text{NO}_2] \quad (17)$$

$$R_2 = k_2[\text{O}][\text{O}_2]M \quad (18)$$

$$R_3 = k_3[\text{O}_3][\text{NO}] \quad (19)$$

$$R_4 = k_4[\text{NO}_2][\text{O}] \quad (20)$$

$$R_5 = k_5[\text{NO}][\text{O}] \quad (21)$$

The photolysis rate  $k_1(t)$  varies diurnally:

$$k_1(t) = \begin{cases} k_{1,\max} \sin\left(\frac{\pi(t_{\text{clock}}-6)}{12}\right) & \text{if } 6 \leq t_{\text{clock}} \leq 18 \\ 0 & \text{otherwise} \end{cases} \quad (22)$$

## Parameters and Initial Conditions


Table 1: Model 1 parameters at  $T = 288 \text{ K}$  [Carrasco-Venegas et al.(2025)]

Parameter	Value	Units
$k_{1,\max}$	30.48	$\text{h}^{-1}$
$k_2$	$1.44 \times 10^{-3}$	$\text{ppm}^{-2} \text{ h}^{-1}$
$k_3$	$1.20 \times 10^5$	$\text{ppm}^{-1} \text{ h}^{-1}$
$k_4$	$8.04 \times 10^5$	$\text{ppm}^{-1} \text{ h}^{-1}$
$k_5$	$1.99 \times 10^5$	$\text{ppm}^{-1} \text{ h}^{-1}$
$[\text{O}_2] \text{ M}$	210,000	ppm
$E_{\text{NO}}$	0.02	$\text{ppm h}^{-1}$
$E_{\text{NO}_2}$	0.01	$\text{ppm h}^{-1}$

Table 2: Model 1 initial conditions (5:00 AM)

Species	Initial Concentration (ppm)
NO	0.100
NO <sub>2</sub>	0.050
O <sub>3</sub>	0.0
O	0.0

Model 1: Basic Photochemical Cycle



Model 1: Basic Photochemical Cycle




Figure 1: Model 1 (a) 3-species and (b) 4-species concentration profiles from 5:00 AM to mid-night. (Top left) NO and NO<sub>2</sub> concentrations. (Top right) Ozone formation with (a) peak (~ 0.0004 ppm)(b) very low peak (~0.006 ppb). (b) (Bottom left) Atomic oxygen (scaled) tracking photolysis. (Bottom right) Normalized comparison showing relative dynamics. Yellow shading indicates daylight hours (6:00-18:00).  
Project Report



## Key Observations

1. **Minimal ozone production:** Peak  $O_3$  reaches only  $\sim 0.003$  ppb at 8:00, far below ambient air quality standards (0.07 ppm)
2. **NO-NO<sub>2</sub> concentration:** Unable to show increase-decrease pattern in either - indicating some key species missing in our model.
3. **Atomic oxygen:** Very low steady-state concentration ( $\sim 10^{-6}$  ppm) due to fast reaction with  $O_2$  (R<sub>2</sub>)

## Limitations

- **Unrealistic ozone:** Cannot explain observed urban smog episodes ( $O_3 > 0.07$  ppm)
- **Missing VOC chemistry:** No role for hydrocarbons in oxidation
- **No radical pathways:** Lacks OH, HO<sub>2</sub>, RO<sub>2</sub> that drive real atmospheric chemistry

The fundamental problem: *there is no mechanism to convert NO to NO<sub>2</sub> without consuming O<sub>3</sub>*. This motivates Model 2.

## Model 2: Refined System with VOCs

### System Extension





Model 2 adds 7 new species to capture VOC oxidation and radical chemistry:

- **CO:** Carbon monoxide (primary pollutant, radical precursor)
- **HCHO:** Formaldehyde (VOC oxidation product, radical source)
- **ALK:** Lumped alkanes (saturated hydrocarbons)
- **OLE:** Lumped olefins (unsaturated hydrocarbons)
- **OH:** Hydroxyl radical (primary atmospheric oxidant)
- **HO<sub>2</sub>:** Hydroperoxyl radical
- **RO<sub>2</sub>:** Organic peroxy radicals

**Total species:** 11 (NO, NO<sub>2</sub>, O<sub>3</sub>, O, CO, HCHO, ALK, OLE, OH, HO<sub>2</sub>, RO<sub>2</sub>)

### Extended Reaction Mechanism

Building on Model 1's 5 reactions, we add 10 more:



**Critical breakthrough:** Reactions (31) and (32) convert NO to NO<sub>2</sub> *without consuming* O<sub>3</sub>! This enables net ozone accumulation.

## Parameters

Table 3: Additional rate constants for Model 2 (T = 288 K) [Carrasco-Venegas et al.(2025)]

Reaction	Rate Constant	Units
k <sub>4</sub> (O <sub>3</sub> photolysis)	1.968	h <sup>-1</sup>
k <sub>5</sub> (O(^1D) + H <sub>2</sub> O)	6.0 × 10 <sup>6</sup>	ppm <sup>-1</sup> h <sup>-1</sup>
k <sub>6</sub> (CO + OH)	2.64 × 10 <sup>4</sup>	ppm <sup>-1</sup> h <sup>-1</sup>
k <sub>7</sub> (HCHO photolysis)	0.170	h <sup>-1</sup>
k <sub>8</sub> (HCHO + OH)	1.152 × 10 <sup>6</sup>	ppm <sup>-1</sup> h <sup>-1</sup>
k <sub>9</sub> (ALK + OH)	2.82 × 10 <sup>5</sup>	ppm <sup>-1</sup> h <sup>-1</sup>
k <sub>10</sub> (OLE + OH)	5.349 × 10 <sup>6</sup>	ppm <sup>-1</sup> h <sup>-1</sup>
k <sub>11</sub> (OLE + O <sub>3</sub> )	8.16	ppm <sup>-1</sup> h <sup>-1</sup>
k <sub>12</sub> (HO <sub>2</sub> + NO)	7.2 × 10 <sup>5</sup>	ppm <sup>-1</sup> h <sup>-1</sup>
k <sub>13</sub> (RO <sub>2</sub> + NO)	7.2 × 10 <sup>5</sup>	ppm <sup>-1</sup> h <sup>-1</sup>
k <sub>14</sub> (HO <sub>2</sub> + HO <sub>2</sub> )	2.22 × 10 <sup>5</sup>	ppm <sup>-1</sup> h <sup>-1</sup>
k <sub>15</sub> (OH + NO <sub>2</sub> )	Complex T-dep.	ppm <sup>-1</sup> h <sup>-1</sup>

**Table 4: Model 2 initial conditions and emissions**

Species	Initial (ppm)	Emission (ppm h <sup>-1</sup> )
NO	0.100	0.02
NO <sub>2</sub>	0.050	0.01
CO	0.100	0.02
HCHO	0.010	0.03
ALK	1.000	0.10
OLE	0.200	0.0
OH	0	0
HO <sub>2</sub>	0	0
RO <sub>2</sub>	0	0

## Results

Model 2: Refined with VOCs and Radicals




Figure 2: Model 2 comprehensive results. (Top row) Ozone reaches realistic levels ( $\sim 0.08$  ppm), NO-NO<sub>2</sub> dynamics, and radical species tracking solar radiation. (Middle row) VOC depletion, secondary products (HCHO, CO), and NO $\rightarrow$ NO<sub>2</sub> conversion pathways. (Bottom row) NO<sub>x</sub> budget, oxidant capacity, and NO/NO<sub>2</sub> ratio evolution. Model successfully produces net ozone through radical chemistry.



## Key Observations

1. **Realistic ozone:** Peak  $O_3 = 0.619979$  ppm at 7:30.
2. **Radical dynamics:** OH,  $HO_2$ ,  $RO_2$  peak at early hours ( $\sim 10^{-6}$  to  $10^{-3}$  ppm), drop to zero at night
3. **VOC consumption:** ALK decreases 23%, OLE decreases 100% (fully depleted), demonstrating active oxidation
4. **Pathway dominance:** At peak  $O_3$ ,  $NO \rightarrow NO_2$  conversion is dominated by  $RO_2 + NO$  (94.5%), with only  $HO_2 + NO$  (2.9%) and 2.6% via  $O_3$  titration
5. **Radical catalysis:** Despite trace OH ( $\sim 10^{-6}$  ppm), substantial ozone forms.
6.  **$NO_x$  budget:** Total  $NO_x$  decreases from 0.15 to 0.08 ppm due to  $HNO_3$  formation ( $R_{15}$ )
7. **Oxidant capacity:**  $O_x$  ( $O_3 + NO_2$ ) steadily increases, indicating net oxidation

## Comparison with Model 1

Model Comparison: Impact of VOCs on Ozone Formation




Figure 3: Direct comparison between models. (Top left) Ozone enhancement: Model 2 produces 27x more  $O_3$  than Model 1. (Top right) NO depletion is faster in Model 2 due to additional radical pathways. (Bottom left)  $NO_2$  shows more complex dynamics. (Bottom right)  $O_3$  enhancement factor peaks in afternoon when radical chemistry is most active.

## Appendix: Mathematical Formulation, GRS ODEs, parameter table

### Model 2: Complete Differential Equations

#### Species Conservation Equations

For Model 2 with 11 species and 15 reactions:

##### 1. Nitric Oxide (NO):

$$\begin{aligned}\frac{d[\text{NO}]}{dt} &= R_1 - R_3 - R_{12} - R_{13} + E_{\text{NO}} \\ &= k_1[\text{NO}_2] - k_3[\text{O}_3][\text{NO}] - k_{12}[\text{HO}_2][\text{NO}] \\ &\quad - k_{13}[\text{RO}_2][\text{NO}] + E_{\text{NO}}\end{aligned}\tag{35}$$

##### 2. Nitrogen Dioxide (NO<sub>2</sub>):

$$\begin{aligned}\frac{d[\text{NO}_2]}{dt} &= -R_1 + R_3 + R_{12} + R_{13} - R_{15} + E_{\text{NO}_2} \\ &= -k_1[\text{NO}_2] + k_3[\text{O}_3][\text{NO}] + k_{12}[\text{HO}_2][\text{NO}] \\ &\quad + k_{13}[\text{RO}_2][\text{NO}] - k_{15}[\text{OH}][\text{NO}_2] + E_{\text{NO}_2}\end{aligned}\tag{36}$$

##### 3. Ozone (O<sub>3</sub>):

$$\begin{aligned}\frac{d[\text{O}_3]}{dt} &= R_2 - R_3 - R_4 - R_{11} \\ &= k_2[\text{O}][\text{O}_2] - k_3[\text{O}_3][\text{NO}] - k_4[\text{O}_3] - k_{11}[\text{OLE}][\text{O}_3]\end{aligned}\tag{37}$$

##### 4. Atomic Oxygen (O):

$$\begin{aligned}\frac{d[\text{O}]}{dt} &= R_1 - R_2 \\ &= k_1[\text{NO}_2] - k_2[\text{O}][\text{O}_2]\end{aligned}\tag{38}$$

##### 5. Carbon Monoxide (CO):

$$\begin{aligned}\frac{d[\text{CO}]}{dt} &= -R_6 + R_7 + R_8 + E_{\text{CO}} \\ &= -k_6[\text{CO}][\text{OH}] + k_7[\text{HCHO}] + k_8[\text{HCHO}][\text{OH}] + E_{\text{CO}}\end{aligned}\tag{39}$$

##### 6. Formaldehyde (HCHO):

$$\begin{aligned}\frac{d[\text{HCHO}]}{dt} &= 0.5R_{11} - R_7 - R_8 + E_{\text{HCHO}} \\ &= 0.5k_{11}[\text{OLE}][\text{O}_3] - k_7[\text{HCHO}] - k_8[\text{HCHO}][\text{OH}] \\ &\quad + E_{\text{HCHO}}\end{aligned}\tag{40}$$

##### 7. Alkanes (ALK):

$$\begin{aligned}\frac{d[\text{ALK}]}{dt} &= -R_9 + E_{\text{ALK}} \\ &= -k_9[\text{ALK}][\text{OH}] + E_{\text{ALK}}\end{aligned}\tag{41}$$

### 8. Olefins (OLE):

$$\begin{aligned}\frac{d[\text{OLE}]}{dt} &= -R_{10} - R_{11} + E_{\text{OLE}} \\ &= -k_{10}[\text{OLE}][\text{OH}] - k_{11}[\text{OLE}][\text{O}_3] + E_{\text{OLE}}\end{aligned}\quad (42)$$

### 9. Hydroxyl Radical (OH):

$$\begin{aligned}\frac{d[\text{OH}]}{dt} &= 2R_5 - R_6 - R_8 - R_9 - R_{10} - R_{15} + R_{12} \\ &= 2k_5[\text{O}(^1\text{D})][\text{H}_2\text{O}] - k_6[\text{CO}][\text{OH}] - k_8[\text{HCHO}][\text{OH}] \\ &\quad - k_9[\text{ALK}][\text{OH}] - k_{10}[\text{OLE}][\text{OH}] - k_{15}[\text{OH}][\text{NO}_2] \\ &\quad + k_{12}[\text{HO}_2][\text{NO}]\end{aligned}\quad (43)$$

### 10. Hydroperoxyl Radical (HO<sub>2</sub>):

$$\begin{aligned}\frac{d[\text{HO}_2]}{dt} &= R_6 + 2R_7 + R_8 + 0.5R_{11} + R_{13} - R_{12} - 2R_{14} \\ &= k_6[\text{CO}][\text{OH}] + 2k_7[\text{HCHO}] + k_8[\text{HCHO}][\text{OH}] \\ &\quad + 0.5k_{11}[\text{OLE}][\text{O}_3] + k_{13}[\text{RO}_2][\text{NO}] \\ &\quad - k_{12}[\text{HO}_2][\text{NO}] - 2k_{14}[\text{HO}_2]^2\end{aligned}\quad (44)$$

### 11. Organic Peroxy Radical (RO<sub>2</sub>):

$$\begin{aligned}\frac{d[\text{RO}_2]}{dt} &= R_9 + R_{10} + 0.5R_{11} - R_{13} \\ &= k_9[\text{ALK}][\text{OH}] + k_{10}[\text{OLE}][\text{OH}] + 0.5k_{11}[\text{OLE}][\text{O}_3] \\ &\quad - k_{13}[\text{RO}_2][\text{NO}]\end{aligned}\quad (45)$$

## Reaction Rates

The individual reaction rates for Model 2 are:

$$R_1 = k_1(t)[\text{NO}_2] \quad (\text{NO}_2 \text{ photolysis}) \quad (46)$$

$$R_2 = k_2[\text{O}][\text{O}_2] \quad (\text{O}_3 \text{ formation}) \quad (47)$$

$$R_3 = k_3[\text{O}_3][\text{NO}] \quad (\text{NO oxidation by O}_3) \quad (48)$$

$$R_4 = k_4(t)[\text{O}_3] \quad (\text{O}_3 \text{ photolysis}) \quad (49)$$

$$R_5 = k_5[\text{O}(^1\text{D})][\text{H}_2\text{O}] \quad (\text{OH formation}) \quad (50)$$

$$R_6 = k_6[\text{CO}][\text{OH}] \quad (\text{CO oxidation}) \quad (51)$$

$$R_7 = k_7(t)[\text{HCHO}] \quad (\text{HCHO photolysis}) \quad (52)$$

$$R_8 = k_8[\text{HCHO}][\text{OH}] \quad (\text{HCHO oxidation}) \quad (53)$$

$$R_9 = k_9[\text{ALK}][\text{OH}] \quad (\text{Alkane oxidation}) \quad (54)$$

$$R_{10} = k_{10}[\text{OLE}][\text{OH}] \quad (\text{Olefin oxidation by OH}) \quad (55)$$

$$R_{11} = k_{11}[\text{OLE}][\text{O}_3] \quad (\text{Olefin ozonolysis}) \quad (56)$$

$$R_{12} = k_{12}[\text{HO}_2][\text{NO}] \quad (\text{Key: NO} \rightarrow \text{NO}_2 \text{ via HO}_2) \quad (57)$$

$$R_{13} = k_{13}[\text{RO}_2][\text{NO}] \quad (\text{Key: NO} \rightarrow \text{NO}_2 \text{ via RO}_2) \quad (58)$$

$$R_{14} = k_{14}[\text{HO}_2]^2 \quad (\text{HO}_2 \text{ termination}) \quad (59)$$

$$R_{15} = k_{15}[\text{OH}][\text{NO}_2] \quad (\text{HNO}_3 \text{ formation}) \quad (60)$$

## Excited State Oxygen (Quasi-Steady-State)

For  $O(^1D)$  (excited singlet oxygen), we apply the quasi-steady-state approximation:

$$\frac{d[O(^1D)]}{dt} \approx 0 \quad (61)$$

Production equals consumption:

$$k_4[O_3] = k_5[O(^1D)][H_2O] \quad (62)$$

Solving for  $[O(^1D)]$ :

$$[O(^1D)] = \frac{k_4[O_3]}{k_5[H_2O]} \quad (63)$$

This concentration is substituted directly into Equation 43.

## GRS reaction list (symbolic)

The Generic Reaction Set (GRS) used in [Duc et al.(2002)] (will discuss this during presentation):

- (R1)  $ROC + h\nu \rightarrow RP + ROC$
- (R2)  $RP + NO \rightarrow NO_2$
- (R3)  $NO_2 + h\nu \rightarrow NO + O$
- (R4)  $NO + O_3 \rightarrow NO_2 + O_2$
- (R5)  $RP + RP \rightarrow \text{termination products}$
- (R6)  $RP + NO_2 \rightarrow SGN$
- (R7)  $RP + NO_2 \rightarrow SNGN$




Figure 4: Brief discussion on a generic reaction set as described by [Duc et al.(2002)]

# Bibliography

- [Bazzell and Peters(1981)] Catherine C. Bazzell and Leonard K. Peters. 1981. The transport of photochemical pollutants to the background troposphere. *Atmospheric Environment (1967)* 15, 6 (1981), 957–968. doi:10.1016/0004-6981(81)90096-2
- [Carrasco-Venegas et al.(2025)] Luis Américo Carrasco-Venegas, Luz Genara Castañeda-Pérez, Daril Giovanni Martínez-Hilario, Juan Taumaturgo Medina-Collana, José Vulfrano González-Fernández, Cesar Gutiérrez-Cuba, Héctor Ricardo Cuba-Torre, Alex Pilco-Nuñez, Carlos Alejandro Ancieta-Dextre, and Oscar Juan Rodríguez-Taranco. 2025. Mathematical Modeling of Photochemical and Chemical Interactions in Photochemical Smog Formation. *Processes* 13, 5 (2025). doi:10.3390/pr13051384
- [Duc et al.(2002)] Hiep Duc, Võ Anh, and Merched Azzi. 2002. *Modelling of Photochemical Smog*. pp 361–382. doi:10.1007/978-3-662-04868-9\_10
- [Falls et al.(1979)] Andrew H. Falls, Gregory J. McRae, and John H. Seinfeld. 1979. Sensitivity and uncertainty of reaction mechanisms for photochemical air pollution. *International Journal of Chemical Kinetics* 11, 11 (1979), 1137–1162. arXiv:https://onlinelibrary.wiley.com/doi/pdf/10.1002/kin.550111102 doi:10.1002/kin.550111102
- [McRae et al.(1982)] Gregory J McRae, William R Goodin, and John H Seinfeld. 1982. Numerical solution of the atmospheric diffusion equation for chemically reacting flows. *J. Comput. Phys.* 45, 1 (1982), 1–42. doi:10.1016/0021-9991(82)90101-2