

Notes for MATH 3210: Foundation of Analysis I

Jing Guo

August 23, 2017

Contents

1	Ring and Field	1
1.1	Ring	1
1.1.1	Addition Axiom	2
1.1.2	Multiplication Axiom	2
1.2	Field	2
2	Basic Topology	2
2.1	Metric Space M	2
2.2	Open Sets	3
2.2.1	Metric Spaces X	3
2.2.2	Topological Spaces (X, U)	3
2.3	Compact Sets	3

1 Ring and Field

1.1 Ring

The set A has two binary operations, addition and multiplication.

For any $a, b \in A$,

$$A * A \rightarrow A \tag{1}$$

$$a, b \rightarrow a + b \tag{2}$$

Notations:

- \mathbb{Z} : The set of integers (ring, not field, has no inverse);
- \mathbb{Q} : The set of rational numbers;
- \mathbb{R} : The set of real numbers (ring and field).

1.1.1 Addition Axiom

1. $a + b = b + a$ (commutative)
2. $(a + b) + c = a + (b + c)$ (associative)
3. There is an element 0 such that $a + 0 = a$ (additive identity)
4. There exists $-a$ (additive inverse) such that $a + (-a) = 0$

1.1.2 Multiplication Axiom

1. $a \cdot b = b \cdot a$
2. $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
3. There is an element 1 such that $a \cdot 1 = a$ and $0 \neq 1$
4. $a \cdot (b + c) = a \cdot b + a \cdot c$ (distributive)

1.2 Field

A is a field if A is a *ring* for any $a \in A$ and $a \neq 0$, there exists a^{-1} (inverse of a) such that $a \cdot a^{-1} = 1$

Example: $x, y \in A, x \neq 0, y \neq 0, xy = 0$

$$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \cdot \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

2 Basic Topology

Metric (distance function) of x_0 and x : $d(x_0, x) = |x - x_0|$

2.1 Metric Space M

$$d: M \times M \rightarrow \mathbb{R}$$

1. $d(x_0, x_1) \geq 0$ where $x_0, x_1 \in M$
2. $d(x_0, x_1) = 0 \iff x_0 = x_1$
3. $d(x_0, x_1) = d(x_1, x_0)$ (irrespective of order)
4. $d(x, z) \leq d(x, y) + d(y, z)$ (triangular inequity)

Euclidean metric on \mathbb{R}^2 : $d(x, y) = \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2}$

Discrete metric: $d(x, y) = \begin{cases} 1 & \text{if } x \neq y \\ 0 & \text{if } x = y \end{cases}$

Open ball of radius ϵ : $B_\epsilon(x_0) = B(x_0, \epsilon) = \{y \in M \mid d(x_0, y) < \epsilon\}$

2.2 Open Sets

2.2.1 Metric Spaces X

$U \subset X$ is open if for any $x \in U$ there exists $\epsilon > 0$ such that $B(x, \epsilon) \subset U$.

Open interval is open set; closed interval is not open set.

2.2.2 Topological Spaces (X, U)

Let U be a family of all sets, X be a set. U is a **topology** on X if

1. \emptyset (Empty set) is always open; X is open. $\Leftrightarrow \emptyset$ and X itself belong to U .
2. F is a collection of open sets, then $\bigcup_{U \in F} U$ is open. \Leftrightarrow Any union of members of U still belongs to U . (Union)
3. F is a *finite* collection of open sets, then $\bigcap_{U \in F} U$ is open. \Leftrightarrow The intersection of any finite number of members of U belongs to U . (Intersection)

Finite case: $x \in \bigcap_{U \in F} U$, $x \in U$, hence $B(x, \epsilon_U) \subset U$

$$\delta = \min \epsilon_U > 0 \quad \text{where } U \in F$$

$$B(x, \delta) \subset B(x, \epsilon_U) \subset U$$

$$\therefore B(x, \delta) \subset \bigcap_{U \in F} U$$

Infinite case: For example, the intersection of all intervals of $(-\frac{1}{n}, \frac{1}{n})$, where n is a positive number, is the set $\{0\}$ which is not open in the real line.

2.3 Compact Sets

In metric space X , compact sets are closed.

Compact \Leftrightarrow closed and bounded (only for Euclidean metric, \mathbb{R}^n)