

order check for separating T and U

zichi zhang

June 2026

1 the problem

In the further quantum mechanics lecture notes, it was claimed that

$$e^{-\frac{i}{\hbar}H(t_n)\Delta t} e^{-\frac{i}{\hbar}H(t_{n-1})\Delta t} \dots e^{-\frac{i}{\hbar}H(t_1)\Delta t}$$

cannot be approximated as

$$e^{-\frac{i}{\hbar} \int_0^t H(t') dt'}$$

While the approximation

$$\prod_{i=n}^1 e^{-\frac{i}{\hbar}(T+V(t_i))\Delta t} \approx \prod_{i=n}^1 e^{-\frac{i\Delta t}{\hbar}T} e^{-\frac{i\Delta t}{\hbar}V(t_i)}$$

is valid.

However, I was confused when checking the orders.
it was concluded in the notes that

$$e^{-\frac{i}{\hbar}(T+V(t_i))\Delta t} = e^{-\frac{i\Delta t}{\hbar}T} e^{-\frac{i\Delta t}{\hbar}V(t_i)} + O(\Delta t^2)$$

, which is true.

However, it is also true that

$$e^{-\frac{i}{\hbar}(H(t_{i+1})+H(t_i))\Delta t} = e^{-\frac{i\Delta t}{\hbar}H(t_{i+1})} e^{-\frac{i\Delta t}{\hbar}H(t_i)} + O(\Delta t^2)$$

Given the same order of error, why the dividing the kinetic and potential energy terms is a valid approximation but neglecting the time-ordering is not?

2 solution: different global errors

We would like to approximate the time-evolution operator by

$$U(t) \simeq \exp\left(-\frac{i}{\hbar} \int_0^t H(t') dt'\right) (?)$$

Let us check the order of the approximation.

$$\prod_{i=n}^1 e^{-\frac{i\Delta t}{\hbar} H(t_i)} = e^{\left(\sum_i -\frac{i\Delta t}{\hbar} H(t_i) + \sum_{i \neq j} \frac{1}{2} \left(-\frac{i\Delta t}{\hbar}\right)^2 [H(t_i), H(t_j)] \right)}.$$

Let

$$N = \frac{t}{\Delta t}.$$

The number of commutator correction terms with $O(\Delta t^2)$ is given by the number of terms in the $i \neq j$ summation:

$$\binom{N}{2} = \frac{N(N-1)}{2} \sim O(\Delta t^{-2}),$$

so

$$\boxed{\Delta U \sim O(1).}$$

This global error does not disappear with $\Delta t \rightarrow 0$, so is not valid.

On the other hand, although the local error of the Trotter decomposition is the same, the global error is different.

local error:

$$e^{-\frac{i\Delta t}{\hbar}(T+V(t_i))} = e^{-\frac{i\Delta t}{\hbar}T} e^{-\frac{i\Delta t}{\hbar}V(t_i)} + O(\Delta t^2),$$

expansion:

$$e^{-\frac{i\Delta t}{\hbar}(T+V(t_n))} \dots e^{-\frac{i\Delta t}{\hbar}(T+V(t_1))} = \left(e^{-\frac{i\Delta t}{\hbar}T} e^{-\frac{i\Delta t}{\hbar}V(t_n)} + O(\Delta t^2) \right) \dots \left(e^{-\frac{i\Delta t}{\hbar}T} e^{-\frac{i\Delta t}{\hbar}V(t_1)} + O(\Delta t^2) \right).$$

still define

$$N = \frac{t}{\Delta t}$$

,since one bracket only contribute one error term, the global error is:

$$\boxed{O(\Delta t^2) \times N = O(\Delta t).}$$

which means the approximation will become precise with Δt substituted by dt .